Runners with mid-portion Achilles tendinopathy have greater triceps surae

intracortical inhibition than healthy controls

Gabriel L. Fernandes; Lucas B. R. Orssatto; Anthony J. Shield; Gabriel S. Trajano

School of Exercise and Nutrition Sciences, Faculty of Health, Queensland University of Technology

(QUT), Brisbane, Australia.

Abstract

Objectives: This study aimed to investigate short-interval intracortical inhibition (SICI) and muscle

function in the Triceps surae of runners with mid-portion Achilles tendinopathy (AT).

Methods: Runners with (n=11) and without (n=13) AT were recruited. Plantar flexor isometric peak

torque and rate of torque development (RTD) were measured using an isokinetic dynamometer. Triceps

surae endurance was measured as single leg heel raise (SLHR) to failure test. SICI was assessed using

paired-pulse transcranial magnetic stimulation during a sustained contraction at 10% of plantar flexor

isometric peak torque.

Results: Triceps surae SICI was 14.3% (95%CI: -2.1 to 26.4) higher in AT than control group (57.9%,

95%CI: 36.2, to 79.6; and 43.6% 95%CI:16.2 to 71.1; p=0.032) irrespective of the tested muscle. AT

performed 16 (95%CI: 7.9 to 23.3, p<0.001) fewer SLHR repetitions on the symptomatic side compared

to controls, and 14 (95%CI: 5.8 to 22.0, p=0.004), fewer SLHR repetitions on the non-symptomatic

compared to controls. We found no between-groups differences in isometric peak torque (p=0.971) or

RTD (p=0.815).

Perspective: Our data suggest greater intracortical inhibition for the Triceps surae muscles for the AT

group accompanied by reduced SLHR endurance, without deficits in isometric peak torque or RTD.

The increased SICI observed in the AT group could be negatively influencing Triceps surae endurance;

thus, rehabilitation aiming to reduce intracortical inhibition should be considered to improve patient

outcomes. Furthermore, SLHR is a useful clinical tool to assess plantar flexor function in AT patients.

Keywords

Achilles tendinopathy; running injury; SICI; single leg heel raise; triceps surae endurance

1. Introduction

Achilles tendinopathy (AT) is an overload injury, most commonly seen in the mid-portion of the

tendon, which affects both recreational and elite athletes in sports that involve running and jumping[1].

AT is a debilitating multifactorial injury, with high recurrency rates[1,2]. Triceps surae weakness is a key

feature in chronic AT and is suggested to be present even before the development of tendinopathy

symptoms[3–5]. Muscle weakness has been reported in individuals with AT as reduced maximal concen-

tric and eccentric strength, and endurance. These deficits in muscle strength and endurance are still

observed after years of rehabilitation[6,7], even with improvements in pain and function[5,8,9]. Thus, it is of

great clinical relevance to understand the mechanisms underpinning chronic muscle weakness to de-

velop better rehabilitation strategies.

Chronic muscle weakness in individuals with AT could have an origin in the central nervous

system. Bilateral sensory and motor deficits (i.e. proprioception and muscle weakness) have been re-

ported in individuals with unilateral tendinopathy, implying involvement of the central nervous sys-

tem[10–12]. The central nervous system controls the motor action through different inhibitory and excita-

tory intracortical mechanisms, which will fine tune and adjust the intensity of the stimuli to achieve the

required level of force and control[13]. Thus, any changes in cortical mechanisms of inhibition could

affect motor performance[13,14]. Short-interval intracortical inhibition (SICI) is a measurement of intra-

cortical mechanisms of inhibition, and it is believed to represent post-synaptic inhibition mediated by

$GABA_A$ receptors[15]. SICI reflects the balance between inhibition and facilitation and can be measured

using transcranial magnetic stimulation (TMS), by delivering a pair of pulses to the primary motor

cortex. Increased SICI has been reported in patellar tendinopathy patients, suggesting an imbalance

within intracortical mechanisms of inhibition; affecting muscle activation and consequently muscle

force[14]. Therefore, it is possible that increased intracortical inhibition is an important mechanism af-

fecting muscle force in AT patients.

Although increased SICI has been observed in patellar tendinopathy[14], it is unclear if this is also

present in chronic mid-portion AT patients. Thus, this study aimed to investigate intracortical inhibitory

mechanisms in runners with mid-portion AT and triceps surae isometric peak torque, rate of torque

development (RTD), and maximal number of single leg heel raises (SLHR) (endurance measure). We

hypothesized that runners with chronic mid-portion AT would present with greater levels of cortical

inhibition in the triceps surae and lower plantar flexor isometric peak torque, RTD, and number of

SLHR than controls.

2. Method

2.1 Study design and participant selection criteria

This was a cross-sectional study comparing runners with and without mid-portion AT. A sample

size of 10 participants was calculated based on the results of Rio et al, 2015[14] (GPower software param-

eters: effect size d= 2.63; α err prob: 0.05; power 0.95; n=10). Twenty-four runners were recruited and

allocated into two groups: mid-portion AT (n=11) and healthy controls (n=13). Participants' self-re-

ported height, body mass, age, gender, running mileage, VISA-A, and active motor threshold are re-

ported in Table 1. All volunteers were endurance runners, recruited from local running clubs in the area

of Southeast Queensland, Australia.

Diagnosis of mid-portion AT was confirmed by an experienced physiotherapist during examina-

tion if patients presented with localized mid-portion Achilles tendon pain for more than three months,

pain provoked by physical activities in a dose dependent way and had pain with palpation at the mid

portion of tendon. Once diagnosis was confirmed, inclusion and exclusion criteria were applied. Inclu-

sion criteria for the AT group were: 1) a diagnosis of mid-portion AT for > 3 months; 2) a running

routine of more than twice weekly for > 4 months; and 3) between 18-60 years of age and able to give

informed consent. Exclusion criteria for the AT group were: 1) insertional AT; 2) previous rupture or

surgery of the Achilles tendon; 3) clinical findings indicating a differential diagnosis for the Achilles

tendon pain, such as tendon tear; regular participation in other sports involving high speed running

(football, rugby, AFL etc.); 4) VISA-A score > 90 points; and 5) any other musculoskeletal injuries of

the lower limb; any neurological disorder; or mental health issues affecting consent. Inclusion criteria

for the control group were: 1) to be free of any current musculoskeletal injuries of the lower leg; 2) a

running routine of more than twice a weekly for > 4 months; and 3) between 18-60 years of age and

able to give informed consent. Exclusion criteria for the control group were: 1) previous injury of the

Achilles tendon; 2) regular participation in other sports involving high speed running (football, rugby,

AFL etc.); 3) VISA-A score < 100 points; and 4) any neurological disorder; or mental health issues

affecting consent. Prior to testing, all participants read and signed a detailed informed consent document

and completed the VISA-A questionnaire[16] and a TMS contra-indication questionnaire. All participants

were free of comorbidities such as cardiac, pulmonary, renal, endocrine of gastrointestinal and were not

taking any medication for tendon pain or that would affect tendon structure[17].

This study was approved by the Queensland University of Technology Human Research and

Ethics Committee in line with the Declaration of Helsinki. Data collection was conducted during the

COVID-19 pandemic and all safety procedures followed local state government policies.

2.2 Neuromuscular assessments

2.2.1 Torque measures and single leg heel raise

Plantar flexor isometric peak torque was measured using an isokinetic dynamometer (Biodex

Medical Systems, Shirley, New York). For the bilateral AT presentations (n=3), the most symptomatic

leg was used and for the control group, the dominant leg was used for testing. Leg dominance was

selected by asking the participants what their preferred leg was, and if they were unsure, they were

asked which leg they would use to kick a ball. Participants were positioned in a seated position (75

degrees of hip flexion) with their knee straight and with the foot perpendicular to shank. Warm up

consisted of 2×4 s isometric contractions of each participant's perceived 20, 40, 60 and 80% maximal

voluntary isometric contraction intensity. After warm-up, participants performed at least 3 maximal

voluntary isometric contractions, until <5% variation was observed between contractions, and the high-

est value was used. Participants were instructed to contract as hard and fast as possible and received

verbal encouragement and visual feedback of force output during the contractions. Absolute RTD was

calculated as the average slope of torque-time curve (0-100 ms) relative to onset of contraction. The

onset of contraction was determined by visually inspecting the torque trace and calculated from the time

immediately after torque increased from baseline. Relative RTD was calculated by dividing the absolute

RTD value by the contraction peak torque. After all neuromuscular and corticospinal assessments were

completed, participants were instructed to perform the SLHR endurance test to failure from a flat sur-

face, with each leg. Participants were instructed to lift the heel as high as possible, and return to flat

position, as many times as possible[18]. Test was ceased once participants started to perform SLHR test

with a reduced range of motion.

2.2.3 Electromyography (EMG)

After skin preparation, bipolar surface electromyography (sEMG) was used to record electrical

signals from soleus (SOL), gastrocnemius medialis (GM) and gastrocnemius lateralis (GL), and a ref-

erence electrode was placed on the lateral malleolus. sEMG signal was collected during the corticospi-

nal assessments (see below), and amplified (x1000), band-passed filtered (10-1000 Hz), digitized at 4

kHz. sEMG signals were recorded with PowerLab (16/35) and analysed with LabChart 8 software (AD

instruments, New Zealand).

2.2.4 Corticospinal assessments

SICI was assessed by measuring motor evoked potential (MEP) amplitudes elicited by TMS over

the primary motor cortex. The vertex (Cz) was located according to the 10-20 International System and

used as the anatomic reference point for stimulator coil placements. Coil was then, positioned over the

motor cortex area related to the lower limb[15,19]. Single and paired pulses were delivered using a double

cone coil connected to a Magstim[2] stimulator (Magstim, Dyfed, UK). Coil was positioned over the

motor cortex area, contralateral to the examined limb and oriented to induce a postero-anterior current

to the motor cortex. The optimal stimulation site for the triceps surae, was identified using a supra-

threshold stimulation intensity with a simultaneous background contraction (10% of peak torque)[20]. The

optimal site was considered the area to produce the largest MEP in SOL[15]. Thereafter, active motor

threshold was determined using a non-parametric algorithm tool for estimating TMS motor threshold,

MTAT2.0[21]. Motor threshold was defined as the minimal intensity of TMS output required to produce

a positive MEP in the target muscle in at least 5 of 10 trials. A positive MEP was considered when

peak-to-peak amplitude was greater than 0.1mV in SOL[15].

SICI was investigated using paired pulse stimulation with a conditioning stimulus of 0.8× active

motor threshold, followed by a test stimulus of 1.2× active motor threshold, using a 3-ms interstimulus

interval[19]. A single pulse was delivered at the same intensity as the test stimulus of 1.2x active motor

threshold, as reference to calculate the percentage of SICI. Participants performed a submaximal con-

traction of 10% of maximal voluntary torque during transcranial stimulation. To ensure correct back-

ground torque participants received real-time visual feedback on a monitor positioned in front of them.

The decision of whether to initiate the protocol with SICI or single pulse stimulation was randomized

for each participant. To ensure MEP reliability across conditions, twenty paired pulses and twenty sin-

gle pulses were delivered. The results were then compared as an index of intracortical inhibition,

whereas the higher the SICI value, the greater the intracortical inhibition. SICI is described as a per-

centage of the averaged peak-to-peak MEP amplitude and it was calculated using the formula: 100 -

(conditioned MEP/unconditioned MEP×100)[19,22]. Testers were not blinded to group during assessment.

2.3 Statistical analysis

Repeated measures ANCOVA was used to compare SICI values between muscles of the triceps

surae and between groups, as the AT was statistically older than the control group ($p<0.01$), age was

used as covariate in the model. ANCOVA was used to compare between group differences of torque

measures, active motor threshold and SLHR using age as covariate to account for statistical difference

found in age between groups. Robust paired t test was used to compare SLHR test between legs for

participants with unilateral AT (n=8). An alpha level of 5% was adopted for all statistical tests. Hedge's

g was calculated for estimates of effect size (presented as g). Hedge's g effect size interpretation is as

follows: ≤ 0.2 small effect; ≤ 0.5 medium effect; ≤ 0.8 large effect[23]. All statistical analyses were con-

ducted using Jamovi software (Version 1.6, Sydney, Australia). Data is presented as means and 95%

confidence intervals and effect size.

Table 1

Demographic data for Achilles Tendinopathy and healthy Control group

Variable	AT (n=11)	Control (n=13)	Mean difference
Age (years) *	44.1 (35.7 to 52.5)	34 (30.0 to 38.2)	10.2 (5.65 to 14.7)
Gender	6 males (54%)	7 males (54%)	-
Height (cm)	172.0 (165.9 to 178.1)	170.6 (165 to 176.2)	2.8 (-6.9 to 12.6)
Body Mass (Kg)	76.2 (65.8 to 86.6)	64.8 (58 to 71.8)	8.8 (-5.8 to 23.5)
Running mileage km/week	37.4 (26.6 to 48.2)	30.4 (22 to 38.8)	8.7 (-6.8 to 24.4)
VISA-A score *	70.7 (63.7 to 77.7)	100 (0)	-30.1 (-38.7 to -21.5)
Active Motor Threshold (stimulator output)	33.9 (29.7 to 38.1)	32.6 (28.8 to 36.4)	1.3 (-4.3 to 8.5)

Data represent Mean and 95%CI in brackets for each variable. * Denotes statistical differences.

3. Results

3.1 Corticospinal assessments

Triceps surae SICI was 14.3% (95%CI: -2.1 to 26.4; F=5.271, p=0.032, η^2p=0.201) higher in AT

than in controls (57.9%, 95%CI: 36.2 to 79.6; and 43.6%, 95%CI:16.2 to 71.1) (Figure 1); irrespective

of the tested muscle (F=0.929, p=0.403, η^2p=0.042); no muscle × group interaction was observed

(F=0.647, p=0.529, η^2p=0.030) and there was no affect from age in between groups (F=0.991, p=0.331,

η^2p=0.045). Mean active motor threshold did not differ between groups -1.3 stimulator output (-4.3 to

8.5; F=0.197, p=0.661, η^2p= 0.009).

Figure 1. **Repeated measures ANCOVA showing SICI individual data for all Triceps surae muscles: Soleus (SOL), Gastrocnemius medialis (GM) and Gastrocnemius lateralis (GL) between groups.** The AT group had ~ 14% more intracortical inhibition than the Control group. SICI was calculated and represents a percentage of inhibition; thus, the higher the SICI value, the greater the cortical inhibition. Data reported as Grand mean. * Denotes statistical difference between groups.

3.2 Neuromuscular assessments

We found no differences between groups in any of the torque measures (Table 2); isometric peak torque (F<0.001, p=0.971, η^2p<0.001); isometric peak torque normalised for body mass (F=1.000, p=0.329, η^2p=0.045); RTD 0-100ms (F=0.055, p=0.815 η^2p<0.001); RTD 0-100ms normalised for % peak torque (F=0.150, p=0.702, η^2p=0.007). For the SLHR test, the AT performed 16 fewer SLHR repetitions on the symptomatic side compared to controls (F=15.64, p<0.001, η^2p=0.427) and 14 fewer SLHR repetitions in the non-symptomatic side compared to controls (F=11.13, p=0.004, η^2p=0.382).

190 When analysing side to side differences of the AT, the symptomatic side performed on average, 2.3 less

191 SLHR repetitions compared to the non-symptomatic (g:-0.2, p=0.019), for this analysis, only unilateral

192 presentations were considered. Figure 2 displays data of SLHR values for visual comparison. No par-

193 ticipants reported pain during peak isometric torque test or SLHR test.

Table 2

Neuromuscular measures between AT and Control

Variable	AT (n=11)	Control (n=13)	Mean difference
Isometric peak Torque (N.m)	110.0 (84.1 to 136)	111.0 (87.1 to 134)	-0.6 (-35.4 to 34.1)
Isometric peak Torque/BM (N.m/BM)	1.4 (1.1 to 1.8)	1.7 (1.3 to 2.1)	-0.2 (-0.7 to 0.3)
RTD 0-100 (N.m)	239.0 (151.0 to 326.0)	224.0 (145.0 to 304.0)	14.3 (-117.2 to 132.8)
RTD 0-100/ peak torque (N.m/ % peak torque)	2.33(1.5 to 3.1)	2.12 (1.4 to 2.8)	0.2 (-0.7 to 1.2)
SLHR (Symptomatic vs Control) (repetitions) *	25.0 (19.3 to 30.7)	40.6 (35.4 to 45.8)	-15.6 (-23.3 to -7.9)
SLHR (Non-symptomatic vs Control) (repetitions) *	26.3 (19.7 to 32.9)	40.6 (35.4 to 45.8)	-13.9 (-22 to -5.8)
Variable	AT (Symptomatic n=8)	AT (Non-symptomatic n=8)	Mean difference
SLHR (AT Symptomatic vs AT Non-symptomatic) (repetitions) *	26.4 (18.4 to 44.4)	29.0 (21 to 37)	-2.3 (-3.63 to -1.0)

Data represent Mean and 95%CI in brackets for each variable.

Figure 2. **Single leg heel raise (SLHR) endurance test – AT vs Control (individual values).** Each point represents individual values of maximal SLHR performed by each participant during endurance test. Both the symptomatic and the non-symptomatic leg of AT group were weaker than the Control group. *AT symptomatic* refers to the most symptomatic leg; *AT Non symptomatic* SLHR used for analysis represent data only from unilateral presentations and for the *Control* group, the dominant leg was used for analysis. Data reported as mean and 95% CI. *; **; *** Denote statistical difference.

4. Discussion

This study aimed to determine whether runners with mid-portion AT had: 1) greater intracortical inhibition (measured as SICI) of the triceps surae muscles than healthy controls and 2) lower levels of muscle function (measured as isometric peak torque, rate of torque development and SLHR endurance test). We found that runners with mid-portion AT had higher SICI than controls, accompanied by re-duced SLHR endurance but without deficits in isometric peak torque or RTD. The data supports the primary hypothesis that runners with mid-portion AT have greater intracortical inhibition than controls; and partially supports our secondary hypothesis as runners with mid-portion AT have reduced triceps surae endurance. Contrary to the hypothesis, runners with AT were not weaker nor slower in developing torque than controls during isometric contractions.

As expected, SICI was higher in the AT group, outlining increased intracortical inhibition of the

triceps surae muscles (SOL, GM, GL), which could be negatively impacting triceps surae endurance.

To our knowledge, no other studies have investigated intracortical inhibitory mechanisms, using SICI

measures, in chronic mid-portion AT. SICI can be investigated with the target muscle at rest or during

a submaximal isometric force (5-10% peak torque); with the latter used in this study, and it measures

the activity of intracortical inhibitory circuits within the primary motor cortex (M1) mediated by

$GABA_A$ receptors,[19] one of the many circuits that control motor output. Increased cortical inhibition,

can reduce the neural drive to the muscle, impacting motor output[24], such as maximal strength[25] and

endurance[26]. We did not find differences in maximal strength measures; but one study in patellar tendi-

nopathy[14] reported increased SICI associated with a reduction in maximal voluntary contraction.

They found a reduction of intracortical inhibition, associated with reduction in tendon pain and

increased in maximal force following exercise; this study, however, did not have a control group to

compare SICI values with. Our results expand these previous findings by demonstrating that patients

with mid-portion AT also have higher intracortical inhibition, compared to controls. Additionally, there

is evidence[27] that GL has a 22.8% reduction in activation during submaximal isometric contractions in

AT; but our study was not structured to investigate individual muscle differences in neural drive. SICI

values can be influenced by age, with an increase in inhibition in older adults[28], for that reason we used

age as a covariate in our statistical model, and still we observed increased SICI in the AT group. This

is an important finding to help and better understand the neurophysiological characteristics present in

runners with Achilles tendinopathy and its potential impact in chronic triceps surae deficits observed in

AT; however, this study cannot address whether increased intracortical inhibition is driving the muscle

deficits observed and causing AT, or if this is an adaptation from long-lasting pain and tendon disfunc-

tion present in chronic AT. Furthermore, this study does not aim to drive changes in tendinopathy treat-

ment. An interesting point to note, triceps surae heavy resistance rehabilitation programs have yield

great improvements in both pain and function in AT patients; heavy resistance training is described in

the literature as effective in reducing cortical inhibition[29,30].

Although there were no differences between groups in measures of isometric peak torque, the

AT group showed a reduced capacity during the SLHR endurance test, suggesting a deficit in triceps

surae endurance. The lack of group difference in plantar flexor isometric peak torque is in line with the

result of other studies[27,31]. In contrast, weakness in plantar flexor dynamic peak torque has been widely

reported in AT[5,9,12]. Perhaps, as observed in other chronic musculoskeletal disorders, such as chronic

hamstring strains[32], chronic AT presents deficits in dynamic but not in isometric peak torque. It is in-

teresting to note that the AT group was on average 8.8 kg heavier that the control group, and yet there

were no differences in torque measures in absolute or in normalised values. To the authors' knowledge,

studies have not yet compared concentric, eccentric, and isometric torque measures from the same group

of AT patients, against a control group. Further research analysing different contraction types is neces-

sary to fully characterize triceps surae`s deficits in AT and help guide better rehabilitation strategies.

Explosive plantar flexor torque, measured as the RTD, was not different between groups in ab-

solute or normalized values, suggesting chronic AT has no impact on this important parameter. A recent

meta-analysis[9] found conflicting evidence about reductions in RTD in AT; perhaps other factors could

be impacting hop distance, rather than explosive force, such as increased tendon compliance observed

in chronic AT[31]. Our study compared RTD from the most symptomatic leg of AT against healthy con-

trols, whereas all other studies investigating RTD in AT, compared RTD between symptomatic and

non-symptomatic legs of the same subjects. As a consequence, there is limited evidence to compare our

data to.

Kinesiophobia (fear of movement) influences endurance recovery in AT patients; patients with

higher levels of kinesiophobia recovered less of their muscle endurance in a 5-year follow-up[2]. This

interesting finding shows how fear of moving in a certain way or using a specific body part can impact

the neural drive to the muscle, possibly involving inhibitory mechanisms within the cortex, as observed

in the AT group.

The SLHR is a common clinical test of triceps surae endurance[18,33] and function in AT patients[33].

Our data showed a reduction of 38% in triceps surae endurance on the symptomatic side and 34% on

the non-symptomatic side in unilateral presentations, compared to controls. Although we found a dif-

ference between sides of the AT group, the effect size was only small. Participants performed SLHR

first on their symptomatic/dominant leg, followed by their non-symptomatic/non-dominant leg. The

non-symptomatic leg of the AT group was performed immediately after the symptomatic leg. Thus, it

is possible that the difference observed between the symptomatic and the non-symptomatic observed in

the AT group were attenuated by mechanisms of central fatigue.

Studies that have adopted the SLHR endurance test[2] or calculated the total work done in dyna-

mometry[5] for endurance measurement have also observed a reduction in endurance capacity compared

to the non-symptomatic side and to controls, respectively. Bilateral deficits in unilateral presentations

have been widely described in chronic AT[9,12]. These deficits on both symptomatic and non-symptomatic

sides, emphasize that the non-symptomatic leg should not be used as comparison for neuromuscular

function in AT. It is possible that the degree of cortical inhibition present in our AT participants, is

insufficient to impact muscle performance during a brief maximal voluntary isometric contraction, but

sufficient to reduce performance during the SLHR test, once additional inhibitory impacts (sensory

inhibitory drive to motor neurons or higher centres) are added. Furthermore, it would also be relevant

to investigate the impact neural mechanisms would have in muscular responses to fatigue in AT.

Our AT group had greater running mileage/week than controls (Table 1). AT is an overloading

injury and higher mileage could be a contributing factor for the development of AT. It is suggested that,

amongst other factors, load magnitude and recovery time, may influence tendon response and help de-

velop tendinopathy[34].

The literature shows no association between tendon structure and levels of tendon pain and

dysfunction[35]; however, not having scanned participants tendons could be considered a limitation of

this study. In this study we had 3 runners with bilateral AT and 8 runners with unilateral AT. It is still

unknown if runners with bilateral AT present specific cortical changes in comparison with runners with

unilateral AT. Futures studies investigating neurophysiological differences between unilateral and

bilateral AT could help to fill this gap.

Unilateral AT presentations displayed bilateral deficits in muscle endurance; yet, without

presenting any symptoms; thus, having scanned participants bilateral tendons, would have helped

profiling our participants for further analysis and understanding. Testers were not blinded for group,

which is also a limitation of this study.

5. Perspective

Runners with mid-portion AT have greater intracortical inhibition to the triceps surae muscles

accompanied by a bilateral reduction in triceps surae endurance, despite no deficits in isometric peak

torque or explosive torque. The increased SICI observed in the AT group may be negatively influencing

triceps surae endurance; thus, rehabilitation strategies aiming to reduce intracortical inhibition should

be considered for better outcomes. Furthermore, the SLHR test, is a useful clinical tool to assess triceps

surae function in patients with mid-portion AT as an outcome measure of improvement.

Conflict of interests

The authors declare no conflict of interest with the present research.

Acknowledgments

The authors declare not having received any funding for this study.

The authors thank the following for their contribution to the development and achievement of

this research:

Miss Charmaine Enculescu who assisted with figure plotting;

All volunteers who participated in this study.

References

1.   Brund RBK, Rasmussen S, Kersting UG, Arendt-Nielsen L, Palsson TS. Prediction of running-

316   induced Achilles tendinopathy with pain sensitivity - A 1-year prospective study. *Scand J Pain*.

317   2019;19(1):139-146. doi:10.1515/sjpain-2018-0084

318   2.   Silbernagel KG, Brorsson A, Lundberg M. The majority of patients with Achilles tendinopathy

319   recover fully when treated with exercise alone: A 5-year follow-up. *Am J Sports Med*.

320   2011;39(3):607-613. doi:10.1177/0363546510384789

321   3.   McAuliffe S, Tabuena A, McCreesh K, et al. Altered strength profile in Achilles tendinopathy:

322   A systematic review and meta-analysis. *J Athl Train*. 2019;54(8):889-900. doi:10.4085/1062-

323   6050-43-18

4.   Mahieu NN, Witvrouw E, Stevens V, Van Tiggelen D, Roget P. Intrinsic risk factors for the

development of Achilles tendon overuse injury: A prospective study. *Am J Sports Med*.

2006;34(2):226-235. doi:10.1177/0363546505279918

5.   O'Neill S, Barry S, Watson P. Plantarflexor strength and endurance deficits associated with mid-

portion Achilles tendinopathy: The role of soleus. *Phys Ther Sport*. 2019;37:69-76.

doi:10.1016/j.ptsp.2019.03.002

6.   Silbernagel KG, Thomeé R, Eriksson BI, Karlsson J. Full symptomatic recovery does not ensure

full recovery of muscle-tendon function in patients with Achilles tendinopathy. *Br J Sports Med*.

2007;41(4):276-280. doi:10.1136/bjsm.2006.033464

7.   Van Der Plas A, De Jonge S, De Vos RJ, et al. A 5-year follow-up study of Alfredson's heel-

drop exercise programme in chronic midportion Achilles tendinopathy. *Br J Sports Med*.

2012;46(3):214-218. doi:10.1136/bjsports-2011-090035

8.   Wang HK, Lin KH, Wu YK, Chi SC, Shih TTF, Huang YC. Evoked spinal reflexes and force

development in elite athletes with middle-portion achilles tendinopathy. *J Orthop Sports Phys*

Ther. 2011;41(10):785-794. doi:10.2519/jospt.2011.3564

9.   Hasani F, Vallance P, Haines T, Munteanu SE, Malliaras P. Are Plantarflexor Muscle

Impairments Present Among Individuals with Achilles Tendinopathy and Do They Change with

Exercise? A Systematic Review with Meta-analysis. *Sport Med - Open*. 2021;7(1).

doi:10.1186/s40798-021-00308-8

10.   Plinsinga ML, Brink MS, Vicenzino B, Van Wilgen CP. Evidence of nervous system

sensitization in commonly presenting and persistent painful tendinopathies: A systematic

review. *J Orthop Sports Phys Ther*. 2015;45(11):864-875. doi:10.2519/jospt.2015.5895

11.   Heales LJ, Lim ECW, Hodges PW, Vicenzino B. Sensory and motor deficits exist on the non-

injured side of patients with unilateral tendon pain and disability - Implications for central

nervous system involvement: A systematic review with meta-analysis. *Br J Sports Med*.

2014;48(19):1400-1406. doi:10.1136/bjsports-2013-092535

12.   Masood T, Kalliokoski K, Bojsen-Møller J, Peter Magnusson S, Finni T. Plantarflexor muscle

function in healthy and chronic Achilles tendon pain subjects evaluated by the use of EMG and

PET imaging. *Clin Biomech*. 2014;29(5):564-570. doi:10.1016/j.clinbiomech.2014.03.003

13. Kandel ER, Schwartz JH, Jessell TM, Siegelbaum SA H. *Principles of Neural Science*.; 2013.

14. Rio E, Kidgell D, Purdam C, et al. Isometric exercise induces analgesia and reduces inhibition

in patellar tendinopathy. *Br J Sports Med*. 2015;49(19):1277-1283. doi:10.1136/bjsports-2014-

094386

15. Rossini PM, Burke D, Chen R, et al. Non-invasive electrical and magnetic stimulation of the

brain, spinal cord, roots and peripheral nerves: Basic principles and procedures for routine

clinical and research application: An updated report from an I.F.C.N. Committee. *Clin*

Neurophysiol. 2015;126(6):1071-1107. doi:10.1016/j.clinph.2015.02.001

16. Martin RL, Chimenti R, Cuddeford T, et al. The VISA-A questionnaire: An index of the severity

of Achilles tendinopathy I. *J Orthop Sports Phys Ther*. 2018;48(5):9-11.

https://pubmed.ncbi.nlm.nih.gov/11579069/

17. Knobloch K. Drug-Induced Tendon Disorders. In: Ackermann PW, Hart DA, eds. *Metabolic*

Influences on Risk for Tendon Disorders. Springer International Publishing; 2016:229-238.

doi:10.1007/978-3-319-33943-6_22

18. Hébert-Losier K, Wessman C, Alricsson M, Svantesson U. Updated reliability and normative

values for the standing heel-rise test in healthy adults. *Physiother (United Kingdom)*.

2017;103(4):446-452. doi:10.1016/j.physio.2017.03.002

19. Di Lazzaro V, Rothwell J, Capogna M. Noninvasive Stimulation of the Human Brain: Activation

of Multiple Cortical Circuits. *Neuroscientist*. 2018;24(3):246-260.

doi:10.1177/1073858417717660

20. Neige C, Grosprêtre S, Martin A, Lebon F. Influence of voluntary contraction level, test stimulus

intensity and normalization procedures on the evaluation of short-interval intracortical

inhibition. *Brain Sci*. 2020;10(7):1-16. doi:10.3390/brainsci10070433

21. Ah Sen CB, Fassett HJ, El-Sayes J, Turco C V., Hameer MM, Nelson AJ. Active and resting

motor threshold are efficiently obtained with adaptive threshold hunting. *PLoS One*.

2017;12(10):1-9. doi:10.1371/journal.pone.0186007

22. Lauber B, Gollhofer A, Taube W. Differences inmotor cortical control of the soleus and tibialis

anterior. *J Exp Biol*. 2018;221(20). doi:10.1242/jeb.174680

23.   Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: A practical

primer for t-tests and ANOVAs. *Front Psychol*. 2013;4(NOV). doi:10.3389/fpsyg.2013.00863

24.   Brownstein CG, Ansdell P, Škarabot J, et al. Motor cortical and corticospinal function differ

during an isometric squat compared with isometric knee extension. *Exp Physiol*.

2018;103(9):1251-1263. doi:10.1113/EP086982

25.   Lahouti B, Lockyer EJ, Wiseman S, Power KE, Button DC. Short-interval intracortical

inhibition of the biceps brachii in chronic-resistance versus non-resistance-trained individuals.

Exp Brain Res. 2019;237(11):3023-3032. doi:10.1007/s00221-019-05649-1

26.   Brownstein CG, Espeit L, Royer N, Lapole T, Millet GY. Fatigue-induced changes in short-

interval intracortical inhibition and the silent period with stimulus intensities evoking maximal

versus submaximal responses. *J Appl Physiol*. 2020;129(2):205-217.

doi:10.1152/japplphysiol.00282.2020

27.   Crouzier M, Tucker K, Lacourpaille L, et al. *Force-Sharing within the Triceps Surae: An*

Achilles Heel in Achilles Tendinopathy. Vol 52.; 2020. doi:10.1249/MSS.0000000000002229

28.   McGinley M, Hoffman RL, Russ DW, Thomas JS, Clark BC. Older adults exhibit more

intracortical inhibition and less intracortical facilitation than young adults. *Exp Gerontol*.

2010;45(9):671-678. doi:10.1016/j.exger.2010.04.005

29.   Beyer R, Kongsgaard M, Hougs Kjær B, Øhlenschlæger T, Kjær M, Magnusson SP. Heavy slow

resistance versus eccentric training as treatment for achilles tendinopathy: A randomized

controlled trial. *Am J Sports Med*. 2015;43(7):1704-1711. doi:10.1177/0363546515584760

30.   Latella C, Goodwill AM, Muthalib M, et al. Effects of eccentric versus concentric contractions

of the biceps brachii on intracortical inhibition and facilitation. *Scand J Med Sci Sport*.

2019;29(3):369-379. doi:10.1111/sms.13334

31.   Child S, Bryant AL, Clark RA, Crossley KM. Mechanical properties of the achilles tendon

aponeurosis are altered in athletes with achilles tendinopathy. *Am J Sports Med*.

2010;38(9):1885-1893. doi:10.1177/0363546510366234

32.   Maniar N, Shield AJ, Williams MD, Timmins RG, Opar DA. Hamstring strength and flexibility

after hamstring strain injury: A systematic review and meta-analysis. *Br J Sports Med*.

2016;50(15):909-920. doi:10.1136/bjsports-2015-095311

33.   Silbernagel KG, Gustavsson A, Thomeé R, Karlsson J. Evaluation of lower leg function in

411   patients with Achilles tendinopathy. *Knee Surgery, Sport Traumatol Arthrosc*.

412   2006;14(11):1207-1217. doi:10.1007/s00167-006-0150-6

413   34.   Cook JL, Rio E, Purdam CR, Docking SI. Revisiting the continuum model of tendon pathology:

414   What is its merit in clinical practice and research? *Br J Sports Med*. 2016;50(19):1187-1191.

415   doi:10.1136/bjsports-2015-095422

416   35.   Docking SI, Rio E, Cook J, Carey D, Fortington L. Quantification of Achilles and patellar tendon

417   structure on imaging does not enhance ability to predict self-reported symptoms beyond grey-

418   scale ultrasound and previous history. *J Sci Med Sport*. 2019;22(2):145-150.

419   doi:10.1016/j.jsams.2018.07.016

420