Changes in non-oscillatory features of the cortical sensorimotor rhythm in Parkinson's disease across age

Authors

Mikkel C. Vinding* 1), Allison Eriksson 1)2), Cassia Man Ting Low 1)3), Josefine Waldthaler 4)5),

Daniel Ferreira 6,7), Martin Ingvar 1), Per Svenningsson 4), Daniel Lundqvist 1).

Affiliations

1. NatMEG, Department of Clinical Neuroscience, Karolinska Institutet, Sweden.

2. Department of Women's and Children's Health, Uppsala University, Sweden.

3. Cognitive Neuroimaging Centre, Lee Kong Chien School of Medicine, Nanyang Technological University, Singapore.

4. Section of Neurology, Department of Clinical Neuroscience, Karolinska Institutet, Sweden.

5. Department of Neurology, University Hospital Marburg, Marburg, Germany.

6. Division of Clinical Geriatrics, Center for Alzheimer's Research, Department of Neurobiology, Care Sciences, and Society, Karolinska Institutet, Stockholm, Sweden.

7. Department of Radiology, Mayo Clinic, Rochester, MN, USA.

Corresponding author

Mikkel C. Vinding

Department of Clinical Neuroscience
Karolinska Institutet
Nobels väg 9, D2
171 77 Stockholm
Sweden

Email: mikkel.vinding@ki.se

Abstract

Parkinson's disease (PD) is associated with changes in neural activity in the sensorimotor alpha and beta bands. Using magnetoencephalography (MEG), we investigated the role of spontaneous neuronal activity within the somatosensory cortex in a large cohort of early- to mid-stage PD patients (N = 78) and age- and sex matched healthy controls (N = 60) using source reconstructed resting-state MEG. We quantified features of the time series data in terms of oscillatory alpha power, beta power, and 1/f broadband characteristics using power spectral density, and also characterised transient beta burst events in the time-domain signals. We examined the relationship between these signal features and the patients' disease state, symptom severity, age, sex, and cortical thickness.

PD patients and healthy controls differed on PSD broadband characteristics, with PD patients showing a steeper 1/f exponential slope and higher 1/f offset. PD patients further showed a steeper age-related decrease in the burst rate. Out of all the signal features of the sensorimotor activity, only burst rate was associated with increased severity of bradykinesia. Our study shows that general non-oscillatory features (broadband PSD slope and offset) of the sensorimotor signals are related to disease state and *oscillatory burst rate* scales with symptom severity in PD.

1 Introduction

Parkinson's disease (PD) is a common neurodegenerative disease characterised by a gradual

accumulation of Lewy bodies and death of dopaminergic neurons.[1,2] The Lewy body pathology of PD

begins long before the manifestation of motor symptoms. Accumulation of Lewy bodies is initially

found in the olfactory bulb and brain stem and then spreads to the substantia nigra pars compacta,

followed by several brain regions, including the basal ganglia and the neocortex.[3] The progressive

structural and neurochemical changes in PD are accompanied by widespread functional changes in

neuronal activity, which in turn lead to worsening clinical signs and symptoms such as tremor, rigidity,

and bradykinesia and co-occurring non-motor symptoms like sleep disorders, depression, fatigue, and

cognitive deficits.[1]

The changes in brain function in PD are particularly prominent in the oscillatory activity of neurons.[4]

In PD, spontaneous oscillatory beta band (13–30 Hz) activity in the sub-thalamic nucleus (STN)

exhibits a systematic disease-related increase in synchronicity that is related to the dopamine level[5-8],

and correlates with the severity of bradykinesia and rigidity symptoms.[9,10] Changes in the beta band

extend beyond the STN through the basal ganglia-thalamic cortical sensorimotor network. The cortical

manifestation of the disease-related changes in the sensorimotor network can be measured non-

invasively from the cortex, using electro- or magnetoencephalography (EEG/MEG). Such non-invasive

neural recordings can potentially provide easily available prospective biomarkers of disease or

symptom-related neural changes in PD. Increased oscillatory beta-band activity in the sensorimotor

cortex has been linked to increased symptom severity, such as rigidity and bradykinesia.[11,12] The role of

dopamine on the cortical beta band is, however, still unclear. There is no consensus on how

dopaminergic medication affects cortical beta-band power, with some studies reporting no effects[11,13-]

[15] and others an increase in beta-band power.[16–18] Deep brain stimulation of the STN in PD patients has

been shown to lead to a decrease in the power of spontaneous activity in the cortical sensorimotor beta

and alpha (8-12Hz) bands[19,20] (but see also[16,21]).

Importantly, there is evidence that the beta-band changes are not in the same direction across the

different stages of PD. For example, there are reports of increased cortical beta-band power in the early

stages of PD[22], whereas the later stages are associated with decreased beta-band power.[23] Further, the

beta-band power is not the *only* feature of the sensorimotor rhythms that is altered in PD. Several

studies have found a shift in the beta-band centre frequency (the frequency at which the power

spectrum density peaks in the beta-band) towards a lower frequency in PD patients compared to

healthy controls.[24–26] The shift towards lower beta-band centre frequency is more pronounced in PD

patients with dementia[27–30] and correlates with reduced cognitive ability.[26,31] Notably, the centre

frequency shift is detectable already in the early stages of PD[25] and does not seem to be affected by

dopaminergic medication.[32] The changes in beta-band power and centre frequency in PD could indicate

that different features of the oscillatory beta-band activity reflect different underlying neural functions

expressed in the measured sensorimotor signals. Changes in beta-band power could be functionally

related to sensorimotor disturbances, and changes in centre frequency could be related to cognitive

function.

The characteristics of neuronal oscillatory activity may hold additional information of disease-related

changes in PD.[33] Both beta-band power and centre frequency reflect a quantification of power spectral

density (PSD). While these features can provide valuable information about disease-related changes in

PD, the PSD quantification of a neural time series provides a static summary of the oscillatory activity

across the entire time series. PSD does not account for inherent dynamics in this activity or changes in

the time series on shorter time scales—as is prevalent in neural time series. The beta-band exhibits a

great degree of variation over time and contains characteristic high-amplitude "bursts" that last about

50-200 ms, both in the cortical and sub-cortical beta-bands.[34–37] Functionally, the transient bursts

appear to play a pivotal role in sensorimotor processing through the basal ganglia-thalamic-cortical

network. For instance, the presence of a beta burst in the sensorimotor cortex close to a tactile

stimulation decreased the likelihood of tactile detection[38], and the rate of beta bursts is shown to

decrease in the time leading up to a movement both in STN[39–41] and in the sensorimotor cortex.[42,43]

In PD, quantification of beta-band burst activity from recordings in the STN has shown that beta-burst

rate and duration are reduced by dopaminergic medication[44,45] and deep brain stimulation.[37]

Furthermore, PD patients exhibit a decrease in the rate of beta burst at the cortical level compared to

healthy controls.[14] This decrease in beta burst rate is inversely related with increased severity of motor

symptoms;[46] particulary bradykinesia and postural-kinetic tremor symptoms, but there is not evidence

pointing to an effect of dopaminergic medication on cortical bursting properties.[14] Notably, the burst

rate showed a higher sensitivity than PSD beta power for discriminating PD patients from healthy

controls, demonstrating that the choice of method for analysing beta-band features influences the

sensitivity of subsequent analyses. This is further complicated by the fact that in addition to disease-

related changes, these features likely differ with age,[43,47] and the fact that most studies on oscillatory

changes in PD come from studies with small sizes.[48] The central challenge is quantifying the measured

neural signals to extract the disease's relevant features from the signals, be it the spectral power, centre

frequencies, or burst-like features.

In the current study, we aimed to compare how different oscillatory features of cortical sensorimotor

activity change in PD to elucidate what oscillatory features in the neural time-series differ between PD

patients and healthy controls and how these features are associated with different motor symptoms in

PD. We extracted the sensorimotor neural resting-state activity from source reconstructed resting-state

MEG signals in the sensorimotor cortex (Figure 1) and quantified the time-series in terms of the PSD in

the canonical mu-band (8-30 Hz).[49,50] In addition to the band-specific analysis, we compared the 1/f

broadband characteristics of the PSD.[51,52] Finally, we compared features of the sensorimotor rhythm in

terms of time-domain analysis of spontaneous transient bursts.[14,38] We tested the hypotheses of altered

functional changes in PD by analysing how these features differed between PD patients and healthy

controls and further investigated the interactions with age and sex. As ageing is associated with

structural and functional changes in the sensorimotor cortex[53,54], we investigated if the potential

changes in sensorimotor activity in PD differed across age. Since both healthy ageing and PD disease

progression are linked to thinning of the cortex[55,56], we further included thickness of the sensorimotor

cortex in the analysis as a potential mediating factor on the sensorimotor activity that potentially also

interacts with disease state.

The central hypothesis was that there would be differences between healthy controls and PD in features

of the sensorimotor signals, but also that different features may be related to different functional

changes. We hypothesised that individual oscillatory features would reflect different underlying neural

functions in the sensorimotor system and thereby show different relationships to the clinical

manifestations of specific motor symptoms in PD. We tested this hypothesis in two steps: first,

examining the inter-relationship between all different measures, and subsequently, examining what

feature—or combination of features—best explained the variation in severity within each motor

symptom.

Figure 1: Overview of the data processing pipeline. Three minutes raw resting-state MEG data was

obtained from each participant. The signals were projected through a minimum-norm source

reconstruction to extract the activity in the sensorimotor cortex. We did a frequency decomposition of

the source reconstructed signal to calculate the PSD to which a 1/f and Gaussian curve were fitted to

extract the PSD features (Table 2). In addition, we quantified sensorimotor bursts in the signal time

109 series in the sensorimotor ROI by thresholding the envelope of the band-pass filtered (8-30 Hz) signal

110 to the mu-beta frequency range.

2 Results

112 To enable a sensitive assessment of disease-related oscillatory changes in PD, we aimed for a relatively

113 large cohort compared to other functional neuroimaging studies of PD patients (N=78) and healthy

114 controls (N=60), balanced across gender and age, and with gender- and age-matched groups (Table 1).

115 Table 1: Group-level summary of the participants included in the analysis. Mean (standard deviation).

Measure	Parkinson's patients	Healthy controls	Statistics
N	78	60	
Sex (female/male)	29/49	27/33	$\chi^2 = 0.57$, p = 0.45
Age	65.6 (9.5)	63.93 (8.4)	Welsh t(138.0) = 1.08, p = 0.28
Disease duration	4.4 (3.7) years	-	-
LEDD	548 (273) mg	-	-
MDS-UPDRS-III	18.9 (10.8)	-	-
MoCA	26.1 (2.8)	26.2 (2.1)	Welsh t(136) = 0.10, p = 0.92

117 Table 2: Summary explanations of the main outcome variables in the analysis

Variable category	Variable	Explanation
PSD	Beta power	The maximum peak in the 13-30 Hz band. Estimated as the height of the Gaussian function fitted to the PSD after regressing out the 1/f spectrum.

	Beta centre frequency (Hz)	The dominant frequency bin in the 13-30 Hz band. Estimated as the centre of the Gaussian function fitted to the 13-30 Hz range of the PSD after regressing out the 1/f spectrum.
	Alpha power	The maximum peak in the 8-12 Hz band. Estimated as the height of the Gaussian function fitted to the PSD after regressing out the 1/f spectrum.
	Alpha centre frequency (Hz)	The dominant frequency bin in the 8-12 Hz band. Estimated as the centre of the Gaussian function fitted to the 8-12 Hz range of the PSD after regressing out the 1/f spectrum.
	1/f offset	The intercept of the log-linear regression line estimated from the full PSD in the 0.5-40 Hz range.
	1/f exponent	The decay exponent ($1/f^x$) of the PSD, corresponding to the slope of the log-linear regression line, estimated from the full PSD in the 0.5-40 Hz range.
Burst	*Rate*	The number of burst events in the time series divided by the length of the time series
	Duration (ms)	The time point from where the signal envelope rise above the threshold until the next time point it drops below the threshold.
	Interval (ms)	The time point from where the signal envelope drops below the threshold until the next time point it rises above the threshold.
	Amplitude	The maximum envelope amplitude within a burst event.

119   For the first analysis, we investigated how features in the resting-state activity from the sensorimotor

120   area quantified by features of the PSD and burst characteristics (see Table 2) differed as a function of

121   the predictors *group* (PD patients/healthy controls; Table 1), *age*, *sex*, and *cortical thickness* as well as

122   the interaction between the predictors.

2.1 PSD features

The PSD suggests an apparent group difference between PD patients and healthy controls in the alpha

and beta bands (Figure 2). However, analysing the oscillatory components of PSD by first adjusting for

the broadband characteristic of the PSD[51] removed the apparent group difference in the mu- and beta

band. Bayesian model comparison was used to test which predictors explained the variation in the PSD

(Bayes Factors (BF) > 3 taken as cutoff for substantial evidence for an effect of a given predictor[57]).

The model comparison showed evidence for group differences on the 1/f offset (BF = 37.77) and 1/f

exponent (BF = 5.92). There were no substantial evidence of an effect beyond the threshold for

substantial evidence on any other PSD features. However, it might be worth noting that there was

anecdotal evidence of an effect of cortical thickness on 1/f exponent (BF = 1.73), an interaction effect

between age and group on alpha centre frequency (BF = 1.43), and only minute evidence for a group

difference on beta power (BF = 1.28). The coefficients of regression models analysing the effect of the

predictors on PSD for all outcome measures are presented in Figure 3.

Figure 2: Group-level PSD. Grand average PSD (mean+standard error) for the PD group (blue) and

healthy control group (red).

Analysing the PSD by regressing out the 1/f contribution to the spectrum, the difference between PD

patients and healthy controls is not so much in the canonical beta- and alpha bands but manifests in the

broadband characteristics of the signal. The 1/f intercept was 23.5% [CI: 11.8:33.7%] higher for PD

patients than healthy controls, and PD patients had on average 11.9% [CI: -30.4:3.8%] steeper 1/f

exponential slope compared to healthy controls.

Figure 3: Regression analysis of PSD features. Standardized regression coefficients for the analyses

of sensorimotor PSD features as a function of group, age, sex, cortical thickness and the interaction

between these factors. * indicate factors with Bayes Factor > 3 in the model comparison.

2.2   Burst features

The view of sensorimotor oscillatory activity has recently changed from a steady oscillating signal to

viewing the activity in the sensorimotor bands occurring in short bursts. We compared features of the

sensorimotor rhythm in terms of time-domain analysis of spontaneous transient bursts in the time

series. Model comparison to test which predictors explained the variation in the burst features gave

evidence for an interaction effect between group and age on the burst rate (BF = 33.57) and a main

effect of sex on the burst rate (BF = 4.34). Model coefficients are displayed in Figure 4.

The age-related effects from the model amount to a change in burst rate of -0.7% [CI: -1.6:0.2] per year

for female PD patients and -1.7% [CI: -3.1:-0.3] change in burst rate per year for male PD patients,

whereas female controls had a relative increase in burst rate of 0.8% [CI: 0.1:1.7] per year and male

controls had a stable trend of -0.2% [CI: -1.0:0.6] change per year.

Figure 4: Regression analysis of burst features. Standardized regression coefficients for the analyses

of burst features as a function of group, age, sex, cortical thickness and the interaction between these

factors. * indicate factors with Bayes Factor > 3 in the model comparison.

No other predictors showed evidence of an effect beyond the threshold for substantial evidence on

neither burst length, the interval between bursts, nor burst amplitude. There was anecdotal evidence

(i.e. $1/3 < BF < 3$) for a group difference in burst amplitude (BF = 2.08) as well as an interaction effect

of age and cortical thickness on the burst rate (BF = 1.43), and an interaction between sex and cortical

thickness on the burst rate (BF = 1.86).

Figure 5: Age-related differences in sensorimotor signal features. Scatterplots of the individual

measures and model predictions over age for (A) burst rate, (B) PSD broadband 1/f intercept, and (C)

PSD broadband 1/f exponent, split between PD patients (blue) and healthy controls (red), and female

(solid) and male (dashed).

2.3   Clinical symptoms and oscillatory features

In the second analysis, we tested for associations between motor symptoms and the features of the

sensorimotor signal in the PD group. All sensorimotor signal features listed in Table 2 were used as

predictors in a multiple regression analysis that further included age, sex, and cortical thickness to

regress out the contribution hereof. The standardised regression coefficients of each predictor variable

on the motor symptoms measured with MDS-UPDRS-III[58] are presented in Figure 6.

Model comparison of multiple regression models showed evidence that burst rate was negatively

associated with upper limbs bradykinesia (BF = 12.70). The negative direction of the effect of burst

rate means that reduced burst rate was associated with increased severity of bradykinesia. The analysis

yielded no substantial evidence for effects of other sensorimotor signal features on symptom ratings for

axial symptoms, rest tremor, rigidity, rest tremor, postural/kinetic tremor, nor lower limb bradykinesia.

The analysis yielded substantial evidence for an effect of age on axial symptoms (BF = 5.79) and

evidence for a difference between male and female patients on axial symptoms (BF = 7.56) and rest

tremor (BF = 6.06). There were anecdotal evidence for an effect of alpha centre frequency on upper

limbs bradykinesia (BF = 2.63), an effect of burst length on axial symptoms (BF = 1.49), as well as

anecdotal evidence for an effect of cortical thickness on upper limbs bradykinesia (BF = 2.63).

Figure 6: Regression analysis of motor symptoms. Standardized regression coefficients for the

regression analyses of sensorimotor signal features on clinical motor symptom ratings in PD. * indicate

factors with Bayes Factor > 3 in the model comparison.

3 Discussion

This study aimed to explore how different features of cortical somatosensory oscillatory activity at rest

differed between PD patients and healthy controls across age and gender and how these features relate

to motor symptoms in PD. The analysis of spontaneous sensorimotor bursts showed an increased age-

related reduction in PD patients compared to healthy controls. Notably, our current results show that

the reduced burst rate in PD is not a static group-level difference but interacts with age with a steeper

age-related reduction in burst rate in PD compared to healthy controls.

We hypothesised that different oscillatory features reflect distinct underlying functional neural

properties and manifest as different motor symptoms in PD. The results showed that a reduction in mu-

beta burst rate in PD was accompanied with an increase in bradykinesia severity, confirming previous

findings from our group.[14] This relationship was exclusive for the bursts rate, as we observed no

statistically evident relationships between other burst features or PSD features and clinical motor

symptoms.

3.1 Characterizing neural time series data

The cortical sensorimotor activity of the PD patients differed from healthy controls, but did not differ

on the PSD in the canonical sensorimotor mu and beta bands. The PD patients showed a steeper

broadband 1/f slope and exponent of the broadband PSD than healthy controls. The observation of a

group difference in the spectral broadband 1/f characteristics of the signal adds to the growing evidence

that a focus on neural activity as narrow-band steady oscillations—e.g. narrowly focusing only on the

beta-band power—could potentially miss essential aspects of the neural signals for understanding

mechanistic changes in disease.[59,60] Widening the quantitative analysis of PSD rather than focusing

exclusively on narrow band activity is of potential clinical value: quantifying only the peaks in the PSD

to differentiate patients from controls can misrepresent the actual oscillatory response at those

frequencies as the peaks are influenced by the broadband offset and 1/f exponent. Any unaccounted-for

systematic differences in either PSD offset or decay exponent can lead to a false conclusion that there

is a difference in the oscillatory response.[51,52] Non-invasive measurements of changes in sensorimotor

activity is—despite the often conflicting findings[48]—a potential useful method to assess disease-related

changes. At the current stage of the field there is, however, a need to further bridge how features in the

signals are linked to disease mechanisms.

Analysis of neural time series by frequency decomposition is a powerful tool to extract and summarise

features of the signal, but the method comes with limitations in what one can infer. In the time domain,

increased oscillatory power can reflect both increased burst duration and change in burst amplitude and

as an expression of true sustained oscillations in the signal.[36,61] The presence of more sustained

oscillations in the sensorimotor rhythm might reflect a higher level of inhibition of sensorimotor

information; as is seen in recordings from STN[5] and, to some extent, also at the cortical level.[38,62]

However, sustained oscillations are not in contrast to the bursting properties of the sensorimotor

rhythm. The neural time-series can express both a degree of sustained oscillations while also exhibiting

variation in the degree of transient bursts—e.g., a signal of steady oscillation with transient high-

amplitude bursts.

3.2 Beta bursts and beta activity

Bursting properties of the cortical sensorimotor neural activity are proposed to occur due to long-range

input through the ascending thalamic-cortical connection to the cortex, leading to an increase in the

local neural excitation and resulting in a burst of synchronous activity.[36] The observed disease-related

changes in spontaneous cortical bursts, in the form of a more rapid decrease in rate over age for PD

patients, could reflect inhibition of these projections along the thalamic-cortical pathways caused by

disturbances in the dopamine-dependent structures projecting to the cortex. Interestingly, we did not

find significant group differences in burst duration in the current study—in line with previously

reported findings on cortical burst in PD[14]—supporting the view that the central mechanisms of the

cortical bursts are not primarily affected in PD—instead, it is the *rate* of bursts that is reduced at the

cortical level. The sub-cortical beta-band activity is influenced by the activity of dopamine-responding

neurons[5,6,45]. The effect of dopamine and dopaminergic medication on the cortical beta-band is likely

mediated by the dopaminergig neurons projecting to the cortex that terminates in the pre-fontal cortex

but also to less extend in the primary sensorimotor cortex.[63]. The differences in the cortical

sensorimotor burst rate in PD might be an indirect effect of the loss of dopamine and changes in the

beta band in the sub-cortical structures projecting to the sensorimotor cortex. The notion that the

cortical sensorimotor activity is indirectly related to dopamine depletion in PD is further supported by

findings from animals studies showing that 6-hydroxydopamine injections lead to exaggerated beta-

band oscillations only after several days had passed, suggesting that oscillatory changes occurred as an

indirect compensatory effect after dopamine depletion rather than a direct consequence of the depletion

itself.[7] The indirect influence of dopamine on the cortical beta-band might also explain the often weak

or even absent effect of dopaminergic medication on cortical beta-band activity.[11,13–15] The current

study cannot directly address the role of dopamine on cortical oscillations since all patients in the study

were tested on medication. However, a recent study found that cortical burst characteristics measured

with MEG were influenced by DBS therapy in PD and normalised the bursting chracteristics during

DBS to resemble the burst characteristics of healthy controls.[46] This further suppots that cortical

bursting activity is mediated by subthalamic projections.

3.3 Age-related differences

We explored how age-related differences in cortical sensorimotor neural activity might interact with

disease-related changes in PD. Age-related effects on spontaneous sensorimotor activity are commonly

dealt with by matching the age distributions of the patient group and the healthy control group—

usually within a narrow age span. The analysis showed age-related differences in burst rate, with PD

patients showing a more considerable reduction of burst as a function of age than healthy controls. The

steeper reduction in burst rate with age in PD seems in accordance with the fact that higher age at PD

onset is associated with a faster disease progression and more rapid decline in motor function[64], though

a longitudinal design is needed to confirm the relation between disease progression, reduction in burst

rate, and age. We did not see a significant "slowing" of the beta PSD centre frequency between groups,

as reported in several previous studies.[48] An explanation might be that such slowing is more

pronounced in PD patients with dementia[27–30] and correlates with cognitive ability.[26] The PD patients in

the current study did not differ in their cognitive ability from the healthy controls. Furthermore, we

focused on the activity in the sensorimotor cortex, whereas the slowing of alpha and beta PSD is

usually found in frontal areas and globally throughout the brain.[25,28,31] We included cortical thickness

measures within the same ROI from which we extracted the functional time-series, as we hypothesised

that age-related effects upon the functional measures might be mediated through the age-related

structural changes in the cortex. However, despite the negative correlation between age and cortical

thickness (see supplementary material), we did not find pervasive evidence that cortical thickness

affected any of the functional measures.

3.4 Sex differences

We also included sex to explore if disease-related changes in sensorimotor oscillatory activity differed

between males and females, as there are well-documented sex differences in the manifestation of

PD.[1,65] Male sex is a risk factor for developing PD, with an average incidence ratio of approximately

2:1 male-female ratio across all stages of the disease.[66] The disease onset is on average two years

earlier in males than females and differs in the initial manifestation of symptoms, with women more

likely to develop tremor specific symptoms and men more likely to develop rigidity.[67] We are not

aware of any previous studies that explicitly included sex as a factor in analysing neural oscillations in

PD. The regression analysis of motor symptoms showed evidence for a difference in midline function

and rest tremor between male and female patients. We did not, however, find widespread sex

differences in the various features of the sensorimotor activity. The analysis of the sensorimotor singal

features only showed evidence for differences between males and females in the burst rate. A possible

factor behind the sex differences in PD is the contribution of sex hormones on the nigrostriatal pathway

and linked to the deterioration of the dopaminergic system, where testosterone levels appear to enhance

dopamine loss, while estrogen has been identified as a neuroprotective agent for PD. Estrogen has been

demonstrated to influence incidence levels of PD while menopause-related variations in estrogen levels

are linked to variations in PD symptom severity.[68] At the current stage, it is unclear if estrogen sex

hormones influences oscillatory bursts in the sensorimotor cortex. These findings illustrate the need for

further studies into sex-specific changes in neural function and how they manifest and relate to PD.

3.5 Limitations and conclusions

The present study quantified the neural time series from the sensorimotor cortex based on pre-defined

summary measures of its PSD and burst properties. We included more PD patients and healthy controls

than similar previously conducted functional studies—typically in the range of 5-30 participants.[48] A

limitation of our study for understanding the extent of changes in oscillatory sensorimotor activity is

the focus on different features within a narrow ROI, which ignores other types of measurements that

are potentiallyrelevant to understanding the development of PD and motor symptoms: for example, the

long-range connectivity between the sensorimotor cortex and other cortical areas and the connections

between the sensorimotor cortex and the basal ganglia and thalamus (though the subcortical structures

are practically invisible in MEG).

Treating the activity in the sensorimotor cortex as single time series also means that we remove the

sensitivity to spatial features of the signals, e.g., focal versus spatially blurred activity in one group or

the other. If the oscillatory activity extends over a larger cortical surface area, that signal will also

manifest as power differences in the measured signal.[61] There are potentially other features to be

uncovered, and future studies may explore how the PSD- and burst features further interact with other

aspects of brain activity in the global function of the brain to fully understand the interaction between

functional and structural changes in PD.

We investigated a relatively large cohort of PD patients and healthy controls (for a neuroimaging

study) to make meaningful inferences about how age and sex interact with the group level difference

between PD patients and healthy controls; however, a limitation is that our study is cross-sectional. We

aim to follow this cohort longitudinally to estimate the development trajectories of the sensorimotor

oscillatory activity in PD compared to healthy ageing.

Sensorimotor activity measured non-invasively with MEG/EEG contains rich information about the

functional state of the sensorimotor system and how it changes in PD. The central challenge is

quantifying the measured neural signals to extract the disease's relevant features from the signals, be it

the spectral power, centre frequencies, or burst-like features. Finding features of neural signals that can

explain disease mechanism or symptoms, even if extracted along with a reduced number of dimensions,

will be helpful if they provide adequate information about the disease- or symptom-state. Further

characterisation of the association between features in the non-invasive brain signals and motor

symptoms can potentially be a valuable tool to aid in diagnosis and treatment evaluation.

Understanding how features in the neural time series are related to motor symptoms in PD will also

help develop non-invasive neural stimulation that can potentially relieve motor symptoms.[37,69]

4   Methods

4.1   Participants

Eighty PD patients (age 44-85; 32 female) and 71 healthy controls (age 46–78; 46 female) participated in the study. The study was approved by the regional ethics committee (Etikprövningsnämden Stockholm, DNR 2019-00542) and followed the Declaration of Helsinki. All participants gave written informed consent before participation.

The PD patients were recruited from the Parkinson's Outpatient Clinic, Department of Neurology, Karolinska University Hospital, Stockholm, Sweden. The healthy controls were recruited by advertising or amongst spouses of PD patients. 22 participants (18 patients, 4 healthy controls) were included from a previous study[14] who were qualified based on the recruitment criteria of the present study and had done the same MEG and MRI procedures as in the present study. All data were reanalysed following the procedure described below.

The inclusion criteria for the PD group were a diagnosis of PD according to the United Kingdom Parkinson's Disease Society Brain Bank Diagnostic Criteria with Hoehn and Yahr stage 1-3.[70] Inclusion criteria for the control group were not having a diagnosis of PD, no form of movement disorder, and no history of neurological disorders, epilepsy, or psychiatric disorders.

Exclusion criteria for both groups were a diagnosis of major depression, dementia, history or presence of schizophrenia, bipolar disorder, epilepsy, or history of alcoholism or drug addiction according to the *Diagnostic and Statistical Manual of Mental Disorders.*[71]

One participant declined to do the MRI scanning, one participant had a scanner malfunction during MRI acquisition, and 11 participants had their MRI scans cancelled due to the Covid-19 pandemic and

were not included in the analysis. In total, two PD patients and 11 healthy controls were excluded from

the analysis. Table 1 is a summary of the participants included in the analysis.

The PD patients participated in the study while on their regular prescribed dose of medication. The

levodopa equivalent daily dose (LEDD) was calculated according to Tomlinson et al.[72] Motor

symptoms in the PD group were assessed using the motor section of the Movement-Disorder Society

Unified Parkinson's Disease Rating Scale (MDS-UPDRS-III).[59] Global cognition was assessed with the

Montreal Cognitive Assessment battery (MoCA).[74]

4.2 MEG recordings

MEG data were recorded with a Neuromag TRIUX 306-channel MEG system, with 102

magnetometers and 102 pairs of planar gradiometers. Data were sampled at 1000 Hz with an online 0.1

Hz high-pass filter and 330 Hz low-pass filter. The MEG scanner was located inside a two-layer

magnetically shielded room (Vacuumschmelze GmbH) with internal active shielding active to suppress

electromagnetic artefacts. The subjects' head position and head movements inside the MEG scanner

were measured during recordings with head-position indicator coils (HPI) attached to subjects' heads.

The HPI location and additional points sampled uniformly across the subjects' head shape were

digitalised with a Polhemus Fastrak motion tracker before the measurements. Horizontal and vertical

electrooculogram (EOG) and electrocardiogram (ECG) were recorded simultaneously with the MEG.

We recorded three minutes of resting-state MEG while the participants sat with their eyes closed. The

participants were instructed to close their eyes and relax. The recordings began after assuring the

participant sat still with their eyes closed.

4.3 MRI acquisition

3D T1-weighted magnetisation-prepared rapid gradient-echo (MPRAGE) sequence structural images (voxel size: `1x1x1 mm`) were obtained on a GE Discovery 3.0 T MR scanner for morphological analysis and creating source spaces for MEG source reconstruction. Multi-echo "FLASH"[75] images were obtained to create volumetric headmodels for MEG source reconstruction (see below).

4.4 MRI processing

The MRI images were processed with Freesurfer[76] (v. 5.3) to get surface reconstructions of the cortical mantle. The surfaces were obtained with the automatic routine for extracting cortical surfaces in Freesurfer from the individual T1-weighted MRI.

We defined the cortical sensorimotor area by segmenting the cortical surface using the anatomical labels provided by Freesurfer automatic labelling.[77] The analysis focused on a region of interest (ROI) consisting of the left pre- and post-central gyri and central sulcus. The pre/postcentral gyri were combined because a biomagnetic source on either sulci wall will leave a trance on the other side due to the close distance and the field spread of MEG signals. The ROI was defined for each subject based on the individual cortical reconstructions. The average cortical thickness in the ROI was estimated with Freesurfer.[78]

4.5 MEG pre-processing

The MEG data was processed by applying temporal signal space separation (tSSS) to suppress artefacts from outside the scanner helmet and correct head movement during the recording.[79] The tSSS had a buffer length of 10 s and a cut-off correlation coefficient of 0.95. Movement correction was done by

shifting the head position to a position based on the median of the continuous head position during the

three-minute recording.

The MEG data processing and source reconstruction was done with MNE-Python[80] in Python 3.8. First,

we marked data segments containing muscle artefacts and SQUID jumps with the automatic artefact

detection in MNE-Python. The data was filtered with a 48 Hz low-pass filter and 50 Hz notch filter to

remove line noise. The continuous data were cut into 1.0 s epochs, and epochs with muscle artefacts or

extreme values (5000 fT for magnetometers and 4000 fT/cm for gradiometers) were rejected. Between

0-65 % (median: 6.0 %) of data was rejected resulting in 63.0-180 s (median: 174.0 s) of useful MEG

data per participant. The remaining data length was not significantly different between groups

(Wilcoxon rank sum test, p = 0.98). We then performed an independent component analysis (ICA)

using the *fastica* algorithm[81] to identify artefacts from blinks and heartbeats. Components showing

correlation with the EOG and ECG were removed from the raw data. Between 0-5 (median 3)

components were removed per participant. The number of removed ICA components was not

significantly different between groups (Wilcoxon rank sum test, p = 0.71).

We then applied source reconstruction using noise weighted minimum-norm estimates.[82] The noise

covariance matrix was estimated from two minutes of empty room MEG data recorded before each

session. The source space consisted of 5124 evenly spaced points sampled across the white matter

surfaces. The inner skull boundary was estimated from the multi-echo MRI to create a single shell

volume conductor model. The time series from the sensorimotor ROI (see Figure. 1) was extracted

from the estimated source time series by singular value decomposition of all source points within the

ROI.

4.6 Power spectral analysis

We analysed the spectral properties of the sensorimotor activity by calculating the PSD from 0.5 to 40 Hz across the entire cleaned ROI time series using Welch's method by segmenting the continuous data into 3.072 s epochs with 50% overlap and averaging the PSD across the segments.

Since the narrow-band beta power in the PSD is dependent on the broader features of the broadband spectrum, we further analysed the 1/f broadband characteristic of the sensorimotor activity as this could play a role in the functional properties of the beta-band and has been shown to differ between healthy control and PD patients.[14] We used the *fitting oscillations & one over f* (FOOOF) toolbox[51] to analyse the 1/f broadband characteristic of the PSD (intercept and exponent) and the oscillatory peaks in the canonically defined beta band (13-30 Hz) and alpha band (8-12 Hz). A log-linear regression is fitted to the PSD and subtracted before fitting Gaussian functions to the peaks in the PSD. The midpoint of the Gaussian function fittied to a given frequency band corresponds to the peak frequency in that frequency band and the height represents the signal power. A new log-linear function is fitted to the PSD after subtracting the Gaussian function to estimate the 1/f characteristic.

All participants showed a discernible beta peak in the PSD. Nine PD patients and nine healthy controls did not show a peak in the PSD alpha band (no difference between groups, $\chi^2(1) = 0.12$; $p = 0.73$).

4.7 Burst analysis

To calculate the burst properties of the sensorimotor activity in the time domain, we band-pass filtered the time-series with an 8-30 Hz band-pass filter using FieldTrip[83] in MATLAB (R2016b; MathWorks Inc.) and calculated the Hilbert envelope of the signal. The burst threshold was defined as two times the median of the signal. The burst onset was defined as the time-point where the signal first reached half the max amplitude of the burst and ended at the time-point where the signal again dropped below half

434   the max amplitude of the burst. The *burst amplitude* was defined as the maximum value of the burst.

435   The burst *duration* was defined as the time from burst onset to burst end. The *burst interval* was

436   defined as the time from the end of a burst to the time-point where the next burst began.

437   4.8   Statistics

438   4.8.1   Analysis of sensorimotor rhythm features

The main analyses tested the effect of *group* (PD patients/healthy controls), *age*, *sex*, and *ROI cortical*

thickness on the features listed in Table 2. For the PSD features, we modelled the outcomes as a linear

function of group (PD patients/healthy controls), age, age squared, sex, and cortical thickness with

linear regression in *R* (v. 4.0.2).[84] The regression models were fitted to the data for each participant

with all factors and up to their three-way interactions between the four predictors. Gaussian regression

models were estimated for each feature, except for the burst rate (burst per minute), which was

modelled with Poisson regression using the same predictor variables. Before fitting the regression

models, burst duration, burst interval, and burst amplitude were log-transformed.

Significance testing was done by removing one predictor from the model and comparing the variance

explained between the full model and the model with a predictor removed. Hypothesis testing was done

by computing the Bayes factor (BF) for the model with a given predictor (H1) versus the model

without the predictor (H0) using the BIC approximation.[85] The BF tells how much more likely the

observed data is under the alternative model (H1) versus the null model (H0). The Bayesian model

comparison, therefore, avoids the multiple comparison problem of frequentist hypothesis testing

(theoretical likelihood of the hypothesis given the observed data). The model comparison approach

furthermore circumvents issues with the interpretation of p-values of individual regression coefficients

due to internal correlation between predictor variables. Following the conventional interpretation of

BFs, we used BF > 3 as a cut-off between anecdotal evidence and substantial effects.[57] The BFs for all

comparisons are presented in Supplementary Table S3.

4.8.2   Clinical scores and sensorimotor oscillatory features

The MDS-UPDRS-III scores were divided into subscales based on symptoms: *midline function*, *rest*

tremor, *rigidity*, *upper-body bradykinesia*, *postural and kinetic tremor*, and *lower limb bradykinesia*;

according to Goetz et al.[86], with the exception that left- and right-side upper-body bradykinesia were

combined into a single factor.

Each symptom score was analysed by multiple regression and modelled as a function of the burst rate,

median burst duration, median bursts interval, median burst amplitude, PSD 1/f intercept, PSD 1/f

exponent, PSD beta power, PSD beta centre frequency, PSD alpha power, and PSD alpha centre

frequency for each PD patient. The models further included the age, sex, and cortical thickness to

regress out the contribution hereof and estimate the relative effect size of each signal feature. All

symptom ratings and continuous predictor variables, except age, were z-transformed to get the

standardised effect size. Significance testing was done by removing one predictor from the model and

calculating the BF between the full model (H1) and the model without the predictor (H0) using the BIC

approximation for BFs. The BFs for all comparisons are presented in Supplementary Table S4.

4.9   Data Availability

473   The full dataset cannot be made publicly available, as the ethical permits for the study does not allow

474   for open data sharing. Parts of the data used in this analysis will be made available as part of an online

475   data repository (Vinding, et al. *in prep*). The scripts used to process the data and run the analysis

476   presented in the paper are available at: https://github.com/natmegsweden/PD_beta_bursts2.

Authorship contribution statement

Mikkel C. Vinding: conceptualization, methodology, software, validation, formal analysis, investigation, data curation, writing - original draft, writing - review & editing, project administration. **Allison Eriksson**: investigation, resources, data curation, writing - review & editing, project administration. **Cassia Man Ting Low**: investigation, writing - review & editing. **Josefine Waldthaler**: investigation, writing - review & editing. **Daniel Ferreira**: supervision, writing - review & editing. **Martin Ingvar**: conceptualization, writing - review & editing, resources, funding acquisition. **Per Svenningsson**: conceptualization, methodology, resources, supervision, data curation, writing - review & editing, Project administration, Funding acquisition. **Daniel Lundqvist**: conceptualization, methodology, formal analysis, resources, supervision, writing - original draft, writing - review & editing, project administration, funding acquisition.

Declaration of competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1.   Kalia, L. V. & Lang, A. E. Parkinson's disease. *The Lancet* **386**, 896–912 (2015).

2.   Rodriguez-Oroz, M. C. *et al.* Initial clinical manifestations of Parkinson's disease: features and

pathophysiological mechanisms. *The Lancet Neurology* **8**, 1128–1139 (2009).

3.   Braak, H. *et al.* Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiology of*

aging **24**, 197–211 (2003).

4.   Jenkinson, N. & Brown, P. New insights into the relationship between dopamine, beta oscillations

and motor function. *Trends in Neurosciences* **34**, 611–618 (2011).

5.   Alonso-Frech, F. *et al.* Slow oscillatory activity and levodopa-induced dyskinesias in Parkinson's

disease. *Brain* **129**, 1748–1757 (2006).

6.   Giannicola, G. *et al.* The effects of levodopa and ongoing deep brain stimulation on subthalamic

beta oscillations in Parkinson's disease. *Experimental Neurology* **226**, 120–127 (2010).

7.   Mallet, N. *et al.* Disrupted Dopamine Transmission and the Emergence of Exaggerated Beta

Oscillations in Subthalamic Nucleus and Cerebral Cortex. *Journal of Neuroscience* **28**, 4795–4806

(2008).

8.   Neumann, W.-J. *et al.* Long term correlation of subthalamic beta band activity with motor

impairment in patients with Parkinson's disease. *Clinical Neurophysiology* **128**, 2286–2291 (2017).

9.   Kühn, A. A., Kupsch, A., Schneider, G.-H. & Brown, P. Reduction in subthalamic 8-35 Hz

oscillatory activity correlates with clinical improvement in Parkinson's disease: STN activity and

motor improvement. *European Journal of Neuroscience* **23**, 1956–1960 (2006).

10. Martin, S. *et al.* Differential contributions of subthalamic beta rhythms and 1/f broadband activity

to motor symptoms in Parkinson's disease. *npj Parkinson's Disease* **4**, 32 (2018).

11. Airaksinen, K. *et al.* Somatomotor mu rhythm amplitude correlates with rigidity during deep brain

stimulation in Parkinsonian patients. *Clinical Neurophysiology* **123**, 2010–2017 (2012).

12. Feldmann, L. K. *et al.* Subthalamic beta band suppression reflects effective neuromodulation in

chronic recordings. *European Journal of Neurology* ene.14801 (2021) doi:10.1111/ene.14801.

13. Hall, S. D. *et al.* GABA-mediated changes in inter-hemispheric beta frequency activity in early-

stage Parkinson's disease. *Neuroscience* **281**, 68–76 (2014).

14. Vinding, M. C. *et al.* Reduction of spontaneous cortical beta bursts in Parkinson's disease is linked

to symptom severity. *Brain Communications* (2020) doi:10.1093/braincomms/fcaa052.

15. Vinding, M. C. *et al.* Attenuated beta rebound to proprioceptive afferent feedback in Parkinson's

disease. *Scientific Reports* **9**, (2019).

16. Cao, C. *et al.* L-dopa treatment increases oscillatory power in the motor cortex of Parkinson's

disease patients. *NeuroImage: Clinical* **26**, 102255 (2020).

17. Heinrichs-Graham, E. *et al.* Hypersynchrony despite pathologically reduced beta oscillations in

patients with Parkinson's disease: a pharmaco-magnetoencephalography study. *Journal of*

Neurophysiology **112**, 1739–1747 (2014).

18. Melgari, J.-M. *et al.* Alpha and beta EEG power reflects L-dopa acute administration in

parkinsonian patients. *Frontiers in Aging Neuroscience* **6**, 302 (2014).

19. Abbasi, O. *et al.* Unilateral deep brain stimulation suppresses alpha and beta oscillations in

sensorimotor cortices. *NeuroImage* **174**, 201–207 (2018).

20. Luoma, J. *et al.* Spontaneous sensorimotor cortical activity is suppressed by deep brain stimulation

in patients with advanced Parkinson's disease. *Neuroscience Letters* **683**, 48–53 (2018).

21. Cao, C.-Y. *et al.* Modulations on cortical oscillations by subthalamic deep brain stimulation in

patients with Parkinson disease: A MEG study. *Neuroscience Letters* **636**, 95–100 (2017).

22. Pollok, B. *et al.* Motor-cortical oscillations in early stages of Parkinson's disease: Suppression of

motor cortical beta oscillations is altered in early PD. *The Journal of Physiology* **590**, 3203–3212

(2012).

23. Olde Dubbelink, K. T. E. *et al.* Resting-state functional connectivity as a marker of disease

progression in Parkinson's disease: A longitudinal MEG study. *NeuroImage: Clinical* **2**, 612–619

(2013).

24. Boon, L. I. *et al.* Motor effects of deep brain stimulation correlate with increased functional

connectivity in Parkinson's disease: An MEG study. *NeuroImage: Clinical* **26**, 102225 (2020).

25. Stoffers, D. *et al.* Slowing of oscillatory brain activity is a stable characteristic of Parkinson's

disease without dementia. *Brain* **130**, 1847–1860 (2007).

26. Vardy, A. N. *et al.* Slowing of M1 activity in Parkinson's disease during rest and movement – An

MEG study. *Clinical Neurophysiology* **122**, 789–795 (2011).

27. Bosboom, J. L. W., Stoffers, D., Wolters, E. Ch., Stam, C. J. & Berendse, H. W. MEG resting state

functional connectivity in Parkinson's disease related dementia. *Journal of Neural Transmission*

116, 193–202 (2009).

28. Bosboom, J. L. W. *et al.* Resting state oscillatory brain dynamics in Parkinson's disease: An MEG

study. *Clinical Neurophysiology* **117**, 2521–2531 (2006).

29. Olde Dubbelink, K. T. E. *et al.* Predicting dementia in Parkinson disease by combining

neurophysiologic and cognitive markers. *Neurology* **82**, 263–270 (2014).

30. Ponsen, M. M., Stam, C. J., Bosboom, J. L. W., Berendse, H. W. & Hillebrand, A. A three

dimensional anatomical view of oscillatory resting-state activity and functional connectivity in

Parkinson's disease related dementia: An MEG study using atlas-based beamforming.

NeuroImage: Clinical **2**, 95–102 (2013).

31. Olde Dubbelink, K. T. E. *et al.* Cognitive decline in Parkinson's disease is associated with slowing

of resting-state brain activity: a longitudinal study. *Neurobiology of Aging* **34**, 408–418 (2013).

32. Stoffers, D., Bosboom, J. L. W., Wolters, E. Ch., Stam, C. J. & Berendse, H. W. Dopaminergic

modulation of cortico-cortical functional connectivity in Parkinson's disease: An MEG study.

Experimental Neurology **213**, 191–195 (2008).

33. Karekal, A., Miocinovic, S. & Swann, N. C. Novel approaches for quantifying beta synchrony in

Parkinson's disease. *Exp Brain Res* **240**, 991–1004 (2022).

34. Feingold, J., Gibson, D. J., DePasquale, B. & Graybiel, A. M. Bursts of beta oscillation

differentiate postperformance activity in the striatum and motor cortex of monkeys performing

movement tasks. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 13687–13692 (2015).

35. Leventhal, D. K. *et al.* Basal Ganglia Beta Oscillations Accompany Cue Utilization. *Neuron* **73**,

523–536 (2012).

36. Sherman, M. A. *et al.* Neural mechanisms of transient neocortical beta rhythms: Converging

evidence from humans, computational modeling, monkeys, and mice. *Proceedings of the National*

Academy of Sciences **113**, E4885–E4894 (2016).

37. Tinkhauser, G. *et al.* The modulatory effect of adaptive deep brain stimulation on beta bursts in

Parkinson's disease. *Brain* **140**, 1053–1067 (2017).

38. Shin, H., Law, R., Tsutsui, S., Moore, C. I. & Jones, S. R. The rate of transient beta frequency

events predicts behavior across tasks and species. *eLife* **6**, (2017).

39. Duchet, B. *et al. Average beta burst duration profiles provide a signature of dynamical changes*

between the ON and OFF medication states in Parkinson's disease.

http://biorxiv.org/lookup/doi/10.1101/2020.04.27.064246 (2020) doi:10.1101/2020.04.27.064246.

40. Khawaldeh, S. *et al.* Subthalamic nucleus activity dynamics and limb movement prediction in

Parkinson's disease. *Brain* **143**, 582–596 (2020).

41. Lofredi, R. *et al.* Beta bursts during continuous movements accompany the velocity decrement in

Parkinson's disease patients. *Neurobiology of Disease* (2019) doi:10.1016/j.nbd.2019.03.013.

42. Little, S., Bonaiuto, J., Barnes, G. & Bestmann, S. Human motor cortical beta bursts relate to

movement planning and response errors. *PLoS Biol* **17**, e3000479 (2019).

43. Brady, B., Power, L. & Bardouille, T. Age-related trends in neuromagnetic transient beta burst

characteristics during a sensorimotor task and rest in the Cam-CAN open-access dataset.

NeuroImage **222**, 117245 (2020).

44. Tinkhauser, G. *et al.* Beta burst coupling across the motor circuit in Parkinson's disease.

Neurobiology of Disease **117**, 217–225 (2018).

45. Tinkhauser, G. *et al.* Beta burst dynamics in Parkinson's disease OFF and ON dopaminergic

medication. *Brain* **140**, 2968–2981 (2017).

46. Pauls, K. A. M. *et al.* Cortical beta burst dynamics are altered in Parkinson's disease but

normalized by deep brain stimulation. *NeuroImage* 119308 (2022)

doi:10.1016/j.neuroimage.2022.119308.

47. Power, L. & Bardouille, T. Age‐related trends in the cortical sources of transient beta bursts

during a sensorimotor task and rest. *NeuroImage* **245**, 118670 (2021).

48. Boon, L. I. *et al.* A systematic review of MEG‐based studies in Parkinson's disease: The motor

system and beyond. *Hum Brain Mapp* **40**, 2827–2848 (2019).

49. Salmelin, R., Hämäläinen, M., Kajola, M. & Hari, R. Functional Segregation of Movement-Related

Rhythmic Activity in the Human Brain. *NeuroImage* **2**, 237–243 (1995).

50. Salmelin, R. & Hari, R. Spatiotemporal characteristics of sensorimotor neuromagnetic rhythms

related to thumb movement. *Neuroscience* **60**, 537–550 (1994).

51. Donoghue, T. *et al.* Parameterizing neural power spectra into periodic and aperiodic components.

Nat Neurosci **23**, 1655–1665 (2020).

52. He, B. J., Zempel, J. M., Snyder, A. Z. & Raichle, M. E. The Temporal Structures and Functional

Significance of Scale-free Brain Activity. *Neuron* **66**, 353–369 (2010).

53. Bardouille, T., Bailey, L. & CamCAN Group. Evidence for age-related changes in sensorimotor

neuromagnetic responses during cued button pressing in a large open-access dataset. *NeuroImage*

193, 25–34 (2019).

54. Wolpe, N. *et al.* Ageing increases reliance on sensorimotor prediction through structural and

functional differences in frontostriatal circuits. *Nature Communications* **7**, (2016).

55. Yau, Y. *et al.* Network connectivity determines cortical thinning in early Parkinson's disease

progression. *Nat Commun* **9**, 12 (2018).

56. Lyoo, C. H., Ryu, Y. H. & Lee, M. S. Cerebral cortical areas in which thickness correlates with

severity of motor deficits of Parkinson's disease. *J Neurol* **258**, 1871–1876 (2011).

57. Wetzels, R. *et al.* Statistical Evidence in Experimental Psychology An Empirical Comparison

Using 855 t Tests. *Perspectives on Psychological Science* **6**, 291–298 (2011).

58. Goetz, C. G. *et al.* Movement Disorder Society-sponsored revision of the Unified Parkinson's

Disease Rating Scale (MDS-UPDRS): Process, format, and clinimetric testing plan. *Movement*

Disorders **22**, 41–47 (2007).

59. Robertson, M. M. *et al.* EEG power spectral slope differs by ADHD status and stimulant

medication exposure in early childhood. *Journal of Neurophysiology* **122**, 2427–2437 (2019).

60. Gao, R., Peterson, E. J. & Voytek, B. Inferring synaptic excitation/inhibition balance from field

potentials. *NeuroImage* **158**, 70–78 (2017).

61. Zich, C., Quinn, A. J., Mardell, L. C., Ward, N. S. & Bestmann, S. Dissecting Transient Burst

Events. *Trends in Cognitive Sciences* S136466132030173X (2020) doi:10.1016/j.tics.2020.07.004.

62. Muralidharan, V. & Aron, A. R. Behavioral Induction of a High Beta State in Sensorimotor Cortex

Leads to Movement Slowing. *Journal of Cognitive Neuroscience* 1–18 (2021)

doi:10.1162/jocn_a_01717.

63. Luft, A. R. & Schwarz, S. Dopaminergic signals in primary motor cortex. *Int. j. dev. neurosci.* **27**,

415–421 (2009).

64. De Pablo-Fernández, E., Lees, A. J., Holton, J. L. & Warner, T. T. Prognosis and Neuropathologic

Correlation of Clinical Subtypes of Parkinson Disease. *JAMA Neurol* **76**, 470 (2019).

65. Göttgens, I. *et al.* The Impact of Sex and Gender on the Multidisciplinary Management of Care for

Persons With Parkinson's Disease. *Frontiers in Neurology* **11**, (2020).

66. Elbaz, A. *et al.* Risk tables for parkinsonism and Parkinson's disease. *Journal of Clinical*

Epidemiology **55**, 25–31 (2002).

67. Georgiev, D., Hamberg, K., Hariz, M., Forsgren, L. & Hariz, G.-M. Gender differences in

Parkinson's disease: A clinical perspective. *Acta Neurol Scand* **136**, 570–584 (2017).

68. Gillies, G. E., Pienaar, I. S., Vohra, S. & Qamhawi, Z. Sex differences in Parkinson's disease.

Frontiers in Neuroendocrinology **35**, 370–384 (2014).

69. Cook, A. J., Pfeifer, K. J. & Tass, P. A. A Single Case Feasibility Study of Sensorimotor Rhythm

Neurofeedback in Parkinson's Disease. *Front. Neurosci.* **15**, 623317 (2021).

70. Hoehn, M. M. & Yahr, M. D. Parkinsonism: onset, progression, and mortality. *Neurology* **17**, 427–

427 (1967).

71. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*.

(American Psychiatric Publishing, 2013).

72. Tomlinson, C. L. *et al.* Systematic review of levodopa dose equivalency reporting in Parkinson's

disease. *Movement Disorders* (2010) doi:10.1002/mds.23429.

73. Goetz, C. G. *et al.* Movement disorder society-sponsored revision of the unified Parkinson's

disease rating scale (MDS-UPDRS): Process, format, and clinimetric testing plan. *Movement*

Disorders (2007) doi:10.1002/mds.21198.

74. Nasreddine, Z. S. *et al.* The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild

cognitive impairment. *Journal of the American Geriatrics Society* **53**, 695–699 (2005).

75. Fischl, B. *et al.* Sequence-independent segmentation of magnetic resonance images. *NeuroImage*

23, S69–S84 (2004).

76. Dale, A. M., Fischl, B. & Sereno, M. I. Cortical Surface-Based Analysis: I. segmentation and

surface reconstruction. *NeuroImage* **9**, 179–194 (1999).

77. Destrieux, C., Fischl, B., Dale, A. & Halgren, E. Automatic parcellation of human cortical gyri and

sulci using standard anatomical nomenclature. *NeuroImage* **53**, 1–15 (2010).

78. Fischl, B. & Dale, A. M. Measuring the thickness of the human cerebral cortex from magnetic

resonance images. *Proceedings of the National Academy of Sciences* **97**, 11050–11055 (2000).

79. Taulu, S. & Simola, J. Spatiotemporal signal space separation method for rejecting nearby

interference in MEG measurements. *Physics in Medicine and Biology* **51**, 1759–1768 (2006).

80. Gramfort, A. *et al.* MEG and EEG data analysis with MNE-Python. *Frontiers in Neuroscience* **7**,

267 (2013).

81. Hyvarinen, A. Fast and robust fixed-point algorithms for independent component analysis. *IEEE*

Transactions on Neural Networks **10**, 626–634 (1999).

82. Dale, A. M. *et al.* Dynamic Statistical Parametric Mapping: Combining fMRI and MEG for High-

Resolution Imaging of Cortical Activity. *Neuron* **26**, 55–67 (2000).

83. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open Source Software for

Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational*

Intelligence and Neuroscience **2011**, 1–9 (2011).

84. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for

Statistical Computing, 2020).

85. Wagenmakers, E.-J. A practical solution to the pervasive problems of p values. *Psychonomic*

Bulletin & Review **14**, 779–804 (2007).

86. Goetz, C. G. *et al.* Movement Disorder Society-sponsored revision of the Unified Parkinson's

Disease Rating Scale (MDS-UPDRS): Scale presentation and clinimetric testing results. *Movement*

Disorders **23**, 2129–2170 (2008).

Figure legends

Figure 1: Overview of the data processing pipeline.

Three minutes raw resting-state MEG data was obtained from each participant. The signals were projected through a minimum-norm source reconstruction to extract the activity in the sensorimotor cortex. We did a frequency decomposition of the source reconstructed signal to calculate the PSD to which a 1/f and Gaussian curve were fitted to extract the PSD features (Table 2). In addition, we quantified sensorimotor bursts in the signal time series in the sensorimotor ROI by thresholding the envelope of the band-pass filtered (8-30 Hz) signal to the mu-beta frequency range.

Figure 2: Group-level PSD.

Grand average PSD (mean+standard error) for the PD group (blue) and healthy control group (red).

Figure 3: Regression analysis of PSD features.

Standardized regression coefficients for the analyses of sensorimotor PSD features as a function of group, age, sex, cortical thickness and the interaction between these factors. * indicate factors with Bayes Factor > 3 in the model comparison.

Figure 4: Regression analysis of burst features.

Standardized regression coefficients for the analyses of burst features as a function of group, age, sex, cortical thickness and the interaction between these factors. * indicate factors with Bayes Factor > 3 in the model comparison.

Figure 5: Age-related changes in sensorimotor signal features.

Scatterplots of the individual measures and model predictions over age for (A) burst rate, (B) PSD

broadband 1/f intercept, and (C) PSD broadband 1/f exponent, split between PD patients (blue) and

healthy controls (red), and female (solid) and male (dashed).

Figure 6: Regression analysis of motor symptoms.

Standardized regression coefficients for the regression analyses of sensorimotor signal features on

clinical motor symptom ratings in PD. * indicate factors with Bayes Factor > 3 in the model

comparison.

Tables

27 Tables

Table 1

28 Table 1

29 Group-level summary of the participants included in the analysis. Mean (standard deviation).

Measure	Parkinson's patients	Healthy controls	Statistics
N	78	60	
Sex (female/male)	29/49	27/33	$\chi^2 = 0.57$, $p = 0.45$
Age	65.6 (9.5)	63.93 (8.4)	Welsh $t(138.0) = 1.08$, $p = 0.28$
Disease duration	4.4 (3.7) years	-	-
LEDD	548 (273) mg	-	-
MDS-UPDRS-III	18.9 (10.8)	-	-
MoCA	26.1 (2.8)	26.2 (2.1)	Welsh $t(136) = 0.10$, $p = 0.92$

31   Table 2

32   Summary explanations of the main outcome variables in the analysis

Variable category	Variable	Explanation
PSD	*Beta power*	The maximum peak in the 13-30 Hz band. Estimated as the height of the Gaussian function fitted to the PSD after regressing out the 1/f spectrum.
	Beta centre frequency (Hz)	The dominant frequency bin in the 13-30 Hz band. Estimated as the centre of the Gaussian function fitted to the 13-30 Hz range of the PSD after regressing out the 1/f spectrum.
	Alpha power	The maximum peak in the 8-12 Hz band. Estimated as the height of the Gaussian function fitted to the PSD after regressing out the 1/f spectrum.
	Alpha centre frequency (Hz)	The dominant frequency bin in the 8-12 Hz band. Estimated as the centre of the Gaussian function fitted to the 8-12 Hz range of the PSD after regressing out the 1/f spectrum.
	1/f offset	The intercept of the log-linear regression line estimated from the full PSD in the 0.5-40 Hz range.
	1/f exponent	The decay exponent ($1/f^x$) of the PSD, corresponding to the slope of the log-linear regression line, estimated from the full PSD in the 0.5-40 Hz range.
Burst	*Rate*	The number of burst events in the time series divided by the length of the time series
	Duration (ms)	The time point from where the signal envelope rise above the threshold until the next time point it drops below the threshold.
	Interval (ms)	The time point from where the signal envelope drops below the threshold until the next time point it rises above the threshold.
	Amplitude	The maximum envelope amplitude within a burst event.