Title: **A prospective study of time-dependent childhood adversity and DNA methylation across childhood and adolescence**

Running title: Time-varying adversity and DNA methylation trajectories

Authors: Alexandre A. Lussier, PhD*[1,2,3]; Yiwen Zhu, MSc[1,4]; Brooke J. Smith, MSc[1]; Janine Cerutti, BSc[1]; Andrew J. Simpkin, PhD[5]; Andrew D.A.C. Smith, PhD[6]; Matthew J. Suderman, PhD[7]; Esther Walton, PhD[8]; Caroline L Relton, PhD[7]; Kerry J. Ressler MD, PhD[2,9]; Erin C. Dunn, ScD**[1,2,3,10]

Affiliations:

[1] Psychiatric and Neurodevelopmental Genetics Unit, Centre for Genomic Medicine, Massachusetts General Hospital, Boston, MA, 02114, USA.

[2] Department of Psychiatry, Harvard Medical School, Boston, MA, 02115, USA.

[3] Stanley Center for Psychiatric Research, The Broad Institute of Harvard and MIT, Cambridge, MA, 02142, USA.

[4] Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, 02114, USA

[5] School of Mathematics, Statistics and Applied Mathematics, National University of Ireland, Galway, H91 H3CY, Ireland.

[6] Mathematics and Statistics Research Group, University of the West of England, Bristol, BS16 1QY, UK.

[7] MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, BS8 1UD, UK.

[8] Department of Psychology, University of Bath, Bath, BA2 7AY, UK.

[9] McLean Hospital, Belmont, MA, 02478, USA.

[10] Center on the Developing Child at Harvard University, Cambridge, MA, 02138, USA.

Corresponding authors contact information

*Alexandre A. Lussier: alussier[at]mgh.harvard.edu; 617-642-0193
**Erin C. Dunn: edunn2[at]mgh.harvard.edu; 617-726-9387
Address: 185 Cambridge Street, Room 6.260
Boston, MA 02114

Keywords: ALSPAC, longitudinal, epigenetic, DNA methylation, childhood adversity, adolescence, trajectories.

Word count: 3381

ABSTRACT

Background: Children exposed to adversity often have epigenetic profiles, including DNA methylation (DNAm) signatures, that differ from children without adversity histories. These signatures may be more common among children exposed during sensitive periods in development. However, it remains unclear if adversity has persistent (versus transient) effects on the epigenome across childhood and adolescence. Thus, we examined the relationship between time-varying adversity and genome-wide DNAm, measured three times from birth to adolescence using prospective data from the Avon Longitudinal Study of Parents and Children.

Methods: We first investigated the timing of exposure to seven types of adversity (measured 5-8 times between ages 0-11) and DNAm at age 15 using a structured life course modeling approach (SLCMA). We also assessed the persistence of adversity-DNAm associations identified from age 7 DNAm into adolescence and the influence of adversity on DNAm trajectories from ages 0-15.

Results: Adversity exposure was associated with differences in age 15 DNAm at 41 loci ($R^2 \geq 0.035$; $p < 1 \times 10^{-5}$; 22 at FDR<0.05). Most loci were associated with adversities (i.e., physical, sexual, or emotional abuse; one-adult households) occurring between ages 3-5. DNAm differences present at age 7 resolved by adolescence; age 15 differences were not apparent in childhood. We also identified six distinct DNAm trajectories that highlighted both immediate and latent effects of adversity. Associations were robust in internal validation analyses using nonparametric bootstrapping.

Conclusions: These findings highlight the immediate and latent effects of childhood adversity on DNAm, providing a potential biological mechanism linking adversity to physical and mental health outcomes across development.

INTRODUCTION

Children exposed to adversity, such as abuse or maltreatment[1], family disruption or dysfunction[2], or poverty[3], frequently have poorer physical and mental health outcomes later in life[4-6]. Epigenetic processes, including DNA methylation (DNAm), are increasingly recognized as potential underlying mechanisms for these associations, as DNAm is responsive to life experiences[7] and may mediate the link between environmental exposures and chronic disease[8]. Several large-scale population-based studies, systematic reviews, and meta-analyses have shown links between childhood adversity, DNAm, and adverse health outcomes across the life course[9-14]. However, prior studies investigating the epigenome of children exposed to adversity have not yet explored two key dimensions of the adversity-DNAm relationship, which are critical to understand the biological risk posed by childhood adversity, identify children at risk for poor health, and improve intervention targets for health promotion and disease prevention.

First, it remains unclear how the *timing* of childhood adversity might shape DNAm. Recent human and animal studies suggest there may be *sensitive periods* for epigenetic programming when physiological and neurobiological systems are primed for external influences, allowing life experiences to impart more enduring effects[15-18]. Few studies have investigated the specific developmental periods when adversity may have greater effects on DNAm[13,19], with none investigating sensitive-period effects on epigenetic patterns in adolescence.

Second, little is known about how the epigenome of children exposed to adversity varies across development and how DNAm variation may shape health. In a recent article, Oh and Petronis[20] argued that the dynamic nature of epigenetic mechanisms is best examined through longitudinal studies that model chrono-epigenetic patterns, or the temporal dynamics of

epigenetic processes. Although previous studies have shown the epigenome is dynamic across

development[21-30], no study has determined how childhood adversity might influence DNAm

trajectories.

To address these gaps, we examined the longitudinal relationship between early-life

adversity and genome-wide DNAm across childhood and adolescence, using data collected over

two decades from a subsample of youth enrolled in the Avon Longitudinal Study of Parents and

Children (ALSPAC) cohort. We examined the associations between exposure to seven types of

childhood adversity, assessed repeatedly between birth and age 11, and DNAm at age 15. Given

the unique availability of three waves of DNAm in ALSPAC (birth, age 7, age 15), we also

examined DNAm trajectories from birth to adolescence.

Our aims were to: 1) determine whether childhood adversity has time-dependent effects

on adolescent DNAm; 2) characterize the developmental trajectories of DNAm linked to

adversity; and 3) evaluate the persistence of previously-identified associations between adversity

and DNAm in childhood[19]. To our knowledge, this study is the first to investigate the time-

varying influences of childhood adversity on adolescent DNAm and DNAm trajectories from

childhood to adolescence. These analyses are a novel extension of our prior work, which for one

of the first times revealed sensitive periods for the effect of childhood adversity on epigenetic

alterations at age 7 in ALSPAC[13,19].

MATERIALS AND METHODS

Sample

ALSPAC is a large population-based birth cohort from Avon, UK of 14,451 children

followed from before birth through early adulthood[31,32]. Blood-based DNAm profiles were

generated for a subsample of ALSPAC mother-child pairs with cord blood at birth (n=905),

whole blood at age 7 (n=970), and peripheral blood leukocytes at age 15 (n=966)[33] (full

ALSPAC details in **Supplemental materials**). Of note, ALSPAC is one of the few longitudinal

studies with both repeated measures of childhood adversity and multiple waves of DNAm data.

Measures of childhood adversity

We examined the effect of seven types of childhood adversity previously associated with

DNAm[34-38]: 1) caregiver physical or emotional abuse; 2) sexual or physical abuse (by anyone);

3) maternal psychopathology; 4) one-adult households; 5) family instability; 6) financial

hardship; and 7) neighborhood disadvantage. These adversities were generated from maternal

reports via mailed questionnaires, collected 5-8 times between birth and age 11 (**Figure 1**; details

in **Table S1**).

DNAm data generation

DNAm was measured at 485,577 CpG sites using the Illumina Infinium

HumanMethylation450 BeadChip microarray (Illumina, San Diego, CA). Laboratory procedures,

preprocessing analyses, and quality control steps performed were described previously[33]. We

removed non-variable CpGs (<5% DNAm difference between children in the 10th and 90th

percentile), resulting in an analytic sample of 302,581 CpGs (details in **Supplemental**

materials).

Covariates

To adjust for potential confounders and be consistent with prior work[13], we controlled for

age of blood collection, sex, race/ethnicity, maternal age at birth, maternal education at birth,

birthweight, number of previous pregnancies, maternal smoking during pregnancy, and cell type

proportions estimated using the Houseman method[39] (details in **Supplemental materials**).

Analyses

Structured Life Course Modeling Approach (SLCMA)

Our primary analyses focused on identifying time-dependent associations between each

type of childhood adversity and DNAm measured in adolescence (age 15). We used the

structured life course modeling approach (SLCMA; pronounced "slick-mah"), a two-stage

method that simultaneously compares *a priori* life course hypotheses explaining exposure-

outcome relationships[40-44]. SLCMA first uses variable selection to identify the life course

hypothesis explaining the greatest proportion of outcome variation. Effect estimates, confidence

intervals, and p-values are then calculated for the selected life course hypothesis using post-

selective inference. Importantly, SLCMA detects time-varying associations with more statistical

power and less bias than traditional epigenome-wide association studies[13,19,45].

We examined time-dependent associations for each adversity among children with

complete data for the timepoints shown in **Figure 1**. We generated variables corresponding to six

separate life course hypotheses, including four sensitive periods hypotheses encoding exposure

to each childhood adversity during: 1) *very early childhood* (ages 0-2), 2) *early childhood* (ages

3-5), 3) *middle childhood* (ages 6-7), 4) *late childhood* (ages 8-11); and two additive hypotheses:

5) *accumulation of exposures* (total exposures across childhood), and 6) *recency of exposures*

(total exposures weighted by age) to determine whether more recent exposures had a stronger

impact than distal exposures.

We tested associations using selective inference[46] and accounted for multiple-testing

using 5% a false-discovery rate (FDR). Details on the SLCMA and functional analyses of top

loci are in **Supplemental materials**.

To our knowledge, there are currently no comparable cohort studies with repeated

measures of childhood adversity and DNAm in which to replicate our findings. Thus, we

assessed the robustness of the SLCMA results through internal validation analyses of our

associations using ordinary nonparametric bootstrapping (**Supplemental materials**). We also

assessed the effects of potential confounders or alternate mediators of the association between

childhood adversity and DNAm at age 15, including exposures to other types of childhood

adversity in the same or different sensitive periods (**Supplemental materials**).

Trajectories of DNAm response to childhood adversity

The three waves of longitudinal DNAm data available in ALSPAC allowed us to

investigate patterns of DNAm across development. Thus, we pursued three additional sets of

analyses, building from the SLCMA findings.

<u>Pre-existence of age 15 associations</u>

We first determined whether DNAm differences identified at age 15 using the SCLMA

emerged earlier in development. Using linear regression, we tested whether exposure to adversity

was associated with DNAm at the same top loci at birth or age 7, adjusting for covariates.

<u>Types of DNAm trajectories across development</u>

We also investigated DNAm patterns beyond the cross-sectional age 15 time point,

studying longitudinal change and stability patterns in our top loci among children from three

distinct exposure groups: 1) children who had adversity exposure *during* the sensitive period

identified from the SLCMA (labeled as exposed-SP); 2) children who had adversity exposure

outside the sensitive period identified from the SLCMA (exposed-other); and 3) children who

were never exposed to adversity. We performed an analysis of variance (ANOVA) of the

statistical interaction between age at DNAm collection and exposure group, controlling for

DNAm repeated measures as fixed effects. Loci showing differences between exposure groups

across time (i.e., group-by-age interactions; FDR<0.05) were carried forward to subsequent

analyses, to identify more granular temporal patterns between exposure groups.

From these loci, we identified different types of DNAm trajectories based on ANOVA

and Tukey post-doc analyses of three main parameters: 1) mean exposure group differences

across ages, 2) mean age differences *across exposure groups*, and 3) exposure group differences

within each age. We then used hierarchical clustering of these three parameters to identify

homogeneous subsets within these longitudinal patterns (**Supplemental Materials**).

<u>Persistence of childhood DNAm differences to adolescence</u>

We assessed whether DNAm alterations linked to childhood adversity, which we

previously identified at age 7[19], persisted to adolescence. Specifically, we performed linear

models between adversity and age 15 DNAm data for these 46 childhood loci (**Supplemental**

materials).

RESULTS

Sample characteristics and prevalence of exposure to adversity

Demographic characteristics did not differ between the ARIES sample and children

exposed to any adversity between ages 0-11 (**Table S2**). The prevalence of exposure to a given

adversity from age 0-11 ranged from 15.1% (sexual/physical abuse) to 34.8% (maternal

psychopathology) (**Figure S1; Table S3**). The correlation of exposure within each adversity

across development ranged from 0.36 (family instability) to 0.786 (one-adult households).

Different types of adversity were weakly correlated (r_{avg}=-0.04-0.16).

Childhood adversity showed time-dependent associations with adolescent DNAm profiles

Across all types of childhood adversity, 41 loci showed significant associations between

exposure to adversity and DNAm levels at age 15 ($\geq$3.5% of DNAm variance explained by

adversity; nominal $p<1\times10^{-5}$; **Table 1; Table S4**). Of these, 22 loci were significant after

multiple-test correction (FDR<0.05). As we and others have recently shown that p-values are

poor metrics of statistical inference when used on their own[47,48], particularly in the context of

time-varying effects[19], we focus on downstream analyses on the subset of CpGs that met both the

R^2 threshold and nominal p-value.

Sensitive periods were the most often selected life course hypothesis by the SLCMA,

with 35 loci showing associations with childhood adversity that occurred during *very early*

childhood (20% or 8 of 41), *early childhood* (56% or 23 of 41), or *late childhood* (10% or 4 of

41) (**Figure 2**). Only 3 loci (7%) showed associations with either the accumulation or recency of

adversity.

Most of these associations were for exposure to one-adult households (20 of 41 loci). We

also identified associations with caregiver physical or emotional abuse (3 loci), sexual or

physical abuse by anyone (4 loci), family instability (1 locus), financial hardship (9 loci),

maternal psychopathology (1 locus), and neighborhood disadvantage (3 loci).

Childhood adversity was mainly associated with a decrease in DNAm (20 of 23 loci). On

average, childhood adversity exposure was linked to a 3.5% absolute difference in DNAm (range

0.9-10.4%). For loci associated with accumulated time living in one-adult households, each

additional timepoint of exposure was associated with a 1% difference in DNAm (range 0.3-

1.4%). For loci associated with the recency of financial hardship, one additional exposure was

linked to a -1.3% to 2.3% change in DNAm per year of age at exposure.

Top loci showed higher representation in regions of lower CpG density, such as

enhancers (χ^2=7.1, p=0.008) and Open Seas (χ^2=13.6, p=0.018) (**Figure S2**). Most loci (28/41) also

had weak, but positive correlations between brain and blood[49] (**Table S5**; **Figure S3**), suggesting

adversity-induced alterations to blood DNAm may reflect similar changes in the central nervous

system. Although no biological processes were significantly enriched in top loci using the

DAVID gene ontology tool[50,51] (**Figure S4**), seven genes linked to sexual/physical abuse (*TAF1*),

family instability (*PKD2*), financial hardship (*FBXL16, XKR6*), or one-adult households (*DSP*,

CUX2, *STK38L*) showed evidence of strong evolutionary constraint (**Table S4; Figure S5**)[52].

Together, these findings suggest different types of childhood adversity may act through diverse

biological processes and hunt at a potential evolutionary role for genes responsive to parental and

social environments (**Supplemental materials**).

Associations between childhood adversity and DNAm at age 15 were internally validated

and specific to each adversity

Internal validation analysis of the top associations yielded nearly identical results to the

initial analyses (largest difference in effect estimates=2.03%); all bootstrap effect estimates were

significant at the 5% level (**Figure S6; Table S6**). Our results also remained stable when

correcting for exposure to other adversities during the sensitive period or across childhood,

suggesting they were not influenced by co-occurring incidences of adversity (**Supplemental**

materials; Figure S7-9). Together, these results point to the robustness and specificity of

associations between time-varying childhood adversity and DNAm at age 15.

DNAm differences at age 15 were not present earlier in childhood

For the 41 loci identified in age 15 DNAm, none showed associations between adversity

and DNAm at birth (**Table S7**) or age 7 (**Table S8**) (FDR<0.05). Notably, the age 7 effect

estimates were *smaller* than the age 15 associations, with consistent directions-of-effect in about

half (20 of 41) (**Figure 3A**). Importantly, the emergence these associations was not explained by

early-life confounders or biological mediators during adolescence (**Supplemental materials;**

Figures S10-16), suggesting some adolescent differences may emerge later in development and

become stronger with time.

Childhood adversity was linked to distinct trajectories of DNAm across development

Moving beyond adolescent DNAm alone, 34 of the 41 loci had significant adversity

exposure group-by-age interactions (FDR<0.05), suggestive of more complex patterns of change

and stability across development. From these loci, we identified five additional types of

longitudinal DNAm trajectories (**Table S9**), which showed distinct DNAm patterns across ages

and adversity exposure groups (**Figures S17-20**), but not between the FDR and R^2 subsets of

CpGs (**Figure S21**). **Table 2** provides a full description and examples of the patterns

distinguishing DNAm trajectories, which included differences that emerged earlier versus later in

development, differences between children exposed during a sensitive period or at other

developmental stages, and differences linked to age at DNAm measurement.

Associations between adversity and childhood DNAm did not persist into adolescence

Of the 46 CpG sites previously showing time-varying associations between adversity and

DNAm at age 7 [19], only one showed an association at age 15 ($p<0.05$; **Table S10**), which did not

pass multiple-test correction. Again, approximately half of loci showed consistent direction-of-

effect between age 7 and 15 (24/46) (**Figure 3B**). These findings suggest some childhood

epigenetic responses to adversity may not persist into adolescence.

DISCUSSION

The main finding from this study is that associations between childhood adversity and

DNAm vary across the life course, manifesting at different developmental stages through distinct

patterns of persistence and latency. To our knowledge, this is the first study to incorporate time-

dependent measures of childhood adversity in the study of human longitudinal epigenetic

patterns.

Our findings point to early childhood – meaning the period between ages 3 to 5 – as a

possible sensitive period for the biological embedding of childhood adversity that manifests in

adolescence. These findings are consistent with multiple prior human[11,13] and animal studies[53,54]

showing that exposures earlier in life may have greater influence on epigenetic patterns[19]. As

early childhood is a time of rapid cognitive, social, emotional, and regulatory development[55],

epigenetic processes may be more malleable during this period[24], resulting in increased

sensitivity to life experiences that shape DNAm levels and trajectories across development.

Importantly, these findings suggest early childhood as a potential period for focused

interventions to limit or prevent the long-term sequelae of childhood adversity.

Of the seven types of adversity examined, exposure to single parent households had the

greatest number of associations to DNAm in adolescence. By contrast, previous research on

DNAm from the same children at age 7 identified no associations with one-adult households[19],

suggesting these effects may be adolescent-specific. Prior studies have shown the effects of

single parent households begin to emerge around puberty, manifesting through shifts in the

timing of puberty[56,57], poorer self-esteem[58], and higher depressive symptoms[59] and externalizing

behaviors[59]. These findings suggest a latency to the effects of one adult households, which may

not become apparent until the rapid biological changes that occur during puberty[60].

Curiously, we observed fewer associations than expected for other adversities, such as

maternal psychopathology and experiences of sexual, physical, or emotional abuse. These

adversities may have subtler influences on the adolescent epigenome, requiring larger sample

sizes or meta-analyses to uncover. Notably, none of our top loci overlapped between different

types of childhood adversity, nor were they present among top loci from a twin study of

adolescent victimization (N=118), where half of the children in the sample were exposed to

severe victimization[22]. However, it remains unknown whether subtypes of adversity might have

differential effects on downstream biological processes and vulnerability to disease. As has been

discussed in ongoing debates surrounding the "lumping or splitting" of childhood adversities in

clinical research[61], it may be that different dimensions of adversity result in different epigenetic

signatures, a hypothesis supported by the finding that adjusting for other types of adversity had

modest influences on the strength of associations. To this end, we investigated potential

differences in sensitive periods for exposures to adversities using a threat vs deprivation

paradigm[62], showing that exposures to deprivation-type adversities may have more influence on

adolescent DNAm than threat-type adversities, particularly when they occur in early childhood

(**Figure S22**).

Arguably the most novel set of findings from our study concerned the patterns of stability

and change in the relationship between adversity and DNAm. Most DNAm trajectories showed

primarily *latent* effects of adversity, meaning they did not emerge until age 15 in youth exposed

to adversity. These findings align with previous longitudinal studies of genome-wide DNAm

from ALSPAC and Project Viva, which have shown that early-life stressors, such as prenatal

maternal smoking[26] and socio-economic disadvantage during childhood[21,27], can have both

immediate and latent effects on DNAm during childhood and adolescence. Subtle

desynchronization of DNAm levels may appear earlier in development, while evading immediate

detection until later in life. These "sleeper" effects may explain why complex diseases unfold

over years of development, rather than immediately after exposures or risk factors[20]. Future

research should investigate whether these latent effects of childhood adversity on the epigenome

persist into adulthood and whether they are more likely to influence physical and mental health

than alterations arising earlier in development.

Similarly, the DNAm differences we previously observed at age 7 did not persist into

adolescence[19]. Studies on early-life stressors[21,27] and markers of prenatal environments, such as

birthweight and gestational age[29], or maternal weight before and during pregnancy[28], parallel

these findings, showing that DNAm differences linked to early-life environments do not

generally persist across time. Whether these patterns resolved naturally or due to active

intervention is unknown and should be investigated to determine whether interventions are

needed to rescue the effects of early-life stressors. Nevertheless, even short-term alterations that

eventually fade over time could still alter the developmental trajectories of cellular pathways or

physiological systems, influencing downstream health and risk for disease[63].

Although this study did not focus on causal relationships, several differentially

methylated genes were implicated in key processes that could influence downstream disease. For

instance, *CUX2* is transcription factor involved in dendrite and synapse formation[64], alterations

to which could influence neurodevelopment and vulnerability to mental disorders. Several top

genes, including *DUSP10*[65], *DSP*[66], and *VEGFA*[67], are also linked to cardiac function, and may

partially reflect mechanisms linking childhood adversity to heart disease[68]. However, additional

research is needed to identify the true health consequences of these differences and determine

whether short- and/or long-term DNAm changes mediate the link between childhood adversity

and health outcomes.

Our study had limitations. First, DNAm data were generated from slightly different tissue

types at each wave. Although we corrected for cell type composition using established methods,

differences in the stability of DNAm differences between waves may have been partially driven

by tissue-based differences and variability. Second, we were unable to replicate our findings in

an independent sample, due to the lack of data available from other cohort studies. We also

cannot fully rule-out the possibility that our findings were influenced by unmeasured or technical

factors specifically influencing age 15 associations. However, our results were robust in internal

validation analyses and when controlling for 14 potential confounders. Nonetheless, future

studies are needed to fully replicate and confirm these longitudinal epigenetic responses to

childhood adversity. Third, our analytic subsample was mainly composed of children from

European descent. This lack of diversity limited the generalizability of our findings. Given the

disparities in childhood adversity exposure and health[69], our findings should be replicated in

more diverse cohorts.

CONCLUSIONS

This study highlights developmental variability in the relationship between childhood

adversity and DNAm trajectories and its potential role in adversity-related health outcomes.

Future studies should continue to investigate longitudinal measures of DNAm to identify the

potential role of latent and persistent epigenetic alterations in driving short- and long-term health

outcomes. Ultimately, this research will help guide intervention strategies and identify individual

who are at higher risk for physical and mental disorders arising from exposure to childhood

adversity.

ACKNOWLEDGEMENTS

This work was supported by the National Institute of Mental Health of the National Institutes of
Health (grant number R01MH113930 awarded to ECD). The content is solely the responsibility
of the authors and does not necessarily represent the official views of the National Institutes of
Health. Dr. Dunn and Dr. Lussier were also supported by a grant from One Mind. We are
extremely grateful to all the families who took part in the ALSPAC study, the midwives for their
help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer
and laboratory technicians, clerical workers, research scientists, volunteers, managers,
receptionists, and nurses. The UK Medical Research Council and Wellcome (Grant ref:
217065/Z/19/Z) and the University of Bristol provide core support for ALSPAC. A
comprehensive list of grants funding is available on the ALSPAC website
(http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf); This
research was specifically funded by grants from the BBSRC (BBI025751/1; BB/I025263/1),
MRC IEU (MC_UU_00011/5), National Institute of Child and Human Development
(R01HD068437), NIH (5RO1AI121226-02), and CONTAMED EU (212502). This publication
is the work of the authors, whom will serve as guarantors for the contents of this paper.
Dr. Walton is funded by CLOSER, whose mission is to maximize the use, value, and impact of
longitudinal studies (www.closer.ac.uk). CLOSER was funded by the Economic and Social
Research Council (ESRC) and the Medical Research Council (MRC) between 2012 and 2017. Its
initial five-year grant has since been extended to March 2021 by the ESRC (grant reference:
ES/K000357/1). The funders took no role in the design, execution, analysis, or interpretation of
the data or in the writing up of the findings. Dr. Walton is also supported by the European
Union's Horizon 2020 research and innovation programme (grant n° 848158).

Finally, we would also like to thank Dr. Garrett Fitzmaurice for his guidance in the

characterization of DNAm trajectories across development. We also thank Elizabeth Leblanc for

her assistance with the prediction of age at pubertal onset using superimposition by translation

and rotation (SITAR).

AUTHOR CONTRIBUTIONS

AAL designed the study, performed all analyses, interpreted the results, and wrote the

manuscript. YZ, BJS, JC, AJS, ADACS, MJS, EW, CLR, and KJR assisted in the design and

interpretation of the study and provided critical input in writing the manuscript. ECD obtained

grant support for this work, designed the study, interpreted the results, and helped write the

manuscript.

COMPETING INTEREST

The authors have no conflicts of interest to declare.

REFERENCES

Slopen, N., Koenen, K. C. & Kubzansky, L. D. Cumulative adversity in childhood and emergent risk factors for long-term health. *Journal of Pediatrics* **164**, 631-638 (2014).

Repetti, R. L., Taylor, S. E. & Seeman, T. E. Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin* **128**, 330-366 (2002).

Brooks-Gunn, J. & Duncan, G. J. Effects of poverty on children. *The Future of Children* **7**, 55-71 (1997).

McLaughlin, K. A. *et al.* Childhood adversities and adult psychiatric disorders in the National Comorbidity Survey Replication II: Associations with persistence of DSM-IV disorders. *Archives of General Psychiatry* **67**, 124-132 (2010).

McLaughlin, K. A. *et al.* Childhood adversities and first onset of psychiatric disorders in a national sample of US adolescents. *JAMA Psychiatry* **69**, 1151-1160 (2012).

Grummitt, L. R. *et al.* Association of Childhood Adversity With Morbidity and Mortality in US Adults: A Systematic Review. *JAMA pediatrics*, doi:10.1001/jamapediatrics.2021.2320 (2021).

Aristizabal, M. J. *et al.* Biological embedding of experience: A primer on epigenetics. *Proceedings of the National Academy of Sciences* **117**, 23261, doi:10.1073/pnas.1820838116 (2020).

Fujii, R., Sato, S., Tsuboi, Y., Cardenas, A. & Suzuki, K. DNA methylation as a mediator of associations between the environment and chronic diseases: A scoping review on application of mediation analysis. *Epigenetics* **ahead-of-print**, 1-27, doi:https://doi.org/10.1080/15592294.2021.1959736 (2021).

Cecil, C. A. M., Zhang, Y. & Nolte, T. Childhood maltreatment and DNA methylation: A systematic review. *Neuroscience & Biobehavioral Reviews* **112**, 392-409, doi:https://doi.org/10.1016/j.neubiorev.2020.02.019 (2020).

Parade, S. H. *et al.* A systematic review of childhood maltreatment and DNA methylation: candidate gene and epigenome-wide approaches. *Translational Psychiatry* **11**, 134, doi:10.1038/s41398-021-01207-y (2021).

Essex, M. J. *et al.* Epigenetic vestiges of early developmental adversity: childhood stress exposure and DNA methylation in adolescence. *Child Development* **84**, 58-75, doi:10.1111/j.1467-8624.2011.01641.x (2013).

Yang, B. Z. *et al.* Child abuse and epigenetic mechanisms of disease risk. *Am J Prev Med* **44**, 101-107, doi:10.1016/j.amepre.2012.10.012 (2013).

Dunn, E. C. *et al.* Sensitive periods for the effect of childhood adversity on DNA methylation: results from a prospective, longitudinal study. *Biological Psychiatry* **85**, 838-849, doi:10.1016/j.biopsych.2018.12.023 (2019).

Klengel, T. *et al.* Allele-specific FKBP5 DNA demethylation mediates gene-childhood trauma interactions. *Nat Neurosci* **16**, 33-41, doi:10.1038/nn.3275 (2013).

Bornstein, M. H. Sensitive periods in development: Structural characteristics and causal interpretations. *Psychological Bulletin* **105**, 179-197 (1989).

Ben-Shlomo, Y. & Kuh, D. A life course approach to chronic disease epidemiology: conceptual models, empirical challenges, and interdisciplinary perspectives. *Int J Epidemiol* **31**, 285-293 (2002).

Knudsen, E. Sensitive periods in the development of the brain and behavior. *J Cogn Neurosci* **16**, 1412-1425 (2004).

Shonkoff, J. P., Boyce, W. T. & McEwen, B. S. Neuroscience, molecular biology, and the childhood roots of health disparities. *JAMA* **301**, 2252-2259 (2009).

Lussier, A. A. *et al.* Updates to data versions and analytic methods influence the reproducibility of results from epigenome-wide association studies. *Epigenetics*, doi:10.1080/15592294.2022.2028072 (2022).

417	20	Oh, E. S. & Petronis, A. Origins of human disease: the chrono-epigenetic perspective. *Nat Rev*
418		*Genet*, doi:10.1038/s41576-021-00348-6 (2021).
419	21	Laubach, Z. M. *et al.* Socioeconomic status and DNA methylation from birth through mid-
420		childhood: a prospective study in Project Viva. *Epigenomics* **11**, 1413-1427, doi:10.2217/epi-
421		2019-0040 (2019).
422	22	Kandaswamy, R. *et al.* DNA methylation signatures of adolescent victimization: analysis of a
423		longitudinal monozygotic twin sample. *Epigenetics*, 1-18, doi:10.1080/15592294.2020.1853317
424		(2020).
425	23	Gui, A. *et al.* Leveraging epigenetics to examine differences in developmental trajectories of
426		social attention: A proof-of-principle study of DNA methylation in infants with older siblings
427		with autism. *Infant behavior & development* **60**, 101409, doi:10.1016/j.infbeh.2019.101409
428		(2020).
429	24	Mulder, R. H. *et al.* Epigenome-wide change and variation in DNA methylation in childhood:
430		trajectories from birth to late adolescence. *Hum Mol Genet* **30**, 119-134,
431		doi:10.1093/hmg/ddaa280 (2021).
432	25	Martino, D. J. *et al.* Evidence for age-related and individual-specific changes in DNA
433		methylation profile of mononuclear cells during early immune development in humans.
434		*Epigenetics* **6**, 1085-1094, doi:10.4161/epi.6.9.16401 (2011).
435	26	Richmond, R. C. *et al.* Prenatal exposure to maternal smoking and offspring DNA methylation
436		across the lifecourse: findings from the Avon Longitudinal Study of Parents and Children
437		(ALSPAC). *Hum Mol Genet* **24**, 2201-2217, doi:10.1093/hmg/ddu739 (2015).
438	27	Alfano, R. *et al.* Socioeconomic position during pregnancy and DNA methylation signatures at
439		three stages across early life: epigenome-wide association studies in the ALSPAC birth cohort.
440		*Int. J. Epidemiol.* **48**, 30-44, doi:10.1093/ije/dyy259 (2019).
441	28	Sharp, G. C. *et al.* Maternal pre-pregnancy BMI and gestational weight gain, offspring DNA
442		methylation and later offspring adiposity: findings from the Avon Longitudinal Study of Parents
443		and Children. *Int J Epidemiol* **44**, 1288-1304, doi:10.1093/ije/dyv042 (2015).
444	29	Simpkin, A. J. *et al.* Prenatal and early life influences on epigenetic age in children: a study of
445		mother–offspring pairs from two cohort studies. *Human Molecular Genetics* **25**, 191-201,
446		doi:10.1093/hmg/ddv456 (2016).
447	30	Martins, J. *et al.* Childhood adversity correlates with stable changes in DNA methylation
448		trajectories in children and converges with epigenetic signatures of prenatal stress. *Neurobiol*
449		*Stress* **15**, 100336, doi:10.1016/j.ynstr.2021.100336 (2021).
450	31	Fraser, A. *et al.* Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC
451		mothers cohort. *Int. J. Epidemiol.* **42**, 97-110, doi:10.1093/ije/dys066 (2013).
452	32	Boyd, A. *et al.* Cohort Profile: the 'children of the 90s'--the index offspring of the Avon
453		Longitudinal Study of Parents and Children. *Int. J. Epidemiol.* **42**, 111-127,
454		doi:10.1093/ije/dys064 (2013).
455	33	Relton, C. L. *et al.* Data Resource Profile: Accessible Resource for Integrated Epigenomic
456		Studies (ARIES). *Int J Epidemiol* **44**, 1181-1190, doi:10.1093/ije/dyv072 (2015).
457	34	Ramo-Fernández, L., Schneider, A., Wilker, S. & Kolassa, I. T. Epigenetic alterations associated
458		with war trauma and childhood maltreatment. *Behavioral sciences & the law* **33**, 701-721 (2015).
459	35	Cunliffe, V. T. The epigenetic impacts of social stress: how does social adversity become
460		biologically embedded? *Epigenomics* **8**, 1653-1669 (2016).
461	36	McLaughlin, K. A. Future directions in childhood adversity and youth psychopathology. *Journal*
462		*of Clinical Child & Adolescent Psychology* **45**, 361-382 (2016).
463	37	Vaiserman, A. M. & Koliada, A. K. Early-life adversity and long-term neurobehavioral
464		outcomes: epigenome as a bridge? *Human genomics* **11**, 34 (2017).
465	38	Eachus, H. & Cunliffe, V. T. in *Epigenetics of Aging and Longevity* 251-270 (Elsevier, 2018).
466	39	Houseman, E. A., Molitor, J. & Marsit, C. J. Reference-free cell mixture adjustments in analysis
467		of DNA methylation data. *Bioinformatics* **30**, doi:10.1093/bioinformatics/btu029 (2014).

40 Ben-Shlomo, Y. & Kuh, D. A life course approach to chronic disease epidemiology: conceptual
models, empirical challenges and interdisciplinary perspectives. *Int. J. Epidemiol.* **31**, 285-293,
doi:10.1093/ije/31.2.285 (2002).
41 Kuh, D. & Ben-Shlomo, Y.   (Oxford University Press, Oxford, 2004).
42 Mishra, G. *et al.* A structured approach to modelling the effects of binary exposure variables over
the life course. *Int. J. Epidemiol.*, doi:10.1093/ije/dyn229 (2009).
43 Smith, A. D. A. C. *et al.* Model Selection of the Effect of Binary Exposures over the Life Course.
Epidemiology, doi:10.1097/EDE.0000000000000348 (2015).
44 Smith, A. D. A. C. *et al.* A structured approach to hypotheses involving continuous exposures
over the life course. *Int. J. Epidemiol.*, doi:10.1093/ije/dyw164 (2016).
45 Zhu, Y. *et al.* A Structured Approach to Evaluating Life Course Hypotheses: Moving Beyond
Analyses of Exposed Versus Unexposed in the Omics Context. *Am. J. Epidemiol.*,
doi:10.1093/aje/kwaa246 (2020).
46 Tibshirani, R. J., Taylor, J., Lockhart, R. & Tibshirani, R. Exact Post-Selection Inference for
Sequential Regression Procedures. *Journal of the American Statistical Association* **111**, 600-620,
doi:10.1080/01621459.2015.1108848 (2016).
47 Amrhein, V. & Greenland, S. Remove, rather than redefine, statistical significance. *Nature
Human Behaviour* **2**, 4-4, doi:10.1038/s41562-017-0224-0 (2018).
48 McShane, B. B., Gal, D., Gelman, A., Robert, C. & Tackett, J. L. Abandon Statistical
Significance. *The American Statistician* **73**, 235-245, doi:10.1080/00031305.2018.1527253
(2019).
49 Hannon, E., Lunnon, K., Schalkwyk, L. & Mill, J. Interindividual methylomic variation across
blood, cortex, and cerebellum: implications for epigenetic studies of neurological and
neuropsychiatric phenotypes. *Epigenetics* **10**, 1024-1032, doi:10.1080/15592294.2015.1100786
(2015).
50 Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. *Nat. Protoc.* **4**, 44-57,
doi:10.1038/nprot.2008.211 (2009).
51 Huang da, W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools: paths toward
the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**, 1-13,
doi:10.1093/nar/gkn923 (2009).
52 Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285-
291, doi:10.1038/nature19057 (2016).
53 Curley, J. P. & Champagne, F. A. Influence of maternal care on the developing brain:
Mechanisms, temporal dynamics and sensitive periods. *Frontiers in neuroendocrinology* **40**, 52-
66, doi:10.1016/j.yfrne.2015.11.001 (2016).
54 Pena, C. J., Neugut, Y. D. & Champagne, F. A. Developmental timing of the effects of maternal
care on gene expression and epigenetic regulation of hormone receptor levels in female rats.
Endocrinology **154**, 4340-4351, doi:10.1210/en.2013-1595 (2013).
55 Shonkoff, J. P. & Phillips, D. A. *From neurons to neighborhoods: The science of early childhood
development*. (National Academy Press, 2000).
56 Aghaee, S. *et al.* Early life household intactness and timing of pubertal onset in girls: a
prospective cohort study. *BMC Pediatrics* **20**, 464, doi:10.1186/s12887-020-02345-w (2020).
57 Ellis, B. J. Timing of pubertal maturation in girls: an integrated life history approach. *Psychol
Bull* **130**, 920-958, doi:10.1037/0033-2909.130.6.920 (2004).
58 Alami, A., Khosravan, S., Sadegh Moghadam, L., Pakravan, F. & Hosseni, F. Adolescents' self-
esteem in single and two-parent families. *Int J Community Based Nurs Midwifery* **2**, 69-76
(2014).
59 Daryanani, I., Hamilton, J. L., Abramson, L. Y. & Alloy, L. B. Single Mother Parenting and
Adolescent Psychopathology. *Journal of Abnormal Child Psychology* **44**, 1411-1423,
doi:10.1007/s10802-016-0128-x (2016).

60 Aylwin, C. F., Toro, C. A., Shirtcliff, E. & Lomniczi, A. Emerging Genetic and Epigenetic
Mechanisms Underlying Pubertal Maturation in Adolescence. *Journal of Research on*
Adolescence **29**, 54-79, doi:https://doi.org/10.1111/jora.12385 (2019).
61 Smith, K. E. & Pollak, S. D. Rethinking Concepts and Categories for Understanding the
Neurodevelopmental Effects of Childhood Adversity. *Perspectives on psychological science : a*
journal of the Association for Psychological Science **16**, 67-93, doi:10.1177/1745691620920725
(2021).
62 McLaughlin, K. A., Sheridan, M. A. & Lambert, H. K. Childhood adversity and neural
development: deprivation and threat as distinct dimensions of early experience. *Neurosci*
Biobehav Rev **47**, 578-591, doi:10.1016/j.neubiorev.2014.10.012 (2014).
63 Walton, E. Epigenome-wide Associations With Attention-Deficit/Hyperactivity Disorder in
Adults: The Need for a Longitudinal Life Course Approach in Epigenetic Psychiatry. *Biological*
Psychiatry **86**, 570-572, doi:https://doi.org/10.1016/j.biopsych.2019.07.021 (2019).
64 Cubelos, B. *et al.* Cux1 and Cux2 regulate dendritic branching, spine morphology, and synapses
of the upper layer neurons of the cortex. *Neuron* **66**, 523-535, doi:10.1016/j.neuron.2010.04.038
(2010).
65 Liu, R. & Molkentin, J. D. Regulation of cardiac hypertrophy and remodeling through the dual-
specificity MAPK phosphatases (DUSPs). *Journal of Molecular and Cellular Cardiology* **101**,
44-49, doi:https://doi.org/10.1016/j.yjmcc.2016.08.018 (2016).
66 Smith, E. D. *et al.* Desmoplakin Cardiomyopathy, a Fibrotic and Inflammatory Form of
Cardiomyopathy Distinct From Typical Dilated or Arrhythmogenic Right Ventricular
Cardiomyopathy. *Circulation* **141**, 1872-1884, doi:10.1161/CIRCULATIONAHA.119.044934
(2020).
67 Liu, D. *et al.* Association of Genetic Polymorphisms on VEGFA and VEGFR2 With Risk of
Coronary Heart Disease. *Medicine* **95**, e3413-e3413, doi:10.1097/MD.000000000003413
(2016).
68 Jakubowski, K. P., Cundiff, J. M. & Matthews, K. A. Cumulative childhood adversity and adult
cardiometabolic disease: A meta-analysis. *Health Psychol.* **37**, 701-715, doi:10.1037/hea0000637
(2018).
69 Slopen, N. *et al.* Racial Disparities in Child Adversity in the U.S.: Interactions With Family
Immigration History and Income. *Am J Prev Med* **50**, 47-56, doi:10.1016/j.amepre.2015.06.013
(2016).
551

TABLES AND FIGURES

Table 1. Top associations between time-dependent exposure to adversity and DNA methylation at age 15.

Adversity	Timing	Age (years)	CpG	DNAm unexp[1]	DNAm SP[2]	Δ DNAm[3]	Effect estimate[4]	SE*	R²	P-value	FDR	Nearest gene	Trajectory class
Caregiver physical or emotional abuse	Early childhood	5	cg14855874	0.091	0.121	0.030	0.030	0.005	0.041	3.32E-07	1.01E-01	BANK1	**Emergent**
			cg15454534	0.885	0.868	-0.017	-0.017	0.003	0.039	6.76E-07	1.02E-01	OR2T1	**Latent**
			cg06215562	0.847	0.826	-0.021	-0.021	0.004	0.035	2.37E-06	1.81E-01		**Latent**
Sexual or physical abuse (by anyone)	Early childhood	3.5	**cg26970800**	**0.902**	**0.847**	**-0.055**	**-0.055**	**0.010**	**0.044**	**8.51E-08**	**2.08E-02**	**CBLIF**	**Emergent**
			cg15723468	**0.822**	**0.779**	**-0.043**	**-0.045**	**0.009**	**0.041**	**1.89E-07**	**2.08E-02**	**GALNT2**	**Latent**
			cg17928317	**0.681**	**0.785**	**0.104**	**0.076**	**0.015**	**0.041**	**2.06E-07**	**2.08E-02**	**MAGEC3**	**Primed**
	Late childhood	8	cg27558057	0.257	0.289	0.032	0.107	0.024	0.036	1.53E-06	1.16E-01	TAF1	**Stable**
Family instability	Very early childhood	2.5	cg02735620	0.877	0.857	-0.021	-0.019	0.004	0.036	2.07E-06	4.63E-01	PKD2	Emergent
Financial hardship	Very early childhood	0.66	cg14455319	0.289	0.339	0.050	0.052	0.011	0.036	3.87E-06	2.00E-01	ANKK1	Time-stable
			cg13204236	0.861	0.824	-0.037	-0.037	0.007	0.036	5.94E-06	2.00E-01	STPG4	Latent
	Early childhood	5	cg15037420	0.780	0.746	-0.035	-0.034	0.007	0.036	3.04E-06	2.00E-01	BSPH1	Latent
			cg06410970	0.860	0.825	-0.035	-0.033	0.006	0.036	5.56E-06	2.00E-01	ANXA11	Overcompensation
	Late childhood	11	cg02011706	0.861	0.799	-0.062	-0.064	0.013	0.036	5.35E-06	2.00E-01	LMF1	Emergent
			cg04659536	0.901	0.873	-0.029	-0.028	0.006	0.035	5.52E-06	2.00E-01	SDK1	Latent
	Recency		cg17670999	0.817	0.807	-0.010	-0.002	0.000	0.041	8.76E-07	2.00E-01	ARHGAP39	Stable
			cg25459301	0.769	0.756	-0.013	-0.003	0.001	0.036	4.24E-06	2.00E-01	XKR6	Overcompensation
			cg06812747	0.837	0.825	-0.012	-0.003	0.001	0.035	4.98E-06	2.00E-01	FBXL16	Stable
Maternal psychopathology	Very early childhood	2.75	**cg16813552**	**0.898**	**0.883**	**-0.015**	**-0.015**	**0.003**	**0.045**	**7.11E-08**	**2.15E-02**	**OGA**	**Stable**
Neighborhood disadvantage	Very early childhood	2.75	cg04288299	0.914	0.905	-0.009	-0.021	0.004	0.039	4.52E-07	7.00E-02	NELFA	**Overcompensation**
			cg25019631	0.201	0.223	0.023	0.044	0.009	0.038	6.16E-07	7.00E-02	CASP9	**Overcompensation**
			cg04224851	0.907	0.894	-0.013	-0.014	0.003	0.038	6.94E-07	7.00E-02	ZFP36L2	**Overcompensation**
One adult in the household	Very early childhood	1.75	**cg05491478**	**0.908**	**0.880**	**-0.028**	**-0.027**	**0.006**	**0.038**	**7.33E-07**	**2.81E-02**	**LRRFIP1**	Overcompensation
	Early childhood	3.9	**cg16907527**	**0.853**	**0.824**	**-0.030**	**-0.032**	**0.005**	**0.060**	**4.17E-10**	**1.26E-04**	**VEGFA**	**Flat emergent**
			cg08818094	**0.847**	**0.798**	**-0.048**	**-0.050**	**0.008**	**0.051**	**8.79E-09**	**1.33E-03**	**TBC1D19**	**Latent**
			cg01060989	**0.824**	**0.794**	**-0.031**	**-0.031**	**0.005**	**0.047**	**6.73E-08**	**6.78E-03**	**DUSP10**	**Latent**
			cg15814750	**0.723**	**0.684**	**-0.039**	**-0.040**	**0.009**	**0.039**	**6.57E-07**	**2.81E-02**	**WDR72**	**Latent**
			cg15783822	**0.868**	**0.848**	**-0.021**	**-0.021**	**0.004**	**0.039**	**8.08E-07**	**2.81E-02**	**PRR4**	**Latent**
			cg15864691	**0.907**	**0.889**	**-0.018**	**-0.018**	**0.004**	**0.038**	**8.36E-07**	**2.81E-02**	**HOXA10**	**Overcompensation**
			cg02584161	**0.661**	**0.603**	**-0.057**	**-0.058**	**0.011**	**0.038**	**1.28E-06**	**3.42E-02**		**Latent**
			cg02810291	**0.840**	**0.818**	**-0.022**	**-0.023**	**0.005**	**0.037**	**1.35E-06**	**3.42E-02**	**AKAP13**	**Overcompensation**
			cg04036644	**0.882**	**0.855**	**-0.027**	**-0.026**	**0.005**	**0.037**	**1.36E-06**	**3.42E-02**	**LOC286083**	**Latent**

			DNAm unexp.	DNAm exp. SP	ΔDNAm		SE		p-value	FDR	Gene	Trajectory
		cg11811897	**0.758**	**0.711**	**-0.047**	**-0.047**	**0.010**	**0.037**	**1.68E-06**	**3.64E-02**	**PKD1L1**	**Latent**
		cg15817130	**0.794**	**0.759**	**-0.036**	**-0.038**	**0.007**	**0.037**	**1.83E-06**	**3.69E-02**	**MYO10**	**Latent**
		cg06711254	**0.686**	**0.631**	**-0.055**	**-0.056**	**0.012**	**0.036**	**2.15E-06**	**3.98E-02**	**FSIP2**	**Flat emergent**
		cg19096460	**0.845**	**0.821**	**-0.024**	**-0.024**	**0.005**	**0.035**	**2.89E-06**	**4.85E-02**	**HERC3**	**Latent**
		cg18980650	0.800	0.760	-0.040	-0.036	0.007	0.035	3.31E-06	5.08E-02	NOX1	Emergent
		cg27504269	0.771	0.733	-0.038	-0.040	0.008	0.036	3.52E-06	5.08E-02	SLCO1A2	Latent
Late childhood	10	**cg12096528**	**0.890**	**0.874**	**-0.016**	**-0.016**	**0.003**	**0.036**	**2.24E-06**	**3.98E-02**	**SLC25A41**	**Overcompensation**
Accumulation		**cg00807464**	**0.052**	**0.057**	**0.006**	**0.003**	**0.001**	**0.040**	**7.56E-07**	**2.81E-02**	**CUX2**	**Stable**
		cg10420609	**0.559**	**0.522**	**-0.037**	**-0.014**	**0.003**	**0.039**	**7.71E-07**	**2.81E-02**	**DSP**	**Latent**
		cg14579651	**0.634**	**0.605**	**-0.028**	**-0.012**	**0.002**	**0.037**	**1.68E-06**	**3.64E-02**	**STK38L**	**Stable**

[1]DNAm unexp. = mean DNA methylation levels in children with no exposure to adversity from ages 0 to 11.

[2]DNAm exp. SP = mean DNA methylation levels in children with exposure to adversity that occurred during the selected sensitive period (SP). Accumulation hypotheses show the mean DNA methylation levels in children with at least one exposure to adversity.

[3]ΔDNAm= difference in mean DNA methylation levels between children exposed to adversity during the selected sensitive period and individuals unexposed to adversity (i.e., DNAm exp. SP – DNAm unexp.)

[4]Effect estimates were calculated using linear regression of exposure to adversity from the theoretical model and DNA methylation, correcting for the covariates described in the methods.

* SE = standard error; bolded loci passed a 5% FDR threshold; Very early childhood = 0-3 years, Early childhood = 3-5 years; Late childhood = 8-11 years.

Table 2. Types of DNAm trajectories and response to childhood adversity.

Trajectory name	# R^2 loci[1]	# FDR loci[2]	Sample trajectory[3]	Primary characteristics
Emergent	5	1		Differences in the exposed-SP group appeared in childhood but did not fully emerge until age 15.
Flat emergent	2	2		Differences in the exposed-SP group were modest throughout childhood and fully emerged by age 15.
Overcompensation	9	4		Cross-over effects in the exposed-SP groups were present from age 7 to age 15, along with differences in DNAm level between ages.
Latent	17	11		Effects of childhood adversity for the exposed-SP group emerged at age 15, with no significant effects observed from exposure at other times. Some CpGs in this cluster showed graded effects between childhood exposed in sensitive periods versus other times.
Primed	1	1		Differences in the exposed groups were apparent from birth but were magnified in the exposed-SP group at age 15.
Stable (no group-by-age interaction)	7	3		Differences in the exposed-SP group were present at age 7 and remained stable until age 15.

[1]Of the top 41 loci identified in the SLCMA of adversity between ages 0-11 and adolescent DNA methylation that passed the R2≥0.035 and $p < 1 \times 10^{-5}$ thresholds.

[2]Of the top 22 loci identified in the SLCMA of adversity between ages 0-11 and adolescent DNA methylation that passed an FDR<0.05 threshold.

[3]Sample trajectories show the DNA methylation (DNAm) levels on the y-axis and the age at DNAm collection on the x-axis. Dashed, red trajectories represent the mean DNAm of children exposed to adversity during the sensitive period identified in the SLCMA (exposed-SP). Dotted, blue trajectories represent the mean DNAm of those exposed to adversity outside the sensitive

period identified in the SLCMA (exposed-other). Solid, black trajectories represent the mean DNAm of children with no exposure to adversity across development (unexposed).

Figure 1. Summary of exposures and outcomes examined in the present study. Seven types of childhood adversity were assessed

5-8 times between the ages of 0 and 11. Children with complete cases across all time points and covariates were included in the

present study (N=609-665). Each filled cell represents the time point when the adversity was measured, along with the prevalence of

children exposed to adversity. Colors represent the four sensitive periods used to define time-dependent exposure to adversity: *very*

early childhood (age 0-3), *early childhood* (age 3-5), *middle childhood* (age 5-7), and *late childhood* (age 8-11). The additional life

course models tested were accumulation and recency, which reflect the total number of exposures across development and exposure to

adversity weighted by time, respectively. Genome-wide DNA methylation (DNAm) data were collected at age 0, 7, and 15.

Figure 2. Life course theoretical models selected by the SLCMA for top loci at age 15. The life course theoretical models were split by sensitive periods (i.e., exposure to adversity during specific childhood periods) or additive models (i.e., accumulation or recency of exposures). Colors represent the different types of adversity. The distribution of theoretical models for top loci was significantly different than random chance, with exposure to adversity during sensitive periods more frequently predicting DNA methylation levels as compared to the additive models. **A)** 22 loci were identified at a false-discovery rate (FDR) <0.05. Most loci were associated with exposure to one adult households during early childhood. **B)** 41 loci were identified at an R2≥0.035 cutoff and p<1x10^{-5} threshold, which again mainly showed associations with adversity occurring during early childhood.

Figure 3. Persistence and stability of associations between childhood adversity and DNA methylation across development. A) The effect estimates of associations between childhood adversity and DNAm at age 7 or age 15 generally showed variable directions of effect for the significant loci identified from the SLCMA at age 15 (20 concordant and 21 non-concordant directionality). Effect estimates for age 7 DNAm data were also smaller than those at age 15, suggesting that these loci showed latent responses to adversity. **B)** The effect estimates of associations between childhood adversity and DNAm at age 7 or age 15 generally showed variable directions of effect for the significant loci identified in a previous study of age 7 DNAm (24 concordant and 22 non-concordant directionality). Effect estimates for age 15 DNAm data were also smaller than those at age 7, suggesting that these loci showed early responses to adversity that resolved by adolescence