A cross-sectional study of the association between frequency of telecommuting and unhealthy dietary habits among Japanese workers during the COVID-19 pandemic

Yoshiko Kubo[1], Tomohiro Ishimaru[1], Ayako Hino[2], Masako Nagata[3], Kazunori Ikegami[4], Seiichiro Tateishi[5], Mayumi Tsuji[6], Shinya Matsuda[7], and Yoshihisa Fujino[1], for the CORoNaWork project

[1] Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan

[2] Department of Mental Health, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan

[3] Department of Occupational Health Practice and Management, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan

[4] Department of Work Systems and Health, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan

[5] Department of Occupational Medicine, School of Medicine, University of Occupational and Environmental Health, Japan

[6] Department of Environmental Health, School of Medicine, University of Occupational and Environmental Health, Japan

[7] Department of Preventive Medicine and Community Health, School of Medicine,

University of Occupational and Environmental Health, Japan

Correspondence to Yoshihisa Fujino, M.D., M.P.H., Ph.D.

Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,

University of Occupational and Environmental Health, Japan

1-1, Iseigaoka, Yahatanishiku, Kitakyushu, 807-8555, Japan

Tel: +81-93-691-7401

Email: zenq@med.uoeh-u.ac.jp

Abstract

Background: As a consequence of the COVID-19 pandemic, telecommuting has become a new way of working that has not only changed individuals' work, but also their health, lifestyle, family and community relationships. However, the impact of telecommuting on diet is not well understood. The aim of this study was to determine the relationship between telecommuting frequency and unhealthy dietary habits among Japanese workers during the COVID-19 pandemic.

Methods: A total of 33,302 Japanese workers completed an Internet survey about telecommuting and dietary habits. Data from 13,468 office workers who telecommuted were analyzed. Telecommuting frequency during the COVID-19 pandemic was extracted from a questionnaire. The odds ratios (ORs) of four types of dietary habits, namely, skipping breakfast, solitary eating, lower meal frequency, and meal substitution associated with telecommuting frequency were estimated using multilevel logistic regression.

Results: The multivariate OR of skipping breakfast was 1.15 (95% CI: 1.03–1.29, p=0.013) for participants who telecommuted in excess of four days per week compared to those who rarely telecommuted. Similarly, the multivariate OR of solitary eating, lower meal frequency and meal substitution were 1.44 (95%CI: 1.28–1.63, p<0.001), 2.39 (95%CI: 1.66–3.44, p<0.001), and 1.26 (95%CI: 1.04–1.51, p=0.015) for those who telecommuted in excess of four days per week compared to those who rarely telecommuted. There was a statistically significant increase in the dose-response trend in ORs of solitary eating (p for trend <0.001), lower meal frequency (p for trend <0.001), and meal substitution (p for trend =0.001) with increasing telecommuting frequency.

Conclusion: Increasing telecommuting frequency was associated with an increasing number of unhealthy dietary habits in Japanese workers. Telecommuting during the COVID-19 pandemic, together with other social changes such as refraining from going out and

socializing, is having a significant impact on workers' daily lives and behaviors. This study suggests that in addition to challenges related to physical activity, loneliness, and mental health, telecommuters may also be developing unhealthy dietary habits, indicating the need for strategies to help telecommuters manage their nutrition and diet.

Trial registration: N.A.

Keywords: Teleworking, COVID-19, Feeding behavior, Breakfast, Meal Times, Japan

Background

Telecommuting is a symbol of the new normal brought about by the global COVID-19 pandemic. Governments around the world have recommended telecommuting as a key measure to restrict the movement of people in an effort to contain the spread of infection.[1] In Japan, many companies have introduced telecommuting at the request of government directives.[2] Although some companies had implemented telecommuting before the COVID-19 pandemic, current telecommuting practices differ from those initiated before the pandemic in many respects: the practice advanced rapidly, including among companies and workers who had no experience with telecommuting; was not based on workers' preference; has been taken up by workers in many types of jobs; and has been introduced even in small and medium-sized companies. As a result, the impact of telecommuting on workers' lives and society has been enormous.

For many workers, telecommuting is a new way of working that has not only changed their work, but also their health, lifestyle, family and community relationships. However, the full impact of telecommuting is not well understood. The change to telecommuting has had an impact on health problems previously identified in office workers, such as back pain, visual display terminal-related symptoms, and eye strain, as well as work productivity.[3,4] There are also reports that telecommuting affects stress, communication, and mental health.[5] In addition, telecommuting is thought to be having a significant effect on lifestyle habits like alcohol consumption, exercise, and sleep.[6–10]

Telecommuting is also expected to significantly change workers' eating habits.[9,11–13] For example, reduced commuting leaves more time for other life activities and preparations; lower frequency of going out and less physical activity due to telecommuting may lead to lower calorie consumption; and more flexible use of time at home may lead to more snacking. Government directives to refrain from going out to prevent the spread of

COVID-19 may also have reduced the frequency of eating out and shopping for food. Such changes brought about by telecommuting will thus likely affect how frequently individuals eat meals, whether or not they skip breakfast, the types of foods they consume as meals, and who they eat with. A systematic review reported that there is an association between dietary habits and various occupational factors, including work hours, shift work and job stress.[14] However, few studies have examined how workers' dietary habits have changed due to telecommuting in the COVID-19 pandemic, when they have been required to refrain from going out.[11]

We hypothesized that skipping breakfast, solitary eating, meal substitution, and lower meal frequency are associated with telecommuting. We thus examined the relationship between the frequency of telecommuting and unhealthy dietary habits among Japanese workers during the COVID-19 pandemic.

Methods

Study Design and Participants

A cross-sectional Internet monitor study was conducted between December 22 and 26, 2020, while Japan was experiencing its third wave of COVID-19 infection. Details of the study protocol are published elsewhere.[15] In brief, workers who indicated employment while participating in the survey were included. We used regional sampling to choose participants according region, job type, and sex because COVID-19 infection rates vary greatly by region in Japan.

Of the 33,302 participants, we excluded 6,266 who provided fraudulent responses, which were defined as follows: extremely short response time (≤6 minutes), extremely low body weight (<30 kg), extremely short height (<140 cm), inconsistent answers to similar questions (e.g., inconsistent answers to questions about marital status and living area), and incorrect answers to a question purposely included to detect fraudulent responses (choose the third largest number from the following five numbers). A total of 13,468 (6,896 men and 6,572 women) of the remaining 27,036 participants who stated that they predominantly perform desk work were included in the analysis.

This study was approved by the ethics committee of the University of Occupational and Environmental Health, Japan (reference No. R2-079 and R3-006). Participants provided informed consent by completing a form on the survey website.

Frequency of telecommuting

Telecommuting frequency was determined using the question, "How frequently do you currently telecommute?" Participants chose from the following options: "more than 4 days per week," "more than 2 days per week," "less than 1 day per week," and "hardly ever."

Dietary habits

The following questions were used to determine whether participants missed breakfast and ate alone: "How often do you usually eat breakfast?" and "How often do you eat all of the meals in a day alone?" Respondents chose from the following five options: "6–7 days per week," "4–5 days per week," "2–3 days per week," "less than 1 day per week," and "almost never."

We also inquired about the number of meals and the number of alternative meals eaten per day to determine meal frequency and adoption of meal substitution using the following questions: "How often do you consume meals a day?" and "How many times a day do you eat only individual food items and packaged foods (yogurt, protein bars, pastries, etc.)?" Respondents chose from the following five options: "more than 4 times per day," "3 times a day," "2 times a day," "1 time a day," and "hardly ever."

Other covariates

We controlled for the following confounding variables: age, sex, education, equivalent income (household income divided by the square root of household size), cohabitation status, smoking status (never; quit smoking more than one year ago; quit

smoking within the past year; started smoking less than one year ago; smoking for more than one year), alcohol consumption (6–7 days a week; 4–5 days a week; 2–3 days a week; less than 1 day a week; hardly ever), and physical activity more than 30 minutes per day (6–7 days a week; 4–5 days a week; 2–3 days a week; less than 1 day a week; hardly ever).

Additionally, the community-level variable was the cumulative COVID-19 infection rate in the participants' prefecture of residence one month before the start of the survey.

Statistical analysis

We determined the age-sex-adjusted and multivariate-adjusted odds ratios (ORs) of unfavorable dietary habits associated with telecommuting frequency. We defined unfavorable dietary habits as skipping breakfast, solitary eating, lower meal frequency, and meal substitution. To control for differences in residential area, we used a multilevel logistic model nested in the prefecture of residence. The multivariate model included sex, age, equivalent household income, education, cohabitation status, smoking status, alcohol consumption, and physical activity. Additionally, we used the COVID-19 infection rate in each prefecture as a prefecture-level variable in all comparisons. A p value <0.05 was considered statistically significant. All analyses were conducted using Stata (Stata Statistical Software: Release 16; StataCorp LLC, TX, USA).

Results

Participants' characteristics by telecommuting frequency are summarized in Table 1. Of the 13,468 office workers included in the analysis, approximately 30% telecommuted at least once a week. Those who telecommuted more often tended to be slightly older and male, and less likely to be living with family. Those who telecommuted more frequently tended to have more unfavorable eating habits. Among workers who hardly telecommuted, 25.5% missed breakfast, 25.9% ate all meals alone, 1% ate less than two meals a day, and 6.6% adopted meal substitution. The corresponding proportions among workers who telecommuted in excess of four days per week were 28.7%, 37.0%, 2.5%, and 8.4%, respectively.

The ORs of the four unhealthy dietary habits associated with telecommuting frequency are summarized in Table 2. The multivariate OR of skipping breakfast was 1.15 (95% CI: 1.03–1.29, $p=0.013$) among participants who telecommuted in excess of four days per week compared to those who rarely telecommuted. Similarly, the multivariate ORs of solitary eating, lower meal frequency, and meal substitution were 1.44 (95%CI: 1.28–1.63, $p<0.001$), 2.39 (95%CI: 1.66–3.44, $p<0.001$), and 1.26 (95%CI: 1.04–1.51, $p=0.015$), respectively. There was a statistically significant increasing dose-response trend in ORs of solitary eating (p for trend <0.001), lower meal frequency (p for trend <0.001), and meal substitution (p for trend $=0.001$) with increasing telecommuting frequency.

Multivariate adjustment slightly attenuated the results of the age-adjusted model, but did not affect the findings. Multivariate adjustment also revealed a marginal dose-response trend (p for trend$=0.068$) in the association between skipping breakfast and telecommuting frequency that was not observed in the univariate analysis ($p=0.248$).

Discussion

We found that telecommuting frequency was associated with four unhealthy dietary habits: skipping breakfast, solitary eating, lower meal frequency, and meal substitution. To our knowledge, this is the first study to examine these associations in the COVID-19 pandemic, during which telecommuting has been recommended around the world to prevent the spread of infection.

Eating breakfast is widely recommended for maintaining good health.[16,17] Numerous studies have demonstrated an association between skipping breakfast and adverse health effects such as overweight, obesity, cardiovascular disease, type 2 diabetes, and poor cognitive performance.[18–20] In this study, participants who worked from home in excess of four or more a week were significantly more likely to skip breakfast compared to those who rarely worked from home. However, contrary to our hypothesis, there was no clear dose-response relationship between telecommuting frequency and frequency of skipping breakfast. Several reasons may explain the lack of a dose-response relationship. First, families need breakfast regardless of a worker's telecommuting status. This is supported by the fact that adjustment for cohabitation status in the multivariate analysis produced a dose-response trend that approached significance. Second, the definition of breakfast is ambiguous.[18,21] While some people define breakfast as the first meal eaten upon waking, others define it as any meal eaten in the morning. Thus, vagueness and misclassification of definitions may have affected the observed associations.

We also found that greater telecommuting frequency was associated with more frequent solitary eating. In Japanese culture, avoidance of solitary eating is considered an important public health practice.[22] One of the goals of Japan's national public health policy is to promote community eating.[23] The declining birthrate, women's participation in society, and fragmentation of family life in Japan are thought to have led to an increase in

solitary eating, which is linked to unbalanced nutrition, disrupted life rhythm, and loneliness.[24] However, there is little formal evidence on the health effects of solitary eating, with very few studies having examined the impact of solitary eating on health.[25] During the COVID-19 pandemic, in addition to urging the public to stay home, the government also suggested that individuals avoid eating with people with whom they do not normally spend time. Frequent telecommuting likely further reduced individuals' opportunity to share meals with others. Given the surging public health challenges associated with loneliness and mental health around the world brought about by the pandemic,[26] our results suggest the need to care for those who are forced to eat alone while telecommuting.

There is currently no consensus on how many meals a person should consume per day for good health. However, studies have shown that, in general, consuming more frequent meals per day is associated with a lower risk of weight gain, obesity, high cholesterol, and type 2 diabetes.[20,27] The number of meals consumed per day differs by country and time. Japanese people generally eat three times a day. The current standard health guidance in Japan recommends balanced nutrition and necessary calorie intake at breakfast, lunch, and dinner, and refraining from placing extreme restrictions on the number of meals consumed per day.[28,29] The present study showed that an increase in telecommuting frequency was associated with an increase in the number of people who ate fewer meals per day. We speculate that there may be several possible explanations for this finding. Increased telecommuting frequency may decrease the need for people to go out for commuting purposes and thus their physical activity. This would reduce calorie consumption, which may lead to lower appetite and less frequent meals. Workers who telecommute may be more likely to spend longer hours at home and snack, which may compensate for the food that would normally be consumed at meals. Requests to refrain from going out to prevent the spread of infection may have led people to refrain from going out for meals and shopping, thus

reducing the number of meals eaten. Alternatively, telecommuting may be affecting individuals' loneliness and mental health, thus affecting their dietary patterns.[30–32]

Additionally, we found that increased telecommuting frequency was associated with more frequent meal substitution with individual food items and processed or packaged foods such as bread, cookies, nutritional supplements, yogurt, and pastries. Eating a few individual items and packaged foods may compensate for the consumption of fewer meals. Directives to refrain from going out are likely to have led individuals to eating meals they can easily prepare and buy food with a longer shelf life. The health effects of such meal substitution are unknown, as they depend on the food and frequency of consumption.

This study showed that telecommuters were more likely to have an unhealthy diet, which is associated with the risk of cardiovascular disease and diabetes.[27] Telecommuters are also reportedly less physically active.[7–9] Together, these results indicate the importance of managing cardiovascular and diabetes risk in telecommuters. While guidelines for the work environment have been published for desk workers who telecommute, including specifications for room illumination, display illumination, room temperature, and ergonomic desks and chairs, [33] strategies for promoting healthy diet and exercise habits among telecommuters form a new challenge in occupational health practice.

There are several limitations to this study. First, while we found an association between poor dietary habits and telecommuting frequency, these relationships may not be solely related to the impact of telecommuting, but the COVID-19 pandemic at large and a variety of other consequences such as staying at home and reduced interaction with people. Second, because this was a cross-sectional study, we cannot prove causality. However, we believe that it is logically unlikely that dietary habits could directly affect telecommuting. Third, we examined both the frequency of telecommuting and eating habits in the questionnaire. Dietary surveys are associated with several well-known disadvantages,

including under-reporting, over-reporting, and bias.[34,35] In addition, we did not specify a time period in our questions. Because of the inter-day variability in dietary intake, the impact of these factors on our results is unclear.[36]

Conclusions

We found that increasing telecommuting frequency was associated with an increasing number of unhealthy dietary habits. Telecommuting during the COVID-19 pandemic, together with other social constraints such as refraining from going out and socializing, is having a significant impact on workers' daily lives and behaviors. This study suggests that in addition to challenges related to physical activity, loneliness, and mental health, telecommuters may also be developing unhealthy dietary habits, indicating the need for strategies to help telecommuters manage their nutrition and diet.

Declarations

Ethics approval and consent to participate: This study was approved by the Ethics Committee of the University of Occupational and Environmental Health, Japan (reference No. R2-079 and R3-006). Informed consent was obtained via a form on the survey website.

Consent for publication: N.A.

Availability of data and material: The datasets generated and/or analyzed during the current study are not publicly available due to ethical restrictions, but are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests

Funding: This study was supported and partly funded by the University of Occupational and Environmental Health, Japan; General Incorporated Foundation (Anshin Zaidan); The Development of Educational Materials on Mental Health Measures for Managers at Small-sized Enterprises; Health, Labour and Welfare Sciences Research Grants; Comprehensive Research for Women's Healthcare (H30-josei-ippan-002); Research for the Establishment of an Occupational Health System in Times of Disaster (H30-roudou-ippan-007); scholarship donations from Chugai Pharmaceutical Co., Ltd., the Collabo-Health Study Group, and Hitachi Systems, Ltd.

Authors' contributions: YK contributed to writing the manuscript; TI contributed to creating the questionnaire, reviewing the manuscript, and advice on interpretation; ST, HE, MT, AO, KM and SM contributed to reviewing the manuscript, advice on interpretation, and

securing funding for the research; YF contributed to the overall research plan, creating the questionnaire, analysis, and drafting the manuscript. All authors read and approved the final manuscript.

Acknowledgements: The current members of the CORoNaWork Project, in alphabetical order, are as follows: Dr. Yoshihisa Fujino (present chairperson of the study group), Dr. Akira Ogami, Dr. Arisa Harada, Dr. Ayako Hino, Dr. Hajime Ando, Dr. Hisashi Eguchi, Dr. Kazunori Ikegami, Dr. Kei Tokutsu, Dr. Keiji Muramatsu, Dr. Koji Mori, Dr. Kosuke Mafune, Dr. Kyoko Kitagawa, Dr. Masako Nagata, Dr. Mayumi Tsuji, Ms. Ning Liu, Dr. Rie Tanaka, Dr. Ryutaro Matsugaki, Dr. Seiichiro Tateishi, Dr. Shinya Matsuda, Dr. Tomohiro Ishimaru, and Dr. Tomohisa Nagata. All members are affiliated with the University of Occupational and Environmental Health, Japan.

References

1. Arefi MF, Poursadeqiyan M. A review of studies on the COVID-19 epidemic crisis disease with a preventive approach. Work. 2020;66:717–29.

2. Goto M HW. Trends in Surveys about Home Teleworking During the COVID-19 Pandemic in Japan. Journal of the Institute of Nuclear Safety System. 2020;27:252–74.

3. Parihar JKS, Jain VK, Chaturvedi P, Kaushik J, Jain G, Parihar AKS. Computer and visual display terminals (VDT) vision syndrome (CVDTS). Armed Forces Med J India. 2016;72:270–6.

4. Waongenngarm P, Areerak K, Janwantanakul P. The effects of breaks on low back pain, discomfort, and work productivity in office workers: A systematic review of randomized and non-randomized controlled trials. Appl Ergon. 2018;68:230–9.

5. Oakman J, Kinsman N, Stuckey R, Graham M, Weale V. A rapid review of mental and physical health effects of working at home: how do we optimise health? BMC Public Health. 2020;20:1825.

6. Hallman DM, Januario LB, Mathiassen SE, Heiden M, Svensson S, Bergström G. Working from home during the COVID-19 outbreak in Sweden: effects on 24-h time-use in office workers. BMC Public Health. 2021;21:528.

7. Koohsari MJ, Nakaya T, McCormack GR, Shibata A, Ishii K, Oka K. Changes in Workers' Sedentary and Physical Activity Behaviors in Response to the COVID-19 Pandemic and Their Relationships With Fatigue: Longitudinal Online Study. JMIR Public Health Surveill. 2021;7:e26293.

8. Henke RM, Benevent R, Schulte P, Rinehart C, Crighton KA, Corcoran M. The Effects of Telecommuting Intensity on Employee Health. Am J Health Promot. 2016;30:604–12.

9. Kishimoto M, Ishikawa T, Odawara M. Behavioral changes in patients with diabetes during the COVID-19 pandemic. Diabetol Int. 2020;1–5.

10. Callinan S, Mojica-Perez Y, Wright CJC, Livingston M, Kuntsche S, Laslett A-M, et al. Purchasing, consumption, demographic and socioeconomic variables associated with shifts in alcohol consumption during the COVID-19 pandemic. Drug Alcohol Rev. 2021;40:183–91.

11. Restrepo BJ, Zeballos E. The effect of working from home on major time allocations with a focus on food-related activities. Rev Econ Househ. 2020;1–23.

12. Lizana PA, Vega-Fernadez G, Gomez-Bruton A, Leyton B, Lera L. Impact of the COVID-19 Pandemic on Teacher Quality of Life: A Longitudinal Study from before and during the Health Crisis. Int J Environ Res Public Health [Internet]. 2021;18. Available from: http://dx.doi.org/10.3390/ijerph18073764

13. Xiao Y, Becerik-Gerber B, Lucas G, Roll SC. Impacts of Working From Home During COVID-19 Pandemic on Physical and Mental Well-Being of Office Workstation Users. J Occup Environ Med. 2021;63:181–90.

14. Tanaka R, Tsuji M, Tsuchiya T, Kawamoto T. Association Between Work-Related Factors and Diet: A Review of the Literature. Workplace Health Saf. 2019;67:137–45.

15. Fujino Y, Ishimaru T, Eguchi H, Tsuji M, Tateishi S, Ogami A, et al. Protocol for a nationwide Internet-based health survey in workers during the COVID-19 pandemic in 2020. medRxiv. 2021;2021.02.02.21249309.

16. Rong S, Snetselaar LG, Xu G, Sun Y, Liu B, Wallace RB, et al. Association of Skipping Breakfast With Cardiovascular and All-Cause Mortality. J Am Coll Cardiol. 2019;73:2025–32.

17. Yokoyama Y, Onishi K, Hosoda T, Amano H, Otani S, Kurozawa Y, et al. Skipping Breakfast and Risk of Mortality from Cancer, Circulatory Diseases and All Causes: Findings from the Japan Collaborative Cohort Study. Yonago Acta Med. 2016;59:55–60.

18. Ma X, Chen Q, Pu Y, Guo M, Jiang Z, Huang W, et al. Skipping breakfast is associated with overweight and obesity: A systematic review and meta-analysis. Obes Res Clin Pract. 2020;14:1–8.

19. Hoyland A, Dye L, Lawton CL. A systematic review of the effect of breakfast on the cognitive performance of children and adolescents. Nutr Res Rev. 2009;22:220–43.

20. Horikawa C, Kodama S, Yachi Y, Heianza Y, Hirasawa R, Ibe Y, et al. Skipping breakfast and prevalence of overweight and obesity in Asian and Pacific regions: a meta-analysis. Prev Med. 2011;53:260–7.

21. O'Neil CE, Byrd-Bredbenner C, Hayes D, Jana L, Klinger SE, Stephenson-Martin S. The role of breakfast in health: definition and criteria for a quality breakfast. J Acad Nutr Diet. 2014;114:S8–26.

22. Ishikawa M, Takemi Y, Yokoyama T, Kusama K, Fukuda Y, Nakaya T, et al. "Eating Together" Is Associated with Food Behaviors and Demographic Factors of Older Japanese People Who Live Alone. J Nutr Health Aging. 2017;21:662–72.

23. Minister of Health, Labour and Welfare, Japan. Healthy Japan 21 [Internet]. 2012 [cited 2021 Jun 9]. Available from: https://www.mhlw.go.jp/file/06-Seisakujouhou-10900000-Kenkoukyoku/0000047330.pdf

24. Kim S. Solitary eating, an inferior alternative? An examination of time-use data in South Korea. Int Sociol. SAGE Publications Ltd; 2020;35:415–32.

25. Stürmer B, Ouyang G, Palazova M, Schacht A, Martín-Loeches M, Rausch P, et al. Lunching for Relaxation or Cognitive Control? After-Effects of Social and Solitary Meals. Adv Cogn Psychol. 2018;14:14–20.

26. Jeste DV, Lee EE, Cacioppo S. Battling the Modern Behavioral Epidemic of Loneliness: Suggestions for Research and Interventions. JAMA Psychiatry. 2020;77:553–4.

27. St-Onge M-P, Ard J, Baskin ML, Chiuve SE, Johnson HM, Kris-Etherton P, et al. Meal Timing and Frequency: Implications for Cardiovascular Disease Prevention: A Scientific Statement From the American Heart Association. Circulation. 2017;135:e96–121.

28. Ministry of Health, Labour and Welfare and Ministry of Agriculture, Forestry and Fisheries. Japanese Food Guide Spinning Top [Internet]. 2005 [cited 2021 Jun 9]. Available from: https://www.mhlw.go.jp/bunya/kenkou/pdf/eiyou-syokuji4.pdf

29. y the Ministry of Education, the Ministry of Health and Welfare, and the Ministry of Agriculture, Forestry and Fisheries. Dietary guidelines for Japanese [Internet]. 2006 [cited 2021 Jun 9]. Available from: https://www.maff.go.jp/j/syokuiku/attach/pdf/shishinn-3.pdf

30. Poggiogalle E, Kiesswetter E, Romano M, Saba A, Sinesio F, Polito A, et al. Psychosocial and cultural determinants of dietary intake in community-dwelling older adults: A Determinants of Diet and Physical Activity systematic literature review. Nutrition. 2021;85:111131.

31. Naeim M, Rezaeisharif A, Kamran A. COVID-19 Has Made the Elderly Lonelier. Dement Geriatr Cogn Dis Extra. 2021;11:26–8.

32. Björnwall A, Mattsson Sydner Y, Koochek A, Neuman N. Eating Alone or Together among Community-Living Older People-A Scoping Review. Int J Environ Res Public Health [Internet]. 2021;18. Available from: http://dx.doi.org/10.3390/ijerph18073495

33. Ministry of Health, Labour and Welfare, Japan. 自宅等でテレワークを行う際の作業環境整備 [Internet]. Improving the work environment when teleworking at home. [cited 2021 Jun 9]. Available from: https://www.mhlw.go.jp/stf/newpage_01603.html

34. Murakami K, Sasaki S, Takahashi Y, Uenishi K, Yamasaki M, Hayabuchi H, et al. Misreporting of dietary energy, protein, potassium and sodium in relation to body mass index in young Japanese women. Eur J Clin Nutr. 2008;62:111–8.

35. Zhang J, Temme EH, Sasaki S, Kesteloot H. Under- and overreporting of energy intake using urinary cations as biomarkers: relation to body mass index. Am J Epidemiol. 2000;152:453–62.

36. Ogawa K, Tsubono Y, Nishino Y, Watanabe Y, Ohkubo T, Watanabe T, et al. Inter- and intra-individual variation of food and nutrient consumption in a rural Japanese population. Eur J Clin Nutr. 1999;53:781–5.

Table 1. Subjects chracteristics according to frequency of telecommuting

Factor	Frequency of telecommuting			
	≥4 days/week n=2042	2-3 days/week n=1058	≤1 day/week n=952	Hardly ever n=9416
Age, mean (SD)	49.1 (10.0)	48.4 (10.3)	48.1 (10.2)	47.3 (10.2)
Sex, male (%)	1159 (56.8%)	591 (55.9%)	613 (64.4%)	4533 (48.1%)
Marriage status, married	991 (48.5%)	650 (61.4%)	625 (65.7%)	5498 (58.4%)
Cohabitation	1512 (74.0%)	828 (78.3%)	770 (80.9%)	7532 (80.0%)
Equivalent income (million JPY)				
40-249	490 (24.0%)	115 (10.9%)	95 (10.0%)	1652 (17.5%)
250-375	433 (21.2%)	214 (20.2%)	176 (18.5%)	2593 (27.5%)
376-499	440 (21.5%)	274 (25.9%)	241 (25.3%)	2456 (26.1%)
≥500	679 (33.3%)	455 (43.0%)	440 (46.2%)	2715 (28.8%)
Education				
Junior high school	20 (1.0%)	8 (0.8%)	3 (0.3%)	58 (0.6%)
High school	346 (16.9%)	111 (10.5%)	129 (13.6%)	2343 (24.9%)
University, graduate school, vocational school, junior college	1676 (82.1%)	939 (88.8%)	820 (86.1%)	7015 (74.5%)
Current smoker	500 (24.5%)	267 (25.2%)	261 (27.4%)	2206 (23.4%)
Alcohol consumption				
6–7 days a week	458 (22.4%)	251 (23.7%)	230 (24.2%)	1967 (20.9%)
4–5 days a week	167 (8.2%)	123 (11.6%)	94 (9.9%)	691 (7.3%)
2–3 days a week	252 (12.3%)	139 (13.1%)	148 (15.5%)	1177 (12.5%)
Less than 1 day a week	353 (17.3%)	192 (18.1%)	190 (20.0%)	1603 (17.0%)
Hardly ever	812 (39.8%)	353 (33.4%)	290 (30.5%)	3978 (42.2%)
Physcial activity more than 30 minutes per day				
6–7 days a week	158 (7.7%)	58 (5.5%)	73 (7.7%)	402 (4.3%)
4–5 days a week	138 (6.8%)	86 (8.1%)	60 (6.3%)	474 (5.0%)
2–3 days a week	263 (12.9%)	165 (15.6%)	158 (16.6%)	1033 (11.0%)
Less than 1 day a week	306 (15.0%)	186 (17.6%)	161 (16.9%)	1270 (13.5%)
Hardly ever	1177 (57.6%)	563 (53.2%)	500 (52.5%)	6237 (66.2%)
Miss breakfast less than 3 days per week	586 (28.7%)	264 (25.0%)	230 (24.2%)	2405 (25.5%)
Eat all meals alone at least 4 days a week	755 (37.0%)	327 (30.9%)	262 (27.5%)	2437 (25.9%)
Consume less than two meals per day	52 (2.5%)	17 (1.6%)	13 (1.4%)	90 (1.0%)
Eat only individual food items or packaged food more than twice a day	171 (8.4%)	102 (9.6%)	70 (7.4%)	618 (6.6%)

Table 2. Odds ratios for unhealthdy dietary habit associated with frequeny of telecommuting

Outcome variable	Hardly ever n=9416	≤1 day/week n=952			2-3 days/week n=1058			≥4 days/week n=2042			p for trend
		OR	95%CI	p	OR	95%CI	p	OR	95%CI	p	
Miss breakfast more than 3 days per week											
model 1[a]	reference	0.89	0.76-1.04	0.142	0.90	0.77-1.05	0.183	1.13	1.02-1.27	0.026	0.248
model 2[b]	reference	0.95	0.81-1.12	0.529	0.97	0.83-1.13	0.686	1.15	1.03-1.29	0.013	0.068
Eat all meals alone at least 4 days a week											
model 1[a]	reference	1.03	0.88-1.20	0.743	1.09	0.95-1.26	0.220	1.52	1.37-1.68	<0.001	<0.001
model 2[b]	reference	1.07	0.90-1.27	0.459	1.16	0.98-1.36	0.081	1.44	1.28-1.63	<0.001	<0.001
Consume less than two meals per day											
model 1[a]	reference	1.48	0.82-2.68	0.189	1.75	1.03-2.97	0.040	2.89	2.03-4.12	<0.001	<0.001
model 2[b]	reference	1.35	0.74-2.45	0.332	1.52	0.88-2.63	0.130	2.39	1.66-3.44	<0.001	<0.001
Eat only individual food items or packaged food more than twice a day											
model 1[a]	reference	1.14	0.88-1.47	0.330	1.56	1.24-1.95	<0.001	1.37	1.14-1.64	0.001	<0.001
model 2[b]	reference	1.10	0.85-1.44	0.466	1.46	1.16-1.84	0.001	1.26	1.04-1.51	0.015	0.001

[a] Model 1 was adjusted for age and sex.

[b] Model 2 was adjusted for age, sex, education, annual equivalent household income, cohabitation, smoking status, alcohol consumption, and physical activity.