Title: Chemosensory dysfunctions induced by COVID-19 can persist

up to 7 months: A study of over 700 healthcare workers

Authors: Nicholas **Bussière**[1]; Jie **Mei**[1], PhD; Cindy **Lévesque-Boissonneault**[1], MSc; Mathieu

Blais[2,3], PhD; Sara **Carazo**[4], MD, PhD; Francois **Gros-Louis**[2,3], PhD; Gaston **De Serres**[4], MD,

PhD; Nicolas **Dupré**[5,6], MD, MSc, FRCP FAAN; Johannes **Frasnelli**[1], MD

[1] Department of Anatomy, Université du Québec à Trois-Rivières, Trois-Rivières, QC, Canada

[2] Axe Médecine régénératrice, CHU de Québec–Université Laval, Québec City, QC, Canada

[3] Faculty of Medicine, Department of Surgery, Université Laval, Québec City, QC, Canada

[4] Institut National de Santé Publique du Québec, Québec City, QC, Canada

[5] Faculty of Medicine, Department of Medicine, Université Laval, Québec City, QC, Canada

[6] Axe Neurosciences, CHU de Québec–Université Laval, Québec City, QC, Canada

Correspondence to be sent to :

Johannes Frasnelli MD,

Université du Québec à Trois-Rivières,

3351, boul. des Forges, C.P. 500,

Trois-Rivières, QC G9A 5H7 Canada

Email: johannes.a.frasnelli@uqtr.ca

Abstract:

Several studies have revealed either self-reported chemosensory alterations in large groups or objective quantified chemosensory impairments in smaller populations of patients diagnosed with COVID-19. However, due to the great variability in published results regarding COVID-19-induced chemosensory impairments and their follow-up, prognosis for chemosensory functions in patients with such complaints remains unclear. Our objective is to describe the various chemosensory alterations associated with COVID-19 and their prevalence and evolution after infection. A cross-sectional study of 704 healthcare workers with a RT-PCR confirmed SARS-CoV-2 infection between 28/2/2020 and 14/6/2020 was conducted 3 to 7 months after onset of symptoms. Data were collected with an online questionnaire. Outcomes included differences in reported chemosensory self-assessment of olfactory, gustatory, and trigeminal functions across time points and Chemosensory Perception Test scores from an easy-to-use at-home self-administered chemosensory test. Among the 704 participants, 593 (84.2%) were women, the mean (SD) age was 42 (12) years, and the questionnaire was answered on average 4.8 (0.8) months after COVID-19. During COVID-19, a decrease in olfactory, gustatory, and trigeminal sensitivities were reported by 81.3%, 81.5% and 48.0% respectively. Three to seven months later, reduced sensitivity was still reported by 52.0%, 41.9% and 23.3% respectively. Chemosensory Perception Test scores indicate that 19.5% of participants had objective olfactory impairment. These data suggest a significant proportion of COVID-19 cases have persistent chemosensory impairments at 3 to 7 months after their infection but the majority of those who had completely lost their olfactory, gustatory, and trigeminal sensitivity have improved.

Keywords: COVID-19, anosmia, parosmia, long-term, taste, trigeminal system

Introduction:

Coronavirus disease-2019 (COVID-19) is an ongoing major public health challenge. Olfactory dysfunction (OD) is a specific symptom that may affect approximately 60% of patients suffering from COVID-19 (Spinato, Fabbris et al. 2020, von Bartheld, Hagen et al. 2020, Whitcroft and Hummel 2020), and is now considered as a stronger indicator of COVID-19 than fever, cough and shortness of breath (Gerkin, Ohla et al. 2021).

OD can be quantitative or qualitative. Quantitative OD is defined by a reduction of olfactory sensitivity which can be either a complete (anosmia) or a partial (hyposmia) loss of olfactory function (Hummel, Whitcroft et al. 2016). Qualitative OD describes an altered perception of olfactory stimuli: For example, parosmia is defined as the perception of qualitatively altered smells, and phantosmia is defined as the perception of a smell in the absence of an objective odorant (Hummel, Whitcroft et al. 2016, Sjölund, Larsson et al. 2017). Overall, the prevalence of OD in the general population is around 20% (Landis, Konnerth et al. 2004, Yang and Pinto 2016), and all different forms of OD are associated with reduced quality of life (Croy, Nordin et al. 2014). In addition to OD, COVID-19 also appears to affect other chemosensory modalities, i.e., gustation and trigeminal function (Cooper, Brann et al. 2020, Parma, Ohla et al. 2020).

Olfactory and other chemosensory dysfunctions may have detrimental effects. First, affected individuals can expose themselves to harmful substances such as smoke, gas or spoiled food (Gonzales and Cook 2007, Schiffman 2007). It may trigger dysfunctional nutritional patterns like increased salt and sugar consumption, or anorexia (Mattes, Cowart et al. 1990, Aschenbrenner, Hummel et al. 2008). Individuals with OD also have higher rates of anxiety and depression (Croy, Nordin et al. 2014, Kohli, Soler et al. 2016). Moreover, a functioning olfactory

system may be a necessity in some workplaces, such as healthcare, where staff are required to

have the ability to detect and qualify the smell of urine, excrement, infected wounds or abnormal

smells of breath (Kelly 2012).

Investigation of the long-term effects of COVID-19 on chemosensory function is

hindered by the recent onset of the pandemic and other challenges: First, many studies on the

prevalence of OD during COVID include a relatively small number of participants (Hintschich,

Wenzel et al. 2020, Le Bon, Pisarski et al. 2020) or participants with severe forms of COVID-19

(Moein, Hashemian et al. 2020, Speth, Singer-Cornelius et al. 2020). Secondly, many studies on

the prevalence of OD during COVID-19 also include participants with an unclear diagnosis of

COVID-19, and/or self-diagnosis (Hopkins, Surda et al. 2020, Parma, Ohla et al. 2020). Lastly,

while individuals with anosmia can usually evaluate their olfactory function with accuracy

(Lötsch and Hummel 2019), this self-assessment is often challenging for individuals with

intermediate forms of OD (e.g., hyposmia) (Landis, Hummel et al. 2003). Finally, studies on

persistent post-COVID-19 OD in the past year have used various designs (objective measures

(Lechien, Chiesa-Estomba et al. 2021), semi-objective (Petrocelli, Cutrupi et al. 2021), or self-

reported (Havervall, Rosell et al. 2021, Hopkins, Surda et al. 2021) and collected data at varying

time intervals after onset of disease. For these reasons, to this date, no consensus has been

reached regarding the prevalence of post-COVID-19 OD (Xydakis, Albers et al.).

To comprehensively understand long-term olfactory, gustatory, and trigeminal alterations

after COVID-19, we analyzed questionnaire responses from a cohort of healthcare workers

infected with SARS-CoV-2 during the first wave of the pandemic (February - June 2020). We

also developed a Chemosensory Perception Test (CPT), a formal test employing common

household odorants and tastants to enable accessible yet accurate self-evaluation of

chemosensory functions remotely on a large scale. The CPT is particularly useful when in-person

testing is unsafe and testing a large group of participants at distance with mailable tests such as

the UPSIT (Doty, Shaman et al. 1984) is costly. Moreover, distance testing has been reported to

accurately monitor disease progression in at risk populations (Vaira, Hopkins et al. 2020, Weiss,

Attuquayefio et al. 2020).

Materials and Methods:

Participants

Participants were recruited from a Quebec healthcare worker cohort who have had SARS-CoV-2

infection between 28/2/2020 and 14/6/2020. They were part of a study from the Institut National

de Santé Publique du Québec and had agreed to be contacted for other research projects(Carazo

2021). Inclusion criteria were (1) RT-PCR confirmed COVID-19 (2) above 18 years of age, (3)

French or English speakers, (4) completed the online questionnaire, and (5) did not report of

other respiratory diseases (bacterial or viral infection, or/and allergies with rhinorrhea) within 2

weeks prior to questionnaire completion or chronic sinusitis (Figure 1).

This study was reviewed and approved by the research ethics board of the CHU

de Québec – Université Laval (MP-20-2021-5228) and all protocols were reviewed by an

independent Scientific Review Committee. This study also complies with the Declaration of

Helsinki for Medical Research Involving Human Subjects. All participants provided an online

informed consent prior to participation. The study received funding from the Fonds de recherche

du Québec-Santé. No compensation or incentive was offered for participation. Data were

collected from August 11 to October 29, 2020. Up to four attempts were made to reach by email

potential participants. At the time of data collection, participants were 3-7 months after the onset

of COVID-19 symptoms.

Online questionnaire:

All participants were asked to complete an online questionnaire which was adapted from the core

questionnaire of the Global Consortium on Chemosensory Research(Parma, Ohla et al. 2020).

<u>Demographic information</u>: In the first part of the questionnaire, demographic information

was collected from all participants. Participants were then instructed to provide medical history

and indicate the presence of specific COVID-19 symptoms (Figure 2).

<u>Chemosensory self-assessment</u>: Participants were asked to self-evaluate and report their

olfactory , gustatory, and trigeminal sensitivity using a 10-point visual analog scale (VAS;

Figure 2) for three timepoints: (1) before SARS-CoV-2 infection, (2) during SARS-CoV-2

infection and (3) at questionnaire completion. The specific definition of each chemosensory

modality was presented prior to self-evaluation of each chemosensory modality as follows:

Olfaction: *The following questions relate to your sense of smell (for example, sniffing flowers or*

soap, or smelling garbage) but not the flavor of food in your mouth; Gustation: *The following*

questions are related to your sense of taste. For example, sweetness, sourness, saltiness,

bitterness experienced in the mouth; Trigeminal: *The following questions are related to other*

sensations in your mouth, like burning, cooling, or tingling. For example, chili peppers, mint

gum or candy, or carbonation. Further, information on the presence of parosmia or phantosmia

following the infection (Landis, Frasnelli et al. 2010) and alterations in the 5 tastes (sweet, salty,

sour, bitter, umami) was collected.

<u>Chemosensory Perception Test (CPT)</u>: Items commonly found in North American

households were used to assess participants' olfactory and gustatory functions, as odor intensity

is the best single predictor to classify individuals with normosmia (Parma, Hannum et al. 2021).

Participants had to smell three substances (peanut butter, jam/jelly, and coffee) and rate odor

intensity on a 10-point VAS (0: no smell at all; 10: very strong smell). We obtained olfactory

scores by averaging these ratings. Pilot data on a total of 93 participants show these scores to

accurately detect OD when compared to the Sniffin' Sticks (cut-off score: 6/10; sensitivity:

0.765; specificity: 0.895; Supplement 3). Participants were asked to prepare saline and sweet

water by dissolving respectively a teaspoon of salt or 3 teaspoons of sugar in a cup (250 mL) of

lukewarm water. Then, they were asked to taste saline and sweet water and to rate taste

intensities on a 10-point VAS. We obtained gustatory scores by averaging these ratings. An

ongoing study is comparing CPT gustatory scores with the Waterless-Empirical Taste Test -

Self-Administered (Doty, Wylie et al. 2021), but too few participants have been recruited to this

to establish its accuracy (Supplement 3).

Statistical Analyses:

A Python script (Python 3.7.5, Python Software Foundation, https://www.python.org) was used

to process raw questionnaire data and to calculate the number of participants reporting COVID-

19 symptoms, chronic conditions and recent respiratory illnesses. Processed data were analyzed

and visualized with SPSS 26.0 (Armonk, NY: IBM Corp), GraphPad Prism 8.3.1 (GraphPad

Prism Software, San Diego, CA) and Raincloud plots(Allen, Poggiali et al. 2021).

Parametric (ANOVA) or non-parametric (Friedman) tests were chosen depending on

whether normality assumption was fulfilled. To evaluate the effects of COVID-19 on *modality*

(olfactory, gustatory, and trigeminal) and *time* (prior to, during and after COVID-19 infection),

for *gender* (women, men), repeated measures (rm) ANOVA with age as a covariate were

computed. To disentangle interactions, separate rmANOVA were carried out for individual

modalities and timepoints with the same factors. Greenhouse-Geisser corrections were used for

sphericity and Tukey's multiple comparisons test were used for post-hoc comparisons.

Friedman's test was followed by Dunn's post-hoc test to correct for multiple comparisons. To

assess the correlation between self-reported olfactory, gustatory, and trigeminal abilities and

results of the CPT, Pearson correlation coefficient or Spearman's rank correlation coefficient

was used. For all statistical tests, alpha was set at 0.05. All results are expressed as mean (SD)

unless otherwise specified.

Results:

Characteristics of participants:

A total of 704 healthcare workers (593 (84.2%) women, mean age of 42.0 (SD:11.7, range 18 –

70) years were included. The questionnaire was completed on average 4.8 (SD: 0.8, range 3-7)

months after symptoms onset. COVID-19 symptoms reported by the 704 participants are listed in

Table 1.

Quantitative disorders:

Before COVID-19, average self-reported score was 9.0 (1.6), 9.2 (1.3) and 8.9 (1.9) of 10 for

olfaction, gustation and trigeminal function, respectively. Among participants, 0.9%, 0.7% and

1.8% respectively reported an absence of olfaction, gustation and trigeminal function (score 0;

Figure 3). During COVID-19, average self-reported score was 2.6 (3.6) for olfaction, 3.4 (3.6)

for gustation, and 7.0 (3.0) for trigeminal sensitivity. In the 704 participants, 51.1%, 33.5% and

5.7% reported absence of olfaction, gustation and trigeminal function. At time of questionnaire

completion, mean scores were 7.4 (2.5), 8.0 (2.2) and 8.5 (2.2) for olfaction, gustation and

trigeminal function respectively and absence of chemical senses was reported respectively by

1.4%, 0.7% and 2.3%. Weak correlations were found between the time since infection and the

self-reported olfactory and gustatory scores at questionnaire completion (olfaction: ρ=0.11;

gustation: ρ=0.14; both P<.001; trigeminal ρ = .06; P=.11).

Compared to the baseline chemosensory functions before COVID-19, 572 (81.3%), 574

(81.5) and 338 (48.0%) reported lower olfactory, gustatory, and trigeminal sensitivity during

COVID-19. Olfactory and gustatory dysfunction were present in similar proportions (χ^2(2,

N=704) =0.02, P=.891) and were different to trigeminal (olfaction: χ^2(2, N=704) = 174.81

P<.001; gustation: χ^2(2, N=704) = 174.56 P<.001). Three to seven months after the infection,

366 (52.0%), 295 (41.9%), 164 (23.3%) reported lower olfactory, gustatory, and trigeminal

sensitivity compared to before COVID-19 (Table 2). These proportions were significantly

different between all three chemosensory systems (χ^2(2, N=704) = 123.46, P<.001).

Overall, there were significant effects of *modality* (F(2,1402)=42.83, P<.001, η_p^2=0.058;

olfactory<gustatory<trigeminal; all P<.001), *time* (F(2,1402)=118.47, P<.001, η_p^2=0.145;

during<after<before; all P<.001), and *gender* (F(1,701)=5.52, P=0.019, η_p^2=0.008; women<men)

and significant interactions between these factors (*modality*time, modality*time*gender*; all

P<.001) on chemosensory self-evaluation. To disentangle these interactions, we analyzed data

separately per chemosensory modality and time points.

<u>Chemosensory modality</u>: With regards to <u>olfactory</u> function, significant main effects of *time*

(F(2,1402)=165.07, P<.001 η_p^2=0.191; during<after<before; all P<.001; Figure 3A), *age*

(F(1,701)=4.42, P=.012, η_p^2=0.009) and *gender* (F(1,701)=4.42, P=.036, η_p^2=0.006; women <

men) were revealed. In addition, we observed significant interactions of *time*age*

(F(2,1402)=23.39, P<.001, η_p^2=0.032) and *time*gender* (F(2, 1402)=21.69 , P<0.001, η_p^2=0.030).

With regards to <u>gustatory</u> function, we observed significant main effects of *time*

(F(2,1402)=102.97, P<.001, η_p^2=0.128; during<after<before; all P<.001; Figure 3B) and *gender*

(F(1, 701)=9.80, P=.002, η_p^2=0.014; women<men), but no effect of *age*. We also observed

significant interactions of *time*age* (F(2, 1402))=5.97, P=.005, η_p^2=0.008) and *time*gender* (F(2,

1402))=20.02 , P<.001, η_p^2=0.028).

With regards to <u>trigeminal</u> function, we observed significant main effects of *time*

(F(2,1402)=3.91, *P*=.020, η_p^2=0.006; during<after<before; all *P*<.001; Figure 3C), and age

((1,701)=5.08, *P*=.025, η_p^2=0.007) but no effect of *gender*. We also identified significant

interactions of *time*age* (F(2, 1402)=4.70, *P*=.016, η_p^2=0.007) and *time*gender* (F(2,

1402)=4.50, *P*=.019, η_p^2=0.006).

<u>Time point</u>: With regards to chemosensory function <u>before</u> infection, we observed a significant

effect of *gender* (F(1,701)=8.52, *P*=.004, η_p^2=0.012; men < women), but not of *modality*, *age* nor

interactions. <u>During</u> COVID-19, we observed a significant effects of *modality* (F(2,

1402)=96.714, *P*<.001, η_p^2=0.121; olfaction<gustation<trigeminal; all *P*<.001), *gender* (F(1,

701)=21.98, *P*<.001, η_p^2=0.030; women<men), and age (F(1, 701)=4.74, *P*=.030, η_p^2=0.007).

Further, we found significant interactions *modality*age* (F(2, 1402)=24.185, *P*<.001, η_p^2=0.033)

and *modality*gender* (F(2, 1402)=6.76, *P*=.002, η_p^2=0.010). Finally, <u>after</u> infection, we observed

a significant effect of *modality* (F(2, 1402)=9.91, *P*<.001, η_p^2=0.014;

olfaction<gustation<trigeminal; all *P*<.015), but not of *gender* or *age*, nor any interaction.

Compared to baseline (before infection), changes in chemosensory function were

correlated for all modalities during infection (olfaction-gustation: ρ=0.69; gustation-trigeminal:

ρ=0.43; olfaction-trigeminal: ρ=0.33; all *P*<.001, Figure 4 A-C) and after infection (olfaction-

gustation: ρ=0.69; gustation-trigeminal: ρ=0.40; olfaction-trigeminal: ρ=0.36; all *P*<.001, Figure

4 D-F).

Qualitative disorders:

Among included participants, 78 (11.1%) reported parosmia, 73 (10.4%) experienced

phantosmia and/or 82 (11.6%) had waxing and waning of olfaction following infection. In

addition, 42 (6.0%) claimed that they experienced other forms of OD (hyposmia to specific

substances, hyperosmia, parosmia only at high concentrations or slow identification times).

Furthermore, 335 (47.6%) participants reported changes to perception of sweet, 338 (48.0%)

salty, 293 (41.6%) sour, 309 (43.9%) bitter and 281 (39.9%) umami. A total of 275 (39.1%)

participants reported alterations in all 5 tastes.

Chemosensory Perception Test:

Among the 704 participants, 137 (19.5%) had a CPT score suggestive of OD. Mean CPT scores

were lower for olfaction than gustation (7.84 (1.78) vs 8.42 (2.31); $Z=8.193$, $P<.001$). Neither

age nor *gender* had an effect on CPT scores. CPT scores correlated with self-reported

chemosensory abilities at testing time (olfaction: $\rho=0.67$; gustation: $\rho=0.51$; $P<.001$ for both).

Discussion:

This study reports chemosensory dysfunction 3 to 7 months following SARS-CoV-2 infection in a large cohort of RT-PCR-confirmed healthcare workers. In addition to confirming the now well-established detrimental effect of acute COVID-19 on all three chemosensory systems (olfactory, gustatory, trigeminal), our major findings are: (1) the detrimental effect of COVID-19 lasts beyond the acute phase after the infection, half of those affected indicated that olfactory function had not returned to the baseline levels 3 to 7 months later, while 20% of infected participants reported scores in a formal test that are consistent with the presentation of hyposmia/anosmia; (2) approximately 10% of the patients exhibit parosmia and/or phantosmia; (3) women are more heavily affected than men.

We observed chemosensory dysfunction in the acute phase of COVID-19, which was most pronounced for olfactory function, but less so for gustatory function and even less for trigeminal function. The proportion of participants describing OD and GD in the acute phase of COVID-19 in this study was comparable to earlier studies (Hajikhani, Calcagno et al. 2020). Although the proportions of participants indicating a decrease in olfaction or gustation were comparable, the olfactory system seems to be more severely impaired. Given the cross-sectional design of the present study, recall bias may have a role to play in the prevalence of OD and GD in similar study populations, but published studies with little to no recall bias also report equivalent prevalence of OD and GD (Andrews, Pendolino et al. 2020, Lechien, Chiesa-Estomba et al. 2020, Petrocelli, Cutrupi et al. 2021). Longitudinal studies are needed to further assess the relationship between OD and GD in COVID-19. Nevertheless, on average 4.8 months after infection and thus well after the acute phase, approximately 50% and 40% of patients reported persistent alterations in olfactory and gustatory function, respectively; these numbers are higher

than what has been reported in some studies (Boscolo-Rizzo, Guida et al. 2021, Capelli and Gatti

2021, Lechien, Chiesa-Estomba et al. 2021) and lower than reported by others (Hopkins, Surda

et al. 2021). The great variability in these results is due to very different study designs (self-

report vs. psychophysical test; prospective vs cross-sectional) and studied populations (of

different ethnicity and under different effects of selection bias), which either influence the

measure of OD and GD in study populations or directly impact the baseline prevalence of OD

and GD during COVID-19, offsetting all prevalence calculated at further points (Mazzatenta,

Neri et al. 2020, von Bartheld, Hagen et al. 2020). For instance, in the study population included

in this study, prevalence of OD decreases to 18.9% of participants when measured using the CPT

at 4.8 (SD: 0.8) months after infection. The difference in these frequencies could be due to a

higher sensitivity of the self-reported alterations compared to the CPT. Participants with milder

forms of persistent hyposmia or with higher baseline olfactory sensitivity may have higher scores

on the semi-objective CPT yet have not recovered entirely. We found a moderate-to-strong

correlation between self-reported olfactory and gustatory changes, which were stronger than with

self-reported trigeminal changes. This could be due to similar pathophysiological alterations in

the olfactory and gustatory systems and their differences from that of the trigeminal system.

Knowing that the general population often mixes up retro-olfaction (perceiving odors

from the substances in the mouth traveling posteriorly and rostrally to the olfactory epithelium)

with taste, an alternative explanation would be a misunderstanding of this nuance by participants

despite the fact that specific definitions for each modality were given (Landis, Frasnelli et al.

2005)(Malaty and Malaty 2013). The latter hypothesis is more probable since the correlation

between gustatory self-report and CPT gustatory scores using strict gustatory stimuli (salt, sugar)

is lower than the correlation between olfactory self-report and CPT olfactory scores. When

tasting strictly gustatory stimuli in the CPT, participants reflect solely on their sense of taste,

without the influence of retronasal sensations. These tests have the potential to be more accurate

than simple subjective measures and could simplify large-scale psychophysical chemosensory

testing. Others have reported the usefulness of similar self-administered chemosensory tests in

the detection and follow-up of COVID-19-induced chemosensory dysfunctions *(Vaira, Salzano*

et al. 2020, Petrocelli, Cutrupi et al. 2021). Different theories have been proposed to explain the

persistence of OD in certain individuals, ranging from olfactory epithelium dysfunction to

central nervous system infection (Bilinska and Butowt 2020, Butowt and von Bartheld 2020,

Solomon 2021). Since cells of the olfactory epithelium possess the ability to regenerate, the re-

establishment of olfactory function is possible in the context of postinfectious OD (Cavazzana,

Larsson et al. 2018), as well as in COVID-19-related OD, where 75-85% of the affected

individuals recovered olfactory function within 60 days (Mullol, Alobid et al. 2020, Lechien,

Chiesa-Estomba et al. 2021). The exact rate of olfactory recovery is still unknown, while post-

COVID-19 OD prevalence ranging from 11%-60% at 6 months according to a recent study

(Xydakis, Albers et al.). In addition to OD and GD, TD has also been reported in patients with

COVID-19 (Cooper, Brann et al. 2020, Parma, Ohla et al. 2020).

Persistent chemosensory dysfunctions may be a sign of chronic central nervous system

alterations (Gori, Leone et al. 2020, Wu, Xu et al. 2020), and there is now evidence that SARS-

CoV-2 can infect olfactory sensory neurons in humans (Meinhardt, Radke et al. 2020, de Melo,

Lazarini et al. 2021). Other viruses, such as the Japanese encephalitis virus, Varicella-Zoster

virus, measles virus, human immunodeficiency virus and CoVs, were shown to invade the CNS

(Koyuncu, Hogue et al. 2013). Febrile seizures, loss of consciousness, convulsions, ataxia, status

epilepticus, encephalitis, myelitis, neuritis and extrapyramidal symptoms are among extra-

pulmonary symptoms that have been described (Bohmwald, Gálvez et al. 2018). However, no

evidence of intra-parenchymal replication has been found yet. Additional findings include the

presence of local immune processes (Saussez, Sharma et al. 2021) and persistence of viral

fragments in the olfactory epithelium (de Melo, Lazarini et al. 2021). Therefore, chronic post-

COVID-19 inflammation in the olfactory pathway (epithelium, bulb) with or without direct

infection is the most probable pathophysiological explanation of post-COVID-19 OD

(Kirschenbaum, Imbach et al. 2020, Vaira, Hopkins et al. 2020, Xydakis, Albers et al.). The

persistence of postinfectious neurological inflammation may contribute to the development or

aggravation of chronic neurological diseases such as Parkinson, multiple sclerosis, or psychiatric

outcomes (Morris 1985, Johnson-Lussenburg and Zheng 1987, Fazzini, Fleming et al. 1992,

Murray, Brown et al. 1992, Stewart, Mounir et al. 1992, Cristallo, Gambaro et al. 1997, Arbour,

Day et al. 2000, Koyuncu, Hogue et al. 2013, Cohen, Eichel et al. 2020, Taquet, Geddes et al.

2021). These patients should be followed up to document the development of neurological

sequalae.

Moreover, approximately 10% reported parosmia and/or phantosmia following SARS-

CoV-2 infection. These qualitative smell disorders usually involve unpleasant olfactory

sensations (rotten eggs, sewage, smoke). While the exact patho-mechanism of parosmia and

phantosmia are still to be elucidated, parosmia is probably linked to altered peripheral

input/central processing of olfactory stimuli (Iannilli, Leopold et al. 2019). Importantly, patients

with postviral OD and parosmia exhibit better recovery rates following olfactory training than

those without parosmia (Liu, Sabha et al. 2021). Follow-ups will determine to what extent

parosmia predicts a better outcome.

Women's chemical senses were more affected than men during and after COVID-19

infection. Women typically have better scores in olfactory testing than men at baseline (Wang,

Zhang et al. 2019). However, in line with our results, studies have revealed that women exhibit a

higher prevalence and a longer persistence of postviral OD (Liu, Pinto et al. 2016, Sorokowski,

Karwowski et al. 2019). Gender differences could be explained by a multitude of

neuroendocrine, social, and cognitive factors (Sorokowski, Karwowski et al. 2019). We also

found that older individuals have lower olfactory and gustatory sensitivities, especially during

the acute phase of COVID-19.

Currently, there is no approved therapy specifically for COVID-19-induced OD, although

experts agree that olfactory training could be prescribed for COVID-induced OD as it has a

significant effect on olfactory function according to studies on other viral infections (Damm,

Pikart et al. 2014, Sorokowska, Drechsler et al. 2017, Doty 2019, Huart, Philpott et al. 2021).

Additionally, oral steroids, intranasal steroids and/or omega-3 supplements may be prescribed on

an individual basis (Hopkins, Alanin et al. 2021). Most importantly, long-term follow-up of these

patients will be necessary to assess other signs of neurological damage or spontaneous recovery,

as recoveries can be possible after a year in other post-viral OD (Lee, Lee et al. 2014).

<u>Limitations</u>

Given the cross-sectional design of the study, a recall bias is possible for all self-reported

peri-SARS-CoV-2 infection values before or during the SARS-CoV-2 infection due to the 3-to-

7-month gap. This study did not control for potential confounding factors like race and level of

education. Finally, the CPT requires further validation for its gustatory and trigeminal

components, and it relies on substances found in participants' homes, which may lead to

variation in test results due to the differences in the brand, quality, or expiration date of

substances and consequently, their ability to trigger equal sensorineural responses.

Conclusions:

Nearly two thirds of SARS-CoV-2 infected patients had chemosensory impairments during their

infection and despite improvements, impairments persist in half of them 3 to 7 months after

COVID-19. Quantitative and qualitative olfactory dysfunction as well as persisting gustatory and

trigeminal deficits were common in the cohort presented in this study. Given the frequency of

these problems and the possible neurological underpinnings of these observations, it will be

critical to understand the underlying mechanisms of these chemosensory dysfunctions, their

evolution, and possible therapeutic options.

Acknowledgments:

We thank Josiane Rivard for preparing the online questionnaire, Cécilia Tremblay, Émilie

Aubry-Lafontaine and Frédérique Roy-Côté for data collection and the validation of the

Chemosensory Perception Test, and all study participants and frontline healthcare workers facing

the COVID-19 pandemic. This work was supported by Fonds de recherche du Québec – Santé

(chercheur boursier junior 2 #283144 to JF). NB and JF had full access to all the data in the study

and take responsibility for the integrity of the data and the accuracy of the data analysis. FGL is

the recipient of a tier-2 Canada research Chair. All authors declare no conflict of interest.

References:

Allen, M., D. Poggiali, K. Whitaker, T. Marshall, J. van Langen and R. Kievit (2021). "Raincloud plots: a multi-platform tool for robust data visualization [version 2; peer review: 2 approved]." Wellcome Open Research **4**(63).

Andrews, P. J., A. L. Pendolino, G. Ottaviano, B. Scarpa, J. Grant, P. Gaudioso, A. Bordin, R. Marchese-Ragona, D. Leoni, A. Cattelan, A. Kaura, S. Gane, N. J. Hamilton, D. Choi and J. A. Andrews (2020). "Olfactory and taste dysfunction among mild-to-moderate symptomatic COVID-19 positive health care workers: An international survey." Laryngoscope investigative otolaryngology **5**(6): 1019-1028.

Arbour, N., R. Day, J. Newcombe and P. J. Talbot (2000). "Neuroinvasion by human respiratory coronaviruses." J Virol **74**(19): 8913-8921.

Aschenbrenner, K., C. Hummel, K. Teszmer, F. Krone, T. Ishimaru, H. S. Seo and T. Hummel (2008). "The influence of olfactory loss on dietary behaviors." Laryngoscope **118**(1): 135-144.

Bilinska, K. and R. Butowt (2020). "Anosmia in COVID-19: A Bumpy Road to Establishing a Cellular Mechanism." ACS Chem Neurosci.

Bohmwald, K., N. M. S. Gálvez, M. Ríos and A. M. Kalergis (2018). "Neurologic Alterations Due to Respiratory Virus Infections." Front Cell Neurosci **12**: 386.

Boscolo-Rizzo, P., F. Guida, J. Polesel, A. V. Marcuzzo, P. Antonucci, V. Capriotti, E. Sacchet, F. Cragnolini, A. D'Alessandro, E. Zanelli, R. Marzolino, C. Lazzarin, M. Tofanelli, N. Gardenal, D. Borsetto, C. Hopkins, L. A. Vaira and G. Tirelli (2021). "Self-reported smell and taste recovery in coronavirus disease 2019 patients: a one-year prospective study." European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery: 1-6.

Butowt, R. and C. S. von Bartheld (2020). "Anosmia in COVID-19: Underlying Mechanisms and Assessment of an Olfactory Route to Brain Infection." Neuroscientist.

Capelli, M. and P. Gatti (2021). "Anosmia in the first coronavirus disease 2019 outbreak in Europe: functional recovery after eight months." The Journal of laryngology and otology: 1-5.

Carazo, S. L. D., Villeneuve J, Martin R, Deshaies P, Denis G, , Adib G, Tissot F, Dionne M, De Serres G (2021). "Characterization and evolution of infection control practices among SARS-CoV-2 infected healthcare workers of acute care hospitals and long-term care facilities in Quebec, Canada, Spring 2020." Infect Contr Hospit Epidemiol.

Cavazzana, A., M. Larsson, M. Münch, A. Hähner and T. Hummel (2018). "Postinfectious olfactory loss: A retrospective study on 791 patients." Laryngoscope **128**(1): 10-15.

Cohen, M. E., R. Eichel, B. Steiner-Birmanns, A. Janah, M. Ioshpa, R. Bar-Shalom, J. J. Paul, H. Gaber, V. Skrahina, N. M. Bornstein and G. Yahalom (2020). "A case of probable Parkinson's disease after SARS-CoV-2 infection." Lancet Neurol **19**(10): 804-805.

Cooper, K. W., D. H. Brann, M. C. Farruggia, S. Bhutani, R. Pellegrino, T. Tsukahara, C. Weinreb, P. V. Joseph, E. D. Larson, V. Parma, M. W. Albers, L. A. Barlow, S. R. Datta and A. Di Pizio (2020). "COVID-19 and the Chemical Senses: Supporting Players Take Center Stage." Neuron **107**(2): 219-233.

Cristallo, A., F. Gambaro, G. Biamonti, P. Ferrante, M. Battaglia and P. M. Cereda (1997). "Human coronavirus polyadenylated RNA sequences in cerebrospinal fluid from multiple sclerosis patients." New Microbiol **20**(2): 105-114.

Croy, I., S. Nordin and T. Hummel (2014). "Olfactory Disorders and Quality of Life—An Updated Review." Chemical Senses **39**(3): 185-194.

Damm, M., L. K. Pikart, H. Reimann, S. Burkert, Ö. Göktas, B. Haxel, S. Frey, I. Charalampakis, A. Beule, B. Renner, T. Hummel and K. B. Hüttenbrink (2014). "Olfactory training is helpful in postinfectious olfactory loss: a randomized, controlled, multicenter study." Laryngoscope **124**(4): 826-831.

de Melo, G. D., F. Lazarini, S. Levallois, C. Hautefort, V. Michel, F. Larrous, B. Verillaud, C. Aparicio, S. Wagner, G. Gheusi, L. Kergoat, E. Kornobis, F. Donati, T. Cokelaer, R. Hervochon, Y. Madec, E. Roze, D. Salmon, H. Bourhy, M. Lecuit and P.-M. Lledo (2021). "COVID-19–related anosmia is associated with viral persistence and inflammation in human olfactory epithelium and brain infection in hamsters." <u>Science Translational Medicine</u> **13**(596): eabf8396.

Doty, R. L. (2019). "Treatments for smell and taste disorders: A critical review." <u>Handb Clin Neurol</u> **164**: 455-479.

Doty, R. L., P. Shaman, C. P. Kimmelman and M. S. Dann (1984). "University of Pennsylvania Smell Identification Test: a rapid quantitative olfactory function test for the clinic." <u>Laryngoscope</u> **94**(2 Pt 1): 176-178.

Doty, R. L., C. Wylie and M. Potter (2021). "Validation of the Waterless Empirical Taste Test (WETT®)." <u>Behavior research methods</u> **53**(2): 864-873.

Fazzini, E., J. Fleming and S. Fahn (1992). "Cerebrospinal fluid antibodies to coronavirus in patients with Parkinson's disease." <u>Mov Disord</u> **7**(2): 153-158.

Gerkin, R., K. Ohla, M. Veldhuizen, P. Joseph, C. Kelly, A. Bakke, K. Steele, M. Farruggia, R. Pellegrino, M. Pepino, C. Bouysset, G. Soler, V. Pereda-Loth, M. Dibattista, K. Cooper, I. Croijmans, A. Di Pizio, M. Ozdener, A. Fjaeldstad and V. Parma (2021). "Recent Smell Loss Is the Best Predictor of COVID-19 Among Individuals With Recent Respiratory Symptoms." <u>Chemical Senses</u> **46**.

Gonzales, G. M. and M. J. Cook (2007). CHAPTER 13 - DISORDERS OF SMELL AND TASTE. <u>Neurology and Clinical Neuroscience</u>. A. H. V. Schapira, E. Byrne, S. DiMauro et al. Philadelphia, Mosby**:** 171-177.

Gori, A., F. Leone, L. Loffredo, B. L. Cinicola, G. Brindisi, G. De Castro, A. Spalice, M. Duse and A. M. Zicari (2020). "COVID-19-Related Anosmia: The Olfactory Pathway Hypothesis and Early Intervention." <u>Frontiers in neurology</u> **11**: 956-956.

Hajikhani, B., T. Calcagno, M. J. Nasiri, P. Jamshidi, M. Dadashi, M. Goudarzi, A. A. Eshraghi and M. Mirsaeidi (2020). "Olfactory and gustatory dysfunction in COVID-19 patients: A meta-analysis study." <u>Physiol Rep</u> **8**(18): e14578.

Havervall, S., A. Rosell, M. Phillipson, S. M. Mangsbo, P. Nilsson, S. Hober and C. Thålin (2021). "Symptoms and Functional Impairment Assessed 8 Months After Mild COVID-19 Among Health Care Workers." <u>JAMA</u> **325**(19): 2015-2016.

Hintschich, C. A., J. J. Wenzel, T. Hummel, M. K. Hankir, T. Kühnel, V. Vielsmeier and C. Bohr (2020). "Psychophysical tests reveal impaired olfaction but preserved gustation in COVID-19 patients." <u>International forum of allergy & rhinology</u> **10**(9): 1105-1107.

Hopkins, C., M. Alanin, C. Philpott, P. Harries, K. Whitcroft, A. Qureishi, S. Anari, Y. Ramakrishnan, A. Sama, E. Davies, B. Stew, S. Gane, S. Carrie, I. Hathorn, R. Bhalla, C. Kelly, N. Hill, D. Boak and B. Nirmal Kumar (2021). "Management of new onset loss of sense of smell during the COVID-19 pandemic - BRS Consensus Guidelines." <u>Clin Otolaryngol</u> **46**(1): 16-22.

Hopkins, C., P. Surda, L. A. Vaira, J. R. Lechien, M. Safarian, S. Saussez and N. Kumar (2021). "Six month follow-up of self-reported loss of smell during the COVID-19 pandemic." <u>Rhinology</u> **59**(1): 26-31.

Hopkins, C., P. Surda, E. Whitehead and B. N. Kumar (2020). "Early recovery following new onset anosmia during the COVID-19 pandemic – an observational cohort study." <u>Journal of Otolaryngology - Head & Neck Surgery</u> **49**(1): 26.

Huart, C., C. M. Philpott, A. Altundag, A. W. Fjaeldstad, J. Frasnelli, S. Gane, J. W. Hsieh, E. H. Holbrook, I. Konstantinidis, B. N. Landis, A. Macchi, C. A. Mueller, S. Negoias, J. M. Pinto, S. C. Poletti, V. R. Ramakrishnan, P. Rombaux, J. Vodicka, A. Welge-Lüessen, K. L. Whitcroft and T. Hummel (2021). "Systemic corticosteroids in coronavirus disease 2019 (COVID-19)-related smell dysfunction: an international view." <u>International forum of allergy & rhinology</u> **11**(7): 1041-1046.

Hummel, T., K. L. Whitcroft, P. Andrews, A. Altundag, C. Cinghi, R. M. Costanzo, M. Damm, J. Frasnelli, H. Gudziol, N. Gupta, A. Haehner, E. Holbrook, S. C. Hong, D. Hornung, K. B. Hüttenbrink, R. Kamel, M.

Kobayashi, I. Konstantinidis, B. N. Landis, D. A. Leopold, A. Macchi, T. Miwa, R. Moesges, J. Mullol, C. A. Mueller, G. Ottaviano, G. C. Passali, C. Philpott, J. M. Pinto, V. J. Ramakrishnan, P. Rombaux, Y. Roth, R. A. Schlosser, B. Shu, G. Soler, P. Stjärne, B. A. Stuck, J. Vodicka and A. Welge-Luessen (2016). "Position paper on olfactory dysfunction." Rhinology **56**(1): 1-30.

Iannilli, E., D. A. Leopold, D. E. Hornung and T. Hummel (2019). "Advances in Understanding Parosmia: An fMRI Study." ORL J Otorhinolaryngol Relat Spec **81**(4): 185-192.

Johnson-Lussenburg, C. M. and Q. Zheng (1987). "Coronavirus and multiple sclerosis: results of a case/control longitudinal serological study." Adv Exp Med Biol **218**: 421-429.

Kelly, M. (2012). "Scent of a patient: an underestimated role in clinical practice?" The British journal of general practice : the journal of the Royal College of General Practitioners **62**(600): 378-378.

Kirschenbaum, D., L. L. Imbach, S. Ulrich, E. J. Rushing, E. Keller, R. R. Reimann, K. B. M. Frauenknecht, M. Lichtblau, M. Witt, T. Hummel, P. Steiger, A. Aguzzi and K. Frontzek (2020). "Inflammatory olfactory neuropathy in two patients with COVID-19." The Lancet **396**(10245): 166.

Kohli, P., Z. M. Soler, S. A. Nguyen, J. S. Muus and R. J. Schlosser (2016). "The Association Between Olfaction and Depression: A Systematic Review." Chem Senses **41**(6): 479-486.

Koyuncu, O. O., I. B. Hogue and L. W. Enquist (2013). "Virus infections in the nervous system." Cell Host Microbe **13**(4): 379-393.

Landis, B. N., J. Frasnelli, I. Croy and T. Hummel (2010). "Evaluating the clinical usefulness of structured questions in parosmia assessment." Laryngoscope **120**(8): 1707-1713.

Landis, B. N., J. Frasnelli, J. Reden, J. S. Lacroix and T. Hummel (2005). "Differences Between Orthonasal and Retronasal Olfactory Functions in Patients With Loss of the Sense of Smell." Archives of Otolaryngology–Head & Neck Surgery **131**(11): 977-981.

Landis, B. N., T. Hummel, M. Hugentobler, R. Giger and J. S. Lacroix (2003). "Ratings of overall olfactory function." Chem Senses **28**(8): 691-694.

Landis, B. N., C. G. Konnerth and T. Hummel (2004). "A study on the frequency of olfactory dysfunction." Laryngoscope **114**(10): 1764-1769.

Le Bon, S.-D., N. Pisarski, J. Verbeke, L. Prunier, G. Cavelier, M.-P. Thill, A. Rodriguez, D. Dequanter, J. R. Lechien, O. Le Bon, T. Hummel and M. Horoi (2020). "Psychophysical evaluation of chemosensory functions 5 weeks after olfactory loss due to COVID-19: a prospective cohort study on 72 patients." Eur Arch Otorhinolaryngol.

Lechien, J. R., C. M. Chiesa-Estomba, E. Beckers, V. Mustin, M. Ducarme, F. Journe, A. Marchant, L. Jouffe, M. R. Barillari, G. Cammaroto, M. P. Circiu, S. Hans and S. Saussez (2021). "Prevalence and 6-month recovery of olfactory dysfunction: a multicentre study of 1363 COVID-19 patients." Journal of Internal Medicine **n/a**(n/a).

Lechien, J. R., C. M. Chiesa-Estomba, E. Beckers, V. Mustin, M. Ducarme, F. Journe, A. Marchant, L. Jouffe, M. R. Barillari, G. Cammaroto, M. P. Circiu, S. Hans and S. Saussez (2021). "Prevalence and 6-month recovery of olfactory dysfunction: a multicentre study of 1363 COVID-19 patients." Journal of Internal Medicine **290**(2): 451-461.

Lechien, J. R., C. M. Chiesa-Estomba, D. R. De Siati, M. Horoi, S. D. Le Bon, A. Rodriguez, D. Dequanter, S. Blecic, F. El Afia, L. Distinguin, Y. Chekkoury-Idrissi, S. Hans, I. L. Delgado, C. Calvo-Henriquez, P. Lavigne, C. Falanga, M. R. Barillari, G. Cammaroto, M. Khalife, P. Leich, C. Souchay, C. Rossi, F. Journe, J. Hsieh, M. Edjlali, R. Carlier, L. Ris, A. Lovato, C. De Filippis, F. Coppee, N. Fakhry, T. Ayad and S. Saussez (2020). "Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study." Eur Arch Otorhinolaryngol.

Lee, D., W. Lee, J. Wee and J.-W. Kim (2014). "Prognosis of postviral olfactory loss: Follow-up study for longer than one year." American Journal of Rhinology & Allergy **28**.

Liu, D. T., M. Sabha, M. Damm, C. Philpott, A. Oleszkiewicz, A. Hähner and T. Hummel (2021). "Parosmia is Associated with Relevant Olfactory Recovery After Olfactory Training." Laryngoscope **131**(3): 618-623.

Liu, J., J. M. Pinto, L. Yang, L. Li, J. Sun, X. Miao, K. Li, G. Chen and Y. Wei (2016). "Gender difference in
Chinese adults with post-viral olfactory disorder:a hospital-based study." Acta Otolaryngol **136**(9): 976-
981.
Lötsch, J. and T. Hummel (2019). "Clinical Usefulness of Self-Rated Olfactory Performance-A Data
Science-Based Assessment of 6000 Patients." Chem Senses **44**(6): 357-364.
Malaty, J. and I. A. Malaty (2013). "Smell and taste disorders in primary care." Am Fam Physician **88**(12):
852-859.
Mattes, R., B. Cowart, M. Schiavo, C. Arnold, B. Garrison, M. Kare and L. Lowry (1990). "Dietary
evaluation of patients with smell and/or taste disorders." The American journal of clinical nutrition **51**:
233-240.
Mazzatenta, A., G. Neri, D. D'Ardes, C. De Luca, S. Marinari, E. Porreca, F. Cipollone, J. Vecchiet, C.
Falcicchia, V. Panichi, N. Origlia and C. Di Giulio (2020). "Smell and Taste in Severe CoViD-19: Self-
Reported vs. Testing." Frontiers in medicine **7**: 589409-589409.
Meinhardt, J., J. Radke, C. Dittmayer, J. Franz, C. Thomas, R. Mothes, M. Laue, J. Schneider, S. Brünink, S.
Greuel, M. Lehmann, O. Hassan, T. Aschman, E. Schumann, R. L. Chua, C. Conrad, R. Eils, W. Stenzel, M.
Windgassen, L. Rößler, H. H. Goebel, H. R. Gelderblom, H. Martin, A. Nitsche, W. J. Schulz-Schaeffer, S.
Hakroush, M. S. Winkler, B. Tampe, F. Scheibe, P. Körtvélyessy, D. Reinhold, B. Siegmund, A. A. Kühl, S.
Elezkurtaj, D. Horst, L. Oesterhelweg, M. Tsokos, B. Ingold-Heppner, C. Stadelmann, C. Drosten, V. M.
Corman, H. Radbruch and F. L. Heppner (2020). "Olfactory transmucosal SARS-CoV-2 invasion as a port
of central nervous system entry in individuals with COVID-19." Nat Neurosci.
Moein, S. T., S. M. Hashemian, B. Mansourafshar, A. Khorram-Tousi, P. Tabarsi and R. L. Doty (2020).
"Smell dysfunction: a biomarker for COVID-19." Int Forum Allergy Rhinol.
Morris, J. A. (1985). "Clinical viral infections and multiple sclerosis." Lancet **2**(8449): 273.
Mullol, J., I. Alobid, F. Mariño-Sánchez, A. Izquierdo-Domínguez, C. Marin, L. Klimek, D.-Y. Wang and Z.
Liu (2020). "The Loss of Smell and Taste in the COVID-19 Outbreak: a Tale of Many Countries." Current
allergy and asthma reports **20**(10): 61-61.
Murray, R. S., B. Brown, D. Brian and G. F. Cabirac (1992). "Detection of coronavirus RNA and antigen in
multiple sclerosis brain." Ann Neurol **31**(5): 525-533.
Parma, V., M. E. Hannum, M. O'Leary, R. Pellegrino, N. E. Rawson, D. R. Reed and P. H. Dalton (2021).
"SCENTinel 1.0: development of a rapid test to screen for smell loss." Chem Senses.
Parma, V., K. Ohla, M. G. Veldhuizen, M. Y. Niv, C. E. Kelly, A. J. Bakke, K. W. Cooper, C. Bouysset, N.
Pirastu, M. Dibattista, R. Kaur, M. T. Liuzza, M. Y. Pepino, V. Schöpf, V. Pereda-Loth, S. B. Olsson, R. C.
Gerkin, P. R. Domínguez, J. Albayay, M. C. Farruggia, S. Bhutani, A. W. Fjaeldstad, R. Kumar, A. Menini,
M. Bensafi, M. Sandell, I. Konstantinidis, A. Di Pizio, F. Genovese, L. Öztürk, T. Thomas-Danguin, J.
Frasnelli, S. Boesveldt, Ö. Saatci, L. R. Saraiva, C. Lin, J. Golebiowski, L.-D. Hwang, M. H. Ozdener, M. D.
Guàrdia, C. Laudamiel, M. Ritchie, J. Havlícek, D. Pierron, E. Roura, M. Navarro, A. A. Nolden, J. Lim, K.
Whitcroft, L. R. Colquitt, C. Ferdenzi, E. V. Brindha, A. Altundag, A. Macchi, A. Nunez-Parra, Z. M. Patel, S.
Fiorucci, C. M. Philpott, B. C. Smith, J. N. Lundström, C. Mucignat, J. K. Parker, M. van den Brink, M.
Schmuker, F. P. S. Fischmeister, T. Heinbockel, V. D. C. Shields, F. Faraji, E. Enrique Santamaría, W. E. A.
Fredborg, G. Morini, J. K. Olofsson, M. Jalessi, N. Karni, A. D'Errico, R. Alizadeh, R. Pellegrino, P. Meyer, C.
Huart, B. Chen, G. M. Soler, M. K. Alwashahi, O. Abdulrahman, A. Welge-Lüssen, P. Dalton, J. Freiherr, C.
H. Yan, J. H. B. de Groot, V. V. Voznessenskaya, H. Klein, J. Chen, M. Okamoto, E. A. Sell, P. B. Singh, J.
Walsh-Messinger, N. S. Archer, S. Koyama, V. Deary, S. C. Roberts, H. Yanik, S. Albayrak, L. M. Novákov, I.
Croijmans, P. P. Mazal, S. T. Moein, E. Margulis, C. Mignot, S. Mariño, D. Georgiev, P. K. Kaushik, B.
Malnic, H. Wang, S. Seyed-Allaei, N. Yoluk, S. Razzaghi, J. M. Justice, D. Restrepo, J. W. Hsieh, D. R. Reed,
T. Hummel, S. D. Munger and J. E. Hayes (2020). "More than smell. COVID-19 is associated with severe
impairment of smell, taste, and chemesthesis." medRxiv: 2020.2005.2004.20090902.

Petrocelli, M., S. Cutrupi, G. Salzano, F. Maglitto, F. A. Salzano, J. R. Lechien, S. Saussez, P. Boscolo-Rizzo,
G. De Riu and L. A. Vaira (2021). "Six-month smell and taste recovery rates in coronavirus disease 2019
patients: a prospective psychophysical study." The Journal of Laryngology & Otology **135**(5): 436-441.
Saussez, S., S. Sharma, A. Thiriad, V. Olislagers, I. Vu Duc, S. D. Le Bon, M. Khalife, S. Hans, G. De Riu, C.
Hopkins, J. R. Lechien, L. A. Vaira and A. Marchant (2021). "Predictive factors of smell recovery in a
clinical series of 288 coronavirus disease 2019 patients with olfactory dysfunction." Eur J Neurol.
Schiffman, S. S. (2007). Smell and Taste. Encyclopedia of Gerontology (Second Edition). J. E. Birren. New
York, Elsevier**:** 515-525.
Sjölund, S., M. Larsson, J. K. Olofsson, J. Seubert and E. J. Laukka (2017). "Phantom Smells: Prevalence
and Correlates in a Population-Based Sample of Older Adults." Chemical senses **42**(4): 309-318.
Solomon, T. (2021). "Neurological infection with SARS-CoV-2 — the story so far." Nature Reviews
Neurology **17**(2): 65-66.
Sorokowska, A., E. Drechsler, M. Karwowski and T. Hummel (2017). "Effects of olfactory training: a meta-
analysis." Rhinology **55**(1): 17-26.
Sorokowski, P., M. Karwowski, M. Misiak, M. K. Marczak, M. Dziekan, T. Hummel and A. Sorokowska
(2019). "Sex Differences in Human Olfaction: A Meta-Analysis." **10**(242).
Speth, M. M., T. Singer-Cornelius, M. Oberle, I. Gengler, S. J. Brockmeier and A. R. Sedaghat (2020).
"Olfactory Dysfunction and Sinonasal Symptomatology in COVID-19: Prevalence, Severity, Timing, and
Associated Characteristics." Otolaryngol Head Neck Surg.
Spinato, G., C. Fabbris, J. Polesel, D. Cazzador, D. Borsetto, C. Hopkins and P. Boscolo-Rizzo (2020).
"Alterations in Smell or Taste in Mildly Symptomatic Outpatients With SARS-CoV-2 Infection." JAMA
323(20): 2089-2090.
Stewart, J. N., S. Mounir and P. J. Talbot (1992). "Human coronavirus gene expression in the brains of
multiple sclerosis patients." Virology **191**(1): 502-505.
Taquet, M., J. R. Geddes, M. Husain, S. Luciano and P. J. Harrison (2021). "6-month neurological and
psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic
health records." Lancet Psychiatry.
Vaira, L. A., C. Hopkins, G. Salzano, M. Petrocelli, A. Melis, M. Cucurullo, M. Ferrari, L. Gagliardini, C.
Pipolo, G. Deiana, V. Fiore, A. De Vito, N. Turra, S. Canu, A. Maglio, A. Serra, F. Bussu, G. Madeddu, S.
Babudieri, A. Giuseppe Fois, P. Pirina, F. A. Salzano, P. De Riu, F. Biglioli and G. De Riu (2020). "Olfactory
and gustatory function impairment in COVID-19 patients: Italian objective multicenter-study." Head &
Neck **42**(7): 1560-1569.
Vaira, L. A., C. Hopkins, A. Sandison, A. Manca, N. Machouchas, D. Turilli, J. R. Lechien, M. R. Barillari, G.
Salzano, A. Cossu, S. Saussez and G. De Riu (2020). "Olfactory epithelium histopathological findings in
long-term coronavirus disease 2019 related anosmia." The Journal of Laryngology & Otology **134**(12):
1123-1127.
Vaira, L. A., G. Salzano, M. Petrocelli, G. Deiana, F. A. Salzano and G. De Riu (2020). "Validation of a self-
administered olfactory and gustatory test for the remotely evaluation of COVID-19 patients in home
quarantine." Head & Neck **42**(7): 1570-1576.
von Bartheld, C. S., M. M. Hagen and R. Butowt (2020). "Prevalence of Chemosensory Dysfunction in
COVID-19 Patients: A Systematic Review and Meta-analysis Reveals Significant Ethnic Differences." ACS
Chem Neurosci **11**(19): 2944-2961.
Wang, X., C. Zhang, X. Xia, Y. Yang and C. Zhou (2019). "Effect of gender on odor identification at
different life stages: a meta-analysis." Rhinology **57**(5): 322-330.
Weiss, J. J., T. Attuquayefio, E. B. White, B. Geng, R. Handoko, R. S. Herz, T. L. White, A. Iwasaki, N. D.
Grubaugh, R. Datta, M. Campbell, R. A. Martinello, A. I. Ko, D. M. Small and S. F. Farhadian (2020). "456.
Implementing an At-Home Smell Test for Early Assessment of COVID-19 in High-Risk Healthcare
Workers." Open Forum Infectious Diseases **7**(Supplement_1): S295-S296.

Whitcroft, K. L. and T. Hummel (2020). "Olfactory Dysfunction in COVID-19: Diagnosis and
Management." <u>JAMA</u> **323**(24): 2512-2514.
Wu, Y., X. Xu, Z. Chen, J. Duan, K. Hashimoto, L. Yang, C. Liu and C. Yang (2020). "Nervous system
involvement after infection with COVID-19 and other coronaviruses." <u>Brain, Behavior, and Immunity</u>.
Xydakis, M. S., M. W. Albers, E. H. Holbrook, D. M. Lyon, R. Y. Shih, J. A. Frasnelli, A. Pagenstecher, A.
Kupke, L. W. Enquist and S. Perlman (2021). "Post-viral effects of COVID-19 in the olfactory system and
their implications." <u>The Lancet Neurology</u>.
Yang, J. and J. M. Pinto (2016). "The Epidemiology of Olfactory Disorders." <u>Current otorhinolaryngology</u>
<u>reports</u> **4**(2): 130-141.

Table 1. COVID-19 symptoms of the 704 participants.

Symptoms at time of SARS-CoV-2 infection	No. (%)
Fever	353 (50.1%)
Dry cough	361 (51.7%)
Cough with mucus	77 (10.9%)
Dyspnea	316 (44.9%)
Chest tightness	201 (28.6%)
Runny nose	226 (32.1%)
Sore throat	330 (46.9%)
Changes in food flavor	471 (66.9%)
Changes in smell	520 (73.9%)
Loss of appetite	323 (45.9%)
Headache	518 (73.6%)
Muscle aches	444 (63.1%)
Fatigue	611 (86.8%)
Diarrhea	259 (36.8%)
Abdominal pain	102 (14.5%)
Nausea	179 (25.4%)

628   **Table 2.** Self-reported chemosensory alterations by age group and gender during and 3 to 7

629   months following COVID-19 (n=704).

Age	Gender	During acute COVID-19			3-7 months after COVID-19		
		Olfaction	Gustation	Trigeminal	Olfaction	Gustation	Trigeminal
		N (%)	N (%)	N (%)	N (%)	N (%)	N (%)
18-29	M (N= 11)	9 (81.8)	9 (81.8)	4 (36.4)	6 (54.5)	5 (45.5)	3 (27.3)
	F (N=115)	106 (92.2)	106 (92.2)	54 (47.0)	68 (59.1)	55 (47.8)	16 (13.9)
30-39	M (N=26)	23 (88.5)	21 (80.8)	12 (46.2)	13 (50.0)	7 (26.9)	4 (15.4)
	F (N=153)	133 (86.9)	129 (84.3)	77 (50.3)	83 (54.2)	63 (41.2)	37 (24.2)
40-49	M (N=33)	23 (69.7)	23 (69.7)	13 (39.4)	12 (36.4)	9 (27.3)	5 (15.2)
	F (N=165)	142 (85.5)	137 (83.0)	83 (50.3)	97 (58.8)	77 (46.7)	42 (25.5)
50-59	M (N=28)	13 (46.4)	14 (50.0)	9 (32.1)	7 (25.0)	7 (25.0)	4 (14.3)
	F (N=128)	100 (78.1)	102 (79.7)	66 (51.6)	62 (48.4)	57 (44.5)	39 (30.5)
60+	M (N=13)	5 (38.5)	6 (46.2)	5 (38.5)	2 (15.4)	1 (7.7)	3 (23.1)
	F (N=32)	19 (59.4)	27 (84.4)	15 (46.9)	16 (50.0)	14 (43.8)	11 (34.4
Total (N=704)		**572 (81.3)**	**574 (81.5)**	**338 (48.0)**	**366 (52.0)**	**295 (41.9)**	**164 (23.2)**

Figure 2. Web-based interface and structure of the online questionnaire. Left, Self-rating of

olfaction and reporting of CPT using VAS through the web-based interface, as viewed by the

participant. Right, Sections of the online questionnaire. VAS: visual analog scale.

Figure 3. Self-reported scores for the chemosensory modalities before, during and after

COVID-19 infection (n=704). Raincloud plot representing self-reported scores for olfaction,

gustation, and trigeminal function before, during and after COVID-19. Ratings from individual

participants are displayed as dots. Boxplots show the first to third quartiles, horizontal line denotes

the median, and whiskers denote 1.5 times interquartile range. Compared to baseline, self-reported

scores of olfaction, gustation and trigeminal function were significantly lower during COVID-19

and have not fully returned to baseline values 5 months after COVID-19.

Figure 4. Correlations between alterations in chemosensory modalities (n=704). Red squares,

correlations between alterations in olfaction, gustation, and trigeminal functions during COVID-

19. Grey hexagons, correlations between alterations in olfaction, gustation, and trigeminal

functions after COVID-19. Darker colors indicate higher occurrence.

648
649

Supplementary 1:

Validation of the Chemosensory Perception Test

This study was reviewed and approved by the research ethics board of the Université du Quebec à

Trois-Rivières (CER-20-268-08-01.04). All participants provided a verbal or written informed

consent prior to participation.

Experiment 1:

Methods

Participants were recruited among previously tested groups. Olfactory testing was performed

using the standardized Sniffin' Sticks test at our laboratory from 2016 to 2019. Exclusion criteria

was any perceived changes of their sense of smell since previous testing. Participants were

distributed into 2 groups based on their Threshold-Discrimination-Identification (TDI) scores.

The first group consists of participants with normal olfactory function (normosmia), defined as

TDI scores above 30.5[61]. The second group had subjective olfactory dysfunction and equivalent

TDI scores. Participants were administered CPT by means of a telephone interview.

Results:

TDI scores in the first group range from 32.5 to 41.5 (N= 19, 9 women and 10 men, age range [60-

78]). TDI scores in the second group ranged from 8 to 30.25 (N= 17, 7 women and 10 men, age

range [57-77]). CPT scores were positively correlated with the Sniffin' Sticks (ρ=0.837, P<0.001).

A cut-off score of 6 at the CPT had a sensitivity of 0.765 and specificity of 0.895.

Experiment 2:

Methods:

Participants were recruited among previously tested groups, participants from this cross-sectional

study and in the public via social media. They were administered the University of Pennsylvania

Smell Identification Test (UPSIT) (which was sent by mail) and the CPT under direct supervision

through videoconferencing. Participants were distributed into 2 groups based on their UPSIT

scores with a score equal or less than 33 in males and 34 in females defining hyposmia[62].

<u>Results:</u>

UPSIT scores in the normosmic group ranged from 34 to 28 (n=29 (21 women), age range [22-

73]). The hyposmic group had UPSIT scores ranging from 9 to 34 (n= 28 (21 women), age range:

[22-72]). CPT scores were significantly correlated with the UPSIT score (ρ=0.377, *P*=0.004) in

the whole group of participants. We found this correlation to be much stronger in the hyposmic

group (ρ=0.702, *P*<0.001).

Conclusions:

The CPT allows for distinction between normosmia and olfactory dysfunction with high sensitivity

and specificity. CPT scores are significantly correlated to UPSIT and Sniffin Sticks scores,

especially in a group of individuals with olfactory dysfunction.