Genomic surveillance reveals the spread patterns of SARS-CoV-2 in coastal Kenya during the first two waves

Charles N. Agoti[1,2,#], Lynette Isabella Ochola-Oyier[1], Khadija Said Mohammed[1], Arnold W. Lambisia[1], Zaydah R. de Laurent[1], John M. Morobe[1], Maureen W. Mburu[1], Donwilliams O. Omuoyo[1], Edidah M. Ongera[1], Leonard Ndwiga[1], Eric Maitha[3], Benson Kitole[3], Thani Suleiman[3], Mohamed Mwakinangu[3], John Nyambu[3], John Otieno[3], Barke Salim[3], Jennifer Musyoki[1], Nickson Murunga[1], Edward Otieno[1], John Kiiru[3], Kadondi Kasera[3], Patrick Amoth[3], Mercy Mwangangi[3], Rashid Aman[3], Samson Kinyanjui[1,2,4], George Warimwe[1,4], My Phan[5], Ambrose Agweyu[1], Matthew Cotten[5,6], Edwine Barasa[1], Benjamin Tsofa[1], D. James Nokes[1,7], Philip Bejon [1,4], George Githinji[1]

Affiliations
1. Kenya Medical Research Institute (KEMRI)-Wellcome Trust Research Programme (KWTRP), Kilifi, Kenya.
2. Pwani University, Kilifi, Kenya.
3. Ministry of Health, Kenya.
4. Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom.
5. Medical Research Centre/ Uganda Virus Research Institute, Entebbe, Uganda.
6. MRC-University of Glasgow Centre for Virus Research, Glasgow, UK.
7. University of Warwick, Coventry, United Kingdom.

[#]**Corresponding author**
Dr Charles N. Agoti (Email: cnyaigoti@kemri-wellcome.org)

ABSTRACT

The transmission networks of SARS-CoV-2 in sub-Saharan Africa remain poorly understood. We analyzed 684 genomes from samples collected across six counties in coastal Kenya during the first two waves (March 2020 - February 2021). Up to 32 Pango lineages were detected in the local sample with six accounting for 88.0% of the sequenced infections: B.1 (60.4%), B.1.1 (8.9%), B.1.549 (7.9%), B.1.530 (6.4%), N.8 (4.4%) and A (3.1%). In a contemporaneous global sample, 571 lineages were identified, 247 for Africa and 88 for East Africa. We detected 262 location transition events comprising: 64 viral imports into Coastal Kenya; 26 viral exports from coastal Kenya; and 172 inter-county import/export events. Most international viral imports (61%) and exports (88%) occurred through Mombasa, a key coastal touristic and commercial center; and many occurred prior to June 2020, when stringent local COVID-19 restriction measures were enforced. After this period, local transmission dominated, and distinct local phylogenies were seen. Our analysis supports moving control strategies from a focus on international travel to local transmission.

INTRODUCTION

By 15th June 2021, at least 175 million cases of coronavirus disease 2019 (COVID-19) and > 3.7 million associated deaths were reported worldwide (1). By the same date, Kenya reported 175,176 laboratory-confirmed cases of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and 3,396 COVID-19 associated deaths (2). Serological surveys indicated that Kenya's COVID-19 epidemic had progressed further than could be discerned from the limited laboratory testing case reports that were available during the period (3, 4).   At the end of Kenya's first wave (August 2020), nationwide anti-SARS-CoV-2 IgG prevalence based on analysis of blood donor samples was estimated to be 9% (5).

By June 2021, Kenya had experienced three waves of SARS-CoV-2 infections. The first wave peaked in July-August of 2020, the second in November 2020 and the third in March-April 2021 (6). Despite this progression, the local SARS-CoV-2 spread patterns remain poorly understood. The analysis presented here examined genome sequences from the first two waves in Coastal Kenya. Overall, the documented SARS-CoV-2 infections in Kenya during these early two waves were concentrated in cities, especially Nairobi (~42%), the capital, and Mombasa (~8%), the country's second largest city on the Indian ocean coast. In addition to Mombasa, there are five other counties that make up Kenya's Coastal region namely, Kilifi, Kwale, Taita Taveta, Tana River and Lamu. The region is a major tourism destination accessible through multiple airports, seaports and land border entry points from Tanzania through Kwale and Taita Taveta counties (7).

Throughout the COVID-19 pandemic, genomic surveillance has been critical for tracking the spread of SARS-CoV-2 and investigating viral evolution (8-11). In Kenya, genome sequencing started soon after the identification of the initial case on 13th March 2020 and has remained a key tool informing public health response decisions. We have previously reported on the genomic analysis of SARS-CoV-2 in Coastal Kenya during the early phase of the pandemic (up to July 2020) revealing multiple lineage introductions but with mainly lineage B.1 establishing local transmission (7). Here we report the genetic composition of the subsequent two waves (up to February 2021) with the aim of documenting patterns of viral introductions, evolution and spread across the six Coastal counties of Kenya.

RESULTS

Infection waves in Coastal Kenya counties

The first cases of COVID-19 in Coastal Kenya were reported in Mombasa and Kilifi counties in March 2020, **Figure 1A**. The timing of the first infection peaks across the six counties differed, with Mombasa showing the earliest surge. By 26th February 2021, Mombasa, Lamu and Taita Taveta had observed two major infection peaks while Kilifi, Kwale and Tana River had experienced only a single peak, with the latter two documenting very few cases, **Figure 1A.** During this period, the Kenya Ministry of Health (MoH) reported a total of 12,307 laboratory

confirmed SARS-CoV-2 cases for the six Coastal counties: Mombasa (n=8,450, 68.7%), Kilifi
(n=2,458, 20.0%), Taita Taveta (n=567, 4.6%), Kwale (n=437, 3.6%), Lamu (n=309, 2.5%) and
Tana River (n=85, 0.7%**), S1 Figure.**

SARS-CoV-2 testing and sequencing at KWTRP

The KEMRI-Wellcome Trust Research Programme (KWTRP) was designated as the government

SARS-CoV-2 testing center for Coastal region soon after identification of the first case in Kenya.

Between 17th March 2020 and 26th February 2021, we tested 82,716

nasopharyngeal/oropharyngeal (NP/OP) swab samples from the six Coastal counties and 6,307

(7.6%) of these were determined as SARS-CoV-2 positive (approximately half of the total

reported by MoH), distributed by month as in **Figure 1B**. Overall, a total of 3,137 (6.8%) tests

at KWTRP were positive from samples taken in Mombasa, 1,142 (8.8%) for Kilifi, 657 (4.5%)

for Taita Taveta, 349 (12.7%) for Lamu, 437 (7.9%) for Kwale and 105 (12.0%) for Tana River.

Figure 1. SARS-CoV-2 testing and sequencing on the Kenyan Coast. Panel A, epidemic

curves for each of the six Coastal Kenya counties derived from the daily positive case numbers

reported by the Ministry of Health. Panel B, monthly count of SARS-CoV-2 RT-PCR tests

undertaken at the KEMRI-Wellcome Trust Research Programme (KWTRP) and those positive

during the study period. Panel C, monthly proportion of samples sequenced from total SARS-

CoV-2 positives detected at KWTRP. Panel D, Distribution of the sequenced 684 samples across

the 6 Coastal Kenya counties.

We sequenced the genomes of 684 (10.8%) RT-PCR positive samples as follows: Mombasa: 337
(10.7%), Kilifi: 118 (8.1%), Taita Taveta: 115 (13.4%), Kwale: 65 (14.9%), Lamu: 40 (11.5%)
and Tana River: 9 (8.5%), **Figure 1C and 1D**. The sequences were spread across the study
period with 272 (39.8%) from the introduction phase (17[th] March to 20[th] May 2020 when
nationally <50 new cases were reported per day), 169 (24.7%) from wave one (21[st] May-15[th]
September 2020) and 243 (35.5%) from wave two (16[th] September 2020 to 26[th] February 2021),
S1 Figure. Overall, our sequencing rate corresponded to approximately one sequence for every
18 confirmed cases.

Lineage dynamics in Coastal Kenya
The 684 genomes were classified into 32 Pango lineages, including four first identified in Kenya
(N.8, B.1.530, B.1.549 and B.1.596.1), **S1 Table**. During the introductions phase we detected 12
lineages while during wave one we detected 16 lineages (11 of which were being identified
locally for the first time) and during wave two we detected 18 lineages (nine of which were
being identified locally for the first time). Thirteen lineages (40.6%) were identified in three or
more samples with the top six accounting for 88.0% of the sequenced infections: B.1 (n=413,
60.4%, European lineage which predominated in the Northern Italy outbreak early in 2020),
B.1.1 (n=61, 8.9%, European lineage), B.1.549 (n=54, 7.9%, Kenyan lineage), B.1.530 (n=44,
6.4%, Kenyan lineage), N.8 (n=30, 4.4%, Kenyan lineage) and A (n=21, 3.1%, lineage root to
the pandemic first observed in China).

Lineage B.1 was the first to be detected in the region and comprised the initial cases identified in
Mombasa, Kilifi, Kwale and Tana River**, Figure 2**. This lineage comprised 71.3% (n=194) of the
viruses we sequenced from the introduction phase, 65.7% (n=111) from wave one and 44.4%
(n=108) from wave two. Overall, B.1 accounted for most of the virus sequences detected in all
counties except Lamu where lineage N.8 (alias lineage B.1.1.33.8) was most frequent (70% of
the cases, first detected there on 23[rd] June 2020). Only two additional sequenced cases of
Lineage N.8 occurred outside Lamu County (one in Mombasa and another in Kilifi).

Figure 2. Biweekly distribution of the common Pango lineages identified on the Kenyan coast from March 2020 to February 2021 across the six counties presented as proportions.
Lineages detected in less than three cases were put together and referred to as "other lineages". This group included 18 lineages namely: A.25, B.1.1.254, B.1.179, B.1.378, B.1.393, B.1.1.111, B.1.1.33, B.1.1.485, B.1.1.7, B.1.1.70, B.1.153, B.1.340, B.1.450, B.1.526, B.1.535, B.1.553, B.1.558 and B.4 (see details in S1 Table).

Although lineage A was also detected in Mombasa and Taita Taveta, most of its cases occurred in Kwale where it comprised 20% of all cases identified in the county, **Figure 2**. Lineage B.1.1 was mostly observed in Mombasa with limited number of cases observed in Kilifi, Kwale and Taita Taveta.

During the wave two period in Coastal Kenya, there was an emergence of three lineages first identified in Kenya: (a) lineage B.1.549 (first observed 26th June 2020) which was responsible for a significant number of the sequenced infections in Kilifi (n=24), Mombasa (n=18), Taita-Taveta (n=5), and Kwale (n=7), (b) lineage B.1.596.1 (first observed 01st August 2020) and detected in Mombasa, Kilifi, Kwale and Taita-Taveta and (c) lineage B.1.530 (first observed 28th October 2020) that predominated infections in Taita-Taveta (n=28) with a fewer cases observed in Mombasa (n=8), Kilifi (n=7) and Lamu (n=1).

Two known variants of concern (VOC) were detected during the study period: the lineage B.1.351 (the Beta variant in 10 samples; and the lineage B.1.1.7 (the Alpha variant in one sample

(12). The earliest B.1.351 infections were identified in international travelers from South Africa
in mid-December 2020 while the earliest detected B.1.1.7 case was in second week of January
2020 in a local with no history of recent international travel. We also detected three cases each of
A.23 and A.23.1 after the peak of wave two (both lineages first identified in Uganda, and are
considered variants of interest (VOI) in the region (13)).

Lineage dynamics with a widening scale of observation.
We compared the temporal prevalence patterns of the lineages identified in Coastal Kenya to
Eastern Africa, Africa and a global sub-sample, **Figure 3**. In these comparative genome sets, we
identified 88 Pango lineages for Eastern Africa, 247 for Africa and 571 for the global sub-
sample.

168
Figure 3. Lineage dynamics at widening scales of observation. Panel A, Biweekly distribution
of detected lineages in Coastal Kenya from the sequenced 684 genomes. Panel B, biweekly
distribution of detected lineages in Eastern Africa from 2,316 contemporaneous genomes from
10 countries whose contemporaneous data were available in GISAID. Lineages not observed in
the Coastal Kenya dataset have been lumped together as "Absent Kenya" but with the those
appearing among top six and not observed in Kenya indicated just below the other Kenya group.
Panel C, weekly distribution of detected lineages in Africa from 12,506 contemporaneous
genomes from 39 countries that were available in GISAID. Panel D, weekly distribution of
detected lineages in a global sub-sample of 9,989 contemporaneous genomes from 112 countries
that were compiled from GISAID (see detail in methods section).

In the Eastern Africa sample (n=2,315), all the top lineages were also observed in the Coastal

Kenya data (B.1.351, A.23.1, B.1, B.1.1, B.1.177) except lineage B.1.380 (n=128, a Rwanda

lineage), **Figure 3B**. In the Africa sample (n=12,506), four of the six top lineages (B.1, B.1.1,

B.1.351 and B.1.1.7) were also identified in Coastal Kenya. The undetected two lineages were

B.1.1.448 (n=392) and C.1 (n=330) both of which were mainly observed South Africa samples.

In the global sub-sample (n=8,909) four of the top six lineages (B.1, B.1.1, B.1.1.7 and B.1.177)

where observed in the samples from Coastal Kenya. The unobserved lineages were D.2 (Alias of

B.1.1.25.2, an Australian lineage) and B.1.1.28 (a Brazilian lineage).

At the time of the second wave in Kenya, the VOC (B.1.1.7 and B.1.351) were already

widespread across Eastern Africa and Africa but there were only sporadic detections our dataset

from Coastal Kenya. Although in the comparison data lineage B.1 occurred in substantial

proportions across the different scales early in the pandemic, its prevalence diminished faster

overtime in the outside Kenya sample sets when compared to the Coastal Kenya dataset set.

Approximately 60% of the lineages comprising infections globally were not seen in the Coastal

Kenya samples.

SARS-CoV-2 diversity in Coastal Kenya

When we reconstructed time-resolved ML phylogeny of the Coastal Kenya genomes while

including a global reference set of genomes (n=1500) we observed that (i) the Coastal Kenya

genomes were represented across many but not all of the major phylogenetic clusters observed

globally, ii) some of the coast clusters expanded after introduction whereas others did not

(observed as singletons) and iii) all counties appeared to have had multiple variants introduced

with some clusters comprising genomes detected across multiple counties, **Figure 4A**. There

was considerable correlation of the root-to-tip genetic distance and the date of sampling

(r^2=0.442) **, Figure 4B**.

Figure 4. SARS-CoV-2 diversity in Coastal Kenya. Panel A, Time-resolved global phylogeny
that combined 639 Coastal Kenya SARS-CoV-2 genomes and 1474 global reference sequences.
Panel B, Roo-to-tip regression analysis coastal Kenya (colored by county) and global (shown in
grey) sequences with determination of co-efficient (r^2). Panel C, map of the six study counties
colored scaled by the population density (number of persons per square kilometer).

For detailed investigation into the local SARS-CoV-2 genetic diversity, we reconstructed time-
resolved lineage specific phylogenetic trees are shown in **Figure 5**. Viruses occurring within
lineage B.1 and lineage B.1.1 showed significant genetic divergence and formed multiple
phylogenetic clusters interspersed with other global sequences, a feature indicative of multiple
viral introductions into a geographic area. Three lineages first identified in Kenya (B.1.530,
B.1.549 and B.1.596.1) were found to (a) possess significant diversity consistent with widescale
spread within Kenya (**Figure 5B-D**), (b) formed multiple county-specific sub-clusters and (c)
show local sequences interspersed with global comparison genomes from the same lineage
implying potentially export (or import) events, **Figure 5C**. However, the picture painted by
lineage N.8 was different. This lineage was mainly detected in Lamu forming a single
monophyletic group (**Figure 5E**) when co-analysed with its precursor lineage B.1.1.33, an
observation consistent with a single introduction then expansion.

Figure 5. Time-resolved lineage-specific phylogenetic trees for the commonly detected lineages

in Coastal Kenya. The Coastal Kenya genomes are indicated with filled circles colored by

county. Panel A, phylogeny for lineage B.1 that combined 369 Coastal Kenya genomes and 1134

global reference sequences. Panel B, phylogeny for lineage B.1.530 that combined 44 Coastal

Kenya genomes and 15 global sequences belonging also to B.1.530. Panel C, phylogeny for

lineage B.1.549 that combined 54 Coastal Kenya genomes and 23 sequences from other

countries of lineage B.1.549. Panel D, phylogeny for lineage B.1.596.1 combines 8 Coastal

Kenya genomes and 19 global reference sequences. Panel E, phylogeny for lineage N.8 that

combined 30 Coastal Kenya genomes of lineage N.8, 1 Coastal Kenya genome of lineage

B.1.1.33 and 99 global sequences of lineage B.1.1.33. Panel F, phylogeny for lineage B.1.1 that

combined 60 Coastal Kenya genomes and 824 global sequences belonging to lineage B.1.1.

Viral imports and export from Coastal Kenya

We used ancestral location state reconstruction of the dated phylogeny **(Figure 4)** to infer the

number of viral import and exports (see details in the methods (14)). In total, between March

2020 and February 2021, we detected 262 location transition events, 64 of which were viral

import events into Coastal Kenya from outside and 26 were viral exports events from Coastal

Kenya to outside populations, **Figure 6A-C**. A total of 172 import/export events were detected

between the six Coastal counties. Virus imports into the region occurred through Mombasa

(n=39, 61%), Kwale (n=10, 16%), Taita Taveta (n=10, 16%), Kilifi (n=3, 5%) and Lamu (n=1,

2%). Virus export from the region occurred through Mombasa (n=23, 88%) and Kwale (12%).

Of the 64 detected viral imports from outside the Coastal Kenya region, 40 (63%) occurred

during the introductions phase, 12 (19%) occurred during wave one and another 12 (19%) during

wave two. Of the 26 detected viral exports, 13 (50%) occurred during introductions phase, 10

(38%) during wave one and three (12%) during wave two, **Figure 6B and D**.

Figure 6. Viral imports and export from Coastal Kenya. Panel A, the number of imports and

exports of viral lineages from Coastal Kenya shown as an alluvium plot. Panel B, biweekly

location transition events observed in the Coastal Kenya and global genomes shown as either

viral export, import or inter-county transmission. Panel C, Biweekly number of import events

into the individual Coastal counties and their origins.

Amino acid changes in the Coastal Kenya genomes

We scanned the entire SARS-CoV-2 genome for amino acid changes from the Wuhan reference

genome in the 684 Coastal Kenya sequences. Below we summarize changes that were identified

in ≥5 genomes. We observed 20 changes in the spike (S) protein meeting the set criterion, 15 in

the nucleocapsid (N) protein, 14 in ORF3a, 13 in ORF1b, nine in the ORF1a, seven in ORF8,

five in ORF6, and one in the envelope (E) protein. No codon changes, meeting of criterion was

observed in the matrix (M) protein, ORF7a, ORF7b or ORF10. The most common amino acid

changes observed in the Coastal Kenya genomes were S: D614G (94.4%), ORF1b: P314L

(92.9%), ORF3a: Q57H (15.5%), ORF1a: T265I (15.4%) and N: R195K (12.9%), **S2 Figure.**

Several lineage-defining mutations were observed for the Kenya-specific lineages as summarized

in **S3 Figure.**

DISCUSSION

We present evidence of continuous introductions of SARS-CoV-2 into Coastal Kenya documenting at least 64 independent viral introduction events into the region during the first year of the pandemic. Multiple variant introductions were observed even at the individual counties level, with extensive inter-county transmission events after the early phase period, re-affirming that Coastal Kenya populations are quite linked epidemiologically. Strikingly, many of the imports and exports from the region occurred through Mombasa, a major commercial, industrial and tourist destination in the region, before dispersing to the other counties. This observation highlights Mombasa's central role as a gateway of viral variants into the region and the significance of enhanced and persistent surveillance in this city for early detection of introduced of variants of concern.

The analyzed counties had both differences and similarities in SARS-CoV-2 transmission patterns. While Mombasa, Taita Taveta and Lamu counties had experienced two waves of infections by February 2021, Kilifi, Tana River and Kwale counties had experienced only a single major wave. This may have been brought about by differences in population density, accessibility, and applied infection control measures. For instance, Mombasa is densely populated (~5,604.64 persons/sq.km, 2019 census), has a large seaport and an international airport while Lamu has a sparsely populated mainland and over 65 islands (total population 143,920, population density, 21.84 persons/sq.km). Taita Taveta and Kwale counties are primarily rural counties bordering Tanzania which had a largely uncontrolled epidemic (15). Tana River is the most remote and sparsely populated (8.9 persons/sq.km) county in the region (**Figure 4C**).

During the early months of the pandemic several Pango lineages were introduced into the region (A, B.1, B.1.1, B.1.340, B.1.535, B.4 and B.4.7). However, it is lineage B.1 which has a European origin, that became predominant. This lineage has the D614G change in its spike protein considered to enhance fitness (16) and this may have boosted its local transmission and early dominance. Lineage N.8 was specific to Lamu County and B.1.530 to Taita Taveta. Lineage N.8 has not been observed outside Kenya and its precursor lineage (B.1.1.33) was observed earlier in South America, mainly Brazil. Lineage N.8 may have arisen from a single introduction event into Lamu county, and then remained localized to Lamu. The N.8 lineage has seven characteristic lineage defining mutations including S: D614G and R203K, G204R and I292T on the nucleocapsid protein.

Kilifi county, which neighbors Mombasa to the North (~1.4 million with a population density of 117.67 persons/sq.km, 2019 census) observed its first peak of infections during the second national wave. The Kilifi first wave was comprised of mainly three lineages: B.1 (53.5%), B.1.530 (8.1%) and B.1.549 (25.6%). Lineages B.1.530 and B.1.549, first identified in Kenya, and may have arisen from local evolution of B.1. Lineage B.1.530 has six characteristic mutations including spike P681H change adjacent to the biologically important furin cleavage

314   site while lineage B.1.549 has seven characteristic mutations, five occurring in the ORF1a or
315   ORF1b. Both two lineages have now been observed in a few other countries albeit in small
316   numbers: seven countries for B.1.530 (Rwanda, Netherlands, Germany, Denmark, USA, Japan
317   and Australia) and three countries for B.1.549 (England, USA, Canada).
318
Near the tail end of the national wave two, we detected only sporadic cases of the VOC B.1.351
and B.1.1.7. Lineage B.1.351 was first to be observed, initially in Kilifi in two asymptomatic
international travelers in mid-December 2020. Lineage B.1.1.7 was subsequently detected in a
local who presented to a Mombasa clinic in the second week of January 2021. In the subsequent
weeks up to the end of the period covered by this analysis (February 2021), no additional B.1.1.7
was detected unlike lineage B.1.351 which continued to be detected sporadically in January and
February 2021, especially in truckers screened at Tanzania border points of entry in Taita Taveta
and Kwale counties.
327
When we compared the local lineage patterns to the global patterns, the VOC were already
extensively spread across Eastern Africa (B.1.351 VOC), Africa (B.1.351 VOC) and worldwide
(B.1.1.7 VOC) in the last quarter of 2020. This implies that a lag occurred in their arrival into
Coastal Kenya perhaps due to public health measures that have remained in place during the
period especially at the international borders. It is also notable several lineages were detected
globally (>500) but only a small fraction (<10%) of this were documented locally.
334
Our study contributes to the limited but growing literature illuminating SARS-CoV-2
transmission patterns in Africa (11, 13, 17-19). The patterns revealed have potential to inform
pandemic mitigation strategies. The key limitations of our analysis include, first, the samples we
sequenced were only those available to us through the rapid response teams (RRTs) whose case
identification protocols were altered at the different study phases following guidance from the
MoH. Second, we only sequenced ~10% of the positive samples identified in our laboratory, the
majority from the introduction phase and were prioritizing samples with a Ct value of <30.0.
Third, the sampling across the six Coastal counties was not uniform, probably in part due to
varied distance from our testing centre located in Kilifi County but also the total number of
positive cases varied between the counties.
345
In conclusion, we show that the first two SARS-CoV-2 waves in Coastal Kenya observed
transmission of both newly introduced variants and potentially locally evolved variants. New
variant introductions appeared to mainly occur through Mombasa city. Strikingly, only a limited
number of the many introduced variants progressed to transmit extensively perhaps due to
ongoing public health interventions e.g., screening at ports of entry, case isolation and
quarantining of contacts of cases. Unlike in the global contemporaneous sample, we did not find
evidence of extensive local transmission of the global VOC during wave two. Thus, we infer that
it is more likely that the relaxation of some of the interventions (e.g., reopening of learning

institutions, airspace, bars and restaurants) that drove the second wave of infections. Overall, our
study shows the importance of detailed local genomic surveillance even in remote and under-
resourced settings to understand origins and spread patterns of SARS-CoV-2 to optimize local
interventions.
358
MATERIALS AND METHODS
Ethical statement
Samples analysed here were collected under the MoH protocols as part of the national response
to the COVID-19 pandemic. The whole genome sequencing study protocol was reviewed and
approved by the Scientific and Ethics Review Committee (SERU), Kenya Medical Research
Institute (KEMRI), Nairobi, Kenya (SERU #4035). Individual patient consent was not required
by the committee for the use of these samples for genomic surveillance to inform public health
response.
367
Study period and population
This study examined samples collected between 17[th] March 2020 to 26[th] February 2021 from six
counties in Coastal Kenya i.e Mombasa, Kilifi, Kwale, Tana River, Taita Taveta and Lamu. We
divided the study period into three phases based on trends in daily reported COVID-19 national
case numbers by the MoH (2): (a) Introduction phase – 17[th] March-20[th] May, 2020 (b) Wave one
– 21[st] May-15[th] September 2020 and (c) Wave two – 16[th] September to 28[th] February 2021, **S1**
Figure. We considered the transition from introduction phase to Wave as the timepoint when the
national daily positives exceeded 50 and the transition from Wave one to Wave two as the
timepoint when a consistent renewed rise of national daily number of positives started after
Wave one peak.
378
Patient samples
This study analyzed SARS-CoV-2 positive NP/OP swab samples that were collected by the MoH
County Department of Health RRTs, across all the six counties of Coastal Kenya. SARS-CoV-2
positive diagnosis was made at the KWTRP in Kilifi County (20). The RRTs delivered the
NP/OP swabs to KWTRP laboratories within 48 hours of collection in cool boxes with ice packs.
The samples were from persons of any age and sampling followed the MOH eligibility criteria
that were revised from time to time (7). Persons sampled included those with acute respiratory
symptoms, those with a recent history of travel to the early COVID-19 hotspots, contacts of
confirmed cases, persons presenting at international border points seeking entry into Kenya,
among other criteria.
389
SARS-CoV-2 detection at KWTRP
Viral RNA was extracted from the NP/OP samples using any one of seven commercial kits that
were available namely, QIAamp Viral RNA Mini Kit, RNeasy ® QIAcube ® HT Kit,
QIASYMPHONY ® RNA Kit, T0IANamp Virus RNA Kit, Da An Gene Nucleic acid Isolation

and Purification Kit, SPIN X Extraction and RADI COVID-19 detection Kit. The extracts were
tested for the presence of SARS-CoV-2 nucleic acid following different protocols depending on
which one was available of the following 7 kits/protocols: 1) the Berlin (Charité) primer-probe
set (targeting envelope (E) gene, nucleocapsid (N) or RNA-dependent RNA-polymerase
(RdRp)), 2) European Virus Archive – GLOBAL (EVA-g) (targeting E or RdRp genes), 3) Da
An Gene Co. detection Kit (targeting N or ORF1ab), 4) BGI RT-PCR kit (targeting ORF1ab), 5)
Sansure Biotech Novel Coronavirus (2019-nCoV) Nucleic Acid Diagnostic real-time RT-PCR
kit or 6) Standard M kit (targeting E and ORF1ab) and 7) TIB MOLBIOL kit (targeting E gene).
Protocol-specific recommended cycle threshold cut-offs were followed in defining SARS-CoV-2
positives.

SARS-CoV-2 genome sequencing
Only samples that had a PCR cycle threshold value of <30.0 were targeted for whole genome
sequencing (7). Viral RNA from positive samples was reextracted using QIAamp Viral RNA
Mini kit following the manufacturer's instructions and reverse transcribed using LunaScript®
RT SuperMix Kit. The cDNA was amplified using Q5® Hot Start High-Fidelity 2x Mastermix
along with the ARTIC nCoV-2019 version 3 primers. The PCR products were run on an 1.5%
agarose gel and for samples whose SARS-CoV-2 amplification was considered successful were
purified using Agencourt AMPure XP beads and taken forward for library preparation.
Sequencing libraries were constructed using Oxford Nanopore Technology (ONT) ligation
sequencing kit and the ONT Native Barcoding Expansion kit as described in the ARTIC
protocol (21). Every MinION (Mk1B) run comprised 23 samples and one negative (no-template)
control.

SARS-CoV-2 genome assembly
Following MinION sequencing, the FAST5 files were base-called using the OTN's software
Guppy v3.5-4.2. Consensus SARS-CoV-2 sequences were derived from the reads using the
ARTIC bioinformatics pipeline. A threshold of ×20 coverage was required for a base to be
included in the consensus genome otherwise it was masked with an N. Only complete or near-
complete genomes with >80% coverage and ×20 or more read depth were taken forward for
phylogenetic analysis.
425

Lineage assignment
The consensus genomes were assigned into Pango lineages via the Phylogenetic Assignment of
named Global Lineages (PANGOLIN) software suite (Pangolin v2.4.2, Pango v1.1.23) (22). The
amino acid changes profiles of the Coastal Kenya genomes from the reference strains was
investigated using the nextclade tool v0.14.2 (23).

Global contextual sequences

We prepared three sets of contextual sequences from GISAID deposited as of May 2021.
Selected sequences were those with non-ambiguous sampling date, sampled during the period
covered by our study and after alignment possessed <1500 ambiguous nucleotides i.e showed
>95% genome completeness.

a) **Set 1:** We compiled an African SARS-CoV-2 genome set which we assigned Pango
   lineages and used to compare the continental lineage temporal patterns to the Coastal
   Kenya lineage distribution. A total of 12,542 genomes were retrieved for this purpose on
   the 11th of May 2021. From these we also created an Eastern Africa subset which
   comprised of 2,316 genomes from 10 countries, namely, Zimbabwe, Zambia, Uganda,
   Rwanda, Renunion (overseas territory), Mozambique, Malawi, Madagascar, Ethiopia and
   Comoros.

b) **Set 2**: We compiled a global reference set of 9,989 genomes collected across all the six
   inhabited continents between 24th December 2019 and 28th February 2021. Between 23
   and 899 genomes were selected for each included month (mean of 667 and median of
   701) randomly selected across all continents from across 112 countries. These genomes
   were used to infer the global temporal patterns of the lineages observed in Coastal Kenya.
   A subset of these genomes (n=1500) was combined with the Kenyan genomes to infer
   their global phylogenetic context.

c) **Set 3**: We compiled a genotype reference for the top six observed lineages in Coastal
   Kenya. For lineages B.1. and B.1.1 which >5000 genomes exist in GISAID were
   prepared we used the sub-sample from Set 2 above. For lineages B.1.530, B.1.549,
   B.1.596.1 and N.8 we retrieved all genomes assigned these lineages available from
   GISAID to infer their global phylogenetic context.

Phylogenetic analysis

Multiple sequence alignments were prepared in Nextalign v 0.1.6 software and using the initial
Wuhan sequence (Accession number: NC_045512) as the reference. The alignment was
manually inspected in SEAVIEW v4.6.4 to spot any obvious misalignments. Quick non-
bootstrapped neighbor joining trees were created in SEAVIEW to identify any aberrant
sequences that were henceforth discarded.

We reconstructed maximum likelihood (ML) phylogenies using IQTREE v2.1.3. The software
initiates tree reconstruction after assessment and selection of the best model of nucleotide
substitution for the alignment. The ML trees were linked to the various metadata in R
programming software v4.0.2 and visualized using R ggTree v2.4.2. TempEst v1.5.3 was used to
assess the presence of a molecular clock signal in analysed data and linear regression of root-to-
tip genetic distances against sampling dates plotted.

Import/export analysis

The global ML tree topology was used to estimate the number of viral transmission events
between Coastal Kenya and the rest of the world as in (17). The software TreeTime was used to
transform the ML tree topology into a dated tree assuming a constant genomic evolutionary rate
of SARS-CoV-2 of 8.0×10^{-4} nucleotide substitutions per site per year (14). Outlier sequences
were identified by TreeTime and excluded during this process. A migration model was
henceforth fitted to the resulting time-scaled phylogenetic tree from TreeTime, mapping the
location status of the genomes from the six counties at both the tips and internal nodes. Using
the date and location annotated tree topology, we counted the number of transitions between and
within Coastal Kenya counties and the rest of the world and plotted this using ggplot2 v3.3.3.
481
482 *Epidemiological data*
483 The Kenya daily case data for the period between March 2020 and February 2021 was
484 downloaded from Our World in Data database
485 (https://ourworldindata.org/coronavirus/country/kenya). Metadata for the Coastal Kenya samples
486 was compiled from the MoH case investigation forms.
487
Kenya COVID-19 response
We derived status of Kenya government COVID-19 interventions using data from Our World in
Data database that has calculated the Oxford Stringency Index (SI). These SI estimations are
composite measure based on nine response indicators implemented by governments including
school closures; workplace closures; cancellation of public events; restrictions on public
gatherings; closures of public transport; stay-at-home requirements; public information
campaigns; restrictions on internal movements; and international travel controls, rescaled to a
value 0-100, with 100 being strictest (24). The Kenya government revised the policy
interventions at approximately monthly intervals (6) and the changes in SI overtime are shown
in **S1 Figure** .
498

Acknowledgements

We thank (a) the members of the six Coastal counties of Kenya RRTs for collecting the samples
analysed here; (b) the members of the COVID-19 KWTRP Testing Team who tirelessly analysed
the samples received at KWTRP to identify positives (see full list of members below); (c) the
KWTRP data entry team, (d) Laboratories that have shared sequence data on GISAID that we
included as comparison data in our analysis (see list in appendix); (e) the KRISP team in South
Africa for sharing the scripts we used in the import/export analysis and AFRICA-CDC for
facilitating Africa genomics training. This paper is published with permission of the Director of
KEMRI.

508
509 *Members of COVID-19 Testing Team at KWTRP*

Agnes Mutiso, Alfred Mwanzu, Angela Karani, Bonface M. Gichuki, Boniface Karia, Brian Bartilol, Brian Tawa, Calleb Odundo, Caroline Ngetsa, Clement Lewa, Daisy Mugo, David Amadi, David Ireri, Debra Riako, Domtila Kimani, Edwin Machanja, Elijah Gicheru, Elisha Omer, Faith Gambo, Horace Gumba, Isaac Musungu, James Chemweno, Janet Thoya, Jedida Mwacharo, John Gitonga, Johnstone Makale, Justine Getonto, Kelly Ominde, Kelvias Keter, Lydia Nyamako, Margaret Nunah, Martin Mutunga, Metrine Tendwa, Moses Mosobo, Nelson Ouma, Nicole Achieng, Patience Kiyuka, Perpetual Wanjiku, Peter Mwaura, Rita Warui, Robinson Cheruiyot, Salim Mwarumba, Shaban Mwangi, Shadrack Mutua, Susan Njuguna, Victor Osoti, Wesley Cheruiyot, Wilfred Nyamu, Wilson Gumbi and Yiakon Sein

Data availability
Coastal Kenya genomes have been deposited to both GISAID (Accession numbers: EPI_ISL_1039223-29; EPI_ISL1440075-109, IPI_ISL_457845-931; EPI_ISL_568695-872; EPI_ISL806548-717; EPI_ISL_855494-548; EPI_ISL_968807-9245).

Funding
This work was supported by the National Institute for Health Research (NIHR) (project references 17/63/82 (PI Prof. James Nokes) and 16/136/33 (PI Prof. Mark Wooolhouse) using UK aid from the UK Government to support global health research, The UK Foreign, Commonwealth and Development Office and Wellcome Trust (grant# 220985). The views expressed in this publication are those of the author (s) and not necessarily those of NIHR, the Department of Health and Social Care, Foreign Commonwealth and Development Office, Wellcome Trust or the UK government. Some members of COVID-19 Testing Team at KWTRP were supported by funding received by Dr Marta Maia (BOHEMIA study funded UNITAID), Dr Francis Ndungu (Senior Fellowship and Research and Innovation Action (RIA) grants from EDCTP) and Prof. Anthony Scott (PCIVS grant from GAVI).

References
1. Dong E, Du H, Gardner L. 2020. An interactive web-based dashboard to track COVID-19 in real time. Lancet Infect Dis 20:533-534.
2. MOH. 2021. Update on COVID-19 in Kenya doi:https://www.health.go.ke/press-releases/. MOH.
3. Uyoga S, Adetifa IMO, Karanja HK, Nyagwange J, Tuju J, Wanjiku P, Aman R, Mwangangi M, Amoth P, Kasera K, Ng'ang'a W, Rombo C, Yegon C, Kithi K, Odhiambo E, Rotich T, Orgut I, Kihara S, Otiende M, Bottomley C, Mupe ZN, Kagucia EW, Gallagher KE, Etyang A, Voller S, Gitonga JN, Mugo D, Agoti CN, Otieno E, Ndwiga L, Lambe T, Wright D, Barasa E, Tsofa B, Bejon P, Ochola-Oyier LI, Agweyu A, Scott JAG, Warimwe GM. 2021. Seroprevalence of anti-SARS-CoV-2 IgG antibodies in Kenyan blood donors. Science 371:79-82.

4.   Etyang AO, Lucinde R, Karanja H, Kalu C, Mugo D, Nyagwange J, Gitonga J, Tuju J,
Wanjiku P, Karani A, Mutua S, Maroko H, Nzomo E, Maitha E, Kamuri E, Kaugiria T,
Weru J, Ochola LB, Kilimo N, Charo S, Emukule N, Moracha W, Mukabi D, Okuku R,
Ogutu M, Angujo B, Otiende M, Bottomley C, Otieno E, Ndwiga L, Nyaguara A, Voller
S, Agoti C, Nokes DJ, Ochola-Oyier LI, Aman R, Amoth P, Mwangangi M, Kasera K,
Ng'ang'a W, Adetifa I, Kagucia EW, Gallagher K, Uyoga S, Tsofa B, Barasa E, Bejon P,
Scott JAG, Agweyu A, Warimwe G. 2021. Seroprevalence of Antibodies to SARS-CoV-
2 among Health Care Workers in Kenya. Clin Infect Dis doi:10.1093/cid/ciab346.
5.   Adetifa IMO, Uyoga S, Gitonga JN, Mugo D, Otiende M, Nyagwange J, Karanja HK,
Tuju J, Wanjiku P, Aman R, Mwangangi M, Amoth P, Kasera K, Ng'ang'a W, Rombo C,
Yegon C, Kithi K, Odhiambo E, Rotich T, Orgut I, Kihara S, Bottomley C, Kagucia EW,
Gallagher KE, Etyang A, Voller S, Lambe T, Wright D, Barasa E, Tsofa B, Bejon P,
Ochola-Oyier LI, Agweyu A, Scott JAG, Warimwe GM. 2021. Temporal trends of
SARS-CoV-2 seroprevalence during the first wave of the COVID-19 epidemic in Kenya.
Nature Communications 12:3966.
6.   Brand SPC, Ojal J, Aziza R, Were V, Okiro EA, Kombe IK, Mburu C, Ogero M,
Agweyu A, Warimwe GM, Nyagwange J, Karanja H, Gitonga JN, Mugo D, Uyoga S,
Adetifa IMO, Scott JAG, Otieno E, Murunga N, Otiende M, Ochola-Oyier LI, Agoti CN,
Githinji G, Kasera K, Amoth P, Mwangangi M, Aman R, Ng'ang'a W, Tsofa B, Bejon P,
Keeling MJ, Nokes DJ, Barasa E. 2021. COVID-19 Transmission Dynamics Underlying
Epidemic Waves in Kenya. medRxiv
doi:10.1101/2021.06.17.21259100:2021.06.17.21259100.
7.   Githinji G, de Laurent ZR, Mohammed KS, Omuoyo DO, Macharia PM, Morobe JM,
Otieno E, Kinyanjui SM, Agweyu A, Maitha E, Kitole B, Suleiman T, Mwakinangu M,
Nyambu J, Otieno J, Salim B, Kasera K, Kiiru J, Aman R, Barasa E, Warimwe G, Bejon
P, Tsofa B, Ochola-Oyier LI, Nokes DJ, Agoti CN. 2020. Tracking the introduction and
spread of SARS-CoV-2 in coastal Kenya. medRxiv
doi:10.1101/2020.10.05.20206730:2020.10.05.20206730.
8.   Geoghegan JL, Ren X, Storey M, Hadfield J, Jelley L, Jefferies S, Sherwood J, Paine S,
Huang S, Douglas J, Mendes FK, Sporle A, Baker MG, Murdoch DR, French N,
Simpson CR, Welch D, Drummond AJ, Holmes EC, Duchêne S, de Ligt J. 2020.
Genomic epidemiology reveals transmission patterns and dynamics of SARS-CoV-2 in
Aotearoa New Zealand. Nature Communications 11:6351.
9.   Worobey M, Pekar J, Larsen BB, Nelson MI, Hill V, Joy JB, Rambaut A, Suchard MA,
Wertheim JO, Lemey P. 2020. The emergence of SARS-CoV-2 in Europe and North
America. Science 370:564-570.
10.   Oude Munnink BB, Nieuwenhuijse DF, Stein M, O'Toole Á, Haverkate M, Mollers M,
Kamga SK, Schapendonk C, Pronk M, Lexmond P, van der Linden A, Bestebroer T,
Chestakova I, Overmars RJ, van Nieuwkoop S, Molenkamp R, van der Eijk AA,
GeurtsvanKessel C, Vennema H, Meijer A, Rambaut A, van Dissel J, Sikkema RS,

Timen A, Koopmans M. 2020. Rapid SARS-CoV-2 whole-genome sequencing and analysis for informed public health decision-making in the Netherlands. Nat Med 26:1405-1410.

11. Bugembe DL, Kayiwa J, Phan MVT, Tushabe P, Balinandi S, Dhaala B, Lexow J, Mwebesa H, Aceng J, Kyobe H, Ssemwanga D, Lutwama J, Kaleebu P, Cotten M. 2020. Main Routes of Entry and Genomic Diversity of SARS-CoV-2, Uganda. Emerg Infect Dis 26:2411-2415.

12. Konings F, Perkins MD, Kuhn JH, Pallen MJ, Alm EJ, Archer BN, Barakat A, Bedford T, Bhiman JN, Caly L, Carter LL, Cullinane A, de Oliveira T, Druce J, El Masry I, Evans R, Gao GF, Gorbalenya AE, Hamblion E, Herring BL, Hodcroft E, Holmes EC, Kakkar M, Khare S, Koopmans MPG, Korber B, Leite J, MacCannell D, Marklewitz M, Maurer-Stroh S, Rico JAM, Munster VJ, Neher R, Munnink BO, Pavlin BI, Peiris M, Poon L, Pybus O, Rambaut A, Resende P, Subissi L, Thiel V, Tong S, van der Werf S, von Gottberg A, Ziebuhr J, Van Kerkhove MD. 2021. SARS-CoV-2 Variants of Interest and Concern naming scheme conducive for global discourse. Nature Microbiology doi:10.1038/s41564-021-00932-w.

13. Bugembe DL, Phan MVT, Ssewanyana I, Semanda P, Nansumba H, Dhaala B, Nabadda S, O'Toole ÁN, Rambaut A, Kaleebu P, Cotten M. 2021. Emergence and spread of a SARS-CoV-2 lineage A variant (A.23.1) with altered spike protein in Uganda. Nature Microbiology doi:10.1038/s41564-021-00933-9.

14. Sagulenko P, Puller V, Neher RA. 2018. TreeTime: Maximum-likelihood phylodynamic analysis. Virus Evol 4:vex042.

15. Rice BL, Annapragada A, Baker RE, Bruijning M, Dotse-Gborgbortsi W, Mensah K, Miller IF, Motaze NV, Raherinandrasana A, Rajeev M, Rakotonirina J, Ramiadantsoa T, Rasambainarivo F, Yu W, Grenfell BT, Tatem AJ, Metcalf CJE. 2021. Variation in SARS-CoV-2 outbreaks across sub-Saharan Africa. Nature Medicine 27:447-453.

16. Baric RS. 2020. Emergence of a Highly Fit SARS-CoV-2 Variant. New England Journal of Medicine 383:2684-2686.

17. Wilkinson E, Giovanetti M, Tegally H, San JE, Lessels R, Cuadros D, Martin DP, Zekri A-RN, Sangare AK, Ouedraogo A-S, Sesay AK, Hammami A, Amuri AA, Sayed A, Rebai A, Elargoubi A, Trotter AJ, Keita AK, Sall AA, Kone A, Souissi A, Gutierrez AV, Page AJ, Iranzadeh A, Lambisia A, Sylverken A, Ibrahimi A, Dhaala B, Kouriba B, Kleinhans B, Brook C, Williamson C, Pratt CB, Akoua-Koffi CG, Agoti CN, Morang'a CM, Nokes DJ, Bridges DJ, Bugembe DL, Baker D, Doolabh D, Ssemwanga D, Tshabuila D, Bassirou D, Amuzu DSY, Goedhals D, Maruapula D, Foster-Nyarko E, Lusamaki EK, Simulundu E, et al. 2021. A year of genomic surveillance reveals how the SARS-CoV-2 pandemic unfolded in Africa. medRxiv doi:10.1101/2021.05.12.21257080:2021.05.12.21257080.

18. Tegally H, Wilkinson E, Lessells RJ, Giandhari J, Pillay S, Msomi N, Mlisana K, Bhiman JN, von Gottberg A, Walaza S, Fonseca V, Allam M, Ismail A, Glass AJ,

629		Engelbrecht S, Van Zyl G, Preiser W, Williamson C, Petruccione F, Sigal A, Gazy I,
630		Hardie D, Hsiao NY, Martin D, York D, Goedhals D, San EJ, Giovanetti M, Lourenço J,
631		Alcantara LCJ, de Oliveira T. 2021. Sixteen novel lineages of SARS-CoV-2 in South
632		Africa. Nat Med 27:440-446.
633	19.	Tegally H, Wilkinson E, Giovanetti M, Iranzadeh A, Fonseca V, Giandhari J, Doolabh D,
634		Pillay S, San EJ, Msomi N, Mlisana K, von Gottberg A, Walaza S, Allam M, Ismail A,
635		Mohale T, Glass AJ, Engelbrecht S, Van Zyl G, Preiser W, Petruccione F, Sigal A,
636		Hardie D, Marais G, Hsiao NY, Korsman S, Davies MA, Tyers L, Mudau I, York D,
637		Maslo C, Goedhals D, Abrahams S, Laguda-Akingba O, Alisoltani-Dehkordi A, Godzik
638		A, Wibmer CK, Sewell BT, Lourenço J, Alcantara LCJ, Kosakovsky Pond SL, Weaver S,
639		Martin D, Lessells RJ, Bhiman JN, Williamson C, de Oliveira T. 2021. Detection of a
640		SARS-CoV-2 variant of concern in South Africa. Nature 592:438-443.
641	20.	Agoti C, Mutunga M, Lambisia A, Kimani D, Cheruiyot R, Kiyuka P, Lewa C, Gicheru
642		E, Tendwa M, Said Mohammed K, Osoti V, Makale J, Tawa B, Odundo C, Cheruiyot W,
643		Nyamu W, Gumbi W, Mwacharo J, Nyamako L, Otieno E, Amadi D, Thoya J, Karani A,
644		Mugo D, Musyoki J, Gumba H, Mwarumba S, M. Gichuki B, Njuguna S, Riako D,
645		Mutua S, Gitonga J, Sein Y, Bartilol B, Mwangi S, O. Omuoyo D, M. Morobe J, de
646		Laurent Z, Bejon P, Ochola-Oyier L, Tsofa B. 2021. Pooled testing conserves SARS-
647		CoV-2 laboratory resources and improves test turn-around time: experience on the
648		Kenyan Coast [version 2; peer review: 3 approved]. Wellcome Open Research 5.
649	21.	Tyson JR, James P, Stoddart D, Sparks N, Wickenhagen A, Hall G, Choi JH, Lapointe H,
650		Kamelian K, Smith AD, Prystajecky N, Goodfellow I, Wilson SJ, Harrigan R, Snutch TP,
651		Loman NJ, Quick J. 2020. Improvements to the ARTIC multiplex PCR method for
652		SARS-CoV-2 genome sequencing using nanopore. bioRxiv
653		doi:10.1101/2020.09.04.283077.
654	22.	Rambaut A, Holmes EC, O'Toole Á, Hill V, McCrone JT, Ruis C, du Plessis L, Pybus
655		OG. 2020. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist
656		genomic epidemiology. Nat Microbiol 5:1403-1407.
657	23.	Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, Sagulenko P,
658		Bedford T, Neher RA. 2018. Nextstrain: real-time tracking of pathogen evolution.
659		Bioinformatics 34:4121-4123.
660	24.	Hale T, Angrist N, Boby T, Cameron-Blake E, Hallas L, Kira B, Majumdar S, Petherick
661		A, Phillips T, Tatlow H, Webster S. December 2020 Noam Angrist. Variation in
662		government responses to COVID-19.
663		doi:https://www.bsg.ox.ac.uk/research/publications/variation-government-responses-
664		covid-19doi:https://www.bsg.ox.ac.uk/research/publications/variation-government-
665		responses-covid-19.

SUPPLEMENTAL MATERIAL

668
S1 Figure. The SARS-CoV-2 epidemic in Kenya. Panel A, Map of Kenya showing county
administrative boundaries and neighboring countries. The map is colored by the number of cases
documented in the counties as at of 26th February 2021. Panel, B. Reported daily new cases in
Kenya from March 2020 to February 2021 showing the first two national SARS-CoV-2 infection
waves. Panel C, daily cases reported across coastal Kenya during the study period. Panel D,
Kenya government intervention levels during the study period as measured by the Oxford
stringency index.

S2 Figure. Position of non-synonymous mutations in the 684 genomes from coastal Kenya

compare with the Wuhan reference genome (Accession number NC_045512). Only changes

occurring in >1% of genomes are shown. The percentage frequency of the changes in the 684

genomes is shown on the y-axis.

S3 Figure. Genomic characteristics of the four lineages first identified in Kenya. Panel A,

root-to-tip regression plots for the four lineages first identified in Kenya. Panel B, Genome maps

of the "Kenya" lineages where the spike region is shown in detail and in color and the rest of the

genome is shown in grey color.

S1 Table. Lineages observed in Coastal Kenya, their global history and VOC/VOI status

Lineage	Frequency (%)	Mombasa	Kilifi	Kwale	Taita Taveta	Tana River	Lamu	Earliest date (Global)	Total Global #	Description
A	21 (3.1)	3	–	13	5	–	–	2019-12-30	2930	Root of the pandemic lies within lineage A.
A.23	3 (0.4)	1	1	1	–	–	–	2020-08-14	97	Ugandan lineage
A.23.1	3 (0.4)	1	–	1	1	–	–	2020-10-21	957	International lineage
A.25	2 (0.3)	2	–	–	–	–	–	2020-06-22	34	Ugandan lineage
B	4 (0.6)	3	1	–	–	–	–	2019-12-24	5214	Second major haplotype
B.1	413 (60.4)	241	69	28	56	9	10	2020-01-24	75776	A large European lineage
B.1.1	61 (8.9)	39	5	4	13	–	–	2020-02-15	46205	European lineage
B.1.1.1	4 (0.6)	1	1	2	–	–	–	2020-03-04	2529	England
B.1.1.111	1 (0.2)	–	–	1	–	–	–	2020-04-04	59	African lineage (Zambia, Zimbabwe)
B.1.1.254	2 (0.3)	1	–	1	–	–	–	2020-03-14	76	Mostly South Africa
B.1.1.33	1 (0.2)	1	–	–	–	–	–	2020-03-01	1830	Brazillian lineage
B.1.1.485	1 (0.2)	–	–	–	1	–	–	2020-04-09	86	Ghanian Lineage
B.1.1.7	1 (0.2)	1	–	–	–	–	–	2020-02-07	694561	Alpha variant of concern, UK first identified
B.1.1.70	1 (0.2)	–	–	–	1	–	–	2020-03-06	2596	European Lineage
B.1.153	1 (0.2)	–	–	1	–	–	–	2020-01-24	267	English lineage
B.1.177	3 (0.4)	–	1	1	–	–	1	2020-02-14	68271	Major mostly European lineage
B.1.179	2 (0.3)	2	–	–	–	–	–	2020-03-09	256	Denmark
B.1.340	1 (0.2)	1	–	–	–	–	–	2020-04-06	136	USA lineage
B.1.351	10 (1.5)	–	3	3	4	–	–	2020-05-11	18425	Beta variant of concern
B.1.378	2 (0.3)	2	–	–	–	–	–	2020-03-14	190	USA
B.1.393	2 (0.3)	1	1	–	–	–	–	2020-07-04	32	Ugandan lineage
B.1.450	1 (0.2)	–	1	–	–	–	–	2020-03-14	126	USA (TX)
B.1.526	1 (0.2)	–	–	1	–	–	–	2020-11-23	18387	Iota variant of interest
B.1.530	44 (6.4)	8	7	–	28	–	1	2020-10-11	60	Kenyan
B.1.535	1 (0.2)	1	–	–	–	–	–	2020-03-25	36	Australian
B.1.549	54 (7.9)	18	24	7	5	–	–	2020-06-26	76	Kenya + England lineage
B.1.553	1 (0.2)	–	1	–	–	–	–	-	-	-
B.1.558	1 (0.2)	1	–	–	–	–	–	2020-04-06	167	USA/ Mexico lineage
B.1.596.1	8 (1.2)	4	2	1	1	–	–	2020-08-03	29	Kenyan lineage
B.4	1 (0.2)	1	–	–	–	–	–	2020-01-18	374	Iranian lineage and global export
B.4.7	3 (0.4)	3	–	–	–	–	–	2020-03-14	68	Africa and UAE lineage
N.8	30 (4.4)	1	1	–	–	–	28	2020-06-23	19	Alias of B.1.1.33.8, Kenyan lineage