Transmission networks of SARS-CoV-2 in coastal Kenya during the first two waves: a retrospective genomic study

Charles N. Agoti[1,2,#], Lynette Isabella Ochola-Oyier[1], Khadija Said Mohammed[1], Arnold W. Lambisia[1], Zaydah R. de Laurent[1], John M. Morobe[1], Maureen W. Mburu[1], Donwilliams O. Omuoyo[1], Edidah M. Ongera[1], Leonard Ndwiga[1], Eric Maitha[3], Benson Kitole[3], Thani Suleiman[3], Mohamed Mwakinangu[3], John Nyambu[3], John Otieno[3], Barke Salim[3], Jennifer Musyoki[1], Nickson Murunga[1], Edward Otieno[1], John Kiiru[3], Kadondi Kasera[3], Patrick Amoth[3], Mercy Mwangangi[3], Rashid Aman[3], Samson Kinyanjui[1,2,4], George Warimwe[1,4], My Phan[5], Ambrose Agweyu[1], Matthew Cotten[5,6], Edwine Barasa[1], Benjamin Tsofa[1], D. James Nokes[1,7], Philip Bejon [1,4], George Githinji[1]

Affiliations
1. Kenya Medical Research Institute (KEMRI)-Wellcome Trust Research Programme (KWTRP), Kilifi, Kenya.
2. Pwani University, Kilifi, Kenya.
3. Ministry of Health, Kenya.
4. Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom.
5. Medical Research Centre (MRC)/ Uganda Virus Research Institute, Entebbe, Uganda.
6. MRC-University of Glasgow Centre for Virus Research, Glasgow, UK.
7. University of Warwick, Coventry, United Kingdom.

[#]**Corresponding author:** Email: cnyaigoti@kemri-wellcome.org)

ABSTRACT
Background: The transmission networks of SARS-CoV-2 in sub-Saharan Africa remain poorly understood.
Methods: We undertook phylogenetic analysis of 747 SARS-CoV-2 positive samples collected across six counties in coastal Kenya during the first two waves (March 2020 - February 2021). Viral imports and exports from the region were inferred using ancestral state reconstruction (ASR) approach.
Results: The genomes were classified into 35 Pango lineages, six of which accounted for 79% of the sequenced infections: B.1 (49%), B.1.535 (11%), B.1.530 (6%), B.1.549 (4%), B.1.333 (4%) and B.1.1 (4%). Four identified lineages were Kenya specific. In a contemporaneous global subsample, 990 lineages were documented, 261 for Africa and 97 for Eastern Africa. ASR analysis identified >300 virus location transition events during the period, these comprising: 69 viral imports into Coastal Kenya; 93 viral exports from coastal Kenya; and 191 inter-county import/export events. Most international viral imports (58%) and exports (92%) occurred through Mombasa City, a key touristic and commercial Coastal Kenya center; and many occurred prior to June 2020, when stringent local COVID-19 restriction measures were enforced. After this period, local virus transmission dominated, and distinct local phylogenies were seen.
Conclusions: Our analysis supports moving control strategies from a focus on international travel to local transmission.

Funding: This work was funded by Wellcome (grant#: 220985) and the National Institute for Health Research (NIHR), project references: 17/63/and 16/136/33 using UK aid from the UK Government to support global health research, The UK Foreign, Commonwealth and Development Office.

INTRODUCTION

By 31st August 2021, at least 218 million cases of COVID-19 and > 4.5 million associated deaths were reported worldwide. By the same date, Kenya, an East Africa nation with a population of ~50 million had reported 235,863 cases of SARS-CoV-2 and 4,726 associated deaths (MOH, 2021). Serological surveys indicated that Kenya's COVID-19 epidemic had progressed further than could be discerned from the limited laboratory testing case reports that were available during the period (Etyang et al., 2021; Uyoga et al., 2021a). In the period after the peak of Kenya's second wave (January-March 2021), nationwide anti-SARS-CoV-2 IgG prevalence based on analysis of blood donor samples was estimated to be 48.5% (Adetifa et al., 2021; Uyoga et al., 2021b).

In August 2021, 18 months after the first report of SARS-CoV-2 occurrence in Kenya, the country had experienced four major waves of SARS-CoV-2 infections (Brand et al., 2021). Despite this progression, the local SARS-CoV-2 spread patterns, like in many sub-Saharan Africa settings, remained poorly understood (Wilkinson et al.). The analysis presented here examined genome sequences from the first two waves in Coastal Kenya. Overall, the documented SARS-CoV-2 infections in the country during these early waves were concentrated in cities, especially Nairobi (~42%), the capital, and Mombasa (~8%), the country's second largest city located on the Indian ocean coast (**Figure 1A-C**). In addition to Mombasa city, there are five other counties that make up Kenya's Coastal region (Kilifi, Kwale, Taita Taveta, Tana River and Lamu) which has a total population size of ~4.3 million.

Throughout the COVID-19 pandemic, genomic surveillance has been critical for tracking the spread of SARS-CoV-2 and investigating its evolutionary changes (Bugembe et al., 2020; Geoghegan et al., 2020; Oude Munnink et al., 2020; Worobey et al., 2020). In-country SARS-CoV-2 genome sequencing started in Kenya soon after the identification of the initial case and has remained a key tool informing public health response and tracking the progress of the local epidemic. We have previously reported on the genomic analysis of SARS-CoV-2 in Coastal Kenya during the early phase of the pandemic (up to July 2020) revealing introduction multiple lineages but with mainly Pango lineage B.1 establishing local transmission (Githinji et al., 2021). Here we investigated the lineage composition and diversity of the subsequent two waves (up to February 2021) and examine the patterns of further viral introductions, local evolution and spread of SARS-CoV-2 across the six Coastal counties of Kenya.

RESULTS

Infection waves in Coastal Kenya counties
The first cases of COVID-19 in Coastal Kenya were reported in Mombasa and Kilifi counties in March 2020. This was after initial detection in Nairobi City on 13th March 2020. Following this, several COVID-19 related containment measures were put in place in March /April 2020 to curtail the spread of the virus including ban of movement into and out of infected areas. The national Oxford Stringency index in Kenya in April 2020 was at 89 (Brand et al., 2021), **Figure 1D**. However, two months later (June 2020), a steady removal of these restriction measures started leading to a SI of 75 by August 2020 and a SI <50 by October 2020. At the peak of the second wave, some of the restriction measures were reinstated increasing the SI to >=50, **Figure 1D**.

Figure 1. The SARS-CoV-2 epidemic in Kenya. Panel A, map showing administrative boundaries of the 47 counties in Kenya and the neighboring countries. The map is colored by the cumulative number of SARS-CoV-2 cases documented in each county as of 26[th] February 2021. Panel B, reported daily new cases, 7-day-rolling average, in Kenya from March 2020 to February 2021 showing the first two national SARS-CoV-2 infection waves. Panel C. reported daily cases, 7-day-rolling average, across coastal Kenya during the study period. Panel D, Kenya government COVID-19 intervention levels during the study period as summarized by the Oxford stringency index (abbreviated SI, see details in materials and methods section).

Mombasa county endured the earliest surge of SARS-CoV-2 cases in coastal Kenya with its first peak occurring in June 2020, **Figure 2A**. This was earlier than the first national peak that occurred in July-August of 2020. By 26[th] February 2021 (the most recent date included in our analysis) three of the six coastal Kenya counties (i.e. Mombasa, Lamu and Taita Taveta) had experienced at least two major infection peaks while the other three (i.e. Kilifi, Kwale and Tana River) had experienced only a single peak**.** During this period covered, the Kenya Ministry of Health (MoH) reported 12,307 laboratory confirmed SARS-CoV-2 cases for all the six Coastal counties: Mombasa (n=8,450, 68.7%), Kilifi (n=2,458, 20.0%), Taita Taveta (n=567, 4.6%), Kwale (n=437, 3.6%), Lamu (n=309, 2.5%) and Tana River (n=85, 0.7%)**, Figure 2A.**

Testing and sequencing at KWTRP
The flow of samples analyzed in this study is provided in **Supplementary-Figure 1.** Between 17[th] March 2020 and 26[th] February 2021, the KEMRI-Wellcome Trust Research Programme (KWTRP) tested at total of 82,716 nasopharyngeal/oropharyngeal (NP/OP) swab samples from the six Coastal counties. Of these, 6,307 (7.6%) were determined as SARS-CoV-2 positive (approximately half of the total reported by MoH), distributed by month as in **Figure 2B**. Overall, a total of 3,137 (6.8%) tests at KWTRP were positive from samples taken in Mombasa, 1,142 (8.8%) for Kilifi, 657 (4.5%) for Taita Taveta, 349 (12.7%) for Lamu, 437 (7.9%) for Kwale and 105 (12.0%) for Tana River.

128
Figure 2. SARS-CoV-2 identified cases on the Kenyan Coast and testing and sequencing
at KEMRI-Wellcome Trust Research Programme (KWTRP). Panel A, epidemic curves
for each of the six Coastal Kenya counties derived from the daily positive case numbers, 7-
day-rolling average, reported by the Ministry of Health. Panel B, monthly count of SARS-
CoV-2 RT-PCR tests undertaken at the KWTRP and those positive during the study period.
Panel C, monthly proportion of samples sequenced from total SARS-CoV-2 positives detected
at KWTRP. Panel D, Distribution of the sequenced 747 samples by study period and across the
6 Coastal Kenya counties.

137
Of the identified 3,137 RT-PCR positive samples, 747 (23.8%) were genome sequenced,
distributed as follows; Mombasa: 367 (11.7%), Kilifi: 119 (10.4%), Taita Taveta: 139 (21.2%),
Kwale: 68 (15.6%), Lamu: 43 (12.4%) and Tana River: 11 (10.5%), **Figure 2C**. The multiple
sequence alignment of these coastal Kenya genomes and observed differences from the
Wuhan-Hu-1 isolate is shown in **Supplementary-Figure 2**. The sequences were spread across
the study period with 272 (36.4%) from the introduction phase (17th March to 20th May 2020
when nationally <50 new cases were reported per day), 223 (29.9%) from wave one (21st May-
15th September 2020) and 252 (33.7%) from wave two (16th September 2020 to 26th February
2021), **Figure 1**. Overall, our sequencing rate corresponded to approximately one sequence for
every 16 confirmed cases.
148

Most of the sequenced isolates were from males (n=443, 59.3%), persons aged between 30-60
years (n=420, 56.2%) and persons of Kenyan nationality (n=509, 68.1%), **Table 1**. The second
largest nationality of the individuals providing sequenced samples was Tanzania (n=34)
followed by South Africa (n=5). A total of 110 samples (14.7%) had been collected from
persons with a recent international travel history (<14 days) but a considerable proportion of
the sequenced samples were obtained from persons whose travel history information was
missing (n=373, 49.9%).
156

Lineage dynamics in Coastal Kenya
As a reference, we adopted the PANGOLIN (Phylogenetic Assignment of Named Global
LINeages) lineage nomenclature (Rambaut et al., 2020). The 747 genomes were classified into
35 Pango lineages, including four first identified in Kenya (N.8, B.1.530, B.1.549 and
B.1.596.1), **Table 2 and Figure 3A**. During the introductions phase we detected 14 lineages
while during wave one we detected 21 lineages (10 of which were being identified locally for
the first time) and during wave two we detected 19 lineages (seven of which were being
identified locally for the first time), **Figure 3B**. Seventeen lineages (48.6%) were identified in
three or more samples with the top six lineages accounting for 78.8% of the sequenced
infections; B.1 (n=369, 49.4%, European lineage which predominated in the Northern Italy
outbreak early in 2020), B.1.535 (n=82, Australian lineage), B.1.530 (n=42, 5.6%, Kenyan
lineage), B.1.530 (n=42, 6%, Kenyan lineage), B.1.549 (n=33, 4.4%, Kenyan lineage), B.1.333
(n=32, 4.3%, Washington USA lineage), B.1.1 (n=31, 4.1%, European lineage).
170

Table 1: Demographics of the sequenced SARS-CoV-2 cases in coastal Kenya

Characteristic	Total cases (%)	Introductions phase	Wave 1	Wave 2
Age category				
0-9	16 (2.1	5	6	5
10-19	40 (5.4)	11	10	19
20-29	139 (18.6)	52	39	48
30-39	195 (26.1)	75	47	73
40-49	137 (18.3)	48	40	49
50-59	88 (11.8)	45	12	31
60-69	31 (4.1)	17	6	8
70-79	12 (1.6)	4	3	5
80+	4 (0.5)	3	-	1
Missing	85 (11.4)	12	60	13
Gender				
Female	203 (27.2)	81	43	79
Male	443 (59.3)	167	118	158
Missing	101 (13.5)	24	62	15
Nationality				
Kenyan	509 (68.1)	174	138	197
Tanzania	34 (4.5)	18	7	9
South Africa	5 (0.1)	0	0	5
Other[#]	20 (2.7)	3	4	13
Missing	179 (24.0)	77	74	28
Travel History[¥]				
Yes	110 (14.7)	55	28	27
No	264 (35.3)	108	81	75
Missing	373 (49.9)	109	114	150

172 [#] The other nationalities included Ethiopia (4), Rwanda(2), Sweden (2), USA (2), Britain (2), India (2), Pakistan (1, Burundi (1), Poland (1), Eritrea (1),

173 Uganda (1) and Zambia (1) ;[¥] Defined as having moved into Kenya in the previous 14 days or sampled at a point of entry (POE) into Kenya.

176 **Table 2.** Lineages observed in coastal Kenya, their county distribution, global history and VOC/VOI status

Lineage	Frequency (%)	Mombasa	Kilifi	Kwale	Taita Taveta	Tana River	Lamu	Earliest date	Number assigned	Description
A	22 (2.9)	4	-	13	5	-	-	2019-12-30	2164	Root of the pandemic lies within lineage A. Many sequences originating from China
A.23	2 (0.3)	-	1	1	-	-	-	2020-06-08	99	Ugandan lineage
A.23.1	6 (0.8)	2	1	1	2	-	-	2020-10-21	1109	International lineage with a number of variants of potential biological concern, including 681R
A.25	2 (0.3)	2	-	-	-	-	-	2020-06-08	46	Ugandan lineage
AL.1	4 (0.5)	3	-	-	1	-	-	2020-02-23	1268	Alias of B.1.1.231.1, Canada lineage
B	11 (1.5)	10	1	-	-	-	-	2019-12-24	5408	Second major haplotype (and first to be discovered)
B.1	369 (49.4)	201	65	19	65	8	11	2020-01-24	71452	A large European lineage the origin of which roughly corresponds to the Northern Italian outbreak early in 2020.
B.1.1	31 (4.1)	12	3	4	12	-	-	2020-02-16	42839	European lineage with 3 clear SNPs `28881GA`,`28882GA`,`28883GC`
B.1.1.1	4 (0.5)	1	1	2	-	-	-	2020-03-02	2796	England
B.1.1.33	13 (1.7)	2	1	-	-	-	10	2020-03-01	2039	Brazillian
B.1.1.7	2 (0.3)	2	-	-	-	-	-	2020-02-07	986233	UK lineage of concern, associated with the N501Y mutation.
B.1.13	1 (0.1)	1	-	-	-	-	-	2020-03-09	398	UK lineage
B.1.153	1 (0.1)	-	-	1	-	-	-	2020-03-01	773	English lineage with some Australian, New Zealand and European sequences
B.1.157	2 (0.3)	2	-	-	-	-	-	2020-02-03	324	UK/ Spanish lineage
B.1.160	2 (0.3)	-	-	1	-	-	1	2020-02-06	24207	Large European lineage, sequences from BeNeLuX, Denmark, Switzerland, Hungary and UK.
B.1.177	1 (0.1)	-	1	-	-	-	-	2020-02-14	69319	Major mostly European lineage, result of opening borders summer 2020
B.1.179	2 (0.3)	2	-	-	-	-	-	2020-03-09	231	Denmark
B.1.222	2 (0.3)	2	-	-	-	-	-	2020-02-24	562	Scottish lineage
B.1.229	2 (0.3)	2	-	-	-	-	-	2020-03-19	104	UK lineage
B.1.281	1 (0.1)	1	-	-	-	-	-	2020-04-08	41	Bahrain lineage
B.1.333	32 (4.3)	27	2	3	-	-	-	2020-03-03	188	USA lineage WA
B.1.351	25 (3.3)	6	4	8	7	-	-	2020-03-27	25467	Lineage of concern detected in South Africa
B.1.393	1 (0.1)	-	1	-	-	-	-	2020-05-29	37	Uganda
B.1.525	1 (0.1)	-	1	-	-	-	-	2020-12-11	6729	International lineage with E484K, del69-70 among other defining mutations.
B.1.530	42 (5.6)	8	7	-	26	-	1	2020-10-20	54	Kenya
B.1.535	82 (11.0)	50	11	10	10	1	-	2020-02-03	2573	Australia

B.1.549	33 (4.4)	11	17	3	2	-	-	2020-07-17	51	Kenya + England
B.1.551	2 (0.3)	-	-	-	2	-	-	2020-07-31	751	USA lineage
B.1.558	1 (0.1)	1	-	-	-	-	-	2020-04-06	190	USA/ Mexico lineage
B.1.593	2 (0.3)	-	-	-	-	2	-	2020-08-27	90	USA lineage
B.1.596.1	15 (2.0)	10	2	2	1	-	-	2020-09-07	54	Kenyan lineage
B.1.612	7 (0.9)	-	-	-	6	-	1	2020-03-11	380	USA lineage, split out from B.1.361
B.4	3 (0.4)	3	-	-	-	-	-	2020-01-18	376	Iranian lineage and global exports
B.4.7	1 (0.1)	1	-	-	-	-	-	2020-03-14	68	Africa and UAE
N.8	20 (2.7)	1	-	-	-	-	19	2020-06-23	14	Alias of B.1.1.33.8, Kenyan

177
178
179

Lineage B.1 was the first to be detected in the region and comprised the initial cases identified
in Mombasa, Kilifi, Kwale and Tana River, **Figure 3A and C**. This lineage comprised 48.2%
(n=131) of the viruses we sequenced from the introduction phase, 58.7% (n=131) from wave
one and 42.5% (n=107) from wave two. Overall, B.1 accounted for most of the virus sequences
detected in all counties except Lamu where lineages B.1.1.33 and N.8 (alias lineage B.1.1.33.8)
were most frequent (together accounting for 67.5% of Lamu cases, especially wave one, first
detected there on 23[rd] June 2020), **Table 2**. Only three additional sequenced cases of lineage
B.1.1.33 occurred outside Lamu County (two Mombasa and one in Kilifi) and only one
additional sequenced case of lineage N.8 occurred outside Lamu County (in Mombasa).
189

190
Figure 3. Lineage introductions and temporal dynamics in Coastal Kenya. Panel A,
SARS-CoV-2 Pango lineages in the sequenced 747 coastal Kenya samples and timing of
detections. Panel B, Lineage detections by study phase. Intr. phase refers to introduction phase.
Panel C, Biweekly distribution of the common lineages identified across the six counties
presented as proportions. Lineages detected in less than three cases were put together and

referred to as "other lineages". This group included 18 lineages namely: A.25, AL.1, B.1.1.1,
B.1.13, B.1.153, B.1.157, B.1.160, B.1.177, B.1.179, B.1.222, B.1.229, B.1.281, B.1.393,
B.1.551, B.1.558, B.1.593, B.4 and B.4.7.
199
During wave one, 18 cases of lineage A were observed. Most (56%) of these were identified
in Kwale county (one of the Kenya counties bordering Tanzania, **Figure 1A**), and the lineage
comprised 19% of all cases identified Kwale county, **Figure 3C.** Only two other counties
observed lineage A cases: Mombasa (n=4) and Taita Taveta (n=5). Lineage B.1.535, which
was the second most common lineage in the region was mostly observed in during wave one
and predominantly in Mombasa County with limited number of cases observed in Kilifi, Kwale
and Taita Taveta, **Table 2**.
207
During the wave two period, there was an emergence of three lineages first identified in Kenya:
(a) lineage B.1.549 (first observed 17th July 2020) which was responsible for a significant
number of the sequenced infections in Kilifi (n=17), Mombasa (n=11), Taita Taveta (n=2), and
Kwale (n=3), (b) lineage B.1.596.1 (first observed 03rd August 2020) and detected in
Mombasa, Kilifi, Kwale and Taita-Taveta and (c) lineage B.1.530 (first observed 15th
September 2020) that predominated infections in Taita Taveta (n=26) with a fewer cases
observed in Mombasa (n=8), Kilifi (n=7) and Lamu (n=1), **Figure 3A** and **B**.
215
Two variants of concern (VOC) were detected during the study period: Beta (i.e. lineage
B.1.351 in 25 samples); and Alpha (i.e lineage B.1.1.7 in two samples (Konings et al., 2021)).
The earliest B.1.351 infections were identified in international travelers from South Africa in
mid-December 2020 while the earliest detected B.1.1.7 case was in second week of January
2020 in a local with no history of recent international travel. We also detected two cases of
A.23 and six cases of A.23.1 after the peak of wave two (both lineages first identified in
Uganda, and A.23.1 was considered a variant of interest (VOI) in the region in early 2021
(Bugembe et al., 2021a)).
224
Lineage dynamics with a widening scale of observation.
We compared the temporal prevalence patterns of the lineages identified in Coastal Kenya to
Eastern Africa, Africa and a global sub-sample, **Figure 4**. In these comparative genome sets,
we identified 97 Pango lineages for Eastern Africa, 261 for Africa and 990 for the global sub-
sample. Note that by May 2021, a total of 1,293 Pango lineages has been assigned worldwide
since the beginning of the pandemic (O'Toole et al., 2021) indicating very high
representativeness in our global subsample.
232
In our Eastern Africa sample (n=2,261), the top lineages were also observed in the Coastal
Kenya data (B.1.351, A.23.1, B.1.1 and B.1) except lineage B.1.351.2 (n=180, sub-lineage of
B.1.351 circulating in Mayotte) and B.1.380 (n=128, a Rwanda lineage (Butera et al., 2021)),
Figure 4B. In the Africa sample (n=12,191), four of the six top lineages (B.1, B.1.1, B.1.351
and B.1.1.7) were also identified in Coastal Kenya. The undetected two lineages were
B.1.1.448 (n=392) mainly observed South Africa samples, and B.1.351.2, **Figure 4C**. In the
global sub-sample (n=24,511) four of the top six lineages (B.1, B.1.1, B.1.1.7 and B.1.351)
were observed in the samples from Coastal Kenya. The unobserved lineages were D.2 (Alias
of B.1.1.25.2, an Australian lineage) and B.1.2 (a USA lineage), **Figure 4D**.
242
243 At the time of the second wave in Kenya, the Alpha and Beta VOC were already widespread
244 across Eastern Africa and Africa but there were only sporadic detections from our Coastal
245 Kenya dataset. Although in the comparison data lineage B.1 occurred in substantial proportions

across the different scales early in the pandemic, its prevalence diminished faster overtime in
the outside Coastal Kenya sample sets when compared to the Coastal Kenya dataset set.
Greater than 95% of the lineages comprising infections globally were not seen in the Coastal
Kenya samples.

a. Coastal Kenya

b. Eastern Africa

c. Africa

d. Global sample

Figure 4. Lineage dynamics at widening scales of observation. Panel A, Biweekly distribution of detected lineages in Coastal Kenya from the sequenced 747 genomes. Panel B, biweekly distribution of detected lineages in Eastern Africa from 2,261 contemporaneous genomes from 10 countries whose contemporaneous data were available in GISAID. Lineages not observed in the Coastal Kenya dataset have been lumped together as "Absent Kenya" but with the those appearing among top six and not observed in Kenya indicated just below the other Kenya group. Panel C, weekly distribution of detected lineages in Africa from 12,191 contemporaneous genomes from 39 countries that were available in GISAID. Panel D, weekly distribution of detected lineages in a global sub-sample of 25,511 contemporaneous genomes from 139 countries that were compiled from GISAID (see detail in methods section).

SARS-CoV-2 diversity in Coastal Kenya

We constructed both a mutational maximum likelihood (ML) (i.e. non-time resolved) and a time-resolved phylogeny of the Coastal Kenya genomes both while including a random global reference set of genomes (n=4000) shown in **Supplementary-Figure 3** and **Figure 5A,** respectively. These analysis showed that (i) the Coastal Kenya genomes were represented across many but not all of the major phylogenetic clusters observed globally, ii) some of the coastal Kenya clusters expanded after introduction whereas others did not (observed as singletons) and iii) all counties appeared to have had multiple lineage/variant introductions with some clusters comprising genomes detected across multiple counties**, Figure 5A**. There was considerable correlation of the root-to-tip genetic distance and the date of sampling dates (r^2=0.49)**, Figure 5B**.

The coastal Kenya genomes had a fewer pairwise nucleotide differences (median 12, interquartile range (IQR): 7-17) compared to the global subsampled comparison dataset median 22, IQR: 15-29)**, Figure 5C.** A similar difference was observed at amino acid level, with the coastal Kenya genomes showing a median of six amino acid substitutions (IQR: 3-10) while the global dataset showed a median of eight amino acid substitutions (IQR: 6-12)**, Figure 5D**.

For detailed investigation into the local SARS-CoV-2 genetic diversity, we reconstructed time-resolved lineage specific phylogenetic trees are shown in **Figure 6**. The three lineages first identified in Kenya (B.1.530, B.1.549 and B.1.596.1) were found to (a) possess significant diversity consistent with widescale spread within Kenya (**Figure 6C-E**), (b) formed multiple county-specific sub-clusters and (c) show local sequences interspersed with global comparison genomes from the same lineage implying potentially export (or import) events, **Figure 6D**. However, the picture painted by lineage N.8 was different. This lineage was mainly detected in Lamu forming a single monophyletic group (**Figure 6H**) when co-analyzed with its precursor lineage B.1.1.33, an observation consistent with a single introduction as B.1.1.33 then local evolution expansion.

293
294 **Figure 5. SARS-CoV-2 diversity in Coastal Kenya.** Panel A, Time-resolved global

phylogeny that combined 732 Coastal Kenya SARS-CoV-2 genomes and 3,958 global
reference sequences (after removal of sequences not fitting molecular clock-like evolution
using Tree-Time). Panel B, Root-to-tip regression analysis coastal Kenya (colored by county)
and global (shown in grey) sequences with determination of correlation co-efficient (r^2). Panel
C, comparison of pairwise nucleotide differences of the 732 coastal Kenya genomes included
on the phylogenetic tree and the 3,958 global genomes. Panel D, comparison of the number of
amino acid substitutions of the coastal Kenya genomes versus the global dataset relative to the
Wuhan-Hu-1 reference genome.

**Figure 6. Time-resolved lineage-specific phylogenetic trees for the commonly detected
lineages in Coastal Kenya.** The Coastal Kenya genomes are indicated with filled circles
colored by county. Panel A, phylogeny for lineage B.1 that combined 369 Coastal Kenya
genomes and 2,319 global reference sequences. Panel B, phylogeny for lineage B.1.535 that
combined 82 Coastal Kenya genomes and 1,867 global sequences belonging also to B.1.535.
Panel C, phylogeny for lineage B.1.530 that combined 42 Coastal Kenya genomes and 15
global sequences belonging also to B.1.530. Panel D, phylogeny for lineage B.1.549 that
combined 33 Coastal Kenya genomes and 23 sequences from other countries of lineage B.1.
549. Panel E, phylogeny for lineage B.1.596.1 combines 15 Coastal Kenya genomes and 36
global reference sequences. Panel F, phylogeny for lineage B.1.1 that combined 31 Coastal
Kenya genomes and 1,629 global sequences belonging to lineage B.1.1. Panel G, phylogeny
for lineage N.8 that combined 20 Coastal Kenya genomes of lineage N.8, 13 Coastal Kenya
genomes of lineage B.1.1.33 and 570 global sequences of lineage B.1.1.33. Panel H, phylogeny
for lineage B.1.333 combines 32 Coastal Kenya genomes and 753 global reference sequences.

In **Figure 7**, we present a comparison of both the number of nucleotide and amino acid substitutions for each major lineage between the coastal Kenya genomes and the global subsample. The divergence of the lineages from the original Wuhan sequence at the global level was closely mirrored at the local coastal Kenya level. The most divergent lineages among the subset we examined were three of the Kenya specific lineages (B.1.530, B.1.549 and B.1.596.1). The least genetically divergence lineages from the original Wuhan SARS-CoV-2 strain were viruses occurring within lineage B.1.333 and lineage B.1.535, **Figure 7**.

Figure 7. Level of genetic divergence of the major detected lineages in coastal Kenya. The sequences were first aligned to the 2019 Wuhan-Hu-1 isolate and the number of changes compared to isolates across the globe of the same lineage. Panel A shows the divergence at nucleotide sequence level. Panel B shows the divergence at amino acid sequence level focusing on synonymous substitutions.

Viral imports and export from Coastal Kenya
We used ancestral location state reconstruction of the dated phylogeny **(Figure 5A)** to infer the number of viral import and exports (Sagulenko et al., 2018). In total, between March 2020 and February 2021, we detected 357 location transition events within and between the six counties

and the outside world. Sixty-nine of these were viral import events into Coastal Kenya from outside and 93 were viral exports events from Coastal Kenya to outside populations, **Figure 8A-C**. A total of 191 import/export events were detected between the six Coastal counties.

Figure 8. Viral imports and export from Coastal Kenya. Panel A, an alluvium plot showing the number of imports and exports of viral lineages from Coastal Kenya shown as an alluvium plot. Panel B, monthly location transition events observed in the Coastal Kenya and global genomes shown as either viral export, import or inter-county transmission. Panel C, monthly number of import events into the individual Coastal counties and their origins.

Virus imports into the region occurred through Mombasa (n=40, 60%), Kwale (n=9, 13%), Taita Taveta (n=13, 19%), Kilifi (n=5, 7%) and Lamu (n=1, 1%) while virus export from the region occurred through Mombasa (n=86, 41%), Kwale (n=6, 6%) and Taita Taveta (n=1, 1%). Of the 69 detected viral imports, 34 (49%) occurred during the introductions phase, 16 (23%) occurred during wave one and 19 (28%) during wave two while of the 93 detected viral exports, 43 (46%) occurred during introductions phase, 31 (33%) during wave one and 19 (20%) during wave two, **Figure 8B and 8D**.

The analysis revealed the top six country sources of importation into coastal Kenya as; USA, (n=19, 31%), England (n=10, 16%), Uganda (n=6, 10%), Japan (n=5, 8%) and Mozambique

360 (n=3, 5%), **Supplementary-Figure 4 and Supplementary-Figure 5.** On the other hand, the
361 top six country destinations of viral export from coastal Kenya were USA (n=22, 17%),
362 Australia (n=16, 13%), India (n=14, 11%), Uganda (n=8, 6%), England (n=7, 6%) and Scotland
363 (n=4, 3%).
364
365 To infer the sensitivity of our analysis, we repeated the import/export analysis using a (a)
366 different global sub-sample (**Supplementary-Figure 4)** and (b) with a normalized subsample
367 of the coastal Kenya genomes accounting for total reported infections per county,
368 **Supplementary-Figure 5**. The reanalysis came to roughly similar conclusions as above and
369 the results are presented in **Supplementary-Table 1.**
370
Amino acid changes in the Coastal Kenya genomes
We scanned the entire SARS-CoV-2 genome for amino acid changes from the Wuhan
reference genome in the 747 Coastal Kenya sequences. Below we summarize changes that
were identified in ≥5 genomes. We observed 20 changes in the spike (S) protein meeting the
set criterion, 15 in the nucleocapsid (N) protein, 14 in ORF3a, 13 in ORF1b, nine in the ORF1a,
seven in ORF8, five in ORF6, and one in the envelope (E) protein. No codon changes, meeting
of criterion was observed in the matrix (M) protein, ORF7a, ORF7b or ORF10. The most
common amino acid changes observed in the Coastal Kenya genomes were S: D614G (94.4%),
ORF1b: P314L (92.9%), ORF3a: Q57H (15.5%), ORF1a: T265I (15.4%) and N: R195K
(12.9%), **Supplementary-Figure 6.** Several lineage-defining mutations were observed for the
Kenya-specific lineages as summarized in **Supplementary-Figure 7.**
382
DISCUSSION
We present evidence of continuous additional introductions of SARS-CoV-2 into Coastal
Kenya during the first year of the pandemic documenting at least 69 independent viral
introduction events into the region. Multiple introductions were observed even at county level,
with extensive inter-county spread observed after the early phase period. Strikingly, many of
the imports and exports from the region occurred through the city of Mombasa, a major
commercial, industrial and tourist destination in the region. This observation highlights
Mombasa's central role as a gateway for virus entry into coastal Kenya and the significance
continuous surveillance in this county for early detection of variants in coastal Kenya
populations.
393
During the early phase of the pandemic several Pango lineages were introduced into the region
(A, B, AL.1, B.1, B.1.1, B.1.1.1, B.1.1.33, B.1.13, B.1.153, b.1.222, B.1.333, B.1.535, B.4 and
B.4.7). However, it is lineage B.1 which has a European origin, that became predominant,
48.2% of the cases (Githinji et al., 2020). This lineage had the D614G change in the spike
protein which has been found to enhance viral fitness (Baric, 2020) and this may have boosted
its local transmission and early dominance. Notably also the import/export analysis found that
the majority of viral imports to coastal Kenya had an Origin in USA, England and neighboring
Uganda. Viral exports from costal Kenya happened to USA, Uganda, Australia among other
destinations.
403
During the national wave one, the Coastal region observed SARS-CoV-2 transmission mainly
in counties of Mombasa, Taita Taveta and Lamu. The key lineages observed then were B.1
(58.7%), N.8 (8.517%), B.1.535 (6.7%), B.1.1 (5.8%). Notably, lineage N.8 was specific to
Lamu County. Lineage N.8 has not been observed outside Kenya and its precursor lineage

(B.1.1.33) was observed earlier in South America, mainly Brazil. Lineage N.8 may have arisen
from an introduction event of B.1.1.33 into Lamu county, and then local evolution and
localization to Lamu. The N.8 lineage has seven characteristic lineage defining mutations
including S: D614G and R203K, G204R and I292T on the nucleocapsid protein.

The national wave two started in October 2020 and led to significant viral transmission in all
the six coastal counties. In Kilifi, Tana River and Kwale counties, this was the first significant
peak. This wave two was majorly comprised mainly three lineages: B.1 (42%), B.1.530 (16%),
and B.1.549 (12%). Lineages B.1.530 and B.1.549, first identified in Kenya, and may have
arisen from local evolution of lineage B.1. Lineage B.1.530 has six characteristic mutations
including spike P681H change adjacent to the biologically important furin cleavage site while
lineage B.1.549 has seven characteristic mutations, five occurring in the ORF1a or ORF1b.
Both two lineages have now been observed in a few other countries albeit in small numbers:
seven countries for B.1.530 (Rwanda, Netherlands, Germany, Denmark, USA, Japan and
Australia) and three countries for B.1.549 (England, USA, Canada).

Near the tail end of the national wave two, we detected cases of the Beta VOC (B.1.351) and
the Alpha VOC (B.1.1.7). Beta was first to be observed, initially in Kilifi in two asymptomatic
international travelers in mid-December 2020. Alpha was subsequently detected in a local who
presented to a Mombasa clinic in the second week of January 2021. In the subsequent weeks
up to the end of the period covered by this analysis (February 2021), only one additional Alpha
case was detected in our sequenced samples unlike lineage Beta which continued to be detected
sporadically in January and February 2021 especially in the counties of Taita Taveta and Kwale
which border Tanzania.

When we compared the local lineage patterns to the global patterns, it was notable that although
several lineages were detected globally (>900), only a small fraction (<10%) of this were
documented locally (O'Toole et al., 2021). The VOC were already extensively spread across
Eastern Africa (Beta VOC), Africa (Beta VOC) and worldwide (Alpha VOC) in the last quarter
of 2020 unlike the findings here for costal Kenya. This implies that a slight lag occurred in the
VOC arrival and large-scale spared in Coastal Kenya perhaps due to remoteness and public
health measures that have remained in place during the period.

While Mombasa, Taita Taveta and Lamu counties had experienced two waves of infections by
February 2021, Kilifi, Tana River and Kwale counties had experienced only a single major
wave. This may have been brought about by differences in population size, density, and
accessibility for these different counties. For instance, Mombasa is densely populated
(~5,604.64 persons/sq.km, 2019 census), has a large seaport and an international airport while
Lamu has a sparsely populated mainland and with several islands (total population 143,920,
population density, 21.84 persons/sq.km). Taita Taveta and Kwale counties are primarily rural
counties bordering Tanzania which had a largely uncontrolled SARS-CoV-2 epidemic during
the study period (Rice et al., 2021). Tana River is the most remote and sparsely populated (8.9
persons/sq.km) county in the region.

Our study contributes to the limited but growing literature illuminating SARS-CoV-2
transmission patterns in sub-Saharan Africa (Bugembe et al., 2021a; Bugembe et al., 2021b;
Tegally et al., 2021a; Tegally et al., 2021b; Wilkinson et al.). The patterns revealed here inform
on frequency, sources and routes of introductions of SARS-CoV-2 into Coastal Kenya and
have implications on future control strategies. The key limitations of our analysis include, first,
the samples we sequenced were only those available to us through the rapid response teams

(RRTs) whose case identification protocols were altered at the different study pandemic
phases. Second, we only sequenced ~20% of the positive samples identified in our laboratory,
the majority from the introduction phase and were prioritizing samples with a Ct value of
<30.0. Third, the sampling across the six Coastal counties was not uniform, probably in part
due to varied distance from our testing centre located in Kilifi County.
463
In conclusion, we show that the first two SARS-CoV-2 waves in Coastal Kenya observed
transmission of both newly introduced and potentially locally evolved variants. New variant
introductions were mainly through Mombasa city and originated USA, England, Uganda
among other predicated sources. Only a limited number of the many introduced variants
progressed to transmit extensively. Unlike in the global contemporaneous sample, we did not
find evidence of extensive local transmission of the global VOC during wave two. Thus, we
infer that it is more likely that the relaxation of some of the interventions (e.g., reopening of
learning institutions, airspace, bars and restaurants) that drove the second wave of infections.
Our study shows the importance of detailed local genomic surveillance even in remote and
under-resourced settings to understand spread patterns of SARS-CoV-2 to optimize local
interventions.
475

MATERIALS AND METHODS

Ethical statement
Samples analysed here were collected under the MoH protocols as part of the national response
to the COVID-19 pandemic. The whole genome sequencing study protocol was reviewed and
approved by the Scientific and Ethics Review Committee (SERU), Kenya Medical Research
Institute (KEMRI), Nairobi, Kenya (SERU #4035). Individual patient consent was not required
by the committee for the use of these samples for genomic surveillance to inform public health
response.
484

Study period and population
This study examined samples collected between 17th March 2020 to 26th February 2021 from
six counties in Coastal Kenya i.e. Mombasa, Kilifi, Kwale, Tana River, Taita Taveta and Lamu.
We divided the study period into three phases based on trends in daily reported COVID-19
national case numbers by the MoH (MOH, 2021): (a) Introduction phase – 17th March-20th
May, 2020 (b) Wave one – 21st May-15th September 2020 and (c) Wave two – 16th September
to 28th February 2021, **Figure 1**. We considered the transition from introduction phase to Wave
one as the timepoint when the national daily positives exceeded 50 and the transition from
Wave one to Wave two as the timepoint when a consistent renewed rise of national daily
number of positives started after Wave one peak.
495

Patient samples
This study analyzed SARS-CoV-2 positive NP/OP swab samples that were collected by the
MoH County Department of Health RRTs, across all the six counties of Coastal Kenya. SARS-
CoV-2 positive diagnosis was made at the KWTRP in Kilifi County (Agoti et al., 2021). The
RRTs delivered the NP/OP swabs to KWTRP laboratories within 48 hours of collection in cool
boxes with ice packs. The samples were from persons of any age and sampling followed the
MOH eligibility criteria that were revised from time to time (Githinji et al., 2020). Persons
sampled included those with acute respiratory symptoms, those with a recent history of travel
to the early COVID-19 hotspots, contacts of confirmed cases, persons presenting at
international border points seeking entry into Kenya, among other criteria.
506

507 *SARS-CoV-2 detection at KWTRP*

Viral RNA was extracted from the NP/OP samples using any one of seven commercial kits that were available namely, QIAamp Viral RNA Mini Kit, RNeasy ® QIAcube ® HT Kit, QIASYMPHONY ® RNA Kit, T0IANamp Virus RNA Kit, Da An Gene Nucleic acid Isolation and Purification Kit, SPIN X Extraction and RADI COVID-19 detection Kit. The extracts were tested for the presence of SARS-CoV-2 nucleic acid following different protocols depending on which one was available of the following 7 kits/protocols: 1) the Berlin (Charité) primer-probe set (targeting envelope (E) gene, nucleocapsid (N) or RNA-dependent RNA-polymerase (RdRp)), 2) European Virus Archive – GLOBAL (EVA-g) (targeting E or RdRp genes), 3) Da An Gene Co. detection Kit (targeting N or ORF1ab), 4) BGI RT-PCR kit (targeting ORF1ab), 5) Sansure Biotech Novel Coronavirus (2019-nCoV) Nucleic Acid Diagnostic real-time RT-PCR kit or 6) Standard M kit (targeting E and ORF1ab) and 7) TIB MOLBIOL kit (targeting E gene). Protocol-specific recommended cycle threshold cut-offs were followed in defining SARS-CoV-2 positives.

SARS-CoV-2 genome sequencing

Only samples that had a RT-PCR cycle threshold value of <30.0 were targeted for whole genome sequencing (Githinji et al., 2020), **Supplementary-Figure 1** . Viral RNA from positive samples was reextracted using QIAamp Viral RNA Mini kit following the manufacturer's instructions and reverse transcribed using LunaScript® RT SuperMix Kit. The cDNA was amplified using Q5® Hot Start High-Fidelity 2x Mastermix along with the ARTIC nCoV-2019 version 3 primers. The PCR products were run on a 1.5% agarose gel and for samples whose SARS-CoV-2 amplification was considered successful were purified using Agencourt AMPure XP beads and taken forward for library preparation.

Sequencing libraries were constructed using Oxford Nanopore Technology (ONT) ligation sequencing kit and the ONT Native Barcoding Expansion kit as described in the ARTIC protocol (Tyson et al., 2020). Every MinION (Mk1B) run comprised 23 samples and one negative (no-template) control.

SARS-CoV-2 genome assembly

Following MinION sequencing, the FAST5 files were base-called and demultiplexed using the OTN's software Guppy v3.5-4.2. For base calling, high accuracy mode was used. Consensus SARS-CoV-2 sequences were derived from the reads using the ARTIC bioinformatics pipeline (https://artic.network/ncov-2019/ncov2019-bioinformatics-sop.html; last accessed 2021-08-03). A threshold of ×20 read depth was required for a base to be included in the consensus genome otherwise it was masked with an N. Only complete or near-complete genomes N count < 5,980 (i.e >80% coverage) were taken forward for phylogenetic analysis.

Lineage assignment and calling of amino acid changes

The consensus genomes were assigned into lineages using the PANGOLIN nomenclature (Rambaut et al., 2020). We used the command line pangolin v3.1.10 software to assign Pango lineages to each consensus sequence with pango v1.2.46 and pangoLEARN model v2021-07-28 (O'Toole et al., 2021). Contextual information about lineages was obtained from the Lineage list available at https://cov-lineages.org/lineage_list.html (last accessed 2021-08-04). Variants of concern (VOC) and variants of interest (VOI) were designated based on the WHO framework as of 31st May 2021 (https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/).

The amino acid changes in the Coastal Kenya genomes from the reference strain Wuhan-Hu-1/2019 was investigated using the nextclade tool v0.14.2 (Hadfield et al., 2018):

558   https://clades.nextstrain.org/; last accessed 2021-08-03. Mutations in the Kenyan lineages were
559   visualized using the tool available on Stanford University CORONAVIRUS ANTIVIRAL &
560   RESISTANCE Database webpage : https://covdb.stanford.edu/page/mutation-viewer/; last
561   accessed 2021-08-03).

Global contextual sequences

We prepared three sets of contextual sequences from GISAID database
(https://www.gisaid.org/). Selected sequences were those with non-ambiguous sampling date,
sampled during the period covered by our study (March 2020 – February 2021) and after
alignment possessed <5,980 ambiguous (N) nucleotides i.e. showed >80% genome
completeness.

a) **Set 1:** We compiled an African SARS-CoV-2 genome set which we assigned Pango
lineages and used to compare the continental lineage temporal patterns to the Coastal
Kenya lineage distribution. A total of 12,191 genomes were included for this purpose
downloaded on the 11[th] of May 2021. From these we also created an Eastern Africa
subset which comprised of 2,261 genomes from 10 countries, namely, Zimbabwe,
Zambia, Uganda, Rwanda, Reunion (a France overseas territory), Mozambique,
Malawi, Madagascar, Ethiopia and Comoros.

b) **Set 2**: We compiled a global reference set of 24,511 genomes collected across all the
six inhabited continents between 17[th] March 2020 and 27[th] February 2021. The
genomes were selected from all available genomes in GISAID database using the R
randomization command: *sample_n()*. Between 1916 and 2174 genomes were selected
for each included month (mean of 2042 and median of 2049) randomly selected across
all continents from across 164 countries. These genomes were used to infer the global
temporal patterns of the lineages observed in Coastal Kenya. A subset of these genomes
(n=4000) was combined with the Kenyan genomes to infer their global phylogenetic
context and in the import/export analysis described below.

c) **Set 3**: We compiled a genotype reference for the top six observed lineages in Coastal
Kenya. For lineages B.1. and B.1.1 which >5000 genomes exist in GISAID were
prepared we used the sub-sample from Set 2 above. For lineages B.1.530, B.1.549,
B.1.596.1 and N.8 we retrieved all genomes assigned these lineages available from
GISAID to infer their global phylogenetic context.

Phylogenetic analysis

Multiple sequence alignments were prepared in Nextalign v 0.1.6 software and using the initial
Wuhan sequence (Accession number: NC_045512) as the reference, command:

nextalign -r NC_045512.fasta -i input.fasta

The alignment was manually inspected in AliView v1.21 to spot any obvious misalignments.
Quick non-bootstrapped neighbor joining trees were created in SEAVIEW v4.6.4 to identify
any aberrant sequences that were henceforth discarded. We reconstructed maximum likelihood
(ML) phylogenies using IQTREE v2.1.3, command:

./iqtree2 -s input.aligned.fasta -nt 4

The software initiates tree reconstruction after assessment and selection of the best model of
nucleotide substitution for the alignment using ModelFinder. For example for the large ML
phylogeny presented in Figure 5A, the Best-fit mode was determined as GTR+F+R3.

The ML trees were linked to the various metadata (lineage, county, source etc) in R programming software v4.0.2 and visualized using R ggTree v2.4.2. TempEst v1.5.3 was used to assess the presence of a molecular clock signal in analysed data and linear regression of root-to-tip genetic distances against sampling dates plotted in RStudio and the correlation coefficient assessed.

Import/export analysis

The global ML tree topology (**Figure 5**) was used to estimate the number of viral transmission events between Coastal Kenya and the rest of the world as in (Wilkinson et al.). The software TreeTime was used to transform the ML tree topology into a dated tree assuming a constant genomic evolutionary rate of SARS-CoV-2 of 8.4×10^{-4} nucleotide substitutions per site per year (Sagulenko et al., 2018), command:

treetime --tre input.aligned.fasta.treefile --aln input.aligned.fasta
-–clock-rate 0.00084 –-dates dates.csv

Outlier sequences were identified by TreeTime and excluded during this process. A migration model was henceforth fitted to the resulting time-scaled phylogenetic tree from TreeTime, mapping the location status of the genomes from the six counties at both the tips and internal nodes, command:

treetime mugration –-tree input.tree.nwk –-states mugration.csv –-attribute Location

Using the date and location annotated tree topology, we counted the number of transitions between and within Coastal Kenya counties and the rest of the world using the python script developed and described by the team at KRISP (https://github.com/krisp-kwazulu-natal/SARSCoV2_South_Africa_major_lineages/tree/main/Phylogenetics ; last accessed 2020-08-04) and plotted this using ggplot2 v3.3.3. This analysis was repeated with different sub-sample of the global background data and with a down sample of the coastal Kenya genomes that was normalized spatial-temporally (using a maximum of 15 samples per county per month).

Epidemiological data

The Kenya daily case data for the period between March 2020 and February 2021 was downloaded from Our World in Data database (https://ourworldindata.org/coronavirus/country/kenya). Metadata for the Coastal Kenya samples was compiled from the MoH case investigation forms delivered alongside the samples.

Kenya COVID-19 response

We derived status of Kenya government COVID-19 interventions using data available from Our World in Data database that has calculated the Oxford Stringency Index (SI). These SI estimations are composite measure based on nine response indicators implemented by governments rescaled to a value 0-100, with 100 being strictest (Hale et al., Noam Angrist). The indicators are: (a) School closures, (b) Workplace closures, (c) Cancellation of public events, (d) Restrictions on public gatherings, (e) Closures of public transport, (f) Stay-at-home requirements, (g) Public information campaigns, (h) Restrictions on internal movements and (d) International travel controls. These are The Kenya government revised the policy interventions at approximately monthly intervals (Brand et al., 2021) and the changes in SI overtime are shown in **Figure 1D**.

Statistical analysis

Statistical data analyses were performed in R v 4.0.5. Summary statistics including proportions, means, median and ranges were calculated and provided where applicable. T-test was used to evaluate a statistical difference in means of the pairwise nucleotide or amino acid differences between the coastal and global sample. The *lm* function in R was used to fit a linear regression model evaluating the relationship between sampling dates and root-to-tip genetic distance in the ML phylogeny. The goodness of fit was inferred from the correlation coefficient.

Data availability

The data and scripts used to generate the figures shown in the manuscript are hosted on Harvard dataverse: DOI: https://doi.org/10.7910/DVN/4ZZYIM.

The Coastal Kenya genomes are available in the GISAID database (Accession numbers: EPI_ISL_1039223-29; EPI_ISL1440075-109, IPI_ISL_457845-931; EPI_ISL_568695-872; EPI_ISL806548-717; EPI_ISL_855494-548; EPI_ISL_968807-9245).

Acknowledgements

We thank (a) the members of the six Coastal counties of Kenya RRTs for collecting the samples analysed here; (b) the members of the COVID-19 KWTRP Testing Team who tirelessly analysed the samples received at KWTRP to identify positives (see full list of members below); (c) the KWTRP data entry team, (d) Laboratories that have shared sequence data on GISAID that we included as comparison data in our analysis (see list in appendix); (e) the KRISP team in South Africa for sharing the scripts we used in the import/export analysis and AFRICA-CDC for facilitating Africa genomics training. This paper is published with permission of the Director of KEMRI.

Members of COVID-19 Testing Team at KWTRP
Agnes Mutiso, Alfred Mwanzu, Angela Karani, Bonface M. Gichuki, Boniface Karia, Brian Bartilol, Brian Tawa, Calleb Odundo, Caroline Ngetsa, Clement Lewa, Daisy Mugo, David Amadi, David Ireri, Debra Riako, Domtila Kimani, Edwin Machanja, Elijah Gicherua, Elisha Omer, Faith Gambo, Horace Gumba, Isaac Musungu, James Chemweno, Janet Thoya, Jedida Mwacharo, John Gitonga, Johnstone Makale, Justine Getonto, Kelly Ominde, Kelvias Keter, Lydia Nyamako, Margaret Nunah, Martin Mutunga, Metrine Tendwa, Moses Mosobo, Nelson Ouma, Nicole Achieng, Patience Kiyuka, Perpetual Wanjiku, Peter Mwaura, Rita Warui, Robinson Cheruiyot, Salim Mwarumba, Shaban Mwangi, Shadrack Mutua, Sharon Owuor, Susan Njuguna, Victor Osoti, Wesley Cheruiyot, Wilfred Nyamu, Wilson Gumbi and Yiakon Sein.

Funding

This work was supported by the National Institute for Health Research (NIHR) (project references 17/63/82 (PI Prof. James Nokes) and 16/136/33 (PI Prof. Mark Wooolhouse) using UK aid from the UK Government to support global health research, The UK Foreign, Commonwealth and Development Office and Wellcome Trust (grant# 220985). The views expressed in this publication are those of the author (s) and not necessarily those of NIHR, the

Department of Health and Social Care, Foreign Commonwealth and Development Office,
Wellcome Trust or the UK government. Some members of COVID-19 Testing Team at
KWTRP were supported by funding received by Dr Marta Maia (BOHEMIA study funded
UNITAID), Dr Francis Ndungu (Senior Fellowship and Research and Innovation Action (RIA)
grants from EDCTP) and Prof. Anthony Scott (PCIVS grant from GAVI), and IAVI (USAID
grant number: AID-OAA-A-16-00032).

709
710

REFERENCES

Adetifa, I.M.O., Uyoga, S., Gitonga, J.N., Mugo, D., Otiende, M., Nyagwange, J., Karanja, H.K.,
Tuju, J., Wanjiku, P., Aman, R., Mwangangi, M., Amoth, P., Kasera, K., Ng'ang'a, W.,
Rombo, C., Yegon, C., Kithi, K., Odhiambo, E., Rotich, T., Orgut, I., Kihara, S., Bottomley,
C., Kagucia, E.W., Gallagher, K.E., Etyang, A., Voller, S., Lambe, T., Wright, D., Barasa, E.,
Tsofa, B., Bejon, P., Ochola-Oyier, L.I., Agweyu, A., Scott, J.A.G., Warimwe, G.M., 2021.
Temporal trends of SARS-CoV-2 seroprevalence during the first wave of the COVID-19
epidemic in Kenya. Nature Communications 12(1), 3966.
Agoti, C., Mutunga, M., Lambisia, A., Kimani, D., Cheruiyot, R., Kiyuka, P., Lewa, C., Gicheru, E.,
Tendwa, M., Said Mohammed, K., Osoti, V., Makale, J., Tawa, B., Odundo, C., Cheruiyot,
W., Nyamu, W., Gumbi, W., Mwacharo, J., Nyamako, L., Otieno, E., Amadi, D., Thoya, J.,
Karani, A., Mugo, D., Musyoki, J., Gumba, H., Mwarumba, S., M. Gichuki, B., Njuguna, S.,
Riako, D., Mutua, S., Gitonga, J., Sein, Y., Bartilol, B., Mwangi, S., O. Omuoyo, D., M.
Morobe, J., de Laurent, Z., Bejon, P., Ochola-Oyier, L., Tsofa, B., 2021. Pooled testing
conserves SARS-CoV-2 laboratory resources and improves test turn-around time: experience
on the Kenyan Coast [version 2; peer review: 3 approved]. Wellcome Open Research 5(186).
Baric, R.S., 2020. Emergence of a Highly Fit SARS-CoV-2 Variant. New England Journal of
Medicine 383(27), 2684-2686.
Brand, S.P.C., Ojal, J., Aziza, R., Were, V., Okiro, E.A., Kombe, I.K., Mburu, C., Ogero, M.,
Agweyu, A., Warimwe, G.M., Nyagwange, J., Karanja, H., Gitonga, J.N., Mugo, D., Uyoga,
S., Adetifa, I.M.O., Scott, J.A.G., Otieno, E., Murunga, N., Otiende, M., Ochola-Oyier, L.I.,
Agoti, C.N., Githinji, G., Kasera, K., Amoth, P., Mwangangi, M., Aman, R., Ng'ang'a, W.,
Tsofa, B., Bejon, P., Keeling, M.J., Nokes, D.J., Barasa, E., 2021. COVID-19 Transmission
Dynamics Underlying Epidemic Waves in Kenya. medRxiv, 2021.2006.2017.21259100.
Bugembe, D.L., Kayiwa, J., Phan, M.V.T., Tushabe, P., Balinandi, S., Dhaala, B., Lexow, J.,
Mwebesa, H., Aceng, J., Kyobe, H., Ssemwanga, D., Lutwama, J., Kaleebu, P., Cotten, M.,
2020. Main Routes of Entry and Genomic Diversity of SARS-CoV-2, Uganda. Emerg Infect
Dis 26(10), 2411-2415.
Bugembe, D.L., Phan, M.V.T., Ssewanyana, I., Semanda, P., Nansumba, H., Dhaala, B., Nabadda, S.,
O'Toole, Á.N., Rambaut, A., Kaleebu, P., Cotten, M., 2021a. Emergence and spread of a
SARS-CoV-2 lineage A variant (A.23.1) with altered spike protein in Uganda. Nature
Microbiology.
Bugembe, D.L., V.T.Phan, M., Ssewanyana, I., Semanda, P., Nansumba, H., Dhaala, B., Nabadda, S.,
O'Toole, Á.N., Rambaut, A., Kaleebu, P., Cotten, M., 2021b. A SARS-CoV-2 lineage A
variant (A.23.1) with altered spike has emerged and is dominating the current Uganda
epidemic. medRxiv, 2021.2002.2008.21251393.
Butera, Y., Mukantwari, E., Artesi, M., D'Arc Umuringa, J., O'Toole, Á.N., Hill, V., Rooke, S.,
Hong, S.L., Dellicour, S., Majyambere, O., Bontems, S., Boujemla, B., Quick, J., Resende,
P.C., Loman, N., Umumararungu, E., Kabanda, A., Murindahabi, M.M., Tuyisenge, P.,
Gashegu, M., Rwabihama, J.P., Sindayiheba, R., Gikic, D., Souopgui, J., Ndifon, W.,
Rutayisire, R., Gatare, S., Mpunga, T., Ngamije, D., Bours, V., Rambaut, A., Nsanzimana, S.,
Baele, G., Durkin, K., Mutesa, L., Rujeni, N., 2021. Genomic Sequencing of SARS-CoV-2 in
Rwanda: evolution and regional dynamics. medRxiv, 2021.2004.2002.21254839.
Etyang, A.O., Lucinde, R., Karanja, H., Kalu, C., Mugo, D., Nyagwange, J., Gitonga, J., Tuju, J.,
Wanjiku, P., Karani, A., Mutua, S., Maroko, H., Nzomo, E., Maitha, E., Kamuri, E., Kaugiria,
T., Weru, J., Ochola, L.B., Kilimo, N., Charo, S., Emukule, N., Moracha, W., Mukabi, D.,

Okuku, R., Ogutu, M., Angujo, B., Otiende, M., Bottomley, C., Otieno, E., Ndwiga, L.,
Nyaguara, A., Voller, S., Agoti, C., Nokes, D.J., Ochola-Oyier, L.I., Aman, R., Amoth, P.,
Mwangangi, M., Kasera, K., Ng'ang'a, W., Adetifa, I., Kagucia, E.W., Gallagher, K., Uyoga,
S., Tsofa, B., Barasa, E., Bejon, P., Scott, J.A.G., Agweyu, A., Warimwe, G., 2021.
Seroprevalence of Antibodies to SARS-CoV-2 among Health Care Workers in Kenya. Clin
Infect Dis.

Geoghegan, J.L., Ren, X., Storey, M., Hadfield, J., Jelley, L., Jefferies, S., Sherwood, J., Paine, S.,
Huang, S., Douglas, J., Mendes, F.K., Sporle, A., Baker, M.G., Murdoch, D.R., French, N.,
Simpson, C.R., Welch, D., Drummond, A.J., Holmes, E.C., Duchêne, S., de Ligt, J., 2020.
Genomic epidemiology reveals transmission patterns and dynamics of SARS-CoV-2 in
Aotearoa New Zealand. Nature Communications 11(1), 6351.

Githinji, G., de Laurent, Z.R., Mohammed, K.S., Omuoyo, D.O., Macharia, P.M., Morobe, J.M.,
Otieno, E., Kinyanjui, S.M., Agweyu, A., Maitha, E., Kitole, B., Suleiman, T., Mwakinangu,
M., Nyambu, J., Otieno, J., Salim, B., Kasera, K., Kiiru, J., Aman, R., Barasa, E., Warimwe,
G., Bejon, P., Tsofa, B., Ochola-Oyier, L.I., Nokes, D.J., Agoti, C.N., 2020. Tracking the
introduction and spread of SARS-CoV-2 in coastal Kenya. medRxiv,
2020.2010.2005.20206730.

Githinji, G., de Laurent, Z.R., Mohammed, K.S., Omuoyo, D.O., Macharia, P.M., Morobe, J.M.,
Otieno, E., Kinyanjui, S.M., Agweyu, A., Maitha, E., Kitole, B., Suleiman, T., Mwakinangu,
M., Nyambu, J., Otieno, J., Salim, B., Kasera, K., Kiiru, J., Aman, R., Barasa, E., Warimwe,
G., Bejon, P., Tsofa, B., Ochola-Oyier, L.I., Nokes, D.J., Agoti, C.N., 2021. Tracking the
introduction and spread of SARS-CoV-2 in coastal Kenya. Nature Communications 12(1),
4809.

Hadfield, J., Megill, C., Bell, S.M., Huddleston, J., Potter, B., Callender, C., Sagulenko, P., Bedford,
T., Neher, R.A., 2018. Nextstrain: real-time tracking of pathogen evolution. Bioinformatics
34(23), 4121-4123.

Hale, T., Angrist, N., Boby, T., Cameron-Blake, E., Hallas, L., Kira, B., Majumdar, S., Petherick, A.,
Phillips, T., Tatlow, H., Webster, S., Noam Angrist. Variation in government responses to
COVID-19. BSG-WP-2020/032 vols. University of Oxford.

Konings, F., Perkins, M.D., Kuhn, J.H., Pallen, M.J., Alm, E.J., Archer, B.N., Barakat, A., Bedford,
T., Bhiman, J.N., Caly, L., Carter, L.L., Cullinane, A., de Oliveira, T., Druce, J., El Masry, I.,
Evans, R., Gao, G.F., Gorbalenya, A.E., Hamblion, E., Herring, B.L., Hodcroft, E., Holmes,
E.C., Kakkar, M., Khare, S., Koopmans, M.P.G., Korber, B., Leite, J., MacCannell, D.,
Marklewitz, M., Maurer-Stroh, S., Rico, J.A.M., Munster, V.J., Neher, R., Munnink, B.O.,
Pavlin, B.I., Peiris, M., Poon, L., Pybus, O., Rambaut, A., Resende, P., Subissi, L., Thiel, V.,
Tong, S., van der Werf, S., von Gottberg, A., Ziebuhr, J., Van Kerkhove, M.D., 2021. SARS-
CoV-2 Variants of Interest and Concern naming scheme conducive for global discourse.
Nature Microbiology.

MOH, 2021. Update on COVID-19 in Kenya. In: coronavirus, N.E.C.o. (Ed.). MOH.

O'Toole, Á., Scher, E., Underwood, A., Jackson, B., Hill, V., McCrone, J.T., Colquhoun, R., Ruis, C.,
Abu-Dahab, K., Taylor, B., Yeats, C., du Plessis, L., Maloney, D., Medd, N., Attwood, S.W.,
Aanensen, D.M., Holmes, E.C., Pybus, O.G., Rambaut, A., 2021. Assignment of
epidemiological lineages in an emerging pandemic using the pangolin tool. Virus Evolution.

Oude Munnink, B.B., Nieuwenhuijse, D.F., Stein, M., O'Toole, Á., Haverkate, M., Mollers, M.,
Kamga, S.K., Schapendonk, C., Pronk, M., Lexmond, P., van der Linden, A., Bestebroer, T.,
Chestakova, I., Overmars, R.J., van Nieuwkoop, S., Molenkamp, R., van der Eijk, A.A.,
GeurtsvanKessel, C., Vennema, H., Meijer, A., Rambaut, A., van Dissel, J., Sikkema, R.S.,
Timen, A., Koopmans, M., 2020. Rapid SARS-CoV-2 whole-genome sequencing and
analysis for informed public health decision-making in the Netherlands. Nat Med 26(9),
1405-1410.

Rambaut, A., Holmes, E.C., O'Toole, Á., Hill, V., McCrone, J.T., Ruis, C., du Plessis, L., Pybus,
O.G., 2020. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic
epidemiology. Nat Microbiol 5(11), 1403-1407.

Rice, B.L., Annapragada, A., Baker, R.E., Bruijning, M., Dotse-Gborgbortsi, W., Mensah, K., Miller,
I.F., Motaze, N.V., Raherinandrasana, A., Rajeev, M., Rakotonirina, J., Ramiadantsoa, T.,

Rasambainarivo, F., Yu, W., Grenfell, B.T., Tatem, A.J., Metcalf, C.J.E., 2021. Variation in
SARS-CoV-2 outbreaks across sub-Saharan Africa. Nature Medicine 27(3), 447-453.
Sagulenko, P., Puller, V., Neher, R.A., 2018. TreeTime: Maximum-likelihood phylodynamic analysis.
Virus Evol 4(1), vex042.
Tegally, H., Wilkinson, E., Giovanetti, M., Iranzadeh, A., Fonseca, V., Giandhari, J., Doolabh, D.,
Pillay, S., San, E.J., Msomi, N., Mlisana, K., von Gottberg, A., Walaza, S., Allam, M., Ismail,
A., Mohale, T., Glass, A.J., Engelbrecht, S., Van Zyl, G., Preiser, W., Petruccione, F., Sigal,
A., Hardie, D., Marais, G., Hsiao, N.Y., Korsman, S., Davies, M.A., Tyers, L., Mudau, I.,
York, D., Maslo, C., Goedhals, D., Abrahams, S., Laguda-Akingba, O., Alisoltani-Dehkordi,
A., Godzik, A., Wibmer, C.K., Sewell, B.T., Lourenço, J., Alcantara, L.C.J., Kosakovsky
Pond, S.L., Weaver, S., Martin, D., Lessells, R.J., Bhiman, J.N., Williamson, C., de Oliveira,
T., 2021a. Detection of a SARS-CoV-2 variant of concern in South Africa. Nature 592(7854),
438-443.
Tegally, H., Wilkinson, E., Lessells, R.J., Giandhari, J., Pillay, S., Msomi, N., Mlisana, K., Bhiman,
J.N., von Gottberg, A., Walaza, S., Fonseca, V., Allam, M., Ismail, A., Glass, A.J.,
Engelbrecht, S., Van Zyl, G., Preiser, W., Williamson, C., Petruccione, F., Sigal, A., Gazy, I.,
Hardie, D., Hsiao, N.Y., Martin, D., York, D., Goedhals, D., San, E.J., Giovanetti, M.,
Lourenço, J., Alcantara, L.C.J., de Oliveira, T., 2021b. Sixteen novel lineages of SARS-CoV-
2 in South Africa. Nat Med 27(3), 440-446.
Tyson, J.R., James, P., Stoddart, D., Sparks, N., Wickenhagen, A., Hall, G., Choi, J.H., Lapointe, H.,
Kamelian, K., Smith, A.D., Prystajecky, N., Goodfellow, I., Wilson, S.J., Harrigan, R.,
Snutch, T.P., Loman, N.J., Quick, J., 2020. Improvements to the ARTIC multiplex PCR
method for SARS-CoV-2 genome sequencing using nanopore. bioRxiv.
Uyoga, S., Adetifa, I.M.O., Karanja, H.K., Nyagwange, J., Tuju, J., Wanjiku, P., Aman, R.,
Mwangangi, M., Amoth, P., Kasera, K., Ng'ang'a, W., Rombo, C., Yegon, C., Kithi, K.,
Odhiambo, E., Rotich, T., Orgut, I., Kihara, S., Otiende, M., Bottomley, C., Mupe, Z.N.,
Kagucia, E.W., Gallagher, K.E., Etyang, A., Voller, S., Gitonga, J.N., Mugo, D., Agoti, C.N.,
Otieno, E., Ndwiga, L., Lambe, T., Wright, D., Barasa, E., Tsofa, B., Bejon, P., Ochola-
Oyier, L.I., Agweyu, A., Scott, J.A.G., Warimwe, G.M., 2021a. Seroprevalence of anti-
SARS-CoV-2 IgG antibodies in Kenyan blood donors. Science 371(6524), 79-82.
Uyoga, S., Adetifa, I.M.O., Otiende, M., Yegon, C., Agweyu, A., Warimwe, G.M., Scott, J.A.G.,
2021b. Prevalence of SARS-CoV-2 Antibodies From a National Serosurveillance of Kenyan
Blood Donors, January-March 2021. Jama.
Wilkinson, E., Giovanetti, M., Tegally, H., San, J.E., Lessells, R., Cuadros, D., Martin, D.P.,
Rasmussen, D.A., Zekri, A.-R.N., Sangare, A.K., Ouedraogo, A.-S., Sesay, A.K., Priscilla,
A., Kemi, A.-S., Olubusuyi, A.M., Oluwapelumi, A.O.O., Hammami, A., Amuri, A.A.,
Sayed, A., Ouma, A.E.O., Elargoubi, A., Anthony, A.N., Victoria, A.F., Kazeem, A., George,
A., Trotter, A.J., Yahaya, A.A., Keita, A.K., Diallo, A., Kone, A., Souissi, A., Chtourou, A.,
Gutierrez, A.V., Page, A.J., Vinze, A., Iranzadeh, A., Lambisia, A., Ismail, A., Rosemary, A.,
Sylverken, A., Femi, A., Ibrahimi, A., Marycelin, B., Oderinde, B.S., Bolajoko, B., Dhaala,
B., Herring, B.L., Njanpop-Lafourcade, B.-M., Kleinhans, B., McInnis, B., Tegomoh, B.,
Brook, C., Pratt, C.B., Scheepers, C., Akoua-Koffi, C.G., Agoti, C.N., Peyrefitte, C.,
Daubenberger, C., Morang'a, C.M., Nokes, D.J., Amoako, D.G., Bugembe, D.L., Park, D.,
Baker, D., Doolabh, D., Ssemwanga, D., Tshiabuila, D., Bassirou, D., Amuzu, D.S.Y.,
Goedhals, D., Omuoyo, D.O., Maruapula, D., Foster-Nyarko, E., Lusamaki, E.K., Simulundu,
E., Ong'era, E.M., Ngabana, E.N., Shumba, E., Fahime, E.E., Lokilo, E., Mukantwari, E.,
Philomena, E., Belarbi, E., Simon-Loriere, E., Anoh, E.A., Leendertz, F., Ajili, F., Enoch,
F.O., Wasfi, F., Abdelmoula, F., Mosha, F.S., Takawira, F.T., Derrar, F., Bouzid, F.,
Onikepe, F., Adeola, F., Muyembe, F.M., Tanser, F., Dratibi, F.A., Mbunsu, G.K., et al., A
year of genomic surveillance reveals how the SARS-CoV-2 pandemic unfolded in Africa.
Science 0(0), eabj4336.
Worobey, M., Pekar, J., Larsen, B.B., Nelson, M.I., Hill, V., Joy, J.B., Rambaut, A., Suchard, M.A.,
Wertheim, J.O., Lemey, P., 2020. The emergence of SARS-CoV-2 in Europe and North
America. Science 370(6516), 564-570.

Supplementary Material for

Transmission networks of SARS-CoV-2 in coastal Kenya during the first two waves: a retrospective genomic study

Agoti et al, 2021

Corresponding author:
Dr Charles N. Agoti
Email : cnyaigoti@kemri-wellcome.org

Contents

Additional available material relating to this manuscript

Supplementary-Figure 1. Flow of samples analysed in this study.

Supplementary-Figure 2. Nucleotide differences in SARS-CoV-2 genome sequences from the coastal Kenya samples compared to the reference genome Wuhan-Hu-1 (Accession #: NC_045512.2). Vertical coloured bars show the nucleotide differences from the reference genome; orange is a change to A, crimson is a change to T, indigo a change to G, and slate blue a change to C. Grey is a nucleotide deletion or an area not sequenced in our labouratory due to amplicon drop-off or inconclusive assembly.

Supplementary-Figure 3. Mutation-resolved global phylogenetic tree showing the genetic closeness of the sequences from coastal Kenya while including a representative global sub-sample. The tree uses the same data as Figure 5A in the main test (time-resolved tree) by combining 3,958 global sequences and 732 coastal Kenya sequences. The coastal Kenya sequences are indicated by a filled circle symbol, coloured by the assigned Pango lineage.

Supplementary-Figure 4. An alluvium plot showing the number of viral imports and exports from coastal Kenya over the study period. This figure, unlike Figure 8A , provides the details of the countries from which import to or exports from coastal Kenya occurred. Exports from coastal Kenya are depicted by a purple colour, imports to costal Kenya are depicted by yellow colour, within coastal Kenya intercounty spread is indicated by the cadet blue colour.

Supplementary-Figure 5. An alluvium plot showing the number of viral imports and exports from coastal Kenya over the study period using a subsample of the coastal Kenya genomes (n=360). The sub-sample was randomly selected aiming to be spatial-temporally representative with up to 15 genomes per county per month over the study period. Exports are depicted by purple colour, imports are depicted by yellow colour, within coastal Kenya intercounty spread is indicated by the cadet blue colour.

Supplementary-Figure 6. Position of non-synonymous mutations in the 747 genomes from coastal Kenya compare with the Wuhan-Hu-1 reference genome (Accession number NC_045512). Only changes occurring in >1% of genomes are shown. The percentage frequency of the changes in the 747 genomes is shown on the y-axis.

Supplementary-Figure 7. Genomic variation characteristics of the four lineages first identified in Kenya. Panel A, root-to-tip regression plots for the four lineages first identified in Kenya. Panel B, Genome maps of the "Kenya" lineages where the spike region is shown in detail and in colour and the rest of the genome is shown in grey colour. Red marks indicate where sequencing of Kenya strains resulted in ambiguous nucleotides. Plots generated using tool available from Stanford University Web-page Coronavirus antiviral and resistance database https://covdb.stanford.edu/page/mutation-viewer/: last accessed on 2021-08-03.

Supplementary-Table 1. Summary output from separate runs of the import/export ancestral state reconstruction (ASR) analysis

Characteristic	Run 1[#]	Run 2[£]	Run 3[$]
Total Location transition events	357	341	217
International imports			
Total	69	65	79
Through Mombasa	40	38	36
Through Kwale	9	5	17
Through Taita Taveta	13	13	20
Through Kilifi	5	5	17
Through Lamu	1	1	1
International exports			
Total	93	87	25
Through Mombasa	86	81	22
Through Kwale	6	3	1
Through Taita Taveta	1	3	1
Through Kilifi	-	-	1
Inter-county transmission events	193	189	113

[#]Import/export analysis results from ASR combining 732 genome sequences from coastal Kenya that passed the molecular clock quality control and a random 3,958 global (outside Kenya) genomes.

[£]Import/export analysis results from ASR combining 728 genomes from coastal Kenya passing the molecular clock quality control and 3,968 global (outside Kenya) genomes.

[$] Import/export analysis results from ASR combining 338 genome sequences from coastal Kenya that passed the molecular clock quality control and a random 3979 global (outside Kenya) genomes. The coastal Kenya sub-sample used for this run 3 was randomly selected aiming to be spatial-temporally representative with up to 15 genomes per county per month over the study period.