Predictors of black fungus fear during the COVID-19 pandemic among the Bangladeshi health workers: a cross-sectional study

Authors:

Md. Kamrul Hasan[1, #, *], Humayun Kabir[1, #], Mamunur Rahman[2], Anjan Kumar Roy[3], Dipak Kumar Mitra[1]

Affiliations:

[1]Department of Public Health, North South University, Dhaka- 1229, Bangladesh.

[2]Department of Pharmacy, East West University, Dhaka-1212, Bangladesh.

[3]Department of Nursing and Health Science, Jashore University of Science and Technology, Jashore-7408, Bangladesh.

# Equal contribution

*Corresponding author:

Md. Kamrul Hasan;
MPH (In course)
Department of Public Health
North South University
Plot 15, Block B, Bashundhara
Dhaka- 1229, Bangladesh
Email: kamrul.hasan11@northosouth.edu

Abstract

Background

Raised of mucormycosis or black fungus cases middle of the COVID-19 pandemic, fear associated with black fungus may turn out to be a terrifying public health issue. This study aimed to assess the association between the fear and knowledge of black fungus and other determinants among healthcare workers in Bangladesh.

Methods

From 25[th] May 2021 to 5[th] June 2021, a cross-sectional study was carried out among healthcare workers. For data collection during the COVID-19 pandemic, a semi-structured online questionnaire was used followed by convenient and snowball sampling methods. A multivariable linear regression model was fitted to assess the association between the fear and knowledge of black fungus and other determinants.

Results

A total of 422 healthcare workers participated in this study. The results indicated that with the increased knowledge score of black fungus, the average score of black fungus fear was significantly increased ($\beta = 0.35$, 95% CI: 0.20, 0.50, $p < 0.001$). Alongside, the respondents having insomnia had a higher score of black fungus fear compare to not having insomnia ($\beta = 3.91$, 95% CI: 2.49, 5.33, $p < 0.001$) significantly. The gender, profession, and death due to COVID-19 of friends and family members had a significant effect on black fungus fear score increment.

Conclusion

Even in the COVID-19 pandemic, the black fungus fear among healthcare workers may hinder their quality of life. Our study findings recommend an emphasis on the mental health aspects and ensure support to the healthcare workers so that they can tackle the ongoing situation with lesser frictions.

Keywords: Mucormycosis, black fungus, COVID-19, fear, knowledge, insomnia, health workers.

1. Introduction

Coronavirus disease 2019 (COVID-19) also known as SARS-Cov-2, the ongoing pandemic which was initiated in December 2019 in China [1], has affected adversely Bangladesh like many other countries [2]. Up to June 4, 2021, 0.8 million cases were recorded along with around 7.5 thousand deaths by COVID-19 which was first diagnosed on 8 March 2020 in Bangladesh according to IEDCR, the country's epidemiology institute [3,4]. Mucormycosis, nowadays popularly known as black fungus in the COVID-19 era, is a treacherous fungal infection that became a new threat recently for COVID-19 survived patients which were first diagnosed in Bangladesh on May 8, 2021 [5,6].

The prevalence of black fungus is eighty times more in India during the COVID-19 pandemic [7]. The black fungus associated with COVID-19 has affected many countries like Bangladesh, Pakistan, Russia, Iran, and Brazil [8–13]. As black fungus infection in India associated with COVID-19 has raised at an alarming rate with a high mortality rate of about 50%, it is a concerning issue for Bangladesh being geographically attached with India [14]. The incidence of

black fungus infection is increasing more rapidly in the second wave than the first wave with at least 14,872 cases as of May 28, 2021 [15].

The black fungus is caused by the order Mucorales, mostly it is due to the intrusion of the genrea *Rhizopus* and *Mucor* [16,17]. People are affected by black fungus through the contact of the fungal spores as inhalation of the spore from the environment can cause infection in the lung and sinus [18]. High-risk black fungus infection determinants include immunosuppression, such as hematologic malignancy, hematopoietic stem cell transplantation (HSCT), solid organ transplantation (SOT), and diabetes mellitus [19]. Underlying disease conditions of the patient considered as a major factor for getting a primary infection of black fungus [20–22]. Nowadays COVID-19 associated black fungus infection is mostly rhino orbital cerebral type infection which is clinically manifested around the nose, eye, and brain [23]. The use of corticosteroid therapy for the management of COVID-19 patients and diabetes mellitus is currently indicated as a risk factor for black fungus infection [24]. A study found that patient suffering from COVID-19 has been affected with pulmonary mucormycosis mostly due to immunocompromised state [25]. The black fungus pathogenesis includes vessel thrombosis and subsequent tissue necrosis which causes mortality around 54% [20,26]. To prevent black fungus infection during COVID-19, avoiding the risk factors is the only option. Though the COVID-19 vaccination has started, there is no vaccine currently available for black fungus. Therefore, early diagnosis and treatment intervention is essential to treat COVID-19 induced black fungus infection effectively.

As front liners, the health care workers are an integral part of the health system who are fighting against the COVID-19 pandemic and emerging black fungus infection. A study found that inadequate hospital infection prevention strategies, workplace safety, and poor control of policies were the reasons for fear among the health care workers during any emerging outbreak [27]. In

addition, safety protocol for Bangladeshi health workers was found to be insufficient [28]. Therefore, fear might be a natural phenomenon for the Bangladeshi health care workers to tackle the COVID-19 pandemic and risen cases of black fungus, simultaneously. However, proper knowledge about the disease and initiatives may guide the health care workers of Bangladesh to eradicate fear.

To our knowledge, there is no study investigated the black fungus fear in Bangladesh among the health care worker to cope with emerging public health phenomenon. Therefore, the present study aims to investigate the association between black fungus fear and knowledge of black fungus and other determinants among health care workers during the COVID-19 pandemic in Bangladesh. To save lives by given the top priority of health care ever, the ongoing pandemic has taught us to be prepared with a capable health system. Addressing the association between black fungus fear, and knowledge of black fungus among the health care workers including the insomnia status, sociodemographic, work-related, COVID-19, and health-related information may help to take appropriate measures to improve their mental health condition.

2. Method

2.1. Study Settings

This cross-sectional study was conducted between 27th May 2021 to 3rd June 2021 in Bangladesh. The inclusion criteria include (a) the registered health care workers; doctor, nurse, and others (technologist, radiologist, and medical assistant), (b) working in clinical settings during the COVID-19 associated black fungus cases were found, (c) willing to participate in this study by an online consent, and (d) responded to all items of the questionnaire.

2.2. Data Collection Procedure

For data collection, a semi-structured online questionnaire was used in this study. We followed convenience and snowball sampling methods, due to the current situation of the COVID-19 pandemic. The data were collected via social media platforms and text messages by using a "Google Form" link. From 468 responses, a total of 422 complete responses were included in the final analysis.

2.3. Background Characteristics

The sociodemographic information such as age, gender, residence, religion, profession, marital status, having children, and living status were collected in this study. Work-related information included job type, working condition, having PPE (personal protective equipment), and directed patient care. We included COVID-19 related information of the participants; COVID-19 positive status, friend and family members' (FNF) COVID-19 positive status, and FNF COVID-19 death status. Additionally, health-related information included comorbidities status, perceived health status, and smoking status.

2.4. Comorbidities Status

The comorbidity status of the respondents was assessed by asking whether they have any comorbidities from the given options; no comorbidity, asthma, diabetes, hypertension, heart disease, cardiovascular disease, chronic kidney disease, chronic liver disease, cancer, and others. The comorbidity status was considered as "no" when the response was "no comorbidity" and the rest of the responses were considered as comorbidity status "yes".

2.5. Perceived Health Status

The World Health Organization (WHO) recommended a single-item scale ("In general, how would you rate your current health status?") that was used to measure the perceived health status in this [29]. The responses of the scale were at five options "very good," "good," "fair," "bad," or "very bad." For statistical analysis, "very good" and "good" were considered as "healthy", whereas "fair" was as "relatively healthy" [30]. Similarly, "bad" and "very bad" were considered as "unhealthy".

2.6. Knowledge Score of Black Fungus

The knowledge score was accessed by six items based on a five-point Likert scale; 1 for strongly disagree to 5 for strongly agree. The overall score was obtained by summing all the items, whereas the score range from 6 to 30. A higher score indicated a higher level of knowledge. The items are given in supplementary file-1.

2.7. Black Fungus Fear Score

A nine items measure was used to assess fear associated with the black fungus (presented in **Table 1**) that was inspired by Wheaton et al. (2012) and Blakey et al. (2015), who assessed the H1N1 and Ebola-related fear, respectively [31,32]. The items were responded from 0 (not at all) to 4 (very much), where the sum score range from 0 to 36. The higher score indicated a higher level of black fungus fear. In this current study, the nine items measure demonstrated good internal consistency with an overall reliability coefficient $\alpha = 0.82$. After deletion of each item, none of the reliability coefficients were found less than 0.78.

2.8. Data Analysis

Descriptive statistics were performed for the background characteristics of the study participants. The overall reliability coefficient of nine items' fear measure was calculated along with reliability coefficients after deletion of each item. A multivariable linear regression model was fitted to find the association between black fungus fear and knowledge of black fungus and with the background characteristics. Four hierarchical regression models were fitted to investigate the contributory role of the studied characteristics among health workers' black fungus fear. The p-value <0.05 was considered statistically significant at a 95% confident interval. Data were analyzed by using statistical software STATA-16.

2.9. Ethical Issue

The Ethical Review Broad of the Faculty of Life Science, North South University, approved this study. The review ID is not available currently. The aim and objective of the study were explained on the first page of the questionnaire and the respondents who were willing to participate by online consent were considered as respondents of this study.

3. Result

3.1. Characteristics of the Study Population

In this study, 422 health care workers participated from Bangladesh. In **Table 2,** the background characteristics of the study participants are presented**.** The mean score of knowledge was 24.67 (SD: 4.57). One-third of the participants (30.57%) were found with present insomnia. The majority of the participants (82.94%) were from age between 24 years to 30 years of old, where age between 24-27 years and 28-30 years old was 39.81% and 34.60%; nearly 4.27% were aged

less than 25 years and 21.33% were age more than 30 years old. In this study majority of them were female (52.37%) participants and the rest of them were male (47.63). Most of the respondents were from urban (88.15%) areas. In terms of religion, most of them were from Islam (80.57%); the rests were from Hindu (17.06%) and other religious backgrounds (2.37%). In this study, 45.26% were doctors, 38.15% were nurses and the rest 16.59% were others health care workers. The majority of them were married (54.27%), although only 25.12% had children. However, only 21.09% of the respondents lived alone. Almost half of the respondents did the private job (51.42%), nearly one-third did govt. job (25.83%); the rest of them did autonomic (8.53%) and NGO (14.2%) jobs. One-third of the respondents worked in COVID-19 condition (31.52%) and the majority of the respondents (65.17%) had PPE. More than half of the respondents (74.88%) involved with direct patient care. Almost one-third of the respondents (27.96%) got COVID-19 positive status; although more than two-thirds (75.53%) of their FNF got COVID-19 positive status. Even, 22.09% of the respondents' FNF died of COVID-19 infection. The number of respondents with having comorbidities status was 20.38%. Interm of perceived health status, more than two-thirds (74.17%) was perceived unhealthy, 20.85% was perceived relatively healthy, where only 4.98% was perceived healthy. In this study, most of the respondents (81.04%) were never smoked, where currents smokers were 9.48% and past smokers were 9.48%.

3.2. Association between Fear and Knowledge of Black Fungus and Other Determinants

Table 3 shows the association between black fungus fear score and knowledge score of black fungus and other determinants of health workers. In the adjusted multivariable linear regression model, the result showed that with the increase of knowledge score of black fungus, the average score of black fungus fear was significantly increased ($\beta = 0.35$, 95% CI: 0.20, 0.50, $p < 0.001$).

In this study, the insomnia of the respondents had a significant effect on the black fungus fear score. The respondents having insomnia had a higher score of black fungus fear compare to not having insomnia (β = 3.91, 95% CI: 2.49, 5.33, p <0.001). In terms of gender, the fear score was found significantly higher among the females compared to the male (β = 2.12, 95% CI: 0.56, 3.57, p <0.008). However, the black fear score was found significantly higher among the nurses compared to the doctors (β = 2.02, 95% CI: 0.14, 3.89, p <0.035). Moreover, the respondents' FNF who died of COVID-19 had found a higher black fungus fear score significantly (β = 2.58, 95% CI: 0.90, 4.26, p <0.003). The rest of the variables i.e., age residence, religion, marital status, having children, living status, job type, working condition, having PPE, direct patient care, COVID-19 positive, FNF COVID-19 positive, comorbidities status, health status, smoking status showed no significant association with the black fungus fear score after adjusting in the multivariable linear regression model.

3.3. Predictive Models of Health Care Workers' Black Fungus Fear

The hierarchical regression models are presented in **table 4**. The four models were fitted; wherein model 1, only the knowledge score, and insomnia status of the respondents were included. The sociodemographic information was included in model 2 and the work-related information was included in model 3. However, COVID-19 and health-related information were added in model 4. All of the models were found to significantly predicted the health care workers' black fungus fear. The knowledge score and insomnia status explained almost 12.20% variance of black fungus fear. With the inclusion of demographic information, the total explained variance was increased by 16.20%. However, after adding the worked-related information, the explained variance was not changed. Moreover, the inclusion of COVID-19 and health-related

information in the 4th model, 2% explained variance was increased and a total 18.20% variance of the black fungus fear was explained by the final model.

4. Discussion

This study found that black fungus fear is significantly associated with the knowledge of black fungus among the health care workers in Bangladesh. Alongside, numerous factors like insomnia status, age, gender, profession, and FNF died of COVID-19 infection were found significantly associated with black fungus fear. Albeit, the fear of infectious diseases during the outbreak has been recorded previously in many studies in various aspects [33–36].

In this study, the knowledge and black fungus fear was found significantly associated in both unadjusted and adjusted linear models. With the increase of knowledge score, the black fungus fear score was significantly increased. Similarly, Govindharaj et al. reported that knowledge of the disease and fear were strongly associated with each other [37]. Earlier in 2013, it was reported that the mortality rate due to mucormycosis was 33% but it has been reported much higher in this COVID-19 era [38]. Alongside, multiple crucial steps as removal of the organs have been reported for the mucormycosis management procedure [14]. Thus, based on the study findings, increment of knowledge of black fungus regarding the severity of black fungus for COVID-19 infection may facilitate black fungus fear.

The study revealed that black fungus fear and insomnia were significantly associated. Similarly, a study found a strong association between insomnia and COVID-19 fear among health workers [39]. Zitting et al. reported that the incidence of insomnia due to COVID-19 during the first 5 months of 2020 has been increased in United State by 58% [40]. In Bangladesh, M. Hossain et al. also reported that during the COVID-19 pandemic the sleep disturbance has been increased

[41]. Therefore, health care workers having insomnia may become prone to black fungus fear during COVID-19.

The senior age groups were found to be less fearful of black fungus compare to the junior age group, though age was found statistically insignificant in this study. This study found that black fungus fear was significantly high among females health workers compare to males. A study revealed that the prevalence of fear is more in females than males in the COVID-19 pandemic in Bangladesh [42].

We found a significant association of black fungus fear with the nursing profession comparing with other healthcare professionals. As nurses are involved more prone to direct contact-related care that may induce more fear among the nurses compared to other health workers. This working nature of the nurses may propel more fear. The Bangladeshi nurses were gone through mental health-related symptoms during the first web of the COVID-19 pandemic [43]. Similarly, Tayyib et al. reported a high level of fear and stress among the nurses during the COVID-19 context [44].

A significant relationship was found between the fear of black fungus and COVID-19 death incidence of family and friends. Family and friends are the two most important aspects of one's life, naturally one related to them thinks about their wellbeing and other aspects [41]. Again, thinking is derived from the information stored in the memory [45]. Consequently, the death memory of friends or family members was found to be associated with fearfulness.

Black fungus fear was found high among the comorbid health workers, though it was not statistically significant. Syndrome et al. reported in a systematic review that having comorbidity

is an independent risk factor of black fungus infection [6]. Therefore, the comorbid health workers may perceive more black fungus fear.

Based on this study finding, the healthcare workers are fearsome toward black fungus. The COVID-19 along with the black fungus cases risen might pose synergistic fear among the health workers, as they confront more risk. Therefore, our study recommended that the mental health of healthcare workers should be more prioritized. The arranging of necessitous programs, training, and workshop to cope with the situation can also be commenced.

5. Strength and Limitation

To our knowledge, this was the first study in Bangladesh that focuses on assessing the association between black fungus fear and knowledge of black fungus so far, which is a strength of this study. The finding of the studies can help the policymakers to improve strategic plans towards COVID-19 associated black fungus emergences in a developing country like Bangladesh.

As the nature of the cross-sectional study, causality between the variables could not explain. However, following convenient and snowball sampling methods might have some selection bias. Hence, it is recommended for further researches to assess black fungus fear on a larger scale and the sample size among the health care workers of Bangladesh to strengthen the health care system during the pandemic.

6. Conclusion

Black fungus is not a new phenomenon but the emergence of black fungus associated with COVID-19 is a concern. The healthcare workers were playing the most pivotal role in

ameliorating the COVID-19 catastrophe. Therefore, the growing concern of black fungus cases in Bangladesh during COVID-19 may induce fear among the health care workers. This study documented the nurses were significantly more feared compare to doctors. The black fungus fear was significantly associated with health care workers' knowledge of black fungus, insomnia status, gender, and friend and family COVID-19 death status. The outcomes of this study finding may help the policymakers to establish a proactive mental health support system for the healthcare workers. These might be bona fide essential to create an ecosystem from the side of healthcare workers toward patients in need of the highest intensive health support.

Funding:

The authors received no funding for this research.

Conflict of Interest:

The authors declare no competing interest regarding this study that may arise from third party.

Acknowledgments:

The authors cordially thank Sabiha Hasan and all the research assistants of the project who helped in data collection.

Reference

[1]   A. Hossain, S.M. Nasrullah, Z. Tasnim, M.K. Hasan, M.M. Hasan, Seroprevalence of SARS-CoV-2 IgG antibodies among health care workers prior to vaccine administration in Europe, the USA and East Asia: a systematic review and meta-analysis, EClinicalMedicine. 33 (2021). https://doi.org/10.1016/j.eclinm.2021.100770.

[2]   D.H. in W. Jeremy Page in Beijing, B. McKay, In hunt for Covid-19 origin, patient zero

points to second Wuhan market - WSJ, Wsj. (2021).

[3]   IEDCR, (n.d.). https://iedcr.gov.bd/ (accessed August 25, 2020).

[4]   Worldometers, Bangladesh COVID: 805,980 cases and 12,724 deaths - Worldometer, (2021).

[5]   M.A.-M. Molla, Two black fungus cases detected in Bangladesh | The Daily Star, Dly. Star. (2021). https://www.thedailystar.net/bangladesh/news/two-black-fungus-cases-detected-2098309.

[6]   A.K. Singh, R. Singh, S.R. Joshi, A. Misra, Mucormycosis in COVID-19: a systematic review of cases reported worldwide and in India, Diabetes Metab. Syndr. Clin. Res. Rev. (2021). https://doi.org/10.1016/j.dsx.2021.05.019.

[7]   A. Skiada, I. Pavleas, M. Drogari-Apiranthitou, Epidemiology and diagnosis of mucormycosis: an update, (n.d.). https://doi.org/10.3390/jof6040265.

[8]   Soutik Biswas, Mucormycosis: The "black fungus" maiming Covid patients in India - BBC News, Bbc. (2021). https://www.bbc.com/news/world-asia-india-57027829.

[9]   S.N. Sakib, Bangladesh reports 1st death by black fungus, Heal. ASIA - PACIFIC. (2021).

[10]   M.W. Bhatti, Cases of black fungus emerge across Pakistan, Thenews. (2021). https://www.thenews.com.pk/print/834117-cases-of-black-fungus-emerge-across-pakistan.

[11]   K. Zykov, Russia confirms rare, deadly 'Black Fungus'' infections seen in India,' The Moscow Times. (2021). https://www.themoscowtimes.com/2021/05/17/russia-confirms-rare-deadly-black-fungus-infections-seen-in-india-reports-a73922.

[12]   Trackpersia, Coronavirus in Iran: Power outages, black fungus, and warnings of a fifth surge - Track Persia, (2021). https://iranwire.com/en/special-features/9632.

[13]   Redação, Casos suspeitos de fungo preto são investigados no Brasil; entenda, Catracalivre. (2021). https://catracalivre.com.br/saude-bem-estar/casos-suspeitos-de-fungo-preto-sao-investigados-no-brasil-entenda/.

[14] S. Biswas, Black fungus: India reports nearly 9,000 cases of rare infection - BBC News, Bbc. (2021). https://www.bbc.com/news/world-asia-india-57217246.

[15] A. Raut, N. Tien Huy, Rising incidence of mucormycosis in patients with COVID-19: another challenge for India amidst the second wave?, (2021). https://doi.org/10.1016/S2213-2600(21)00265-4.

[16] T.L. Ray, Cutaneous manifestations of infection in the immunocompromised host, 1996. https://doi.org/10.1016/s0190-9622(96)90304-7.

[17] N.A.R.G. Joseph Heitman, Pedro W. Crous, Timothy Y. James, Barbara J. Howlett, Eva H. Stukenbrock, The fungal kingdom, 2017.

[18] Cdc, People at risk for mucormycosis | Mucormycosis | CDC, (n.d.). https://www.cdc.gov/fungal/diseases/mucormycosis/risk-prevention.html (accessed June 29, 2021).

[19] A. Serris, F. Danion, F. Lanternier, Disease entities in mucormycosis, J. Fungi (Basel, Switzerland). 5 (2019). https://doi.org/10.3390/jof5010023.

[20] M.M. Roden, T.E. Zaoutis, W.L. Buchanan, T.A. Knudsen, T.A. Sarkisova, R.L. Schaufele, M. Sein, T. Sein, C.C. Chiou, J.H. Chu, D.P. Kontoyiannis, T.J. Walsh, Epidemiology and outcome of zygomycosis: a review of 929 reported cases, Clin. Infect. Dis. 41 (2005) 643–653. https://doi.org/10.1086/432579.

[21] W. Jeong, C. Keighley, R. Wolfe, W.L. Lee, M.A. Slavin, D.C.M. Kong, S.C.A. Chen, The epidemiology and clinical manifestations of mucormycosis: a systematic review and meta-analysis of case reports, Clin. Microbiol. Infect. 25 (2019) 26–34. https://doi.org/10.1016/j.cmi.2018.07.011.

[22] A. Skiada, L. Pagano, A. Groll, S. Zimmerli, B. Dupont, K. Lagrou, C. Lass-Florl, E. Bouza, N. Klimko, P. Gaustad, M. Richardson, P. Hamal, M. Akova, J.F. Meis, J.L. Rodriguez-Tudela, E. Roilides, A. Mitrousia-Ziouva, G. Petrikkos, Zygomycosis in Europe: analysis of 230 cases accrued by the registry of the European confederation of medical mycology (ECMM) working group on zygomycosis between 2005 and 2007,

Clin. Microbiol. Infect. 17 (2011) 1859–1867. https://doi.org/10.1111/j.1469-0691.2010.03456.x.

[23]   M. Sen, S.G. Honavar, N. Sharma, M.S. Sachdev, COVID-19 and eye: a review of ophthalmic manifestations of COVID-19, Indian J. Ophthalmol. 69 (2021) 488–509. https://doi.org/10.4103/ijo.IJO_297_21.

[24]   K. Ahmadikia, S.J. Hashemi, S. Khodavaisy, M.I. Getso, N. Alijani, H. Badali, H. Mirhendi, M. Salehi, A. Tabari, M. Mohammadi Ardehali, M. Kord, E. Roilides, S. Rezaie, The double-edged sword of systemic corticosteroid therapy in viral pneumonia: a case report and comparative review of influenza-associated mucormycosis versus COVID-19 associated mucormycosis, Mycoses. (2021). https://doi.org/10.1111/myc.13256.

[25]   K. Alekseyev, L. Didenko, B. Chaudhry, Rhinocerebral mucormycosis and COVID-19 pneumonia, J Med Cases. 12 (2021) 85–89. https://doi.org/10.14740/jmc3637.

[26]   A.S. Ibrahim, B. Spellberg, T.J. Walsh, D.P. Kontoyiannis, Pathogenesis of mucormycosis, Clin. Infect. Dis. 54 (n.d.) S16–S22. https://doi.org/10.1093/cid/cir865.

[27]   Ogolodom MP, Knowledge, attitudes and fears of health care workers towards the Corona Virus Disease (COVID-19) pandemic in South-South, Nigeria, Heal. Sci. J. 14 (2020). https://doi.org/10.36648/1791-809X.S1.002.

[28]   K.N. Uddin, Physician safety in work place, BIRDEM Med. J. 10 (2019) 1–6. https://doi.org/10.3329/birdem.v10i1.44739.

[29]   Health interview surveys towards international harmonization, WHO Reg. Publ. (1996). https://www.euro.who.int/en/publications/abstracts/health-interview-surveys.-towards-international-harmonization (accessed June 21, 2021).

[30]   S. Wu, R. Wang, Y. Zhao, X. Ma, M. Wu, X. Yan, J. He, The relationship between self-rated health and objective health status: a population-based study, BMC Public Health. 13 (2013) 1–9. https://doi.org/10.1186/1471-2458-13-320.

[31]   M.G. Wheaton, J.S. Abramowitz, N.C. Berman, L.E. Fabricant, B.O. Olatunji,

Psychological predictors of anxiety in response to the H1N1 (swine flu) pandemic, Cognit. Ther. Res. 36 (2012) 210–218. https://doi.org/10.1007/s10608-011-9353-3.

[32]   S.M. Blakey, L. Reuman, R.J. Jacoby, J.S. Abramowitz, Tracing "Fearbola": psychological predictors of anxious responding to the threat of Ebola, Cognit. Ther. Res. 39 (2015) 816–825. https://doi.org/10.1007/s10608-015-9701-9.

[33]   G. Pappas, I.J. Kiriaze, P. Giannakis, M.E. Falagas, Psychosocial consequences of infectious diseases, Clin. Microbiol. Infect. 15 (2009) 743–747. https://doi.org/10.1111/j.1469-0691.2009.02947.x.

[34]   M.A. Mamun, N. Sakib, D. Gozal, A.K.M. Israfil, S. Hossain, F. Al, I. Hosen, M. Binte, A. Hasnat, A. Sarker, I. Rayhan, T. Sikder, The COVID-19 pandemic and serious psychological consequences in Bangladesh : a population-based nationwide study, J. Affect. Disord. 279 (2021) 462–472. https://doi.org/10.1016/j.jad.2020.10.036.

[35]   S. Singh, D. Roy, K. Sinha, S. Parveen, G. Sharma, G. Joshi, Impact of COVID-19 and lockdown on mental health of children and adolescents: a narrative review with recommendations, Psychiatry Res. 293 (2020) 113429. https://doi.org/10.1016/j.psychres.2020.113429.

[36]   M. Iorga, R. Iurcov, L.-M. Pop, The relationship between fear of infection and insomnia among dentists from Oradea metropolitan area during the outbreak of Sars-CoV-2 pandemic, J. Clin. Med. 10 (2021) 2494. https://doi.org/10.3390/jcm10112494.

[37]   P. Govindharaj, S. Srinivasan, J. Darlong, Perception toward the disease of the people affected by leprosy, Int. J. Mycobacteriology. 7 (2018) 247–250. https://doi.org/10.4103/ijmy.ijmy_66_18.

[38]   H.L. Hong, Y.M. Lee, T. Kim, J.Y. Lee, Y.S. Chung, M.N. Kim, S.H. Kim, S.H. Choi, Y.S. Kim, J.H. Woo, S.O. Lee, Risk factors for mortality in patients with invasive mucormycosis, Infect. Chemother. 45 (2013) 292–298. https://doi.org/10.3947/ic.2013.45.3.292.

[39]   K.A. Cawcutt, R. Starlin, M.E. Rupp, Fighting fear in healthcare workers during the

COVID-19 pandemic, Infect. Control Hosp. Epidemiol. 41 (2020) 1192–1193. https://doi.org/10.1017/ice.2020.315.

[40] K.M. Kirsi-Marja Zitting, H.M. Lammers-Van Der Holst, R.K. Yuan, W. Wang, S.F. Quan, J.F. Duffy, Google trends reveals increases in internet searches for insomnia during the 2019 coronavirus disease (COVID-19) global pandemic, J. Clin. Sleep Med. 17 (2021) 177–184. https://doi.org/10.5664/JCSM.8810.

[41] T. Ara, M. Rahman, M.A. Hossain, A. Ahmed, J. Lai, R.M. Garcia, T. Jain, Identifying the associated risk factors of sleep disturbance during the COVID-19 lockdown in Bangladesh: a web-based survey, Article. 11 (2020) 1. https://doi.org/10.3389/fpsyt.2020.580268.

[42] M.A. Hossain, M.I.K. Jahid, K.M. Amran Hossain, L.M. Walton, Z. Uddin, M.O. Haque, M.F. Kabir, S.M. Yasir Arafat, M. Sakel, R. Faruqui, Z. Hossain, Knowledge, attitudes and fear of COVID-19 during the rapid rise period in Bangladesh, PLoS One. 15 (2020) e0239646. https://doi.org/10.1371/journal.pone.0239646.

[43] S.R. Chowdhury, T.C. Sunna, D.C. Das, H. Kabir, A. Hossain, S. Mahmud, S. Ahmed, Mental health symptoms among the nurses of Bangladesh during the COVID-19 pandemic, Middle East Curr. Psychiatr. 28 (2021) 23. https://doi.org/10.1186/s43045-021-00103-x.

[44] N.A. Tayyib, F.J. Alsolami, Measuring the extent of stress and fear among registered nurses in KSA during the COVID-19 outbreak, J. Taibah Univ. Med. Sci. (2020) 410–416. https://doi.org/10.1016/j.jtumed.2020.07.012.

[45] A. Cuncic, What happens to your body when you're thinking?, Verywellmind. (2019). https://www.verywellmind.com/what-happens-when-you-think-4688619.

Tables

Table 1: Item Properties of the Black Fungus Measure

Items	Mean	SD	α if item deleted
1. To what extent are you concerned about black fungus?	2.59	1.13	0.79
2. To what extent do you believe that black fungus could become a ''epidemic'' in Bangladesh?	2.37	1.15	0.80
3. How likely is it that you could become infected with black fungus?	1.52	1.12	0.81
4. How likely is it that someone you know could become infected with black fungus?	1.68	1.26	0.81
5. How much exposure have you had to information about black fungus?	1.73	1.03	0.84
6. If you did become infected with black fungus, to what extent are you concerned that you will be severely ill?	2.69	1.13	0.80
7. To what extent has the threat of black fungus influenced your decisions to be around people?	2.02	1.29	0.80
8. To what extent has the threat of black fungus influenced your travel plans?	2.28	1.38	0.78
9. To what extent has the threat of black fungus influenced your use of safety behaviors (e.g., hand sanitizer)?	2.33	1.33	0.80

Table 2: Characteristics of the Study Participants (n=422)

Sample characteristics	Sample distribution (n=frequency)	Percentage (%)/Mean (SD)
Black fungus fear score	422	19.21 (6.98)
Knowledge score	422	24.67 (4.57)
Insomnia status		
Yes	129	30.57
No	293	69.43
Age		
<24	18	4.27
24-27	168	39.81
28-30	146	34.60
>30	90	21.33
Gender		
Male	201	47.63
Female	221	52.37
Residence		
Urban	372	88.15
Rural	50	11.85
Religion		
Islam	340	80.57
Hindu	72	17.06
Others	10	2.37
Profession		
Doctor	191	45.26
Nurse	161	38.15
Others	70	16.59
Marital status		
Unmarried	193	45.73
Married	229	54.27
Having children		
Yes	106	25.12
No	316	74.88
Living status		
Alone	89	21.09
With others	333	78.91
Job type		
Autonomic	36	8.53
Govt.	109	25.83
NGO	60	14.2
Private	217	51.42
Working condition		
COVID-19	133	31.52
Non-COVID-19	289	68.48
Having PPE		
Yes	275	65.17
No	147	34.83
Direct patient care		
Yes	316	74.88
No	106	25.12
COVID-19 positive		
Yes	118	27.96
No	304	72.04
FNF COVID-19 positive		
Yes	318	75.53

No	103	24.47
FNF COVID-19 death		
Yes	93	22.09
No	328	77.91
Comorbidities status		
Yes	86	20.38
No	336	79.62
Health status		
Healthy	21	4.98
Relatively healthy	88	20.85
Unhealthy	313	74.17
Smoking status		
Current smoker	40	9.48
Never smoker	342	81.04
Past smoker	40	9.48

Table 3: Association between Fear and Knowledge of Black Fungus and with Other Determinants (n=422)

Sample Characteristics	Unadjusted β (95% CI)	p-value	Adjusted β (95% CI)	p-value
Knowledge score	0.36 (0.22, 0.50)	**<0.001**	0.35 (0.20, 0.50)	**<0.001**
Insomnia status				
Yes	3.94 (2.53, 5.34)	**<0.001**	3.91 (2.49, 5.33)	**<0.001**
No	Reference		Reference	
Age				
<24	Reference		Reference	
24-27	-1.59 (-5.00, 1.82)	0.359	-2.31 (-5.61, 0.99)	0.169
28-30	-1.56 (-4.99, 1.88)	0.374	-2.03 (-5.52, 1.46)	0.254
>30	-0.80 (-4.35, 2.75)	0.658	-1.87 (-5.66, 1.92)	0.332
Gender				
Male	Reference		Reference	
Female	2.78 (1.47, 4.10)	**<0.001**	2.12 (0.56, 3.67)	**0.008**
Residence				
Urban	1.97 (-0.10, 4.03)	0.061	1.63 (-0.42, 3.68)	0.119
Rural	Reference		Reference	
Religion				
Islam	Reference		Reference	
Hindu	1.17 (-0.61, 2.94)	0.197	0.26 (-1.49, 2.01)	0.771
Others	4.08 (-0.31, 8.47)	0.068	3.25 (-0.96, 7.45)	0.130
Profession				
Doctor	Reference		Reference	
Nurse	0.48 (-0.99, 1.95)	0.521	2.02 (0.14, 3.89)	**0.035**
Others	-1.08 (-3.10, 0.84)	0.268	1.99 (-0.211, 4.19)	0.076
Marital status				
Unmarried	Reference		Reference	
Married	-0.09 (-1.44, 1.25)	0.891	-0.20 (-1.82, 1.42)	0.805
Having children				
Yes	0.06 (-1.49, 1.60)	0.944	0.32 (-1.51, 2.14)	0.734
No	Reference		Reference	
Living status				
Alone	Reference		Reference	
With others	0.53 (-1.11, 2.17)	0.528	0.70 (-.89, 2.30)	0.346
Job type				
Autonomic	Reference		Reference	
Govt.	0.22 (-2.42, 2.85)	0.873	-0.32 (-3.05, 2.41)	0.815
NGO	-1.73 (-4.62, 1.17)	0.241	-0.60 (-3.47, 2.28)	0.684
Private	-0.40 (-2.87, 2.07)	0.748	-0.18 (-2.57, 2.21)	0.881
Working condition				
COVID-19	Reference		Reference	
Non-COVID-19	-0.35 (-1.79, 1.09)	0.636	0.24 (-1.32, 1.81)	0.761
Having PPE				
Yes	0.94 (-0.46, 2.34)	0.186	0.77 (-.76, 2.30)	0.325
No	Reference		Reference	
Direct patient care				
Yes	0.58 (-0.97, 2.12)	0.464	0.14 (-1.44, 1.71)	0.864
No	Reference		Reference	
COVID-19 positive				
Yes	0.25 (-1.25, 1.74)	0.747	-0.16 (-1.66, 1.35)	0.838
No	Reference		Reference	

FNF COVID-19 positive				
Yes	-0.672 (-2.23, 0.89)	0.396	-1.00 (-2.59, 0.57)	0.209
No	Reference		Reference	
FNF COVID-19 death				
Yes	2.48 (0.89, 4.07)	**0.002**	2.58 (0.90, 4.26)	**0.003**
No	Reference		Reference	
Comorbidities status				
Yes	0.13 (-1.53, 1.79)	0.881	0.09 (-1.57, 1.76)	0.912
No	Reference		Reference	
Health status				
Healthy	Reference		Reference	
Relatively healthy	0.68 (-2.66, 4.02)	0.689	0.94 (-2.28, 4.16)	0.566
Unhealthy	0.03 (-3.07, 3.13)	0.984	1.50 (-1.56, 4.57)	0.336
Smoking status				
Current smoker	Reference		Reference	
Never smoker	1.21 (-1.09, 3.50)	0.301	-0.83 (-3.25, 1.58)	0.499
Past smoker	0.35 (-2.72, 3.42)	0.823	-0.09 (-3.06, 2.88)	0.953

Table 4: Predictive Models of Black Fungus Fear (n=422)

Variables	Model 1 (F = 29.01, R^2 = 0.122, adjusted R^2 = 0.118, p <0.001) β	95% CI	Model 2 (F =7.92, R^2 = 0.162, adjusted R^2 = 0.141, p <0.001) β	95% CI	Model 3 (F = 5.72, R^2 =0.164, adjusted R^2 = 0.136, p <0.001) β	95% CI	Model 4 (F =4.44, R^2 =0.182, adjusted R^2 =0.141, p <0.001) β	95% CI
Knowledge score	0.36	0 .22, 0.49	0.36	0.22, 0.51	0.36	0.21, 0.50	0.34	0.19, 0.48
Insomnia status[a]	-3.90	-5.26, -2.54	-3.82	-5.18, -2.46	-3.87	-5.24, -2.50	-3.99	-5.39, -2.58
Age[b]			-0.06	-0.94, 0.82	-0.10	-1.00, 0.81	-0.23	-1.16, 0.70
Gender[c]			-2.05	-3.35, -0.76	-1.98	-3.29, -0.66	-1.83	-3.15, -0.51
Residence[d]			1.67	-0.28, 3.61	1.66	-0.31, 3.62	1.66	-0.32, 3.64
Religion[e]			0.75	-0.63, 2.13	0.72	-0.66, 2.11	0.93	-0.46, 2.33
Profession[f]			0.79	-0.17, 1.75	0.71	-0.28, 1.69	0.87	-0.16, 1.90
Marital status[g]			-0.44	-2.00, 1.12	-0.40	-1.97, 1.16	-0.35	-1.92, 1.21
Having children[h]			0.44	-1.29, 2.17	0.50	-1.25, 2.24	0.57	-1.17, 2.32
Living status[i]			0.82	-0.74, 2.38	0.85	-0.72, 2.42	0.88	-0.70, 2.46
Job type[j]					-0.03	-0.65, 0.59	-0.11	-0.74, 0.51
Working condition[k]					0.24	-1.29, 1.77	0.11	-1.42, 1.64
Having PPE[l]					0.77	-0.70, 2.24	0.57	-0.93, 2.06
Direct patient care[m]					0.33	-1.18, 1.86	0.23	-1.29, 1.76
COVID-19 positive[n]							-0.23	-1.71, 1.26
FNF COVID-19 positive[o]							-0.96	-2.52, 0.61
FNF COVID-19 death[p]							2.40	0.76, 4.04
Comorbidities status[q]							0.21	-1.42, 1.83
Health status[r]							0.65	-0.54, 1.84
Smoking status[s]							-0.01	-1.46, 1.48

[a, h, l, m, n, o, p]1= Yes, 2= No. [b]1= <25, 2= 24-27, 3= 28-30, 4= >30. [c]1= Male, 2= Female, [d]1= Urban, 2= Rural. [e]1= Islam, 2= Hindu, 3= Others. [f]1= Doctor, 2= Nurse, 3= Others. [g]1= Unmarried, 2= Married. [i]1= Alone, 2= With others. [j]1= Autonomic, 2= Govt., 3= NGO, 4= Private. [k]1= COVID-19, 2= Non-COVID-19. [q]1= Healthy, 2= Relatively healthy, 3= Unhealthy. [r]1= Current smoker, 2 = Never smoker, 3= Past smoker.