1

Relating SARS-CoV-2 shedding rate in wastewater to daily positive tests data:

A consistent model based approach

M. Petala[1], M. Kostoglou[2], Th. Karapantsios[2], Ch. Dovas[3], Th. Lytras[4,5], D. Paraskevis[6], E. Roilides[7], A. Koutsolioutsou-Benaki[8], , G. Panagiotakopoulos[4], V. Sypsa[6], S. Metallidis[9], A. Papa[10], E. Stylianidis[11], A. Papadopoulos[12], S. Tsiodras[6], N. Papaioannou[3]

[1]Laboratory of Environmental Engineering & Planning, Department. of Civil Engineering, Aristotle University of Thessaloniki, Thessaloniki, 54 124, Greece

[2]Laboratory of Chemical and Environmental Technology, Dept. of Chemistry, Aristotle University of Thessaloniki, 54 124 Thessaloniki, 54124, Greece

[3]Faculty of Veterinary Medicine, School of Health Sciences, Aristotle University of Thessaloniki, Thessaloniki, 54124, Greece

[4]National Public Health Organization, Athens, Greece

[5]European University Cyprus, Nicosia, Cyprus

[6]National and Kapodistrian University of Athens, Athens, Greece

[7]Infectious Diseases Unit and 3rd Department of Pediatrics, Aristotle University School of Health Sciences, Hippokration Hospital, Thessaloniki, 54642, Greece

[8]Department of Environmental Health, Directory of Epidemiology and Prevention of Non Communicable Diseases and Injuries, National Public Health Organization, Athens, Greece

[9]Department of Haematology, First Department of Internal Medicine, Faculty of Medicine, AHEPA General Hospital, Aristotle University of Thessaloniki, Thessaloniki, 54636, Greece

[10]Department of Microbiology, Medical School, Aristotle University of Thessaloniki, Thessaloniki, 54124 Greece

[11]School of Spatial Planning and Development, Faculty of Engineering, Aristotle University of Thessaloniki, 54124, Greece

[12]EYATH S.A., Thessaloniki Water Supply and Sewerage Company S.A., Thessaloniki, 54636, Greece

ABSTRACT

During the COVID-19 pandemic, wastewater-based epidemiology (WBE) has been engaged to

complement medical surveillance and in some cases to also act as an early diagnosis indicator of viral

spreading in the community. Most efforts worldwide by the scientific community and commercial

companies focus on the formulation of protocols for SARS CoV-2 analysis in wastewater and approaches

addressing the quantitative relationship between WBE and medical surveillance are lacking. In the

present study, a mathematical model is developed which uses as input the number of daily positive

medical tests together with the highly non-linear shedding rate curve of individuals to estimate the

evolution of virus shedding rate in wastewater along calendar days. A comprehensive parametric study

by the model using as input actual medical surveillance and WBE data for the city of Thessaloniki

(~700,000 inhabitants, North Greece) during the outbreak of November 2020 reveals the conditions

under which WBE can be used as an early warning tool for predicting pandemic outbreaks. It is shown

that early warning capacity is different along the days of an outbreak and depends strongly on the

number of days apart between the day of maximum shedding rate of infected individuals in their

disease cycle and the day of their medical testing. The present data indicate for Thessaloniki an average

early warning capacity of around 2 days. Moreover, the data imply that there exists a proportion

between unreported cases (asymptomatic persons with mild symptoms that do not seek medical advice)

and reported cases. The proportion increases with the number of reported cases. The early detection

capacity of WBE improves substantially in the presence of an increasing number of unreported cases.

For Thessaloniki at the peak of the pandemic in mid-November 2020, the number of unreported cases

reached a maximum around 4 times the number of reported cases.

HIGHLIGHTS

• Model estimates viral load evolution in wastewater from infected people dynamics

• Identifying actual conditions for which WBE can be used as an early warning tool

• Early warning capacity increases with an increasing number of unreported cases

• In Thessaloniki Nov20 outbreak, the early warning capacity of WBE was about 2 days

• In Thessaloniki Nov20 outbreak, unreported cases were up to 4 times reported cases

INTRODUCTION

In the COVID-19 pandemic, wastewater-based epidemiology (WBE) has become an important

tool, supplementing public health surveillance, in the hands of scientists, medical experts, and state

officials to evaluate the spread of SARS-CoV-2 in the community [la Rosa et al., 2020; Medema et al., 2020b; Wurtzer1 et al.,

n.d.]. This was more so after the Centers for Disease Control and Prevention (CDC) adopted wastewater

disease surveillance as a valid monitoring tool and provided guidance and recommendations for the

selection and application of testing methods (CDC, 2021). The latter cover issues from sample collection

and sample processing to RNA measurement and use of laboratory controls for the estimation of the

performance of the applied methods and of data quality. Laboratory controls deal with difficulties

arising from the chemical and biological variability of wastewater quality across different places and

over time, e.g. due to season, weather, human activities etc., and also cope with problems of proper

RNA extraction and quantification, elimination of inhibition and contamination of reagents.

The so-called matrix recovery controls suggested by CDC refer exclusively to the amount of virus

lost during sample processing. However, it is well known that viruses strongly adsorb, and so get

inaccessible ("lost"), in the pores of solid particles suspended in wastewater in sewerage networks

(Sellaoui et al., 2020; Ye, 2018). Therefore, if WBE studies aim to quantify the virus shedding rate at the

source (households) from analysis of samples taken at the entrance of wastewater treatment plants

(WTP) then it is of paramount importance to appraise the amount of virus lost in sewerage networks.

Determining the extent of recovery, as suggested by CDC, by spiking techniques in samples obtained at

the entrance of WTP is incapable of describing the amount of virus lost in sewerage networks. This is

because such samples have traveled in sewerage pipes for hours and so have (most of) their adsorption

sites saturated with adsorbed species. Desorption of spiked material would have been completely

different if a sample had been taken fresh at a shedding spot, i.e., from the sanitary plumbing system of

a building.

Recently, a mathematical model has been proposed to account for virus loss in sewerage

networks by means of adsorption of virus on porous particles suspended in wastewater(Petala et al.,

2021). To do so, the model rationalizes the SARS-CoV-2 concentration in samples taken at the entrance

of WTP with respect to certain quality characteristics of wastewater samples. Rationalization is based on

rigorous physicochemical phenomena in adsorption (and not on a statistical dependence of virus

concentration on wastewater parameters), also including the effect of large-scale topological complexity

of actual sewage networks. At the examined period of time (April to June 2020), the rationalized

decreasing shedding rate was in agreement with the observed clinical conditions, contrary to the non-

rationalized data which showed a different picture. Yet, even with rationalized data it is still difficult to

make the critical step ahead and associate the virus shedding rate with the number of cases.

There are several reasons for the discrepancy between viral shedding rate in wastewater and

number of infected people (cases) reported by public health surveillance. Limitations in the capacity of

medical testing, strict criteria for the application of medical testing and parts of the population being

reluctant to seek medical care are responsible for underdiagnosis of cases, mainly asymptomatic cases

or patients with mild symptoms who are not tested. These factors lead to reporting a lower number of

cases than the actual one. On the other hand, poor recovery in the collected wastewater samples and

virus loss in sewage networks, along with ineffectiveness to properly rationalize for the latter, lead to a

smaller measured concentration of virus in the collected samples and, thus, to a lower estimated viral

shedding rate than the actual one.

The time delay between the onset of patients' viral shedding in infected people and symptoms

onset that prompts patients being tested, is another reason for discrepancy. Several reports in literature

indicate a time delay up to 8 days between wastewater signal and infected cases reported in public

health surveillance system (e.g.(D'Aoust et al., 2021; Nemudryi et al., 2020; Peccia et al., 2020)). This is

in line with the reported incubation period of around 6 days from exposure/infection to onset of

symptoms of COVID-19 (Guan et al., 2020; Lauer et al., 2020; Li et al., 2020). Therefore, the discrepancy

between the onset of viral shedding and the onset of symptoms is actually an advantage that might

explain the early detection capacity of WBE.

Another important reason of discrepancy between daily reports of wastewater viral titers and

daily reports of new confirmed cases is the cumulative character of viral shedding rate in wastewater

during the disease. Several clinical studies 14–16 showed that viral shedding is not uniform during the

disease, but there is a maximum shedding rate in the very acute phase of infection, probably even

before respiratory symptoms appear, being followed by an exponential decline in subsequent days. If

one considers that shedding lasts at least 3-4 weeks after the inception of symptoms 14,15. it is

apparent that daily wastewater viral titers correspond more to the cumulative shedding of infected

people rather than to the shedding of the daily new cases. It must be stressed here that the evidence

from clinical studies is limited, and only for samples collected from hospitalized patients, thus after

symptoms onset, so one should be extremely careful in estimating shedding rates at the first days right

after infection.

If one aims to associate wastewater surveillance data with public health surveillance data, the

different sources of bias between these data should be also taken into account. Laboratory practice

shows that variation in wastewater data is considerably high from one day to the next because of the

many experimental parameters that are involved, e.g., individual's shedding rate (orders of magnitude

variation among infected people(Zheng et al., 2020); sampling protocol, sample concentration

techniques, RNA extraction methodology, normalization with respect to population indicators and flow

rate, inhibition assessment in RNA recovery, RNA quantification, etc. On the other hand, public health

surveillance is susceptible to a systematic bias because of limitations in the capacity of medical testing

along with the unpredictable human behavior under stressful conditions. In principle, one should be

concerned about the correctness of both wastewater viral measurements and clinical testing data.

The goal of this work is to setup a methodology for the consistent comparison between the

number of infected people reported by clinical surveillance and SARS-CoV-2 RNA concentration in

wastewater samples. To do so, a model is developed for the estimation of the evolution of virus

shedding rate in a sewage system based on the daily positive medical tests. Apart from corrections

based on the laboratory controls suggested by CDC, virus shedding rate data are further rationalized

with respect to physicochemical parameters of wastewater to account for virus loss by adsorption to

sewage solids (Petala et al., 2021). To increase accuracy, clinical testing data are based on the date of

specimen collection and not on the reporting date. The model is first developed in a generalized

continuous time regime and then it is transformed to its discrete counterpart dictated by the daily basis

data.

METHODS

Sampling

Wastewater samples were collected at the exit of Thessaloniki's (a city at North Greece) main sewerage

pipe, right before the entrance at the wastewater treatment plant of the city, as described

elsewhere(Petala et al., 2021). This plant serves an estimated population of about 700,000 people. The

present work reports 24-hours composite samples (1L each) taken three times per week (Monday-

Wednesday-Friday) from October 5[th], 2020 until January 6[th], 2021. This period includes the second wave

burst of COVID-19 for Thessaloniki in November 2020 which was the worst ever in the whole country.

Typical range of values of the physicochemical parameters of wastewater samples for the examined

period are displayed in Table 1 (supplementary materials). These parameters were employed in the

rationalization of the measured viral concentration with respect to quality characteristics of wastewater

according to Petala et al. (Petala et al., 2021).

Virus concentration

Upon wastewater collection, 200 mL of each wastewater sample was centrifuged at 4000xg for 30 min

to remove particles and pH of the supernatant was adjusted to 4 using a solution of 2.0 M HCl. An

aliquot of 40 mL supernatant was then passed through an electronegatively charged surface of 0.45 μm-

Ø47 mm cellulose nitrate HA membrane ((HAWP04700; Merck Millipore Ltd., Tullagreen, Ireland).

Filtration was performed using a magnetic funnel mounted on a glass filtration flask (Pall Corporation)

Filtration step was conducted in triplicate and membranes were stored in 15 mL falcon tubes for further

processing of RNA extraction and virus quantification.

RNA extraction and virus quantification

For each sewage sample, three electronegative membranes were individually subjected to phenol-

chloroform-based RNA extraction process (Chaintoutis et al., 2019) coupled with magnetic bead binding.

Each membrane filter was rolled into a Falcon™ 15 mL conical centrifuge tube with the top side facing

inward. The following were added sequentially, a) 900 μL of guanidinium isothiocyanate-based "Lysis

buffer I" [5M guanidinium isothiocyanate, 25mM EDTA, 25mM sodium citrate (pH 7.0), 25mM

phosphate buffer (pH 6.6)] containing 1% N-Lauroylsarcosine, 2% Triton X-100, 2% CTAB and 2% PVP, b)

18μl β-mercaptoethanol, c) 300μl H2O, mixed thoroughly by inversion, and each tube was incubated at

4 °C on horizontal rotator (50 rpm) for 10-30 min. 1200 µl of "Lysis buffer II" were added [prepared by

mixing 152.5gr guanidinium hydrochloride, 31.25 ml of 2M acetate buffer (pH 3.8) and water-saturated

phenol stabilized (pH4), up to 500ml final volume], followed by incubation 10 min / RT on horizontal

rotator (150 rpm). The liquid phase was transferred on a 2-mL microcentrifuge tube and clarified by

centrifugation (21,000×g, 5 min, 4 °C). 1600 µl were transferred to a new tube, 200 µl chloroform-

isoamyl alcohol (24:1) were added and shaken vigorously for 30 s followed by incubation (–20 °C, 30

min) and centrifugation (21,000×g, 10 min, 4 °C). 800 µl of the upper aqueous phase were transferred

and mixed with 667µl isopropanol and 20µl of magnetic beads (IDEXX Water DNA/RNA Magnetic Bead

Kit), followed by incubation 15 min / RT on horizontal rotator (150 rpm). The beads were washed

according to the manufacturer's protocol, using the magnetic extraction reagents. Elution of RNA was

done in 100 µl buffer followed by filtering using OneStep PCR inhibitor removal kit (Zymoresearch).

SARS-CoV-2 quantified RNA (1x 106.3 genomic copies) from human clinical samples was added to a

subset of concentrates to estimate the recovery efficiency and reproducibility of the RNA extraction

procedure. Similarly, heat inactivated SARS-CoV-2 from human clinical samples was spiked (1x 107 viral

particles) in a subset of sewage samples to assess the recovery and reproducibility of the virus

concentration procedure.

Each RNA extract was subjected to real-time RT-PCR for SARS-CoV-2 quantification in triplicates. Two

primer/probe sets were utilized: the N2 set from CDC that targets the nucleocapsid (N) gene and the set

targeting the genomic region that encodes the E protein (Corman et al., 2020). The assays were

performed on a CFX96 Touch™ Real-Time PCR Detection System (Bio-Rad Laboratories, Hercules, CA,

USA). Reactions were considered positive if the cycle threshold was below 40 cycles. Calibration curves

were generated using the synthetic single-stranded RNA standard "EURM-019" (Joint Research Centre,

European Commission). The possible presence of RT-PCR inhibitors in each RNA extract was assessed in

duplicates using spiked EURM-019 included at ~1000 copies in additional RT-PCR reactions. Inhibition

was expressed as % reduction in reported copy number, compared to the sum of spiked EURM-019

copies and the mean value of measured SARS-CoV-2 genomic copies in the non-spiked RNA. SARS-CoV-2

viral load in each sewage sample was expressed as mean ± standard deviation genome copies per liter,

after correcting for RT-PCR inhibition if present, and recovery efficiencies (virus concentration and RNA

extraction). Precision of each individual sewage sample quantification was assessed using the coefficient

of variation (CV) of the estimated SARS-CoV-2 viral load of the three electronegative membranes

processed. We set our precision threshold at 35% CV for each sewage sample measurement.

Epidemiological data

Daily numbers of COVID-19 infected people reported in the city of Thessaloniki, adjusted to specimen

collection date, were obtained from the National Public Health Organization (National Public Health

Organization, 2020). These data reflect residents found positive when tested in public and private

laboratories. Data are presented for the period between September 1st, 2020 and January 6th, 2021

(original data are provided in supplementary materials).

Problem formulation

A model is developed for estimating the evolution of virus shedding rate to sewage system based on the

official (announced by the state) results of daily positive tests (cases), registered by the date of

specimen collection and not by the date of public reporting. There is typically a delay of few days

between specimen collection and public reporting. The model is first developed in a generalized

continuous time regime and then it is transformed to its discrete counterpart dictated by the daily basis

data.

Let us denote as f(t) (t: calendar day) the evolution of positive test counts density with f(t)dt being the

number of positive tests in the time period between t and t+dt. The first step is to transform the

function f(t) to the function F(t) which denotes the total number of reported infected people at time t. It

is essential to stress here that F(t) includes people that at time t either had already a positive test or

they are in the first days after infection, so do not have symptoms, but still they shed virus and will be

tested positive later after their onset of symptoms. The effect of infected but unreported people, such

as asymptomatic ones and those not tested because of mild symptoms, is dealt with later.

In order to proceed let us consider the course of the disease of an infected person: Infection starts

(disease incidence) at day $\tau=1$, detection occurs (specimen collection) at day $\tau=\tau_d$ and end of viral

shedding occurs at day τ_e. The number of days for detection and end of shedding are in general not the

same among cases but they show a dispersion which can be described by the respective probability

density functions $P_d(\tau_d)$ and $P_e(\tau_e)$. It is understood that both functions take non-zero values only in

restricted domains of their arguments. The limits of these domains are denoted as τ_{d1}, τ_{d2} and τ_{e1}, τ_{e2},

respectively. The functions P_d and P_e satisfy the conditions:

$$\int_{\tau_{d1}}^{\tau_{d2}} P_d(\tau_d)d\tau_d = 1 \qquad (1a)$$

$$\int_{\tau_{e1}}^{\tau_{e2}} P_e(\tau_e)d\tau_e = 1 \qquad (1b)$$

$$\int_{\tau_{d1}}^{\tau_{d2}} \tau_d P_d(\tau_d)d\tau_d = \tau_{d,av} \qquad (1c)$$

$$\int_{\tau_{e1}}^{\tau_{e2}} \tau_e P_e(\tau_e)d\tau_e = \tau_{e,av} \qquad (1d)$$

where the subscript "av" denotes the average value of the corresponding variable. Taking into account

the above definitions, it can be shown that the functions F(t) and f(t) can be related to each other as:

$$F(t) = \int\limits_{\tau_{d1}}^{\tau_{d2}} \int\limits_{\tau_{e1}}^{\tau_{e2}} P_d(\tau_d) P_e(\tau_e) \int\limits_{t+\tau_d-\tau_e}^{t+\tau_d} f(x) dx d\tau_e d\tau_d \qquad (2)$$

This relation is actually the mathematical expression of the statement that infected persons at time t

include those already detected and those to be detected in the next days. In principle, a multiplication

of this number by the average shedding rate per person would give the required global shedding rate

evolution function R(t). Such averaging over all infected individuals would have been sufficient for data

reduction purposes only if the shedding rate of individuals during their disease cycle had a constant

value. However, as already explained, this is not true since there is a strong variation in virus

concentration in stool not only among infected individuals but also across the days of their disease cycle.

Let us denote this variable function of the daily shedding rate per person as $S(\tau)$, (τ: day of the disease

onset). It has been shown that not only S is not a constant, but on the contrary, it is a strong function of

its argument τ. In order to incorporate the effect of function $S(\tau)$ to the global shedding rate R(t),

knowledge of the distribution of disease days among the infected population is required. The

corresponding function is denoted as $F^*(\tau,t)$ and represents the density function of the distribution of

disease days, τ, at time t. It is fruitful to decompose the function $F^*(\tau,t)$ into the product of the total

number of infected people and the probability density function of the days of infection as

$F^*(\tau,t)=F(t)g(t,\tau)$ where $g(t,\tau)$ is a probability density function with respect to τ and satisfies the

condition:

$$\int\limits_{0}^{\tau_{e2}} g(\tau,t) d\tau = 1 \qquad (3)$$

251   It can be shown that the function g can be derived as

252
$$g(\tau,t) = \frac{1}{F(t)} \int\limits_{\tau_{d1}}^{\tau_{d2}} \int\limits_{\tau_{e1}}^{\tau_{e2}} P_d(\tau_d) P_e(\tau_e) f(t+\tau_d-\tau) U(\tau_e-\tau) d\tau_e d\tau_d \qquad (4)$$

where U is a step function taking the values 0 for negative and 1 for positive argument. The domain of

definition of the above expression is from $t=\tau_{e2}-\tau_{d1}$ to $t=T-\tau_{d2}$ in the case of $f(t)$ known from t=1 to t=T.

Parametrization of the function S(τ)

In order to proceed, one should know the function of virus shedding rate in stool per person and per day

of the disease. A first question is how the onset of infection (i.e., time τ=1) is defined. There are several

possibilities for this definition but in the present context the most relevant one is that τ=1 day is the first

day of non-zero shedding of virus in the stool of a person.

There are only a few available clinical studies in literature (e.g. Wölfel et al.(Wölfel et al., 2020), Huang

et al.(Huang et al., 2020); Tan et al.(Tan et al., 2020)) referring to the kinetics of SARS-CoV-2 shedding

rate in stool during the course of disease (from τ=1 to τ_e). To our best knowledge, from those studies

only the work of Wölfel et al.(Wölfel et al., 2020) reports actual viral concentrations whereas all other

studies refer to Ct values from molecular analysis. It must be mentioned that these studies present data

exclusively from hospitalized patients, thus from people presenting moderate to severe symptoms.

Apparently, it is not easy to collect stool samples from infected persons prior the onset of symptoms,

and as a result, asymptomatic or mild cases are not registered in clinical studies. A first common

observation in these studies is that virus concentration in stool among infected persons vary by several

orders of magnitude. Therefore, here an average shedding function per person will be employed and the

significance of variations among individuals will be discussed on statistical terms. Interestingly, many

past studies assumed in addition a uniform with respect to time shedding rate despite the orders of

magnitude variability across the course of the disease (Ahmed et al., 2020; Gonzalez et al., 2020;

Medema et al., 2020a; Saththasivam et al., 2021). This assumption simplifies the algebra of the problem

enormously but it is incorrect.

A thorough fitting procedure of the virus titers data in stool of Wölfel et al.(Wölfel et al., 2020), led to a

peculiar two-steps model(Miura et al., 2021): a first step with zero viral shedding, where virus load

simply accumulates in infected hosts up to a maximum concentration reached at the day of symptoms

onset, and a subsequent second step characterized by viral shedding at an exponentially decreasing

concentration over the days of the disease. This fitting yields a Gamma function which degenerates to

an exponential one. Evidently, it is quite arbitrary to assume that there is an initial viral accumulation

period up to a maximum concentration without any shedding at all.

Here, an even more general parameterization is introduced. The average over the infected persons

function is represented as a product of the following factors: (1) the average stool amount produced by

a person per day, A (g_{stool}/day), (2) the maximum in time (averaged over infected individuals) virus

concentration in stool, B (g_{virus}/g_{stool}) and (3) a time distribution function s(τ) of this concentration. This

distribution is assumed to consist of an exponential increase from a minimum initial value up to a

maximum value being followed by an exponential decrease down to a minimum final value. These

minimum initial and final values of the distribution are assumed to be 1% of the maximum value. As a

result, the whole distribution spans shedding rates over two orders of magnitude. This assumption may

be changed to any other option, e.g., to 1‰ (three orders of magnitude span), but the approach

remains the same. However, a two orders of magnitude span represents adequately most of the

shedding of infected individuals during a typical course of the disease, so it is adopted herein. A new

parameter is introduced, τ_a, which denotes the value of τ days at which the maximum virus

concentration in stool, i.e. maximum shedding rate, occurs. Summarizing, the above the function

$S(\tau;\tau_a,\tau_e)$ is given as:

$S(\tau;\tau_a, t_e)=ABs(\tau;\tau_a,\tau_e)$ (5)

where $s(\tau;\tau_a,\tau_e)$ is

$s=0.01\exp(4.6\tau/\tau_a)$ $\tau<\tau_a$ (6α)

$s=\exp(4.6(\tau-\tau_a)/(\tau_e-\tau_a))$ $\tau>\tau_a$ (6b)

The shape of the function $s(\tau)$ for several pairs of (τ_a,τ_e) - (6,32), (12,32), (9,25) - is shown in **Figure 1**. The

particular parametrization of the function S proposed in the present work is more realistic than the

Miura et al. (Miura et al., 2021) model, as it incorporates viral shedding even in the early days of the

disease and before the peak maximum (peak) viral concentration in stool is reached. Such early

shedding in stool is in line with clinical studies reporting patients with shedding in oropharyngeal swabs

(He et al., 2020) and gastrointestinal symptoms (Siegel et al., 2020) a few days before the appearance of

respiratory symptoms. Nevertheless, one should keep in mind that respiratory shedding is only a proxy

for fecal shedding. In addition, the proposed function S is very attractive because there is one to one

correspondence between the parameters and major features of the function. The height (amplitude) of

S is determined by the product AB, its time length is determined by the parameter τ_e and its skewness

by the parameter τ_a. In particular, the closer τ_a is to $\tau_e/2$ the smaller the skewness is. It must be stressed

that the proposed parametrization of S does not require explicit information on the duration of the

incubation period after infection nor on the day of symptoms onset. What actually matters in the

context of the present analysis is the day of the maximum shedding rate as this is described by the

parameter τ_a. Some clinical studies indicated that this maximum might occurs at the end of the

incubation period which is taken also as the day of symptom onset (Huang et al., 2020; Miura et al.,

2021; Tan et al., 2020). However, in the present formulation this day is flexible and can be anytime from

the moment of infection to the end of the shedding period (i.e., the active days of the disease).

Figure 1. Time distribution of normalized shedding rate per person in stool at different shedding

durations and shedding peak days.

The final expression for the global shedding rate is taken by integrating the individual's shedding rate

over the days of the disease.

$$R(t) = \int_{0}^{\tau_{e2}} \int_{\tau_{d1}}^{\tau_{d2}} \int_{\tau_{e1}}^{\tau_{e2}} P_d(\tau_d) P_e(\tau_e) f(t+\tau_d-\tau) S(\tau;\tau_a,\tau_e) U(\tau_e-\tau) d\tau_e d\tau_d d\tau \qquad (7)$$

The development up to now is rather complex and includes several unknown probability density

functions. In the absence of any information about them, it is convenient to consider them as Dirac delta

functions. In this way, we assign to each distribution its average value. This approach not only simplifies

considerably the mathematical problem but it also allows -through sensitivity analysis- to assess bounds

on the effect of using a different distribution than the Dirac delta function. This is based on the principle

that any secondary feature of a distribution has much smaller effect on the result than its average value.

By considering the relations $P_d(\tau_d)=\delta(\tau_d-\tau_{d,av})$ and $P_e(\tau_e)=\delta(\tau_e-\tau_{e,av})$ (where δ denotes the Dirac delta

function) and substituting them in equations (2), (4), (7) leads after some algebra to (the subscript "av"

is dropped in the following for clarity):

$$F(t) = \int_{t+\tau_d-\tau_e}^{t+\tau_d} f(x)dxd\tau_e d\tau_d \qquad (8)$$

$$g(\tau,t) = \frac{f(t+\tau_d-\tau)U(\tau_e-\tau)}{F(t)} \qquad (9)$$

$$R(t) = F(t)\int_0^{\tau_e} g(\tau,t)S(\tau;\tau_a,\tau_e)d\tau \qquad (10)$$

The above set of equations is discretized to be compatible with the present data. These data refer to

daily values of virus shedding rate so a finite volume discretization is followed here. Let us denote as f_i

the number of positive medical tests at day i (measurement period i=1 to N), F_i is the total number of

infected people (that is daily cases) at day i (with the specimen of their positive test collected at day i), R_i

the shedding rate at day i and $g_{i,j}$ the probability of being at the j-th day of the disease at the calendar

day i. The governing equations take the form:

$$F_i = \sum_{j=i-m}^{i+n} f_j \qquad (11)$$

$$g_{i,j} = \frac{f_{j-m+i}}{F_i} \qquad (12)$$

$$R_i = \sum_{j=i-m}^{i+n} g_{i,j} S_j \qquad (13)$$

for i>m and i<N-n

where S_j is the shedding rate per person at the j-th day of the disease (m=τ_e-τ_d, n=τ_d)

RESULTS

Figure 2 displays medical surveillance data reported for the city of Thessaloniki (~ 700,000 inhabitants)

(National Public Health Organization, 2020). More specifically, it presents the daily number of infected

people versus the date of their specimen collection for medical testing. The date of specimen collection

is back-dated by 1 to 4 days (median of 3 days) from the date of reporting by the Hellenic National

Public Health Organization. The difference between the two characteristic dates is small but the

epidemiological data adjusted to the date of specimen collection are more appropriate to compare with

wastewater measurements because they are better associated with the date of infection and the date

of symptoms onset. Nevertheless, if early warning by wastewater measurements is the stake then

comparisons should be made with medical surveillance data as announced, i.e., based on the date of

reporting.

Figure 2. Medical surveillance data reported for the city of Thessaloniki from September 1st, 2020 to

January 6th, 2021 and corresponding Bayesian model fit curve.

The presented data cover the period from September, 1st 2020 to January 6th, 2021. A fitting curve is

fitted over the raw data to smooth out the noise. Most of the parametric study that follows is

performed with respect to this curve, to skip the noise. Following the loose atmosphere of Summer

2020, in September and October there was no strict quarantine in the city and only a modest rule for

social distancing was active along with a rule for limited number of people in confined places, like stores

and restaurants. In September and the first 10 days of October the daily reported cases were always

well below 20 and often even below 10. At around mid-October 2020 the number of infected cases

started to escalate exponentially. In just two weeks, the number of cases increased about ten-fold in just

two weeks. This dramatic rise was consistent with the about 2.7 days doubling time of the epidemic

reported for European countries in the absence of control measures (SET-C Steering Committee, 2020).

A strict lock-down was imposed on the city on November 2^{nd} meant to suppress the outburst of the

disease. Between Nov 3^{rd} and Nov 17^{th} the number of infected cases fluctuated approximately from 800

to 300 cases between weekdays and weekends. Similar fluctuations were noticed also in other studies

(Peccia et al., 2020). On December 12^{th} some of the strict measures were released, e.g., retail stores

opened to serve people but only at their entrance door and only through pre-scheduled appointments.

As a result, after mid-November the number of infected cases gradually went down until it became

pretty stable at below 100 daily cases in the first week of 2021.

Figure 3 presents the experimentally determined relative shedding rate of viral RNA copies,

$r_1(t)=R_{exp}/R_{expo}$, in sewage from October 5^{th}, 2020 to January 6^{th}, 2021 for Thessaloniki. October 5^{th} is the

last day after summer that the measured viral RNA copies in wastewater fluctuated around the

experimental limit of quantification of the employed technique (~10 viral RNA copies/mL). R_{exp}

represents the daily value of shedding rate, whereas R_{expo} is a reference shedding rate. Presenting

experimental shedding data in the form of a ratio reduces the effect of determination uncertainty. R_{expo}

has been defined as the average R_{exp} value across the days of the first week of October which was a

period with calm epidemiological conditions in the city (less than 10 daily reported cases).

Figure 3. Experimental relative shedding rate in sewage expressed as the ratio of the measured daily

shedding rate (R_{exp}) over the reference shedding rate (R_{expo}) from October 5[th], 2020 to January 6[th], 2021.

There are certain similarities and differences between the medical surveillance data in **Figure 2** and the

wastewater data in **Figure 3**. They both show an initial abrupt ascend followed by a later gradual

decline. They both show a peak in the first half of November 2020. Yet, the peak in wastewater data is

sharp at around November 13[th] whereas the peak in medical surveillance, despite the intense noise,

looks like a plateau between November 3[rd] and 12[th]. The initial increasing slope in October's data in

these Figures is a bit steeper for wastewater data implying that wastewater surveillance may act as an

early warning indicator for medical surveillance. More about this below.

Next, the time series of the reported daily infected cases in the city of Thessaloniki is analyzed using the

procedure in the previous section to determine the functions F, g, R. It is reminded that these three

functions correspond respectively to (i) the total number of infected people with positive test at a

calendar day, t, (those already registered as positive to day t but also those that will be tested in the

next days and their test result will be also registered to day t), (ii) the distribution of the number of

infected people with respect to the days of the disease, τ, at a calendar day, t, and (iii) the total shedding

rate with respect to the calendar day, t. It is important to note that since the temporal discretization

quantity is one day, the continuous and the discrete forms of the above functions are arithmetically

similar. Thus, the presented results can be read both as continuous functions (per day) or as discrete

values (i.e., histograms).   The essence of the proposed approach is that it accounts for the time

dependence of the shedding rate during the disease. Yet, this is important only when the distribution of

the daily shedding rates is not uniform among the population, but it evolves along the days of the

disease.

Before a parametric analysis is performed it is useful to identify realistic range of values of the model

parameters in order to examine their influence. It is reminded that the three model parameters are (i)

the total number of shedding days counted from the infection day, τ_e, (ii) the day of the maximum

shedding rate, τ_a, and (iii) the day of detection (specimen collection), τ_d. The analysis of the data of

Wölfel et al.(Wölfel et al., 2020) by Miura et al(Miura et al., 2021) indicated an average shedding period

of 26 days after the onset of symptoms. This is a bit longer than the 17 and 18 days, reported by Huang

et al.(Huang et al., 2020) and Tan et al.(Tan et al., 2020), respectively. But both the latter studies

reported a high variance in their average shedding period plus they mentioned that in some individuals

shedding exceeded five weeks. Apart from this, Tan et al.(Tan et al., 2020), indicated a 6 day median

incubation period from the day of infection until the day of symptoms onset. If one considers that the

day of symptoms onset is a good candidate for the average day of maximum shedding rate then τ_a= 6.

For Greek patients it is realistic to assume that specimen collection for medical testing takes place on

the average about 3 days after the onset of symptoms. This is in line with information in literature that

the median time period from symptom onset to hospital admission is 3 days(Huang et al., 2020).

Therefore, a realistic average day of detection is τ_d=8. The above information combined implies a total

number of shedding days τ_e=32 (6+26) and this is the value adopted herein. Summarizing, the base case

parameter values around which parametric analysis is performed are τ_a= 6, τ_d=8 and τ_e=32.

Figure 4 shows the computed values of g(τ,t) for selected values of calendar days, namely, September

25[th], October 10[th], October 23[th], December 9[th] and December 29[th], 2020. The small oscillation appeared

is due to the oscillations in the curve f(t). However, as f(t) – the positive tests count density- increases

new patients are added to the number of infected people. The addition occurs at a higher rate than the

rate of withdrawal of cured people. This leads to the accumulation of infected people at early shedding

days, τ, so the function g(τ) acquires the characteristic, decreasing with τ, profiles shown for October

10[th] and October 23[th]. In the time regime of the maximum f(t), e.g. curve for December 9[th], the function

g(τ) tends fast towards uniformity, like at the early shedding days. Finally, during the decreasing period

of f(t), e.g. curve for December 29[th], the withdrawal of cured people rate overwhelms the entry of new

patients so g(t) is an almost linearly increasing function of τ. The above description manifests that the

function g(τ) varies a lot with calendar time t and it is far from uniform along the shedding days so the

present analysis is necessary for the estimation of shedding rate across calendar days. It is clear that at a

particular calendar day, infected people are at a different stage of the disease, and thus, at a different

stage of viral shedding.

Figure 4. The probability density distribution of the number of infected people with respect to the days

of the disease.

The early warning capacity of wastewater surveillance with respect to medical surveillance is estimated

by comparing in **Figure 5** the dynamics of functions f(t): daily reported infected people, F(t): total

reported infected people, and R(t): experimentally determined viral shedding rate in sewage. More

specifically, **Figure 5a** shows this comparison for the base case τ_a= 6, τ_d=8, τ_e=32; **Figure 5b** for τ_a= 2,

τ_d=8, τ_e=32; **Figure 5c** for τ_a= 12, τ_d=8, τ_e=32 (all values in days). Curves in **Figure 5** stop at December 29[th],

2020, and not at January 6[th], 2021. This is because calculations of F(t) -and accordingly of R(t)- can be

performed only up to December 29[th], 2020, as this function requires for every daily value available data

for the subsequent τ_d=8 days. To avoid experimental noise in the parametric analysis, calculations in

Figure 5 are based on the Bayesian fitted curve in **Figure 2**. For the comparison it is imperative to

express these three functions in a similar scale. This is done by transforming them in their probability

density function counterparts. For the present case, this is done simply by dividing each function with

the sum of its values for the period of interest. The function F(t) always resides after f(t) which is typical

for a cumulative type of function. The temporal distance apart between these two functions depends on

the values of the parameters τ_d and τ_e. This distance increases as τ_d decreases and τ_e increases. The

dynamics of the function R(t) depend also on τ_a. In case of $\tau_a<\tau_d$ then R(t) precedes f(t). This condition is

extremely important as it demonstrates that for wastewater surveillance to precede medical

surveillance the day of the peak of viral shedding rate in stool during the disease days must lie before

the day of specimen collection for medical testing. In all other cases, the curve R(t) resides between f(t)

and F(t), with the latter being the approximate long time bound for the dynamics of R(t). The previous

statement can be confirmed by observing the three aforementioned functions for the three set of

parameters in **Figure 5**. Depending on the calendar day, function R(t) precedes f(t) roughly from 0 to 4

days (**Figure 5b**). For the base case in **Figure 5a** the difference between R(t) and f(t) appears marginal.

Yet, one must recall that in **Figure 2** the ascending part of the Bayesian curve precedes raw data, so in

reality even for the base case sewage data lie earlier than medical data. In literature there is

experimental evidence of identifying SARS-CoV-2 in wastewater earlier than medical reporting by

several days, e.g., 2 days in Ottawa Canada(D'Aoust et al., 2021), 2-4 days in Montana USA(Nemudryi et

al., 2020). In summary, the viral shedding rate evolution curve measured in wastewater lies from a few

days before the curve of the number of daily infected people up to the curve of the total number of

infected people. The exact position of the curves depends on the relation between the day of maximum

shedding rate in stool and the day of specimen collection for medical testing.

Figure 5. Comparison of the relative viral shedding rate in sewage with the number of the daily and total reported infected people per surveillance day. In all cases shedding duration in stool is set as 32days, laboratory test is assumed at DAY 8, whereas the day of shedding peak in stool is at: DAY 6 (a), DAY 2 (b) and DAY 12 (c).

For appraising in real life the early diagnostic capacity of wastewater surveillance it is important to compare wastewater data with medical surveillance data based on the date of reporting (and not on the date of specimen collection). Peccia et al.(Peccia et al., 2020) found in New Haven (USA) that although wastewater data was ahead by only 0-2 days of positive test results by the date of specimen collection, it was 6-8 days ahead of positive test results by the reporting date. In Thessaloniki (National Public Health Organization, 2020) the date of reporting follows the date of specimen collection by about 2 days on the average and so even the base case parameters in **Figure 5a** permit an early diagnosis capacity of wastewater surveillance of around 2 days.

A critical issue in the disease dynamics is the effect of the number of unreported infected people (unreported cases). Let us denote as U the ratio of unreported to reported infected people. If U is constant along the days of the shedding period, the analysis is exactly the same with that shown above for the reported infected people, F, since the probability density functions do not change. However, if the number of unreported people varies along the shedding period, then this may yield a time lag between wastewater and medical data. **Figure 6** compares the relative medical surveillance data of the

smoothed (Bayesian fit) daily reported cases, $f(t)/f_o$, versus the theoretically estimated relative shedding

rate in wastewater, $r_2(t)=R(t)/R_o$. Normalization parameters, f_o and R_o are average values of the

respective parameters over the same reference first week of October. Comparisons are for the base

case $\tau_a= 6$, $\tau_d=8$, $\tau_e=32$. The ratio U is assumed to vary proportionally with the number of reported

infected people. The parameter U_{max} is used to parameterize U designating the maximum value of U for

the case considered. The parameter U_{max} takes the value 0.5, 2 and 4 in **Figure 6**. Apparently, for

$U_{max}=0.5$ (unreported cases becomes at most 50% of reported cases) wastewater and medical data

coincide at the ascent of the curves but at the descent wastewater lags behind. This changes for $U_{max}=2$

(unreported cases becomes at most 200% of reported cases) and for $U_{max}=4$ (unreported cases becomes

at most 400% of reported cases) where wastewater data clearly go before medical data at the ascent of

the curve. Summing up, a rising number of unreported cases as the shedding rate increases leads to

earlier wastewater signal than medical surveillance. Furthermore, the higher the maximum number of

unreported cases the more in advance the wastewater data from the medical data.

Figure 6. Comparison of the theoretical relative shedding rate in sewage with the relative medical

surveillance data of the smoothed (Bayesian fit) daily reported cases for different scenarios of

unreported daily cases. Unreported daily cases vary proportionally with the number of reported daily

cases starting from 0 and reaching a maximum value of 0, 50, 200, 400% of reported cases at the date of

maximum shedding rate.

Next, the experimental relative shedding rate, $r_1(t)=R_{exp}(t)/R_{expo}$, shown in **Figure 3** is compared with the

estimated total number of infected people, including both reported and unreported ones, **Figure 7.** Red

circles denote increasing shedding rates whereas green squares decreasing shedding rates, respectively.

It is apparent that from low to moderate relative shedding rates ($R_{exp}(t)/R_{expo}$ <100) and total number of

infected people (F<15000) there is a roughly linear relationship between the two quantities. This

changes dramatically for higher values of either quantities. Moreover, during the increasing phase of the

relative shedding rate, i.e., outbreak of pandemic wave, the same number of infected people sheds a

higher viral load than during the decreasing phase. This is so because at the outbreak phase of the

pandemic most infected people are at the early phase of their disease when shedding rate is higher, as

shown in **Figure 1**. In line with this, when the relative shedding rate starts to fall right after its peak, the

total number of infected people continues to rise to about 10% more cases. This happens because the

turning point (peak) of the curve designates the moment when the daily new infected cases begin to

reduce, **Figure 2**, so the majority of the total number of infected people are at later phase of their

disease when shedding rate is lower.

Figure 7. Experimental relative shedding rate in sewage, $R_{exp}(t)/R_{expo}$, versus the estimated total number

of infected people (reported and unreported), F. The latter is estimated assuming that, on the average,

the shedding duration in stool is 32 days and medical tests are taken at DAY 8 during the disease. Red

circles denote increasing shedding rates whereas green squares denote decreasing shedding rates.

Let us now try to correlate the theoretically derived function R(t) with the actual one found from

experimental wastewater analysis $R_{exp}(t)$. The latter was shown as a relative shedding rate,

$r_1(t)=R_{exp}(t)/R_{expo}$, in **Figs 3** and **7**. Contrary to what was done in **Figure 6**, now the theoretically estimated

relative shedding rate $r_2(t)=R(t)/R_o$ is computed from the noisy medical surveillance raw data in **Figure 2**

for the base case parameter values $\tau_a= 6$, $\tau_d= 8$ and $\tau_e=32$. As a result, $r_2(t)$ data are also noisy. Therefore,

$\beta(t)=r_1(t)/r_2(t)$ is a ratio reflecting the total number of reported and unreported infected people, F(1+U),

over the total number of reported infected people, F, at every calendar day. In other words, β=1+U,

that is, β-1 stands for U, the ratio of the total unreported over the total reported cases. Thus, the

minimum value of β is 1 and corresponds to zero unreported cases. As before, the values of R_{expo} and R_o

both refer to the same reference first week of October. Selection of the reference week at an

epidemiologically calm period allows assuming that $R_{expo}/R_o≈1$ so the normalization in β can be ignored.

The evolution of β with the calendar days appears in **Figure 8.** Considering the intense day-by-day

scatter, the ratio β takes values roughly between 1 and 5, following a qualitatively similar trend with the

wastewater shedding data in **Figure 3**. Interestingly, these results imply that there are essentially no

unreported cases at periods of low shedding but unreported cases rise up to four times the reported

ones (U/F≈4) when shedding is maximum. Analysis of seroprevalence data in Qatar indicated that

diagnosed infections represented about 10% of actual cases(Saththasivam et al., 2021). Another

serological-testing study(Havers et al., 2020) showed that the actual number of infections could be from

6 to 24 times the number of the reported cases. Overall, the presently estimated values of U/F from ~0

to 4 is close to the above published values. To our knowledge, this is the first time in the SARS-CoV-2

pandemic that wastewater data are employed to indicate a proportion between unreported and

reported infected cases. Moreover, the present study shows that the proportion of unreported to

reported cases is not constant during the outbreak of a disease wave but as the number of infected

people (and shedding rate) increases, it attains higher values and its scatter decreases.

Figure 8. Evolution of the ratio of the total number of infected people (reported + unreported) over the

number of reported infected people. The ratio is calculated from unsmoothed raw data in Figures 2 and

3. Red circles denote increasing shedding rates whereas green squares denote decreasing shedding

rates.

Based on the above, an effort is made to correlate β to the total number of the reported infected cases

F. The result is shown in **Figure 9a** where once more red circles denote increasing shedding rates and

green squares denote decreasing shedding rates. Alike in **Figure 7**, β starts decreasing before the

maximum F is reached. In addition, at high F values there is a hysteresis in β between increasing and

decreasing branches of the curves. This might imply that virus spread among unreported people is a bit

higher in the declining phase of the disease either because of the cumulatively higher virus prevalence in

the community or because more people undergo testing during the ascending spreading period of the

disease than in the milder descending period. **Figure 9b** presents β versus the daily reported number of

the infected cases, f. Interestingly there is an almost linear relation between the two parameters up to

about 400 daily reported cases. After that point, spreading of the virus in the community is too high so β

depends on the total number infected people F and not on the daily reported cases.

Figure 9. Variation of the ratio of the total number of infected people (reported + unreported) over the

number of reported infected people with regards to (a) the total number of reported infected people

and (b) the daily number of infected people. The ratio is calculated from unsmoothed raw data in

Figures 2 and 3. Red circles denote increasing shedding rates whereas green squares denote decreasing

shedding rates.

It must be mentioned that β may depend also on other quantities (apart from F and the

ascending/descending mode), like the value of the F slope, but at present there are not enough data to

support this. Having an estimation of the total number of infected people through β, a crude estimation

of the product AB (average maximum virus shedding rate per person) is obtained. The estimated value

of AB is at about $2 \cdot 10^{11}$ day^{-1}. This value is in fair proximity with that estimated by the wastewater

analysis of Wu et al.(Wu et al., 2020). However, more work is needed to allow judging on the

correctness of this value.

CONCLUSIONS

A mathematical model is developed that permits estimation of SARS-CoV-2 shedding rate in wastewater

from the number of daily infected people (daily cases) announced by medical surveillance. The problem

is complex as it requires a cumulative function of the total number of infected people at every calendar

day and a function of the average shedding rate among infected individuals at every day along the

course of the disease. Based on the limited available evidence in literature, a realistic function for the

shedding rate of SARS-CoV-2 in stool of infected individuals is proposed which calls for an exponential

increase of shedding rate from the day of infection to the day of symptoms onset, being followed by an

exponential decay until the end of the disease. Three characteristic times describe this function: the day

of maximum shedding rate, the day of detection (specimen collection) and the end day after the onset

of the disease. Applying this model to the public health surveillance data for the city of Thessaloniki

(~700,000 inhabitants, North Greece) a thorough parametric study is performed. It is shown that for

WBE to afford an early warning capacity, the day of maximum shedding rate must precede the day of

detection by a few days. In particular, at the beginning of an outbreak curve, a 4-day early WBE signal

requires these two characteristic times to be apart by 6 days. For Thessaloniki where on the average

these two characteristic times are apart by just 2 days but, additionally, the day of reporting follows the

date of specimen collection also by about 2 days, the early diagnosis capacity of wastewater surveillance

is around 2 days. Furthermore, comparison of wastewater surveillance and public health data indicates

the existence of a number of unreported infected people. For Thessaloniki in the outbreak of November

2020, this number was negligible at epidemiologically calm days with low shedding rate but went up to

four times the number of reported people when shedding rate reached a climax at mid-November 2020.

Interestingly, the presence of an increasing number of unreported cases with shedding rate enhances

the early warning capacity of WBE. To this end, the present model is an essential tool to investigate the

dynamics of virus spreading based on wastewater measurements. When such knowledge is adequately

acquired then the inverse problem of estimating the number of cases from wastewater data can be

attempted.

REFERENCES

Ahmed, W., Angel, N., Edson, J., Bibby, K., Bivins, A., O'Brien, J.W., Choi, P.M., Kitajima, M., Simpson,

S.L., Li, J., Tscharke, B., Verhagen, R., Smith, W.J.M., Zaugg, J., Dierens, L., Hugenholtz, P., Thomas,

K. v., Mueller, J.F., 2020. First confirmed detection of SARS-CoV-2 in untreated wastewater in

Australia: A proof of concept for the wastewater surveillance of COVID-19 in the community.

Science of the Total Environment 728, 138764. https://doi.org/10.1016/j.scitotenv.2020.138764

CDC, 2021. CDC [WWW Document]. Center for Disease Control and Prevention, CDC. URL

https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/wastewater-surveillance/public-

health-interpretation.html (accessed 3.10.21).

Chaintoutis, S.C., Papadopoulou, E., Melidou, A., Papa, A., Dovas, C.I., 2019. A PCR-based NGS protocol

for whole genome sequencing of West Nile virus lineage 2 directly from biological specimens.

Molecular and Cellular Probes 46, 101412. https://doi.org/10.1016/j.mcp.2019.06.002

Corman, V.M., Landt, O., Kaiser, M., Molenkamp, R., Meijer, A., Chu, D.K.W., Bleicker, T., Brünink, S.,

Schneider, J., Schmidt, M.L., Mulders, D.G.J.C., Haagmans, B.L., van der Veer, B., van den Brink, S.,

Wijsman, L., Goderski, G., Romette, J.L., Ellis, J., Zambon, M., Peiris, M., Goossens, H., Reusken, C.,

Koopmans, M.P.G., Drosten, C., 2020. Detection of 2019 novel coronavirus (2019-nCoV) by real-

time RT-PCR. Eurosurveillance 25. https://doi.org/10.2807/1560-7917.ES.2020.25.3.2000045

D'Aoust, P.M., Mercier, E., Montpetit, D., Jia, J.J., Alexandrov, I., Neault, N., Baig, A.T., Mayne, J., Zhang,

X., Alain, T., Langlois, M.A., Servos, M.R., MacKenzie, M., Figeys, D., MacKenzie, A.E., Graber, T.E.,

Delatolla, R., 2021. Quantitative analysis of SARS-CoV-2 RNA from wastewater solids in

communities with low COVID-19 incidence and prevalence. Water Research 188, 116560.

https://doi.org/10.1016/j.watres.2020.116560

Gonzalez, R., Curtis, K., Bivins, A., Bibby, K., Weir, M.H., Yetka, K., Thompson, H., Keeling, D., Mitchell, J.,

Gonzalez, D., 2020. COVID-19 surveillance in Southeastern Virginia using wastewater-based

epidemiology. Water Research 186, 116296. https://doi.org/10.1016/j.watres.2020.116296

Guan, W., Ni, Z., Hu, Yu, Liang, W., Ou, C., He, J., Liu, L., Shan, H., Lei, C., Hui, D.S.C., Du, B., Li, L., Zeng,

G., Yuen, K.-Y., Chen, R., Tang, C., Wang, T., Chen, P., Xiang, J., Li, S., Wang, Jin-lin, Liang, Z., Peng,

Y., Wei, L., Liu, Y., Hu, Ya-hua, Peng, P., Wang, Jian-ming, Liu, J., Chen, Z., Li, G., Zheng, Z., Qiu, S.,

Luo, J., Ye, C., Zhu, S., Zhong, N., 2020. Clinical Characteristics of Coronavirus Disease 2019 in China.

New England Journal of Medicine 382, 1708–1720. https://doi.org/10.1056/nejmoa2002032

Havers, F.P., Reed, C., Lim, T., Montgomery, J.M., Klena, J.D., Hall, A.J., Fry, A.M., Cannon, D.L., Chiang,

C.F., Gibbons, A., Krapiunaya, I., Morales-Betoulle, M., Roguski, K., Rasheed, M.A.U., Freeman, B.,

Lester, S., Mills, L., Carroll, D.S., Owen, S.M., Johnson, J.A., Semenova, V., Blackmore, C., Blog, D.,

Chai, S.J., Dunn, A., Hand, J., Jain, S., Lindquist, S., Lynfield, R., Pritchard, S., Sokol, T., Sosa, L.,

Turabelidze, G., Watkins, S.M., Wiesman, J., Williams, R.W., Yendell, S., Schiffer, J., Thornburg, N.J.,

2020. Seroprevalence of Antibodies to SARS-CoV-2 in 10 Sites in the United States, March 23-May

12, 2020. JAMA Internal Medicine 180, 1776–1786.

https://doi.org/10.1001/jamainternmed.2020.4130

He, X., Lau, E.H.Y., Wu, P., Deng, X., Wang, J., Hao, X., Lau, Y.C., Wong, J.Y., Guan, Y., Tan, X., Mo, X.,

Chen, Y., Liao, B., Chen, W., Hu, F., Zhang, Q., Zhong, M., Wu, Y., Zhao, L., Zhang, F., Cowling, B.J.,

Li, F., Leung, G.M., 2020. Temporal dynamics in viral shedding and transmissibility of COVID-19.

Nature Medicine 26, 672–675. https://doi.org/10.1038/s41591-020-0869-5

Huang, J., Mao, T., Li, S., Wu, L., Xu, X., Li, H., Xu, C., Su, F., Dai, J., Shi, J., Cai, J., Huang, C., Lin, X., Chen,

D., Lin, X.L., Sun, B., Tang, S., 2020. Long period dynamics of viral load and antibodies for SARS-CoV-

2 infection: An observational cohort study. medRxiv.

https://doi.org/10.1101/2020.04.22.20071258

la Rosa, G., Iaconelli, M., Mancini, P., Bonanno Ferraro, G., Veneri, C., Bonadonna, L., Lucentini, L.,

Suffredini, E., 2020. First detection of SARS-CoV-2 in untreated wastewaters in Italy. Science of the

Total Environment 736. https://doi.org/10.1016/j.scitotenv.2020.139652

Lauer, S.A., Grantz, K.H., Bi, Q., Jones, F.K., Zheng, Q., Meredith, H.R., Azman, A.S., Reich, N.G., Lessler, J.,

2020. The incubation period of coronavirus disease 2019 (CoVID-19) from publicly reported

confirmed cases: Estimation and application. Annals of Internal Medicine 172, 577–582.

https://doi.org/10.7326/M20-0504

Li, Q., Guan, X., Wu, P., Wang, X., Zhou, L., Tong, Y., Ren, R., Leung, K.S.M., Lau, E.H.Y., Wong, J.Y., Xing,

X., Xiang, N., Wu, Y., Li, C., Chen, Q., Li, D., Liu, T., Zhao, J., Liu, M., Tu, W., Chen, C., Jin, L., Yang, R.,

Wang, Q., Zhou, S., Wang, R., Liu, H., Luo, Y., Liu, Y., Shao, G., Li, H., Tao, Z., Yang, Y., Deng, Z., Liu,

B., Ma, Z., Zhang, Y., Shi, G., Lam, T.T.Y., Wu, J.T., Gao, G.F., Cowling, B.J., Yang, B., Leung, G.M.,

Feng, Z., 2020. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected

Pneumonia. New England Journal of Medicine 382, 1199–1207.

https://doi.org/10.1056/nejmoa2001316

Medema, G., Been, F., Heijnen, L., Petterson, S., 2020a. Implementation of environmental surveillance

for SARS-CoV-2 virus to support public health decisions: Opportunities and challenges. Current

Opinion in Environmental Science and Health 17, 49–71.

https://doi.org/10.1016/j.coesh.2020.09.006

Medema, G., Heijnen, L., Elsinga, G., Italiaander, R., Brouwer, A., 2020b. Presence of SARS-Coronavirus-2

RNA in Sewage and Correlation with Reported COVID-19 Prevalence in the Early Stage of the

Epidemic in The Netherlands. Environmental Science & Technology Letters 7, 511–516.

https://doi.org/10.1021/acs.estlett.0c00357

Miura, F., Kitajima, M., Omori, R., 2021. Duration of SARS-CoV-2 viral shedding in faeces as a parameter

for wastewater-based epidemiology: Re-analysis of patient data using a shedding dynamics model.

Science of the Total Environment 769, 144549. https://doi.org/10.1016/j.scitotenv.2020.144549

National Public Health Organization, 2020. Daily reports COVID-19 2020. URL

https://eody.gov.gr/epidimiologika-statistika-dedomena/ektheseis-covid-19/imerisies-ektheseis-

covid-19-2020/ (accessed 5.14.21).

Nemudryi, A., Nemudraia, A., Wiegand, T., Surya, K., Buyukyoruk, M., Cicha, C., Vanderwood, K.K.,

Wilkinson, R., Wiedenheft, B., 2020. Temporal Detection and Phylogenetic Assessment of SARS-

CoV-2 in Municipal Wastewater. Cell Reports Medicine 1, 100098.

https://doi.org/10.1016/j.xcrm.2020.100098

Peccia, J., Zulli, A., Brackney, D.E., Grubaugh, N.D., Kaplan, E.H., Casanovas-Massana, A., Ko, A.I., Malik,

A.A., Wang, D., Wang, M., Warren, J.L., Weinberger, D.M., Arnold, W., Omer, S.B., 2020.

Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. Nature

Biotechnology 38, 1164–1167. https://doi.org/10.1038/s41587-020-0684-z

Petala, M., Dafou, D., Kostoglou, M., Karapantsios, T., Kanata, E., Chatziefstathiou, A., Sakaveli, F.,

Kotoulas, K., Arsenakis, M., Roilides, E., Sklaviadis, T., Metallidis, S., Papa, A., Stylianidis, E.,

Papadopoulos, A., Papaioannou, N., 2021. A physicochemical model for rationalizing SARS-CoV-2

concentration in sewage. Case study: The city of Thessaloniki in Greece. Science of the Total

Environment 755, 142855. https://doi.org/10.1016/j.scitotenv.2020.142855

Saththasivam, J., El-Malah, S.S., Gomez, T.A., Jabbar, K.A., Remanan, R., Krishnankutty, A.K., Ogunbiyi,

O., Rasool, K., Ashhab, S., Rashkeev, S., Bensaad, M., Ahmed, A.A., Mohamoud, Y.A., Malek, J.A.,

Abu Raddad, L.J., Jeremijenko, A., Abu Halaweh, H.A., Lawler, J., Mahmoud, K.A., 2021. COVID-19

(SARS-CoV-2) outbreak monitoring using wastewater-based epidemiology in Qatar. Science of the

Total Environment 774, 145608. https://doi.org/10.1016/j.scitotenv.2021.145608

Sellaoui, L., Badawi, M., Monari, A., Tatarchuk, T., Jemli, S., 2020. Since January 2020 Elsevier has

created a COVID-19 resource centre with free information in English and Mandarin on the novel

coronavirus COVID- 19 . The COVID-19 resource centre is hosted on Elsevier Connect , the company

' s public news and information .

SET-C Steering Committee, 2020. Reproduction number (R) and growth rate (r) of the COVID-19

epidemic in the UK 1–86.

Siegel, A., Chang, P.J., Jarou, Z.J., Paushter, D.M., Harmath, C.B., Arevalo, J. ben, Dachman, A., 2020.

Lung base findings of coronavirus disease (COVID-19) on abdominal CT in patients with

predominant gastrointestinal symptoms. American Journal of Roentgenology 215, 607–609.

https://doi.org/10.2214/AJR.20.23232

Tan, W., Lu, Y., Zhang, J., Wang, J., Dan, Y., Tan, Z., He, X., Qian, C., Sun, Q., Hu, Q., Liu, H., Ye, S., Xiang,

X., Zhou, Y., Zhang, W., Guo, Y., Wang, X.H., He, W., Wan, X., Sun, F., Wei, Q., Chen, C., Pan, G., Xia,

J., Mao, Q., Chen, Y., Deng, G., 2020. Viral kinetics and antibody responses in patients with COVID-

19. medRxiv. https://doi.org/10.1101/2020.03.24.20042382

Wölfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Müller, M.A., Niemeyer, D., Jones,

T.C., Vollmar, P., Rothe, C., Hoelscher, M., Bleicker, T., Brünink, S., Schneider, J., Ehmann, R.,

Zwirglmaier, K., Drosten, C., Wendtner, C., 2020. Virological assessment of hospitalized patients

with COVID-2019. Nature 581, 465. https://doi.org/10.1038/s41586-020-2196-x

Wu, F., Zhang, J., Xiao, A., Gu, X., Lee, W.L., Armas, F., Kauffman, K., Hanage, W., Matus, M., Ghaeli, N.,

Endo, N., Duvallet, C., Poyet, M., Moniz, K., Washburne, A.D., Erickson, T.B., Chai, P.R., Thompson,

J., Alm, E.J., 2020. SARS-CoV-2 Titers in Wastewater Are Higher than Expected from Clinically

Confirmed Cases. mSystems 5. https://doi.org/10.1128/mSystems.00614-20

Wurtzer[1], S., Marechal, V., Mouchel[4], J.M., Maday, Y., Teyssou[6], R., Richard[1], E., Almayrac[7], J.L., Moulin[1],

L., n.d. Evaluation of lockdown effect on SARS-CoV-2 dynamics through viral genome quantification

in waste water 1. https://doi.org/10.2807/1560-7917

Ye, Y., 2018. The Detection and Fate of Enveloped Viruses in Water Environments 149.

Zheng, S., Fan, J., Yu, F., Feng, B., Lou, B., Zou, Q., Xie, G., Lin, S., Wang, R., Yang, X., Chen, W., Wang, Q.,

Zhang, D., Liu, Y., Gong, R., Ma, Z., Lu, S., Xiao, Y., Gu, Y., Zhang, J., Yao, H., Xu, K., Lu, X., Wei, G.,

Zhou, J., Fang, Q., Cai, H., Qiu, Y., Sheng, J., Chen, Y., Liang, T., 2020. Viral load dynamics and

disease severity in patients infected with SARS-CoV-2 in Zhejiang province, China, January-March

2020: Retrospective cohort study. The BMJ 369. https://doi.org/10.1136/bmj.m1443