Transmission dynamics of COVID-19 in Ghana and the impact of public health interventions

Sylvia K. Ofori,[1] Jessica S. Schwind,[1] Kelly L. Sullivan,[1] Benjamin J. Cowling,[2] Gerardo

Chowell,[3] Isaac Chun-Hai Fung[1] [*]

Affiliations:

1   Department of Biostatistics, Epidemiology and Environmental Health Sciences, Jiann-

Ping Hsu College of Public Health, Georgia Southern University.

2   WHO Collaborating Centre for Infectious Disease Epidemiology and Control, School of

Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong.

3   Department of Population Health Sciences, School of Public Health, Georgia State University.

*Correspondence should be addressed to Isaac Chun-Hai Fung, Ph.D., P.O. Box 7989,

Department of Biostatistics, Epidemiology and Environmental Health Sciences, Jiann-Ping Hsu

College of Public Health, Georgia Southern University, Statesboro, GA 30458-7989, USA.

Email: cfung@georgiasouthern.edu. Tel: +1 912 4785079.

Keywords: Ghana, time-varying reproduction number, COVID-19, nonpharmaceutical
interventions

Manuscript type: Short Report

Number of tables: 1 + 5 supplementary tables

Number of figures: 2 + 6 supplementary figures

Supplementary materials attached

Running head: Transmission dynamics of COVID-19 in Ghana

Word count, 1534.

Abstract (149 words)

The study characterized COVID-19 transmission in Ghana in 2020-21 by estimating the time-

varying reproduction number (R_t) and exploring its association with various public health

interventions at the national and regional levels. Ghana experienced four pandemic waves with

epidemic peaks in July 2020, and January, August and December, 2021. The epidemic peak was

the highest nationwide in December 2021 with $R_t \geq 2$. Throughout 2020-21, per-capita cumulative

case count by region increased with population size. Mobility data suggested negative correlation

between R_t and staying home in the first 90 days of the pandemic. The relaxation of movement

restrictions and religious gatherings were not associated with increased R_t in the regions with lower

case burdens. R_t decreased from above 1 when schools reopened in January 2021 to below 1 after

vaccination rollout in March 2021. Findings indicated most public health interventions were

associated with R_t reduction at the national and regional levels.

Main text

As of January 15, 2022, 153,514 cases of COVID-19 were confirmed in Ghana, with 1,343 deaths

and 9,020 active cases.[1] Among African countries, Ghana has controlled COVID-19 transmission

with a substantial package of public health and social measures (Oxford stringency index of 62.04

as of May 7, 2020), [2,3] including a 14-day mandatory quarantine for all persons who entered the

country, school and church closures, a lockdown of major cities, and internal movement

restrictions.[4]

This study described the COVID-19 transmission dynamics in Ghana by estimating the

time-varying reproduction number (R_t) using Cori et al method[5] and Generalized Growth Model

(GGM) (Supplementary Materials)[6] to assess the impact of interventions at the national and

regional levels and to explore the association between population size and the COVID-19

cumulative incidence. The correlation between population mobility and incident case count, and

between mobility and R_t was also assessed.

The daily number of new infections and daily cumulative incidence data by date of report

for Ghana and each of its sixteen regions was obtained through the Johns Hopkins University

COVID-19 dashboard from March 12, 2020, to December 31, 2021 (Table S1, Figures S1 and S2).

To account for testing delay (three days) and incubation period (six days), the time series was

shifted by nine days to approximate the date of infection.[8] Using three-day moving averages of

interpolated daily incident case count data, we used the EpiEstim package in R version 4.0.3 to

estimate the R_t using the 7-day sliding window and the nonoverlapping time window between

interventions.[5] To compare the R_t before, during, and after policies were implemented and assess

their association with COVID-19 transmission, specific time points at which a bundle of

interventions began were selected for the latter analysis and the average R_t estimates over the

period between two policy change time points were estimated (Table 1).

The power-law relationship between cumulative case number and population size was

investigated at ten time points in 2020-21, using linear regression of $\log_{10}$-transformed per-capita

cumulative case number and $\log_{10}$-transformed population size.[9] The relationship between 7-day

moving average of mobility changes and the 3-day moving average of daily number of new

infections and that between 7-day moving average of mobility changes and the 7-day sliding

window R_t in the first 90 days of the pandemic were assessed using the time-lagged cross-

correlation with data from Google Mobility Report.[10] The Georgia Southern University

Institutional Review Board determined a non-human subject status for this project (H20364) under

the G8 exemption category.

Ghana and its regions experienced four pandemic waves in 2020-21 with epidemic peaks

in July 2020, and January, August and December, 2021. At the national level and in the Greater

Accra and Ashanti regions, the R_t fluctuated around 1, increased to >1 before the epidemic peaks,

and dropped <1 afterwards (Figure 1). The epidemic peak was the highest nationwide and in

Greater Accra region in December 2021 with an estimated R_t of ≥2. Ghana's December 2021

epidemic peak was largely driven by the Greater Accra region, while in other regions, case counts

were increasing fast. Similar patterns in epidemic curves and R_t estimates were observed in

Central, Eastern, Volta and Western regions, with R_t ≥3 in December 2021 (Figure S3).

The R_t estimates obtained using nonoverlapping windows suggested varying associations

between government policies and interventions and the increase and decrease of COVID-19

transmission in Ghana and across regions. The restriction of social gatherings and travel bans

implemented on March 15th, 2020, were associated with insignificant changes in R_t at the national

level and the Greater Accra region. However, when all borders were closed to human traffic

nationwide a week later, the R_t lowered to around 1 accounting for a 32.63% (95% credible

interval, CrI: 22.87%, 41.19%) decrease for Ghana and a 27.41% (95% CrI: 16.68%, 36.26%)

decrease in Greater Accra region. In contrast, the R_t for Ashanti increased by over 100%. On April

27th, 2020, the mandatory wearing of masks at all businesses and organizations was implemented

nationwide which was associated with a decrease in R_t by 19.97% (95% CrI: 18.05%, 21.81%) at

the national level. Ashanti and Greater Accra regions also observed significant declines in R_t. The

relaxation of restrictions on social gatherings was associated with a slight increase in R_t by 4.03%

(95% CrI: 2.61%, 5.49%) for Ghana at the national level and by 6.02% (95% CrI: 4.05%, 7.78%)

in the Greater Accra region, however the other regions observed a decline in R_t by >5%. Reopening

of schools in January 2021 was associated with an increase in transmission in Eastern and Central

regions only. Vaccination rollout was associated with a decline in R_t to <1 in Ghana and the regions

except Greater Accra region which observed about a 3% increase. Overall, the behavioral change

due to Christmas festivities in 2021 was associated with sustained transmission in all regions.

Details of percent changes in R_t are in Table S2.

The assessment of population size and cumulative incidence showed that Ghanaian regions

with larger populations experienced higher COVID-19 attack rates (Figure 2 and Table S3).

Furthermore, there was a weak correlation between mobility changes and COVID-19 incidence in

the first 90 days of the pandemic in Ghana (Tables S4 and S5), e.g., the daily number of new

infections was positively correlated with mobility changes to retails and recreation, grocery and

pharmacy and workplaces 3 days later (r>0, p<0.05 in all 3 cases), but no significant correlation

was observed with residential mobility changes. R_t was negatively correlated with mobility

changes to residence (r=-0.328, p=0.002).

Overall, R_t >1 showed sustained transmission across the country, but some interventions

implemented during the study period were associated with reduced R_t in regions with relatively

higher case counts. Interestingly, relaxation of restrictions at social gatherings was not associated

with an increase in R_t.

The early estimates of R_t for Ghana were lower than the reported values for other countries

like Nigeria,[11] Egypt,[12] and Kenya.[13] The lower R values for Ghana suggests that public health and

social measures were effective in containing the epidemic at the initial stage. The estimates also

differ slightly from Dwomoh and colleagues (2021), but may be explained by the differences in

the methods including the fact that these authors used data from the first 60 days.[14] Nevertheless,

R_t remained greater than 1 at the national level and in most regions indicating sustained

transmission. It is therefore imperative that public health measures are strengthened throughout

the country and efforts are prioritized especially in regions with larger population sizes as the

disparity in the case burden across regions was reported in multiple studies.[15,16] This disparity may

be explained by the difficulty in practicing social distancing due to overcrowding stemming from

high commercial activities, slum areas facilitating disease spread and urban residency. Hence, such

regions will be required to implement more stringent preventive measures in order to decrease

transmission.[17]

The reopening of schools was associated with a surge in cases in other jurisdictions.[18] This

finding supports the need for routine surveillance, case investigation, better protocols for isolation

and quarantine, and deployment of protective personal equipment to schools. Although the

correlation between changes in mobility and transmission intensity was weak in our study, decline

in trips to grocery and pharmacy outlets and workplaces have been reported to be associated with

lower transmission rates in most countries before and after interventions were relaxed.[19] The

difference in correlation between mobility changes and transmission intensity may be due to

cultural differences, economic status, and variations in the types of interventions implemented.

It was unexpected that the relaxation of restrictions on social gatherings was not associated

with increase in the R_t estimates, although such policies make it difficult for social distancing

especially in enclosed places like churches or restaurants. This finding may be observed due to the

residual effect of the prior mask mandates and the reluctance of prominent churches to resume in-

person religious activities.[20] In addition, mobility changes to retail and recreation centers remained

below baseline even after the restrictions were relaxed (Figure S5).

Our study is not without limitations. First, we used publicly available data, which was

subject to underreporting or reporting delays. Second, data were only available by the date of the

report and unavailable by the date of symptom onset. We, therefore, accounted for this by shifting

the data by nine days to approximate the time of infection. It is possible that this method of

approximating the date of symptom affected the results of the cross-correlation analysis, which

suggested that changes in the daily number of new infections were correlated with mobility

changes 3 days later. Third, given the use of aggregate data, individual-level assumptions cannot

be made. Fourth, we cannot rule out the possibility of ceiling effects during months when testing

capacity was limited (Figure S6). Finally, data on socioeconomic variables was limited at the

regional level hence further exploration of the case burden by region could not be performed.

In conclusion, most of the interventions implemented by the Ghanaian government were

followed by an overall decrease in R_t estimates at the national scale, but it did not display a decline

across all regions. This highlights the importance of a sustained, multi-faceted response at the

national level to help mitigate the varying regional effects observed during this pandemic.

Co-Author Contact Information:

Sylvia K. Ofori, Georgia Southern University: so01935@georgiasouthern.edu

Jessica S. Schwind, Georgia Southern University: jschwind@georgiasouthern.edu

Kelly L. Sullivan, Georgia Southern University: ksullivan@georgiasouthern.edu

Benjamin J. Cowling, The University of Hong Kong: bcowling@hku.hk

Gerardo Chowell, Georgia State University: gchowell@gsu.edu

Declaration of competing interests:

Sylvia Ofori declares that she was a paid intern at Ionis Pharmaceuticals and the financial

relationship does not affect the content of the article. Dr. Isaac C. H. Fung declares that he has

invested in equity in Alphabet, Inc. (GOOGL). Prof. Benjamin Cowling declares that he was a

consultant for Roche and Sanofi Pasteur. All other co-authors declare no competing interest.

Table 1: Public health and social measures interventions against COVID-19 implemented in Ghana[1,21-23]

Date	Label	Intervention	Location
15th March 2020	A	• Restricting all air travels to Ghana. • Suspension of all public gatherings including workshops, funerals, and religious activities.	National
16th March, 2020		• Closure of all universities, senior high schools, and basic schools.	National
17th March, 2020		• Mandatory 14-day quarantine for all travelers entering the country. • Travel bans for all travelers who are not Ghanaian citizens or hold resident permits from countries with at least 200 cases. • Isolation and testing of symptomatic individuals. • Activation of contact tracing.	National
22nd March 2020	B	• Closure of all borders to human traffic for two weeks and suspension of passport services.	National
26th March 2020		• Director-General of the Ghana Health Service recalled staff who were on study leave.	National
28th March 2020		• Special health insurance coverage was offered to all frontline workers.	National
30th March 2020	L	• Lockdown of major cities in two regions	Regional
3rd April 2020		• The Director-General of the Ghana Health Service announced that nose masks were going to be manufactured locally. • Disinfection of markets in Northern, North East, Savannah, and Eastern Regions.	Regional
5th April 2020		• Extension of border closure for another two weeks	National
27th April 2020	C	• Mandatory wearing of a mask at all businesses and organizations.	National
11th May 2020		• Hotels, bars, and restaurants are permitted to operate following social distancing measures.	National

Date		Event	Level
		• Extension of the ban of social gathering till the end of May	
5th June 2020	D	• Relaxation of restrictions at social gatherings including church services with mandatory mask-wearing and maximum attendance of 100.	National
15th June 2020		• The president announced a fine for those who refused to wear a mask in public places. • Final year students in tertiary institutions went back to school.	National
21st June 2020		• The president approved the construction of hospital-related facilities in the Greater Accra and Ashanti regions.	Regional
6th July 2020	E	• Deployment of personnel to monitor COVID-19 cases in high schools.	National
18th July 2020		• Two-phase fumigation exercise.	National
26th July 2020		• COVID-19 related restrictions on transport operators and tourist sites were lifted.	National
1st August 2020		• Ease of restrictions on religious activities as the duration of service increased from 1 hour to 2 hours.	National
1st September 2020	F	• Reopening of international air border with mandatory testing of all passengers.	National
7th September 2020		• Relief package announced for private schools affected by school closures.	National
9th January 2021	G	• Re-opening of schools	National
1st March 2021	V	• Vaccine rollout	National
15th December 2021	H	• Behavioral change due to Christmas festivities	National

REFERENCES

1.   Ghana Health Service. COVID-19 Ghana's Outbreak Response Management Updates. Accessed January 15, 2022, 2022. https://ghanahealthservice.org/covid19/archive.php

2.   Sibiri H, Zankawah S, Prah D. Coronavirus diseases 2019 (COVID-19) response: Highlights of Ghana's scientific and technological innovativeness and breakthroughs. *Ethics Med Public Health*. 2020;14:100537.

3.   Hale T, Angrist N, Goldszmidt R, et al. A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). *Nature Human Behaviour*. 2021;5(4):529-538.

4.   Asamoah JKK, Owusu MA, Jin Z, Oduro FT, Abidemi A, Gyasi EO. Global stability and cost-effectiveness analysis of COVID-19 considering the impact of the environment: using data from Ghana. *Chaos Solitons Fractals*. 2020;140:110103-110103. doi:10.1016/j.chaos.2020.110103

5.   Cori A, Ferguson NM, Fraser C, Cauchemez S. A New Framework and Software to Estimate Time-Varying Reproduction Numbers During Epidemics. *American Journal of Epidemiology*. 2013;178(9):1505-1512. doi:10.1093/aje/kwt133

6.   Chowell G. Fitting dynamic models to epidemic outbreaks with quantified uncertainty: A primer for parameter uncertainty, identifiability, and forecasts. *Infectious Disease Modelling*. 2017;2(3):379-398.

7.   Johns Hopkins University Center for Systems Science and Engineering. Novel Coronavirus (COVID-19) Cases Data. https://data.humdata.org/dataset/novel-coronavirus-2019-ncov-cases?force_layout=desktop

8.   Centers for Disease Control and Prevention. COVID-19 Pandemic Planning Scenarios. Accessed November 12, 2020.

. https://www.cdc.gov/coronavirus/2019-ncov/hcp/planning-scenarios.html.

9.   Chowell G, Bettencourt LM, Johnson N, Alonso WJ, Viboud C. The 1918–1919 influenza pandemic in England and Wales: spatial patterns in transmissibility and mortality impact. *Proceedings of the Royal Society B: Biological Sciences*. 2008;275(1634):501-509.

10.   Google LLC. Google COVID-19 Community Mobility Reports. Accessed April 27, 2021, https://www.google.com/covid19/mobility

11.   Adegboye OA, Adekunle AI, Gayawan E. Early Transmission Dynamics of Novel Coronavirus (COVID-19) in Nigeria. *Int J Environ Res Public Health*. 2020;17(9):3054. doi:10.3390/ijerph17093054

12.   Masoud M, Gewaifel G, Gamaleldin N. Transmissibility and mortality trends of COVID-19 epidemic in Egypt. *Alexandria Journal of Medicine*. 2020/01/01 2020;56(1):189-195. doi:10.1080/20905068.2020.1845442

13.   Brand SP, Aziza R, Kombe IK, et al. Forecasting the scale of the COVID-19 epidemic in Kenya. *medRxiv*. 2020;doi:https://doi.org/10.1101/2020.04.09.20059865

14.   Dwomoh D, Iddi S, Adu B, et al. Mathematical modeling of COVID-19 infection dynamics in Ghana: Impact evaluation of integrated government and individual level interventions. *Infectious Disease Modelling*. 2021;6:381-397.

15.   Jahangiri M, Jahangiri M, Najafgholipour M. The sensitivity and specificity analyses of ambient temperature and population size on the transmission rate of the novel coronavirus (COVID-19) in different provinces of Iran. *Science of the total environment*. 2020;728:138872.

16.   Lulbadda KT, Kobbekaduwa D, Guruge ML. The impact of temperature, population size and median age on COVID-19 (SARS-CoV-2) outbreak. *Clinical Epidemiology and Global Health*. 2021/01/01/ 2021;9:231-236. doi:https://doi.org/10.1016/j.cegh.2020.09.004

17.   Ribeiro HV, Sunahara AS, Sutton J, Perc M, Hanley QS. City size and the spreading of COVID-19 in Brazil. *PLoS One*. 2020;15(9):e0239699-e0239699. doi:10.1371/journal.pone.0239699

18.   Quakyi NK, Agyemang Asante NA, Nartey YA, Bediako Y, Sam-Agudu NA. Ghana's COVID-19 response: the Black Star can do even better. *BMJ global health*. 2021;6(3):e005569. doi:10.1136/bmjgh-2021-005569

19.   Nouvellet P, Bhatia S, Cori A, et al. Reduction in mobility and COVID-19 transmission. *Nature Communications*. 2021/02/17 2021;12(1):1090. doi:10.1038/s41467-021-21358-2

20.   Jemima Esinam Kuatsinu. Ghana: Easing of Restrictions On Religious Activities - Churches Reluctant to Congregate for Worship. *The Ghanaian Times*. 4 JUNE 2020. Accessed 05/01/2021. https://allafrica.com/stories/202006040675.html

21.   Nyabor J. Coronavirus: Government bans religious activities, funerals, all other public gatherings. *CITI NEWSROOM*. March 15, 2020. Accessed May 01, 2021. https://citinewsroom.com/2020/03/government-bans-church-activities-funerals-all-other-public-gatherings/

22.   Donald Ato Dapatem CN. Ghana closes its borders for 2 weeks to fight Coronavirus. *GRAPHIC ONLINE*. Accessed 09/25/2020. https://www.graphic.com.gh/news/general-news/ghana-news-coronavirus-borders-closed-directive-will-last-2-weeks.html

23.   Frimpong ED. All Ghana Health Service staff on study leave recalled for Coronavirus 'fight'. *Graphic online*. March 27, 2020. Accessed 3/4/2021. https://www.graphic.com.gh/news/general-news/all-ghana-health-service-staff-on-study-leave-recalled-for-coronavirus-fight.html

Figure legends

Figure 1: The daily number of new cases (left panels), 7-day sliding window R_t (middle panels) and nonoverlapping window R_t (right panels) estimated using the in the 'EpiEstim' package in Ghana, Greater Accra, and the Ashanti region, March 12, 2020—December 31, 2021. The government policies represented by the alphabets in the figure are: A = restrictions of all air travels to Ghana, suspension of social gatherings, school closure, mandatory 14-day quarantine for all travelers, B = closure of all borders to human traffics, L= lockdown of major cities, C = mandatory wearing of a mask at all businesses and organizations, D = relaxation of restrictions at social gatherings, E = deployment of personnel to monitor COVID-19 cases in high schools, F = reopening of international borders, G = Re-opening of schools, V = vaccination rollout, H= Christmas festivities in 2021.

*Greater Accra and Ashanti regions are highlighted because they are the most populous and had the highest case burden in Ghana

Figure 2: Linear regression models between $\log_{10}$-transformed per capita cumulative case count and $\log_{10}$-transformed population size of the sixteen regions of Ghana (grey areas representing the 95% confidence intervals of the regression lines) at 10-time points: June 30, August 31, October 31, December 31, 2020, and February 28, April 30, June 30, August 31, October 31, and December 31, 2021.

Transmission dynamics of COVID-19 in Ghana and the impact of public health interventions

Sylvia K. Ofori,[1] Jessica S. Schwind,[1] Kelly L. Sullivan,[1] Benjamin J. Cowling,[2] Gerardo Chowell,[3] Isaac Chun-Hai Fung[1] *

Online Supplementary Materials

Supplementary text for Methods

Data interpolation and moving averages

The data from obtained from the Johns Hopkins University was interpolated to estimate the values of missing data. The interpolated values were used to estimate the 3-day moving averages to reduce fluctuations in analysis.

Timeline for regions

At the regional level, the time series began when the first cases were reported consecutively and ended after cases were consecutively reported. For example, for Central Region, the time series began on 7th May 2020. This was done to avoid estimating the R_t too early in the pandemic and to minimize error and uncertainty around the R_t. Six out of the sixteen regions were included in the analysis based on having more than 4,000 cumulative cases as of December 31, 2021, to avoid the fluctuations in the estimates (see Figures S1 and S2).

The reproduction number, R$_t$, from daily incidence using the generalized growth model (GGM)

The daily number of new infections for the first 15 days of the epidemic and selected regional levels are calibrated using the Generalized Growth Model (GGM).[1] The GGM uses a simple generalized model for the ascending case of an epidemic using the equation:

$$\frac{dC(t)}{dt} = C'(t) = rC(t)^p$$

where $C'(t)$ characterizes the epidemic curve (incident case count) over the time t and the solution C(t) which is the cumulative incidence at a given time t, r is a positive number defined as the growth rate (1/t), and $p \in [0, 1]$ is the "deceleration of growth parameter". At a p of zero, the equation describes a constant incidence and at $p=1$, the equation outlines an exponential growth dynamics. Values of p between 0 and 1 describe sub-exponential growth patterns.[2]

We first assume a gamma distribution for the generation interval of SARS-CoV-2 with a mean of 4.60 days and a standard deviation of 5.55 days,[3, 4] then estimate the growth rate parameter r, and the deceleration of growth parameter, p. The GGM model simulates the growth of total cases (both imported and local cases) at I_i, and estimates the reproduction using the renewal equation:[5]

$$R_{t_i} = \frac{I_i}{\sum_{j=0}^{i}(I_{i-j} + \alpha J_{i-j})\rho_j}$$

The imported cases at a given time t are given as J_i; the local incidence is I_i at calendar time t_i; and ρ_j represents the discretized probability distribution of generation interval. The factor α measures the relative contribution of imported cases to the transmission process and is assumed to be 0.15. The total number of new cases is the numerator at a given time $t= I_i$, and the denominator is the total number of primary cases that generated the secondary cases. Thus, the reproduction number is the average number of new cases generated by a primary case at a given calendar time. The uncertainty around the R_t is derived from the parameter estimates (r, p). The R_t is estimated from 300 simulations assuming a negative binomial structure with a mean which is assumed to be a third of the variance.[2] The R_t for the national curve is estimated from the first fifteen days of infection (March 12th – March 26, 2020).

Time-varying reproduction number R_t, using the Instantaneous Reproduction Number method (Cori et al. method).

The R_t is used because it can be estimated easily in real-time and when control measures are implemented.

The instantaneous reproduction number method in the R package EpiEstim developed by Cori et al and Thompson et al. was used in the analysis.[6, 7] The EpiEstim package uses a Bayesian framework assuming a gamma prior distribution for the posterior distribution for R_t.[3, 6, 8] The parametric method was used in which the serial interval was assumed to follow a gamma distribution with a mean of 4.60 days and a standard deviation of 5.55.[3, 4]

R_t using 7-day sliding windows.

The instantaneous reproduction number was estimated over a 7-day sliding window given the high variation in daily R_t estimates. The R_t was assumed to be constant in each time window and the average of R_t estimates over 7 days was estimated with its credible intervals.

R_t using non-overlapping windows to assess the impact of various interventions.

This method is used to compare the R_t before, during, and after a policy is implemented to assess its impact on transmission.[7] An average was taken of the R_t estimates over the period between two policy change time points. The data on the major interventions and newly implemented policies were obtained from the Ghana Health Service website as well as the websites of the major news media. The following data were collected: travel bans and border closures, restriction of social activities, school closure, the mandatory wearing of masks, relaxation of restrictions on social and religious activities, deployment of personnel to monitor COVID-19 in senior high schools, reopening of air borders, and reopening of schools. A detailed list of interventions and their

references can be found in Table 1. Bundled policies were implemented in close succession; hence there is difficulty in assessing the individual effects. **Regression analysis**

The power-law relationship between cumulative case count, C, and population size, N (i.e., $C \sim N^g$ where g is an exponent) was transformed into a relationship between $\log_{10}$-transformed per-capita cumulative case count and $\log_{10}$-transformed population size as follows:[9]

$$\frac{\log_{10}(\text{cumulative case count})}{\log_{10}(\text{population size})} = g$$

$$\frac{\log_{10}(\text{per capita cumulative case count} \times \text{population size})}{\log_{10}(\text{population size})} = g$$

$$\frac{\log_{10}(\text{per capita cumulative case count})}{\log_{10}(\text{population size})} + \frac{\log_{10}(\text{population size})}{\log_{10}(\text{population size})} = g$$

$$\frac{\log_{10}(\text{per capita cumulative case count})}{\log_{10}(\text{population size})} = g - 1 = m$$

The per capita cumulative case count was calculated by dividing the total cumulative case count by the population size of the nation or region at the specified time point. A linear regression model was fit to the data of $\log_{10}$-transformed per capita cumulative case count and $\log_{10}$-transformed population size. The slope *m* of the regression line (where *m=g-1*) can be interpreted as follows: if *m=0*, there is a homogeneity of per capita cumulative case count across regions; if *m<0*, low-population regions have higher per capita cumulative case count than high-population regions; and if *m>0*, low-population regions have lower per capita cumulative case count than high-population regions.[9] The 2020 population estimates for Ghana and the regions were obtained from the Ghana Statistical Service.[10]

Google mobility data

The mobility data was sourced from Google to analyze changes in the number of visits to places in the following categories: (a) grocery and pharmacy, (b) parks, (c) transit stations, (d) retail and recreation, (e) residential, and (f) workplaces.[11] The data provides information for how visits and duration of staying at different places changes compared to baseline values which is the median value for the corresponding day of the week, during the five weeks of January 3 — February 6, 2020. The data is representative of users who opted for location history. The relationship between the 7-day moving average of mobility changes in retails and recreation, grocery and pharmacy, and workplace, and the 3-day moving average of the daily number of new infections (or R_t estimates from a 7-day sliding window) is assessed using the time-lagged cross-correlation using the assumed dates of infection.

Supplementary text for Results

Epidemic curves

Four waves were observed in the epidemic curves for Ghana. The daily number of new cases of COVID-19 in Ghana, the Greater Accra Region, and the Ashanti region decreased from mid-May to mid-June and surged in mid-July, 2020. For all three locations, the daily number of new cases dropped again to less than 500 per day and spiked again in January 2021 (Figure 1). Relatively few new cases were observed between March and July 2021, before rising again in late July. Very high numbers of new cases were recorded from December 2021 due to Christmas festivities. The Western region's daily new cases were highest in June and July 2020, and January-February 2021 (Figure S3). The daily number of new cases for the Central, Volta, and Eastern regions fluctuated

throughout the study period but followed a similar trajectory with that of Ghana as a whole in the peak periods (Figures S3).

Comparing R_t Estimates using GGM and Cori et al. method.

Using the GGM, the COVID-19 R_t estimate for Ghana was estimated at 1.8 (95% CI: 1.7, 2) for the first 15 days of the epidemic (Figure S4) compared to that estimated using the Cori et al. method, which estimated the R_t at 1.85 (95% CI: 1.42, 2.27).

The difference in estimates may be because in the Cori et al. method, fluctuations in daily number of new infections are treated as true reflections of the underlying epidemic while the GGM method treats these fluctuations as noise by assuming an error structure underneath. Hence, the estimated value of R_t using the Cori et al. method may provide a better reflection of how early interventions and policy changes affected the R_t at the national level for this study.

REFERENCES

1.   Viboud C, Simonsen L, Chowell G. A generalized-growth model to characterize the early ascending phase of infectious disease outbreaks. *Epidemics*. 2016;15:27-37.
2.   Chowell G. Fitting dynamic models to epidemic outbreaks with quantified uncertainty: A primer for parameter uncertainty, identifiability, and forecasts. *Infectious Disease Modelling*. 2017;2(3):379-398.
3.   Fung IC-H, Hung YW, Ofori SK, Muniz-Rodriguez K, Lai P-Y, Chowell G. SARS-CoV-2 Transmission in Alberta, British Columbia, and Ontario, Canada, December 25, 2019, to December 1, 2020. *Disaster Medicine and Public Health Preparedness*. 2021:1-10. doi:doi:10.1017/dmp.2021.78
4.   You C, Deng Y, Hu W, et al. Estimation of the time-varying reproduction number of COVID-19 outbreak in China. *International Journal of Hygiene Environmental Health*. 2020;228:113555.

5.   Nishiura H, Chowell G. Early transmission dynamics of Ebola virus disease (EVD), West Africa, March to August 2014. *Eurosurveillance*. 2014;19(36):20894.
6.   Cori A, Ferguson NM, Fraser C, Cauchemez S. A New Framework and Software to Estimate Time-Varying Reproduction Numbers During Epidemics. *American Journal of Epidemiology*. 2013;178(9):1505-1512. doi:10.1093/aje/kwt133
7.   Thompson R, Stockwin J, van Gaalen RD, et al. Improved inference of time-varying reproduction numbers during infectious disease outbreaks. *Epidemics*. 2019;29:100356.
8.   Fraser C, Cummings DA, Klinkenberg D, Burke DS, Ferguson NM. Influenza transmission in households during the 1918 pandemic. *American journal of epidemiology*. 2011;174(5):505-514.
9.   Chowell G, Bettencourt LM, Johnson N, Alonso WJ, Viboud C. The 1918–1919 influenza pandemic in England and Wales: spatial patterns in transmissibility and mortality impact. *Proceedings of the Royal Society B: Biological Sciences*. 2008;275(1634):501-509.
10.   Ghana Statistical Service. Ghana's population by region, 2020. Accessed 04/07/2021, 2021. https://www.statsghana.gov.gh/index.php?id=MjYzOTE0MjAuMzc2NQ==/webstats/4238n0op4p
11.   Google LLC. Google COVID-19 Community Mobility Reports. Accessed April 27, 2021, https://www.google.com/covid19/mobility

Table S1: Cumulative number of cases and cumulative incidence rate by region as of December 31, 2021 (N= 142,986)

Region	Population	Cumulative Number of Cases	Cumulative Number of Cases per 100,000
Ghana (national)	32,000,000	142,986	446.83
Ahafo	613,000	1,074	175.20
Ashanti	5,924,500	21,646	365.36
Bono	1,168,800	2,119	181.30
Bono East	1,133,800	2,667	235.23
Central	2,605,500	5,143	197.39
Eastern	3,318,900	6,661	200.70
Greater Accra	5,055,900	79,567	1573.75
North East	588,800	283	48.06
Northern	1,948,900	1,787	91.69
Oti	759,800	850	111.87
Savannah	594,700	263	44.22
Upper East	1,302,700	1,484	113.92
Upper West	868,500	1,543	177.66
Volta	1,907,700	5,621	294.65
Western	2,214,700	7,817	352.96
Western North	949,100	1,049	110.53

Table S2: Median estimates and 95% credible intervals of nonoverlapping window R_t for each period between each policy change (left) and percent changes in nonoverlapping window R_t and associated 95% credible intervals with given policies.

Policy change	R_t Median	95% Credible Interval	Percentage change (%) Median	95% Credible Interval
National				
Before A	1.92	1.54, 2.34		
A to B	1.54	1.42, 1.67	-20.19	-34.75, +3.91
B to L	1.05	0.94, 1.16	-32.63	-41.19, -22.87
L to C	1.31	1.29, 1.34	+25.62	+13.60, +40.42
C to D	1.05	1.04, 1.06	-19.97	-21.81, -18.05
D to E	1.09	1.09, 1.10	+4.03	+2.61, +5.49
E to F	0.92	0.91, 0.93	-15.95	-16.86, -15.10
F to G	1.04	1.03, 1.05	+13.01	+11.74, +14.25
G to V	1.03	1.03, 1.04	-0.60	-1.58, 0.49
V to H	0.98	0.98, 0,99	-4.80	-5.51, -4.09
Beyond H	1.09	1.08, 1.10	+10.57	+9.62, +11.47
Regions				
Greater Accra				
Before A	1.75	1.35, 2.21		
A to B	1.62	1.47, 1.77	-7.48	-27.84, +20.48
B to L	1.17	1.06, 1.29	-27.41	-36.26, -16.68
L to C	1.27	1.23, 1.29	+7.70	-2.70, +19.31
C to D	1.00	0.99, 1.02	-20.84	-22.88, -18.75
D to E	1.06	1.05, 1.07	+6.02	+4.05, +7.78
E to F	0.95	0.94, 0.96	-10.07	-11.45, -8.68
F to G	1.03	1.02, 1.04	+8.24	+6.87, +9.64
G to V	0.98	0.97, 0.99	-5.30	-6.47, -4.30
V to H	1.01	1.00, 1.02	+3.43	+2.35, +4.46

Beyond H	1.47	1.45, 1.49	+45.15	+43.33, +46.99
Ashanti				
Before A	1.84	1.20, 2.69		
A to B	0.61	0.29, 1.11	-66.67	-85.65, -32.54
B to L	2.15	1.50, 2.96	+250.86	+84.23, +681.67
L to C	1.40	1.27, 1.54	-35.34	-52.66, -6.37
C to D	1.13	1.10, 1.16	-19.34	-26.19, -11.69
D to E	1.03	1.02 1.05	-8.49	-11.08, -5.82
E to F	0.95	0.93, 0.96	-8.43	-10.45, -6.53
F to G	1.00	0.95, 1.05	+5.75	+0.35, +10.98
G to V	1.03	1.01, 1.05	+2.87	-2.34, +8.52
V to H	0.98	0.97, 0.99	-4.84	-6.67, -2.27
Beyond H	1.45	1.39, 1.50	+47.31	+41.77, +52.84
Central				
Before C	1.47	1.04, 1.99		
C to D	1.16	1.10, 1.22	-20.86	-42.14, +14.15
D to E	1.06	1.00, 1.09	-9.65	-15.07, -3.46
E to F	0.94	0.91, 0.97	-9.99	-14.28, -5.25
F to G	0.94	0.88, 0.99	-0.58	-7.17, +6.52
G to V	1.16	1.12, 1.20	+23.31	+15.66, +32.75
V to H	0.95	0.93, 0.97	-17.86	-21.02, -14.39
Beyond H	1.24	1.00, 1.51	+29.70	+5.24, +59.34
Eastern				
Before C	1.21	1.09, 1.34		
C to D	1.21	1.11, 1.32	+0.63	-12.97, +15.60
D to E	1.05	1.01, 1.08	-13.89	-21.62, -4.77
E to F	0.99	0.96, 1.01	-5.54	-9.76, -1.18
F to G	0.88	0.84, 0.94	-10.15	-15.58, -4.52
G to V	1.05	1.02, 1.07	+17.86	+10.66, +26.23

V to H	0.98	0.96, 0.99	-6.73	-9.74, -3.58
Beyond H	1.55	1.35, 1.76	+58.30	+40.16, +80.74
Volta				
Before C	1.73	1.40, 2.12		
C to D	1.43	1.33, 1.54	-18.29	-31.95, +3.86
D to E	0.85	0.80, 0.91	-40.35	-45.64, -34.66
E to F	0.90	0.84, 0.96	+4.92	-4.71, +15.57
F to G	1.06	0.97, 1.16	+18.83	+5.92, +31.69
G to V	1.08	1.04, 1.12	+1.06	-7.43, +11.28
V to H	0.97	0.95, 0.99	-10.08	-12.68, -6.92
Beyond H	2.20	2.02, 2.39	+127.92	+109.66, +148.11
Western				
Before C	1.53	1.24, 1.87		
C to D	1.26	1.22, 1.30	-18.77	-33.71, +0.40
D to E	0.97	0.94, 0.99	-23.26	-26.55, -19.97
E to F	0.85	0.82, 0.88	-11.71	-15.88, -7.83
F to G	1.12	1.05, 1.19	+31.58	+22.22, +41.45
G to V	1.04	1.02, 1.06	-7.49	-13.28, -1.24
V to H	0.95	0.93, 0.97	-8.37	-11.05, -5.64
Beyond H	1.91	1.79, 2.03	+101.03	+88.67, +114.64

Table S3: Linear regression analysis between the $\log_{10}$-transformed per capita cumulative case count and $\log_{10}$-transformed population size using cumulative incidence for sixteen regions.

Time point	Slope	95% confidence interval	P value
June 30, 2020	1.59	0.91, 2.28	0.0002
August 31, 2020	1.17	0.53, 1.82	0.0016
October 31, 2020	1.19	0.54, 1.84	0.0014
December 31, 2020	1.21	0.57, 1.85	0.0011
February 28, 2021	1.10	0.62, 1.58	0.000247
April 30, 2021	0.94	0.53, 1.35	0.000216
June 30 2021	0.94	0.52, 1.35	0.00024
August 31 2021	0.91	0.48, 1.34	0.00050
October 31 2021	0.86	0.45, 1.28	0.00052
December 31 2021	0.90	0.47, 1.33	0.00048

Table S4: Time-lag correlation coefficients between the 3-day moving average of daily number of new COVID-19 cases, and 7-day moving average of relative mobility (percentage change from baseline) for trips to retails and recreation, grocery and pharmacy, workplace, and residential categories over the first 90 days of the pandemic in Ghana using the assumed date of infection.[b]

Lag (days)	Retails and recreation		Grocery and pharmacy		Workplace		Residential	
	r	P value	r	P value	r	P value	r	P value
-3	0.042	0.693	0.172	0.103	0.118	0.263	0.067[a]	0.528
-2	0.069	0.516	0.205	0.052	0.149	0.159	0.044	0.629
-1	0.096	0.363	0.237	0.024	0.178	0.091	0.022	0.836
0	0.123	0.242	0.271	0.010*	0.209	0.047*	0	0.999
1	0.167	0.113	0.307	0.004*	0.248	0.019*	-0.042	0.687
2	0.208	0.048*	0.341	0.001*	0.291	0.006*	-0.085	0.418
3	0.243[a]	0.021*	0.366[a]	0.001*	0.328[a]	0.002*	-0.124	0.240

*p<0.05

[a] The model with the largest (absolute value) correlation coefficient in each category.

[b] Data for the first 90 days was used to observe the impact of relaxation of social restrictions on mobility changes.

Table S5: Time-lag correlation coefficients between median 7-day sliding window R_t of COVID-19 and 7-day moving average of relative mobility (percentage change from baseline) for trips to retails and recreation, grocery and pharmacy, workplace, and residential categories over the first 90 days of the pandemic in Ghana using the assumed date of infection.[b]

Lag (days)	Retails and recreation		Grocery and pharmacy		Workplace		Residential	
	r	P value	r	P value	r	P value	r	P value
-3	0.204[a]	0.053	0.061[a]	0.562	0.137	0.192	-0.295	0.005*
-2	0.203	0.054	0.055	0.602	0.139[a]	0.188	-0.310	0.003*
-1	0.197	0.062	0.042	0.691	0.130	0.216	-0.317	0.003*
0	0.197	0.061	0.031	0.766	0.124	0.240	-0.328[a]	0.002*
1	0.164	0.119	0.004	0.972	0.099	0.346	-0.297	0.005*
2	0.133	0.207	-0.020	0.851	0.072	0.497	-0.264	0.012*
3	0.108	0.304	-0.033	0.755	0.051	0.631	-0.238	0.024*

*p<0.05[a] The model with the largest (absolute value) correlation coefficient in each category.

[b] Data for the first 90 days was used to observe the impact of relaxation of social restrictions on mobility changes.

Supplemental Figure Captions and Legends

Figure S1: Cumulative case count as of December 31, 2021 in Ghana by region.

Figure S2: Cumulative case count per 10,000 population as of December 31, 2021 in Ghana by region.

Figure S3: The daily number of new cases (left panel), 7-day sliding window R_t (middle panel) and nonoverlapping window R_t (right panel) estimated using the Cori et al. method in the 'EpiEstim' package, in the Central, Eastern, Volta and Western regions, March 12, 2020— December 31, 2021. The government policies represented by the alphabets in the figure are: A = restrictions of all travels to Ghana, suspension of social gatherings, school closure, mandatory 14-day quarantine for all travelers; B = closure of all borders to human traffics; C = mandatory wearing of facemasks at all businesses and organizations; D = relaxation of restrictions at social gatherings; E = deployment of personnel to monitor COVID-19 cases in high schools; F = reopening of international borders; G= reopening of schools, V = vaccination rollout, H = Christmas festivities in 2021.

Figure S4: Results from the Generalized Growth Model. Upper panel shows the estimated reproduction number of 1.8 (95% CI: 1.7, 2) for Ghana with 95% CI from March 12, 2020, to March 26, 2020. The growth rate, *r*, 0.84 (95% CI: 0.61, 1.1), deceleration growth parameter, *p*, 0.8 (95% CI: 0.74, 0.87). The lower panel shows the GGM fit of the daily reported number of new infections using the first 15 days of data.

Figure S5: Google mobility trends by type of mobility: retail and recreation (first panel), grocery and pharmacy (second panel), workplaces (third panel) and residential (fourth panel) from February 15, 2020, to December 31, 2021. Both the original data (bar) and the 7-day moving average of mobility data (red line) are displayed.

Figure S6: COVID-19 testing positivity rate in Ghana by date of report from May 5, 2020 to October 30, 2021.