Late surges in COVID-19 cases and varying transmission potential partially due to public health policy changes in 5 Western states, March 10, 2020-January 10, 2021

3 4

5 Xinyi Hua[†], MPH; Aubrey R. D. Kehoe[†], MPH; Joana Tome[†], MPH; Mina Motaghi[†], MD,

- 6 MPH; Sylvia K. Ofori, MPH; Po-Ying Lai, MS; Sheikh Taslim Ali, PhD; Gerardo Chowell,
- 7 PhD; Anne C. Spaulding, MD, MPH; Isaac Chun-Hai Fung*, PhD.
- 8

9 Affiliations:

- Department of Biostatistics, Epidemiology and Environmental Health Sciences, Jiann Ping Hsu College of Public Health, Georgia Southern University, Statesboro, GA, USA
 (X. Hua, A. R. D. Kehoe, J. Tome, M. Motaghi, S. K. Ofori, I.C.H. Fung)
- Department of Biostatistics, School of Public Health, Boston University, Boston, MA,
 USA (P.Y. Lai)
- WHO Collaborating Centre for Infectious Disease Epidemiology and Control, Division of Epidemiology and Biostatistics, School of Public Health, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region, China (S.T. Ali)
 - 4. Laboratory of Data Discovery for Health, Hong Kong Science and Technology Park, Hong Kong Special Administrative Region, China (S.T. Ali)
 - 5. Department of Population Health Sciences, School of Public Health, Georgia State University, Atlanta, GA, USA (G. Chowell)
 - 6. Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA, USA (A.C. Spaulding)
- 25 *Corresponding author: Isaac Chun-Hai Fung, PhD. Email: cfung@georgiasouthern.edu
- 26 † Xinyi Hua, Aubrey R. D. Kehoe, Joana Tome and Mina Motaghi serve as co-first authors.
 27
- 28

19

20

21

22

23

24

29

30 Word count of the main text: 3906

31

32 **Conflict of interest statement:** ACS reports grants through her institution from the National 33 Science Foundation, Cellex and Gilead Sciences. Via a subcontract with Einstein School of 34 Medicine, she has received funding from the National Institutes of Health. She has received 35 personal fees from, and served on an advisory board for, Gilead Sciences. She reports honoraria 36 through third parties funded by Gilead, AbbVie, and Merck; personal fees from National 37 Commission on Correctional Health Care, National Sheriff's Association, Harris County 38 Sheriff's Office and California Department of Corrections and Rehabilitation; and travel 39 expenses from bioLytical Sciences and Guardian. 40

41

42

43 Abstract (200 words, max. 200 words)

- 44 **Objective:** This study investigates how the SARS-CoV-2 transmission potential varied in North Dakota,
- 45 South Dakota, Montana, Wyoming, and Idaho from March 2020 through January 2021.
- 46 Methods: Time-varying reproduction numbers, R_t , of a 7-day-sliding-window and of non-overlapping-
- 47 windows between policy changes were estimated utilizing the instantaneous reproduction number
- 48 method. Linear regression was performed to evaluate if per-capita cumulative case-count varied across
- 49 counties with different population size.
- 50 **Results:** The median 7-day-sliding-window R_t estimates across the studied region varied between 1 and
- 51 1.25 during September through November 2020. Between November 13 and 18, *R_t* was reduced by
- 52 14.71% (95% credible interval, CrI, [14.41%, 14.99%]) in North Dakota following a mask mandate;
- 53 Idaho saw a 1.93% (95% CrI [1.87%, 1.99%]) reduction and Montana saw a 9.63% (95% CrI [9.26%,
- 54 9.98%]) reduction following the tightening of restrictions. High-population counties had higher per-capita
- 55 cumulative case-count in North Dakota at four time points (June 30, August 31, October 31, and
- 56 December 31, 2020). In Idaho, North Dakota, and South Dakota, there was a positive correlation between
- 57 population size and per-capita weekly incident case-count, adjusted for calendar time and social
- 58 vulnerability index variables.
- 59 **Conclusions:** R_t decreased after mask mandate during the region's case-count spike suggested reduction
- 60 in SARS-CoV-2 transmission.
- 61
- 62 Keywords: Ecological study, Facemask, COVID-19, Reproduction number, SARS-CoV-2
- 63

64 Abbreviation list:

- 65 1. Coronavirus disease 2019 = COVID-19
 - 2. Severe acute respiratory syndrome coronavirus 2 = SARS-CoV-2
 - 3. Social Vulnerability Index = SVI
 - 4. United States = U.S.
- 68 69

66

67

70

71 INTRODUCTION

72 A case of coronavirus disease 2019 (COVID-19) reported in the state of Washington on 73 January 21, 2020 heralded the arrival of the pandemic in the United States (U.S.), but 74 neighboring states were spared for over a month. While Idaho borders Washington, it and four 75 other contiguous western states-North Dakota, South Dakota, Montana, and Wyoming-did not report a single case until 49 days later.¹ All five states are sparsely populated.² Due to 76 77 demographic heterogeneity, and the division of power between federal and state governments, 78 the reaction to various phases of the pandemic have differed in type and timing by region, 79 resulting in dissimilar patterns of disease spread around the country.³ More regional studies are needed to assess the variation in the spatial heterogeneities of the transmission of the severe 80 81 acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and the distribution of disease burden. 82 In this paper, we will analyze the disease pattern in North Dakota, South Dakota, Idaho, Montana 83 and Wyoming to provide insight for epidemiologists and to help policy makers with future 84 decision-making processes.

85 Orders from the executive branch of federal and state governments and state health 86 officer orders regarding the COVID-19 pandemic included social distancing, guarantine and 87 isolation, mask mandates, closure of businesses, and COVID-19 testing requirements. These 88 were implemented in different time frames across this 5-state region. Evaluating the existing data 89 to explore the impact of policy implementation on COVID-19 pandemic may provide insight on 90 the policies with high impact in reducing the transmission of SARS-CoV-2 infection, morbidity and mortality. Quantifying epidemiologic characteristics of the COVID-19 pandemic in these 91 92 states, so that we can document the potential effect of policies and non-pharmaceutical 93 interventions that reduce COVID-19 transmission and mortality, may make us better prepared 94 for the emergence of future infectious disease epidemics.

95 Central to the description of an epidemic's transmission potential is the reproduction 96 number. The basic reproduction number, also called R_0 , shows the transmissibility of an 97 infectious agent at the beginning of an outbreak; it is calculated as the average number of 98 secondary cases generated by a primary case in a completely susceptible population, prior to any

behavioral changes or public health interventions.⁴ On the other hand, the time-varying reproduction number, also known as R_t , is a time-dependent estimate of the average number of secondary cases that are generated from one case at time t, after there are behavioral changes, depletion of the susceptible population, and implementation of disease control policies.^{5,6}

103 An R_t larger than one indicates sustained transmission and the epidemic is expected to 104 expand in the population. An R_t less than one indicates that the epidemic will tend to decline. 105 Therefore, it is used as an indication of the effectiveness for infection control measures.^{4,5} 106 Calculating R_t over the course of pandemic, from March 2020 through January 2021, this study 107 aims to investigate the time-dependent variability in transmission potential of SARS-CoV-2 in 108 these five states in different time periods, and explore their relationship with the changes in the 109 states' public health policies.

110

111 METHODS

We used time-series data for the COVID-19 pandemic during March 10, 2020 – January 10, 2021, in the states of North Dakota, South Dakota, Idaho, Montana, and Wyoming. A detailed list and description of all counties (North Dakota's 53 counties, South Dakota's 66

counties, Idaho's 44 counties, Montana's 56 counties, and Wyoming's 23 counties) are provided
 in Supplementary Table 1.

117

118 DATA ACQUISITION

119 We downloaded the cumulative confirmed case count during March 10, 2020 – January 10, 2021, for all five states, including the counties located in each state from the New York 120 121 Times GitHub data repository.⁷ The first case of each state was reported during the same week, on March 10 (South Dakota), March 11 (North Dakota and Wyoming), and March 13 (Idaho and 122 123 Montana) respectively. Our cutoff point for all five states was January 10, 2021. Our timeframe 124 covered nearly ten months from the first reported case in those states. We obtained the daily 125 number of newly confirmed COVID-19 cases from the reported cumulative case count numbers 126 (Appendix A). We also retrieved 2019 county-level estimated population data for all five states 127 from the U.S. Census Bureau.⁸

128

We collected and assessed the executive orders from the governors' offices of the five states and identified the timing of the orders to implement and the announcements permitting relaxation of public health interventions in each state respectively (**Supplementary Table 2**).

133 STATISTICAL ANALYSIS

134 We estimated the time-varying reproduction number, R_t , using the instantaneous 135 reproduction number method with parametric definition of the serial intervals as proposed by 136 Cori et al.⁶ as implemented in the R package 'EpiEstim' version 2.2-3. The instantaneous 137 reproduction number is one of a few definitions of R_t . It is an estimate of the transmissibility of 138 the disease at current time t, assuming that it is the same as the transmissibility of prior cases that 139 result in the number of their secondary cases at current time t. An average R_t can be estimated 140 using a fixed sliding window or non-overlapping time windows defined by the user. R_t is a time 141 varying measure of transmissibility and defined as the ratio between the number of incident cases 142 at the time t, and the total infectiousness of all infected individuals at the time t accounted during 143 their infectious periods. As county-level data may give information of health inequities, or 144 crucial information in differences in COVID-19 transmission rate,⁹ we conducted our analysis 145 based on county level data. We first reconstructed the date of infection according to Gostic et al.,¹⁰ by shifting the time series by nine days backward (assuming a mean incubation period of 6 146 days and a median delay to testing of 3 days).¹¹ We assumed the serial interval distribution with 147 a mean of 4.60 days and a standard deviation of 5.55 days.¹² Besides using the default 7-day 148 149 sliding window, we also estimated R_t by the non-overlapping time periods when different 150 combinations of non-pharmaceutical interventions (i.e., face masking, social distancing, school 151 and business closure, etc.) have been implemented (we call them policy change R_t thereafter). 152 The policy change R_t is the average of the daily R_t over the non-overlapping time period between 153 two major policy changes.

154

155 Percentage change is often utilized to help identify the magnitude and direction of change 156 in a statistic. We calculated percentage change for both the 7-day sliding window R_t and the 157 policy change R_t (**Table 1**). This was calculated utilizing percentage change $=\frac{t_2-t_1}{t_1} \times 100$, at

158 the date of each policy implementation and face-to-face school resumption. For 7-day sliding

- 159 window R_t each date of interest is considered time 2 (t₂) and the previous 7-day period was
- 160 utilized as time 1 (t_1); for the policy change R_t each time window was compared to the previous

161 window. We utilized EpiEstim "sample from the posterior R distribution" function to sample 162 1000 estimates of R_t for each t_1 and t_2 then estimate the 95% credible intervals (2.5 and 97.5

- 163 percentile) of the calculated percentage change through bootstrapping.
- 164 165 As a note, R_t percentage change was calculated for the fall implemented interventions of 166 North Dakota on November 13 (Mask order), Idaho on November 14 (Stay Health Order-Stage 167 2: Gatherings of no more than 10 people, and vulnerable population were strongly recommended 168 to stay at home), and Montana on November 18 (Additional mitigation measures were 169 implemented including reduced size of gatherings to no more than 25). Since neither South 170 Dakota nor Wyoming ever ordered interventions during the same time period, for comparison 171 purposes we calculated the 7-day-sliding-windows R_t percentage change for both states with 172 November 13-18, 2020 as t₂ and November 6-11, 2020 as t₁.
- 173

174 We assessed the power-law relationship between the cumulative case count of COVID-175 19 and the population size of the counties. Cumulative case count would be exactly proportional 176 to population size if the per capita cumulative case count remained the same across all counties 177 of different population size (Appendix B). We ran a linear regression model between $\log_{10^{-10}}$ 178 transformed per capita cumulative case count and log₁₀-transformed population size. Counties 179 with lower population sizes would have a higher per capita cumulative case count if the slope of 180 the regression line was negative, and a lower per capita cumulative case count if the slope was positive.¹³⁻¹⁶ We conducted the log-linear regression considering four different dates: June 30th, 181 182 August 31st, October 31st and December 31st, 2020.

183

184 To further explore the relationship between the per capita weekly incident case count and 185 county's population size, log linear analysis was performed as univariable and multivariable 186 analysis. Multivariable models were adjusted for time (more specifically a categorical variable of 187 each 7-day period from March 2020 to January 2021) and other factors that may impact this 188 association. Those other factors were variables that we chose from the CDC Social Vulnerability 189 Index (SVI), which was created by Geospatial Research, Analysis & Services Program to 190 identify and map the most vulnerable communities with a need for support in emergency 191 events.¹⁷ CDC utilizes the U.S. census data to estimate the social vulnerability for every census 192 tract and every county in each state. The SVI is composed of many variables which are 193 categorized in four different themes (socio-economic status, household composition and 194 disability, race/ethnicity/languages, and housing type/transportation). In times of COVID-19, it 195 becomes meaningful to account for social vulnerability variables when assessing the relationship 196 between weekly incident case count per capita and population size. Those variables may partially 197 explain the variability of weekly incident case count per capita. Among many variables included 198 in the SVI database, we chose those variables we thought were more representative and had less 199 collinearity with each other. The chosen variables were the percentage of people living below the 200 poverty threshold, the percentage of people without a high school diploma, the percentage of 201 people 65 and older, the percentage of the civilian non-institutionalized population with a 202 disability, the percentage of minorities (all people excluding white, non-Hispanic) estimate, and 203 the percentage of occupied housing units with more people than rooms estimate. We obtained 204 our data from the Agency for Toxic Substances and Disease Registry.¹⁸

All statistical analyses were performed using R version 4.0.3 (R Core Team, R
 Foundation for Statistical Computing, Vienna, Austria). Maps (Figure 1 and Figure 2) were

207 created using R version 3.5.1 (R Core Team, R Foundation for Statistical Computing, Vienna, 208 Austria). The significance level was set a priori at α =0.05.

209

210 ETHICS

The Georgia Southern University Institutional Review Board made a non-human subject determination for this project (H20364) under the G8 exemption category according to the Code of Federal Regulations Title 45 Part 46.

- 214215 <u>RESULTS</u>
- 215

217 *Rt* ESTIMATES AT THE STATE LEVEL

218 From March 10, 2020 to January 10, 2021, the daily number of new cases showed at least 219 one peak in the Fall across all five states, except Idaho which had a summer peak (Figure 3, left 220 panel). All five states had a very similar qualitative trajectory. For instance, as of January 10, 221 2021, Idaho had reported 149,742 cases, the highest cumulative numbers of COVID-19 cases 222 among five states. South Dakota had reported 113,318 cases, North Dakota 94,724 cases, 223 Montana 86,324 cases, and Wyoming 46,832 cases. Figure 2 presents the geospatial dynamics 224 of cumulative cases number and cumulative case number per 1000 population (incidence) by 225 county in five states at four different dates of report between March 10, 2020 and January 10, 226 2021: June 30th, August 31st, October 31st, and December 31st, respectively. 227

228 The 7-day-sliding-window R_t estimates were between 2 and 3, in the beginning of the 229 pandemic across all five states. In Idaho, Montana, and Wyoming, the 7-day-sliding-window R_t 230 estimates dropped below 1 in the early-April; and in North Dakota and South Dakota, the 7-day-231 sliding-window R_t estimates briefly drop below 1 in mid-May. Then the 7-day-sliding-window R_t 232 estimates steadily stayed above 1 between September and December which correspond to the 233 fall/winter surges in five states. At the end of the study period, the 7-day-sliding-window R_t 234 estimates were slightly above 1 (Idaho, Montana, and Wyoming) or around 1 (North Dakota and 235 South Dakota), demonstrating the extensive community transmission of SARS-CoV-2 in those 236 states.

As of September 7, 2020 all students in K-12 grades resumed face-to-face instruction in all five states. Compared to the week prior, the 7-day-sliding-window R_t increased in Montana, Wyoming, Idaho, South Dakota; whereas North Dakota saw a reduction in R_t (**Table 1 & Figure 3, right panel**). Thus, these five states experienced an average of 11.55% increase in the 7-daysliding-window R_t .

243

237

Following the region's fall surge in cases, North Dakota implemented a mask mandate on November 13th (Policy E) which contributed to a 14.71% decrease in R_t . Idaho saw a 1.93%, and Montana a 9.63% decrease in R_t following the implementation of stricter policies (Policy E respectively). Meanwhile in mid-November 2020, neither Wyoming nor South Dakota implemented any additional restrictions seeing differing effects on R_t ; Wyoming's R_t decreased by 13.22% where South Dakota's R_t increased by 2.8%.

250

251 Percentage change in policy change R_t were computed by comparing each time window 252 (between each major policy change) to the previous (**Table 1**). Detailed description of R_t for 253 each state can be found in **Appendices C and D**.

255 POWER-LAW RELATIONSHIP BETWEEN CUMULATIVE CASE NUMBER AND 256 POPULATION SIZE

257 **Figure 4** presents a linear regression relationship between the log_{10} -transformed per 258 capita cumulative case number and the log_{10} -transformed population size for a total of 242 259 counties in North Dakota, South Dakota, Idaho, Montana, and Wyoming. Each panel corresponds to an assessed date (date of report), June 30th, August 31st, October 31st, and 260 261 December 31st, 2020, respectively; each regression line represents a state in a specific assessed 262 date. The slope, *m*, and its 95% confidence internal of every regression line are presented in 263 **Supplemental Table 3**. North Dakota was the only state that had significant slopes at all four 264 time points (m=0.2758, 0.2171, 0.0729, 0.0986; p=0.0034, 0.0018, 0.046, 0.0024; respectively). 265 Overall, the slopes of regression lines at four different assessed dates were found slightly 266 positive in June and August assessments but very close to zero in October and December assessments as the pandemic unfolded, except that of Montana on June 30th, 2020. This meant 267 268 towards the end of 2020, there was no heterogeneity of per capita cumulative case count across 269 242 counties with different population size, suggesting an extensive community transmission of 270 SARS-CoV-2 as the pandemic progressed. However, as we mentioned previously, the slope of 271 Montana's June regression line (-0.1423) was slightly less than zero. When m < 0, it suggests 272 counties with lower population sizes would experience a higher per capita cumulative case count. 273 In other words, the rural counties in Montana experienced a slightly higher impact of the 274 COVID-19 pandemic than other counties in the same state by mid-2020. Figures S1-S5 275 (Appendix E) present the same regression analyses, separately for each state, with outliers in 276 log-transformed per capita cumulative case number highlighted.

277

254

278 MULTIPLE REGRESSION ANALYSIS BETWEEN PER CAPITA WEEKLY INCIDENT CASE 279 COUNT AND POPULATION SIZE, ADJUSTED FOR SVI VARIABLES

280 Linear regression results for evaluating the relationship between \log_{10} -transformed per 281 capita weekly incident case count and log_{10} -transformed population size is shown in **Table 2**. 282 Each ten-fold increase in population size, after adjusting for the time of data collection (therefore 283 adjusting for different policies running in the state or different stages of the pandemic), was 284 associated with an increase of 0.06-0.27 in the log₁₀-transformed per capita weekly incident 285 case count, in the studied states and all these changes were statistically significant. After 286 controlling for SVI confounders, in addition to time, a positive trend was found in the association 287 between population size and per capita weekly incident case count in Idaho, North Dakota, and 288 South Dakota. However, these coefficients were not statistically significant for Montana and

Wyoming.

290 **DISCUSSION**

From March 2020 through January 2021, North Dakota, South Dakota, Idaho, Montana, and Wyoming have taken various approaches on implementing both policy and health guidelines to reduce the community transmission of SARS-CoV-2. Utilizing both the 7-day-sliding window R_t and policy change R_t , we estimate the change in transmission potential over time and after

295 each major policy change. Throughout the initial spring and summer surges across the U.S. 296 coastal and major metropolitan areas, limited community spread of the virus was observed in the 297 five states. Although we did not observe a corresponding surge in reported case numbers in 298 South Dakota in August, the 7-day-sliding-window Rt has a small peak, suggesting there was 299 community transmission related to Sturgis Motorcycle Rally, that was held in Meade County, South Dakota, August 7-16, 2020. Dave et al.,¹⁹ and Carter et al.,²⁰ assessed the nationwide 300 transmission of SARS-CoV-2 following the Sturgis Motorcycle Rally and identified it as a 301 302 superspreading event. However, entering fall 2020, this region began to see an uptick in case 303 counts ultimately leading a peak seen in late November. During the summer, reopening orders 304 were given and between mid-August and September 1st, all five states resumed face-to-face K-12 305 schooling. Many of the universities also began in-person instruction. By September 7th the R_t in 306 all five states had increased by an average of 11.55%.

307

308 This paper used 7-day sliding window R_t estimates and policy change R_t estimates to 309 assess and evaluate the various non-pharmaceutical interventions and policy changes at the state 310 level for North Dakota, South Dakota, Idaho, Montana, and Wyoming. A few studies have 311 accessed the effectiveness of mitigation measures, especially regarding face mask mandate and 312 school reopening.²¹⁻²³ Among the five states, North Dakota, Montana and Wyoming 313 implemented the mask mandate and followed with the R_t reduction, which echoed with the current existing body of research on the efficacy of face masking in preventing the transmission 314 of SARS-CoV-2.²⁴⁻²⁶ However, South Dakota and Idaho also experienced the R_t reduction 315 316 without adopting the mask mandate, perhaps due to voluntary adoption of facemasks, which will 317 require further investigation.

318

319 Furthermore, we observed an increase in R_t after schools (K-12 schools, colleges, and 320 universities) reopened in South Dakota, Idaho, Montana, and Wyoming, except North Dakota. 321 The detailed R_l description for all five states regarding school reopening are provided in 322 Appendix C. School closure and suspended in-person instruction were considered to be one of 323 the major public health interventions, and its effectiveness was widely discussed and researched. 324 Our research results suggest that school reopening is correlated with an increase in R_t estimates; 325 however, the results contradicted with the existing studies that the in-person instruction posed a low risk of transmission of SARS-CoV-2.²⁷⁻²⁹ This might reflect a concurrent change in social 326 contact patterns when parents returned to workplaces after their children went back to schools. 327 328 School closure as a mitigation strategy requires further research on its effectiveness since it is 329 highly related to school-aged children, adolescents, and young adults' physical health, mental 330 health, and quality of life.³⁰

331

North Dakota had a higher cumulative case count per capita in the more densely populated counties; however, the reason of why North Dakota was the only state of the 5 states under study that had statistically significant slopes (*m*) at all four assessed time points requires further investigation. Additional analysis found that 8 out of 20 slopes were very close to 0 (0 < m < 0.1), and the per capita cumulative case count did not vary by different population sizes.

When adjusting for SVI variables, our study found a positive significant association between log₁₀-transformed per capita weekly incident case count and log₁₀-transformed population size for Idaho, North Dakota, and South Dakota, with borderline significance for

8

341 Wyoming, and no significance for Montana. Concretely, a 10-fold increase in population size

342 was associated with a 0.106 increase in log₁₀-transformed per capita incident case count in Idaho,

343 0.195 increase in North Dakota, and 0.127 increase in South Dakota. This positive significant

344 association means that high-population counties have a higher per capita incident case count than

low-population counties. This finding is supported by Wong & Li³¹ study where U.S. counties

with more people were more likely to have larger numbers of cases especially in late Spring and early Summer 2020, or by McLaughlin et al.³² who emphasized the association of COVID-19

- 348 case rate with densely populated counties, urban counties and crowded housing.
- 349

350 **LIMITATIONS**

351 There are some limitations in our study. First, upticks in cases may be due to external 352 events attracting large crowds, such as a July 4, 2020 outdoor rally by the then President Trump 353 at Mt. Rushmore, South Dakota, and a motorcycle rally in Sturgis, South Dakota in August 2020. 354 It is impossible to distinguish local cases that were associated with the Sturgis Motorcycle Rally 355 from those that were not using the aggregate data. Others have found evidence suggesting that 356 the motorcycle rally might be a superspreading event, leading to at least 649 cases nationally that 357 were associated with transmission chains traced back to the event, and Meade County (where the 358 Rally was) experiencing a faster rate of growth in case rate than the rest of South Dakota, a week after the close of the Rally.^{19,20} The latter event occurred just before school opening and its 359 360 independent effect on R_t increase may be difficult to tease out. Second, the date of symptom 361 onset or the date of (unobserved) infection was not available for our dataset. Only the date of 362 report was available. To correct for the time lag, we shifted the epidemic curve by nine days in 363 order to correct for the incubation period and the delay to test results. Third, our dataset lacked 364 information to distinguish between local and foreign imported cases. However, such distinction 365 was mostly important in the early stages of the epidemic and since April 2020 community 366 transmission was the main driver for the epidemic. Therefore, we argue that the absence of this 367 distinction would not have significantly affected R_t estimates since April 2020. Fourth, we used aggregated numbers of reported cases by political jurisdictions instead of separate data on 368 369 different facilities or settings, while each setting might demonstrate a different dynamic than that 370 of community transmission. Fifth, underreporting due to the limits on testing capacity was 371 especially acute at the beginning of the pandemic. The majority of the states issued orders to 372 increase testing capacities between March and April 2020, and thus overcome this challenge. 373 However, mild and asymptomatic cases are unlikely to get tested and confirmed. Thus, state-374 reported data cannot be used to measure the extent to which asymptomatic spread has progressed 375 during the pandemic. Lastly, government orders to undertake a mitigation practice, and actual 376 compliance, may differ, which is a recognized limitation in the data.^{33,34}

377 <u>CONCLUSIONS</u>

378 In North Dakota, South Dakota, Idaho, Montana, and Wyoming, new cases of 379 COVID-19 started to rise in November and peaked in November-December 2020. From March 380 2020 to January 2021, the R_t for North Dakota, South Dakota, Idaho, Montana, and Wyoming 381 fluctuated around one (with a range of 0.5 to 1.5 starting from June). Various social distancing 382 policies including stay-at-home order and closing businesses and other protective interventions 383 such as mask requirements appeared to be associated with a reduction in the spread of SARS-384 CoV-2 and keeping the R_t at a low level in the states studied in this paper. On May 13, 2021, the 385 Centers for Disease Control and Prevention updated their guidelines regarding face coverings 386 indicating that fully vaccinated people do not have to wear masks or practice physical distancing

387 except where required by federal, state, local, tribal, or territorial laws, rules, and regulations.³⁵

- 388 Since individuals fully vaccinated with COVID-19 vaccine still have a small chance becoming
- infected, this updated guideline is controversial among some healthcare professionals and
- 390 requires further investigation.³⁶ While we await the vaccination of more people so that the herd
- immunity threshold can be reached, government agencies may consider the continuation of the
- above-mentioned policies to manage transmission and keep the R_t below 1.0.
- 393
- 394395 References:
- 396
- Johns Hopkins University. COVID-19 United States Cases by County
 <u>https://coronavirus.jhu.edu/us-map</u>. Published 2021. Accessed February 28, 2021.
 The United States Census Bureau. 2020 Census: 2020 Resident Population for the 50
 States, the District of Columbia, and Puerto Rico. United States Census Bureau.
 <u>https://www.census.gov/library/visualizations/2021/dec/2020-resident-population-</u>
 map.html. Published 2021. Accessed June 30, 2021.
- 4033.Bergquist S, Otten T, Sarich N. COVID-19 pandemic in the United States. Health Policy404Technol. 2020;9(4):623-638.
- 405 4. Vynnycky E, White RG. *An Introduction to Infectious Disease Modelling*. Oxford
 406 University Press; 2010.
- Thompson RN, Stockwin JE, van Gaalen RD, et al. Improved inference of time-varying
 reproduction numbers during infectious disease outbreaks. *Epidemics*. 2019;29:100356.
- 409
 6. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-varying reproduction numbers during epidemics. *Am J Epidemiol.*411
 2013;178(9):1505-1512.
- 412 7. New York Times/covid-19 data. <u>https://github.com/nytimes/covid-19-data</u> Published
 413 2021. Accessed January 11, 2021.
- The United States Census Bureau. County population totals 2010-2019. United States
 Census Bureau. <u>https://www.census.gov/data/tables/time-series/demo/popest/2010s-</u>
 counties-total.html. Published 2019. Accessed January 11, 2021.
- 417 9. Richmond HL, Tome J, Rochani H, Fung IC, Shah GH, Schwind JS. The Use of
 418 Penalized Regression Analysis to Identify County-Level Demographic and
 419 Socioeconomic Variables Predictive of Increased COVID-19 Cumulative Case Rates in
- 420 the State of Georgia. *Int J Environ Res Public Health*. 2020;17(21):8036.
- 421 10. Gostic KM, McGough L, Baskerville EB, et al. Practical considerations for measuring the
 422 effective reproductive number, Rt. *PLoS Comput Biol.* 2020;16(12):e1008409.
- 11. Centers for Disease Control and Prevention. COVID-19 Pandemic Planning Scenarios.
 https://www.cdc.gov/coronavirus/2019-ncov/hcp/planning-scenarios.html. Published
 2020. Accessed November 12, 2020.
- 426 12. You C, Deng Y, Hu W, et al. Estimation of the time-varying reproduction number of
 427 COVID-19 outbreak in China. *Int J Hyg Environ Health.* 2020;228:113555.
- 428 13. Chowell G, Bettencourt LM, Johnson N, Alonso WJ, Viboud C. The 1918-1919
 429 influenza pandemic in England and Wales: spatial patterns in transmissibility and 430 mortality impact. *Proc Biol Sci.* 2008;275(1634):501-509.
- 431 14. Fung IC-H, Zhou X, Cheung C-N, et al. Assessing Early Heterogeneity in Doubling
 432 Times of the COVID-19 Epidemic across Prefectures in Mainland China, January–
 433 February, 2020. *Epidemiologia*. 2021;2(1):95-113.

15.	Ogwara CA, Mallhi AK, Hua X, et al. Spatially refined time-varying reproduction numbers of COVID-19 by health district in Georgia, USA, March - December, 2020.
	<i>Epidemiologia</i> . 2021;2(2):179-197.
16.	Politis MD, Hua X, Ogwara CA, et al. Spatially refined time-varying reproduction
	numbers of SARS-CoV-2 in Arkansas and Kentucky, March - November, 2020.
	Unpublished Work. In:2021.
17.	Agency for Toxic Substances and Disease Registry. CDC Social Vulnerability Index.
	https://www.atsdr.cdc.gov/placeandhealth/svi/index.html. Published 2020. Accessed
	March 27, 2021.
18.	Agency for Toxic Substances and Disease Registry. CDC SVI Data and Documentation
	Download.
	https://www.atsdr.cdc.gov/placeandhealth/svi/data_documentation_download.html.
	Published 2018. Accessed April 11, 2021.
19.	Carter RJ, Rose D, Sabo R, et al. Widespread SARS-CoV-2 Transmission Among
	Attendees at a Large Motorcycle Rally and their Contacts, 30 US Jurisdictions, August-
	September, 2020. Clin Infect Dis. 2021.
20.	Dave D, McNichols D, Sabia JJ. The contagion externality of a superspreading event:
	The Sturgis Motorcycle Rally and COVID-19. South Econ J. 2020.
21.	Muniz-Rodriguez K, Chowell G, Schwind JS, et al. Time-varying Reproduction Numbers
	of COVID-19 in Georgia, USA, March 2, 2020 to November 20, 2020. The Permanente
	Journal 2021;25(20):232.
22.	Wang Y, Siesel C, Chen Y, et al. Transmission of COVID-19 in the state of Georgia,
	United States: Spatiotemporal variation and impact of social distancing. In.
	MedRxiv2020.
23.	Leung K, Wu JT, Leung GM. Real-time tracking and prediction of COVID-19 infection
	using digital proxies of population mobility and mixing. Nat Commun. 2021;12(1):1501.
24.	Herstein JJ, Degarege A, Stover D, et al. Characteristics of SARS-CoV-2 Transmission
	among Meat Processing Workers in Nebraska, USA, and Effectiveness of Risk
	Mitigation Measures. Emerg Infect Dis. 2021;27(4):1032-1038.
25.	Chu DK, Akl EA, Duda S, et al. Physical distancing, face masks, and eye protection to
	prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic
	review and meta-analysis. <i>Lancet</i> . 2020;395(10242):1973-1987.
26.	Liang M, Gao L, Cheng C, et al. Efficacy of face mask in preventing respiratory virus
	transmission: A systematic review and meta-analysis. <i>Travel Med Infect Dis.</i>
	2020;36:101751.
	Fricchione MJ, Seo JY, Arwady MA. Data-Driven Reopening of Urban Public Education
	Through Chicago's Tracking of COVID-19 School Transmission. J Public Health Manag
	Pract. 2021;27(3):229-232.
28.	Gillespie DL, Meyers LA, Lachmann M, Redd SC, Zenilman JM. The Experience of 2
-	Independent Schools With In-Person Learning During the COVID-19 Pandemic. J Sch
	Health. 2021;91(5):347-355.
	Zimmerman KO, Akinboyo IC, Brookhart MA, et al. Incidence and Secondary
-	Transmission of SARS-CoV-2 Infections in Schools. <i>Pediatrics</i> .
	2021;147(4):e2020048090.
	 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28.

- 478 30. McGuine TA, Biese KM, Petrovska L, et al. Mental Health, Physical Activity, and 479 Quality of Life of US Adolescent Athletes During COVID-19-Related School Closures 480 and Sport Cancellations: A Study of 13 000 Athletes. J Athl Train. 2020;56(1):9. 481 31. Wong DWS, Li Y. Spreading of COVID-19: Density matters. PLoS One. 482 2020;15(12):e0242398. 483 32. McLaughlin JM, Khan F, Pugh S, et al. County-Level Predictors of COVID-19 Cases and 484 Deaths in the United States: What Happened, and Where Do We Go from Here? Clin 485 Infect Dis. 2020;XX(XX):ciaa1729. 486 Nivette A, Ribeaud D, Murray A, et al. Non-compliance with COVID-19-related public 33. 487 health measures among young adults in Switzerland: Insights from a longitudinal cohort 488 study. Soc Sci Med. 2021;268:113370. 489 34. Crane MA, Shermock KM, Omer SB, Romley JA. Change in Reported Adherence to 490 Nonpharmaceutical Interventions During the COVID-19 Pandemic, April-November 491 2020. JAMA. 2021;325(9):883-885. 492 35. Centers for Disease Control and Prevention. Guidance for Fully Vaccinated People. 493 https://www.cdc.gov/coronavirus/2019-ncov/vaccines/fully-vaccinated-guidance.html. 494 Published 2021. Accessed May 19, 2021. 495 36. Rabin RC. A major nurses' union condemns the C.D.C.'s new mask advice for 496 vaccinated people. New York Times. 497 https://www.nytimes.com/2021/05/15/world/nurses-union-cdc-mask-vaccinated.html. 498 Published 2021. Accessed May 15, 2021. 499 500 501
- 501

12

State		ate (2020)		Percentage Cha		5% CrI
			7-Da	y Sliding Window	Non-O	verlapping Window
North Dakota	16-Mar 30-Mar 26-May 7-Sep	Policy A Policy B Policy C School Open**	8.86% -10.96% 6.98% -8.03%	[-22.86%, 52.67%] [-15.3%, -10.46%] [-1.61%, 16.12%] [-8.23%, -7.83%]	-18.01% -21.43% 4.30%	[-39.68%, 10.12%] [-32.18%, -8.84%] [0.47%, 8.19%]
South Dakota	21-Sep 13-Nov 21-Dec 13-Mar	Policy D Policy E Policy F Policy A	0.17% -14.71% 32.44% 94.16%	[0.01%, 0.33%] [-14.99%, -14.41%] [26.53%, 38.09%] [-9.05%, 404.87%]	0.82% -27.27% 20.85% 89.99%	[-0.82, 2.49%] [-28.24%, -26.09%] [16.86%, 25.10%] [7.91%, 285.91%]
	6-Apr 28-Apr 13-Aug 7-Sep 25-Sep	Policy B Policy C Policy D School Open** Policy E	-38.85% 37.34% 16.47% 12.36% 18.84%	[-43.80%, -33.50%] [24.20%, 51.22%] [7.06%, 26.37%] [8.07%, 16.72%] [18.19%, 19.48%]	-42.62% 9.64% 7.59% -10.38%	[-47.18%, -38.20%] [7.00%, 12.28%] [7.09%, 8.08%] [-12.00%, -8.86%]
Idaho	25-Mar 1-May 13-Jun 7-Sep	Average* Policy A Policy B Policy C School Open**	2.80% -59.94% 11.14% 8.68% 13.14%	[-7.85%, 34.52%] [-60.57%, -59.31%] [10.38%, 11.98%] [-0.84%, 19.05%] [12.52%, 13.73%]	-62.67% 41.97% -5.02%	[-65.75%, -59.49%] [35.64%, 48.05%] [-9.42%, -0.83%]
	27-Oct 14-Nov	Policy D Policy E	2.60% -1.93%	[2.27%, 2.91%] [-1.99%, -1.87%]	-2.67% -6.24%	[-4.09%, -1.24%] [-7.64%, -4.83%]
Montana	13-Mar 7-May 15-Jul 1-Sep 7-Sep	Policy A Policy B Policy C Policy D School Open**	-3.99% -14.92% -12.43% 34.83% 20.73%	[-38.99%, 49.24%] [-65.10%, 107.88%] [-12.55%, -12.31%] [33.82%, 35.85%] [10.90%, 30.98%]	-51.24% 22.60% -16.17% 8.52%	[-66.21%, -27.83%] [10.14%, 35.95%] [-17.27%, -15.08%] [5.54%, 11.28%]
Wyoming	18-Nov 13-Mar 28-Apr 10-Jun 13-Jul 7-Sep	Policy E Policy A Policy B Policy C Policy D School Open**	-9.63% 15.46% -29.99% 1.17% 0.30% 19.56%	[-9.98%, -9.26%] [-29.23%, 91.29%] [-46.75%, -6.23%] [-1.03%, 3.64%] [-0.76%, 1.43%] [1.66%, 39.27%]	-17.58% -32.95% -5.13% 5.69% 1.96%	[-18.75%, -16.38%] [-54.24%, 2.84%] [-5.79%, -4.48%] [-4.57%, 16.99%] [-2.99%, 6.41%]
		Policy E 3 Average*	-1.98% -13.22%	[-2.64%, -1.33%] [-29.76%, 21.92%]	-8.84%	[-11.03%, -6.78%]
	7-Dec	Policy F	10.58%	[10.34%, 10.83%]	-13.25%	[-15.30%, -10.97%]

Table 1: Percentage change of R_t and 95% credible intervals (CrI) at policy implementation and important dates calculated with both 7-day sliding window and non-overlapping window.

*Note: neither South Dakota nor Wyoming had any policy change during this time-period of interest so bootstrapping was performed on the average of the 6-day range of policy implementation from the other three states. **Note: As of September 7th, 2020, all states had resumed face-to-face K-12 education.

Policy labels: North Dakota: A = Schools closed; B = Promote physical distancing; C = Testing order; D = Lifted travel and quarantine order; E = Mask order; F = Allow all restaurants and bars to resume normal hours. South Dakota: A = State of Emergency declared; B = Minnehaha and Lincoln stay at home order; C = All citizens shall implement 'back to normal plan'; D = All K-12 schools were mandated to start in-person classes again; E = long-term facilities began to relax visitor restrictions. Idaho: A = Statewide stay home order; B = Stay Healthy Order-Stage 1. Businesses and governmental agencies resumed operations at physical locations; C = Stay Health Order-Stage 4. Gatherings of any size were allowed and non-essential travel could resume; D = Stay Health Order-Stage 3. Patrons of bars, nightclubs, and restaurants must remain seated. Face coverings were required at long-term care facilities; E = Stay Health Order-Stage 2. Gatherings of no more than 10 people, and vulnerable population were strongly recommended to stay at home. Montana: A = State of Emergency declared; B = School reopened; C = Statewide mask requirement; D = School's Fall semester started; E = Additional mitigation measures were implemented including reduced size of gatherings to no more than 25. Wyoming: A = All businesses and schools were closed; B = Gymnasiums and childcare facilities were reopened; C = K-12 schools, colleges, universities, and trade schools resumed on-site instructions; D = Removed some conditions and restrictions were applicable to restaurants; E = Additional mitigation measures were implemented including reduced size of gatherings to no more than 50; F =Required face coverings in certain places with exceptions.

Table 2. The linear regression analysis between log₁₀-transformed per capita weekly incident case count and log₁₀-transformed population size for North Dakota, South Dakota, Idaho, Montana, and Wyoming, March 2020 -January 2021.*

	Unadju	usted Model	Adjuste	d for Weeks	Adjusted fo	r all Variables†
	Parameter Estimate	(95% CI)	Parameter Estimate	(95% CI)	Parameter Estimate	(95% CI)
North Dakota	0.066	(-0.009,0.141)	0.190	(0.148, 0.233)*	0.195	(0.131, 0.259)*
South Dakota	-0.018	(-0.092,0.056)	0.143	(0.099, 0.187)*	0.127	(0.069, 0.185)*
Idaho	0.005	(-0.071,0.080)	0.155	(0.108, 0.201)*	0.106	(0.049, 0.164)*
Montana	-0.222	(-0.299, -0.145)*	0.061	(0.017, 0.105)*	0.029	(-0.021, 0.079)
Wyoming	0.212	(0.054, 0.369)*	0.271	(0.182, 0.360)*	0.153	(-0.007, 0.313)

*p<0.05 †Note: Adjusted for below poverty variable, the percentage of people without high school diploma, the percentage of minority, the percentage of people with a disability, the percentage of crowding, the percentage of people 65 years and older, and for weeks.

504

505

506

507 Figure Titles and Legends

508

509 Figure 1. County-level population map (2019) of North Dakota, South Dakota, Idaho, Montana,510 and Wyoming, USA.

511

512 **Figure 2.** County-level maps of North Dakota, South Dakota, Idaho, Montana, and Wyoming by 513 cumulative case count (top four maps), and cumulative case counts per 1000 population (bottom

four maps) on June 30, August 31, October 31, and December 31, 2020 (date of report).

515

Figure 3. The daily number of new cases by their assumed date of infection (left panel), 1-week sliding window R_t estimates (middle panel), and non-overlapping window R_t estimates by policy change (right panel) of North Dakota, South Dakota, Idaho, Montana, and Wyoming, USA,

519 March 10, 2020 – January 10, 2021 (date of report).

520 North Dakota: A = Schools closed; B = Promote physical distancing; C = Testing order; D =

521 Lifted travel and quarantine order; E = Mask order; F = Allow all restaurants and bars to resume

522 normal hours. **South Dakota**: A = State of Emergency declared; B = Minnehaha and Lincoln

523 stay at home order; C = All citizens shall implement 'back to normal plan'; D = All K-12 schools

524 were mandated to start in-person classes again; E = long-term facilities began to relax visitor

525 restrictions. Idaho: A = Statewide stay home order; B = Stay Healthy Order-Stage 1. Businesses

and governmental agencies resumed operations at physical locations; C = Stay Health Order-

527 Stage 4. Gatherings of any size were allowed and non-essential travel could resume; D = Stay

528 Health Order-Stage 3. Patrons of bars, nightclubs, and restaurants must remain seated. Face

529 coverings were required at long-term care facilities; E = Stay Health Order-Stage 2. Gatherings

530 of no more than 10 people, and vulnerable population were strongly recommended to stay at 531 home. **Montana**: A = State of Emergency declared; B = School reopened; C = Statewide mask

requirement; D = School's Fall semester started; E = Additional mitigation measures were

implemented including reduced size of gatherings to no more than 25. Wyoming: A = All

businesses and schools were closed; B = Gymnasiums and childcare facilities were reopened; C

535 = K-12 schools, colleges, universities, and trade schools resumed on-site instructions; D =

536 Removed some conditions and restrictions were applicable to restaurants; E = Additional

537 mitigation measures were implemented including reduced size of gatherings to no more than 50;

538 F = Required face coverings in certain places with exceptions.


539

540 **Figure 4.** Linear regression between log₁₀-transformed per capita cumulative case number (ccn)


and log₁₀-transformed population size by county (grouped in states) for North Dakota, South


542 Dakota, Idaho, Montana, and Wyoming on June 30th, August 31st, October 31st, and December


543 31^{st} , 2020 (date of report).



Longitude















50.0 -











Supplementary material: Late surges in COVID-19 cases and varying transmission potential partially due to public health policy changes in 5 Western states, March 10, 2020-January 10, 2021

Xinyi Hua[†], MPH; Aubrey R. D. Kehoe[†], MPH; Joana Tome[†], MPH; Mina Motaghi[†], MD, MPH; Sylvia K. Ofori, MPH; Po-Ying Lai, MS; Sheikh Taslim Ali, PhD; Gerardo Chowell, PhD; Anne C. Spaulding, MD, MPH; Isaac Chun-Hai Fung^{*}, PhD.

Affiliations:

- 1. Department of Biostatistics, Epidemiology and Environmental Health Sciences, Jiann-Ping Hsu College of Public Health, Georgia Southern University, Statesboro, GA, USA (X. Hua, A. R. D. Kehoe, J. Tome, M. Motaghi, S. K. Ofori, I.C.H. Fung)
- 2. Department of Biostatistics, School of Public Health, Boston University, Boston, MA, USA (P.Y. Lai)
- 3. WHO Collaborating Centre for Infectious Disease Epidemiology and Control, Division of Epidemiology and Biostatistics, School of Public Health, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region, China (S.T. Ali)
- 4. Laboratory of Data Discovery for Health, Hong Kong Science and Technology Park, Hong Kong Special Administrative Region, China (S.T. Ali)
- 5. Department of Population Health Sciences, School of Public Health, Georgia State University, Atlanta, GA, USA (G. Chowell)
- 6. Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA, USA (A.C. Spaulding)

*Corresponding author: Isaac Chun-Hai Fung, PhD. Email: cfung@georgiasouthern.edu

† Xinyi Hua, Aubrey R. D. Kehoe, Joana Tome, and Mina Motaghi serve as co-first authors.

Appendix A. Data cleaning: handling negative incident case counts

Dates with negative case counts were identified (i.e., when public health agencies made corrections to their cumulative case counts at a specific date or dates). Daily case counts were adjusted to remove negative counts that were created when states adjusted their positive cases; for instance, a state would record negative cases in order to account for a previous over-count. To remove these negative counts all data was reviewed, and negative counts were transformed into a zero count by removing this number from the previous days with positive cases until there was no discrepancy. We adjusted the negative daily case counts at both state and county levels in all five states in this study.

Appendix B. Assessing power-law relationship using linear regression of log-transformed per capita cumulative case count and log-transformed population size.

If we assume that a power-law relationship exists between the cumulative case count (C) and the population size (N), with an exponent g, i.e., $C = N^{A}g$, and that per capita cumulative case count, A = C/N, then,

$$g = \frac{\log C}{\log N} = \frac{\log (AN)}{\log N} = \frac{\log A}{\log N} + \frac{\log N}{\log N} = \frac{\log A}{\log N} + 1$$

Thus,

$$\frac{\log A}{\log N} = g - 1 = m$$

Where *m* is the slope of the regression line between the log_{10} -transformed per capita cumulative case count and the log_{10} -transformed population size.

Appendix C. The detailed description for R_t in North Dakota, South Dakota, Idaho, Montana, and Wyoming.

North Dakota. The daily reported cases of COVID-19 in North Dakota started to increase since July 2020, reaching a peak of 2,270 new cases in early November and 2,340 new cases in early December (Figure 3). The 7-day-sliding-window R_t stayed above 1 almost all year around and went below 1 in late May and in December. The governor's executive orders and State Health Officers' orders (Supplemental Table 2) throughout 2020 were chosen to create the policy change R_t throughout the year (Figure 3). From March 16, 2020 till April 4, 2020, the 7-day-sliding-window R_t decreased toward 1 but was still above 1 meaning that the promotion of physical distance was working. On April 18, 2020, the State Health Officer released the quarantine order for the Grand Forks county, and subsequently on May 14, 2020, they issued the testing order for Grand Forks county. In late May, the 7-day-sliding-window R_t fell below 1 where the coverage of the testing order expanded further. From June till November both the 7-day-sliding-window R_t and policy change R_t maintained a level above 1 which corresponded to the lifted quarantine order, the rescinded close contact quarantine order, and the permission of health care workers who are asymptomatically infected with SARS-CoV-2 to work in care facilities. From mid-November, the 7-day-sliding-window R_t and policy change R_t fell below 1, which corresponded to the release of the first mask order, and the allowance of team practice, bars and restaurant to resume normal hours.

South Dakota. From March 1, 2020 through January 1, 2021 South Dakota had 103,318 confirmed COVID-19 cases cumulatively. The daily reported cases of South Dakota had one late surge, peaking in November 2020 (Figure 3). Within South Dakota, there were eight policies or recommendations for COVID-19 precautions were implemented, five major ones can be found in Supplemental Table 2. The 7-day-sliding window R_t estimate fell below 1 on April 4, 2020 following the closure of the Smithfield pork processing plant in Sioux Falls, Minnehaha, South Dakota.¹ The policy change R_t estimate remained below 1 until April 28, 2020 when South Dakota implemented their *Back to Normal Plan* which largely focuses on good hygiene practices and limiting persons in large numbers. From August 13 till September 25, 2020 the policy change R_t estimates increased to above 1 when South Dakota K-12 school reopened in person. The 95% credible intervals (CrI) of 7-day-sliding window R_t estimates in March 2020 were wide due to the small numbers of daily new cases. Meanwhile, during the periods between April 28, 2020 and August 13, 2020 and between September 25, 2020 and January 10, 2021 there were very few policy changes; the policy change R_t estimates may not accurately depict the true R_t estimates during those periods.

Idaho. From March 1, 2020 to January 1, 2021, Idaho's daily reported case number revealed two major surges of new cases in July and November 2020 (Figure 3). The policy change R_t estimates based on statewide policy changes demonstrate the effectiveness of the non-pharmaceutical interventions (face masking, social distancing, prohibited gathering of more than 10 people, etc.) of controlling the spread of COVID-19 within the state of Idaho. The policy change R_t estimates dropped below 1 after the Statewide Stay Home order was in place between March 25, 2020 and April 30, 2020. The statewide policy change R_t estimates increased above 1 between May 1, 2020 and October 26, 2020, with the Stay Healthy Order relaxation of the controlling measures from stage 1 (encourage vulnerable individuals to stay at home,

avoid public and private gathering, and avoid and minimize non-essential travels) to stage 4 (open all business, allow public gathering with any size and resume non-essential travel). With the Stay Healthy Order stage 3 in effect on October 27, 2020, the policy change R_t estimates dropped again and continued to drop below one, after the Stay Healthy Order moved from stage 3 back to stage 2 with more restrictions on November 14, 2020. Overall, Idaho demonstrated an extensive community transmission of SARS-CoV-2 transmission in the study period.

Montana. The number of incident cases in Montana had a surge in November 2020 with up to 1,622 daily cases in mid-November (Figure 3). The 7-day-sliding-window R_t curve and the policy change R_t curve show some significant changes over time parallel with implementing different COVID-19 related policies (Figure 3). There was a drastic drop in the 7-day-sliding-window R_t estimates after closing businesses and the *stay-at-home* orders at the end of March, leading to R_t estimates under 1 from late March until early May. An increase in 7-day-sliding-window R_t was observed in the first days of May as the state lifted the *stay-at-home* order, reopened schools and started the first phase of reopening businesses (such as some retails, restaurants, casinos, and bars). There was a decrease in policy change R_t estimates in mid-July after the statewide mask requirement was applied, with a slight rise in September, probably due to the start of schools' Fall semester. In November after applying new restrictions on gatherings and also placing time and capacity limitations for bars and restaurants, the policy change R_t estimates dropped again below 1 and remained low, mostly till late December.

Wyoming. From March 1, 2020 to January 1, 2021, Wyoming's daily reported case number revealed one major surge of new cases in October and November 2020 (Figure 3). The policy change R_t estimates based on statewide policy changes demonstrated the necessity of the non-pharmaceutical interventions (face masking, social distancing, reduction in the size of public gathering, etc.) of controlling the spreading of COVID-19 within the state of Wyoming. For instance, the R_t estimates stayed around and above 1 while the state opened up all businesses without face mask mandate and public gathering restrictions. The 7-day-sliding-window R_t estimates then dropped below 1 after the Seventeenth Continuation and Modification Statewide Public Health Order regarding the public gathering with no more than 25 people on November 19, 2020. With the first public health order regarding mandatory face covering in effect on December 7, 2020, the policy change R_t estimates maintained below one. Overall, Wyoming demonstrated an extensive community transmission of SARS-CoV-2 transmission in the study period.

Appendix D: Detailed *R_t* description for North Dakota, South Dakota, Idaho, Montana, and Wyoming regarding School reopening.

North Dakota. North Dakota decided to reopen schools and universities on June 1st, 2020.² There are around 21 universities in the state located in Cass, Grand Forks, Ward, Burleigh, Stark, Stutsman, Trail, Mountrail, Bottineau, Benson, Barnes, Pierce, Dickey, Sioux, and Rolette counties.³ During the year of 2020, in most of the counties, the case number starts to increase at the end of the summer, peaking in November. Concretely, in Ward, Burleigh, Stark, Trail, and Sioux counties, the reported cases started to increase in July; in Cass, Grand Forks, Stutsman, Barnes counties, the reported cases started to increase in August; in Mountrail, Bottineau, and Benson counties the reported case started to increase in September, while for the Pierce, Dickey, and Rolette counties, the daily reported case count started to increase in October, peaking in November. For most of the counties, the *R*_t stays at one during all the summer-fall period, except for some fluctuations (above and below one) in Grand Forks, and an *R*_t greater than one for Burleigh (where the state capital is located), and Dickey or Pierce (over one till the end of October). This description may be related to college students returning to school because the increase in the daily

reported case count corresponds to the start of the fall semester in North Dakota. However, after November, new cases infected with SARS-CoV-2 and R_t starts to fall, which correspond to the issued mask order.

South Dakota. Some South Dakota schools began to offer in-person classes in June, but most schools did not begin face-to-face instruction for the mid-August fall semester.⁴ The state is home to four technical institutes, six state universities, six tribal colleges and universities, and six private colleges and universities; of these, six located in Minnehaha county, three in Pennington county, two schools located in Davison and Brown counties respectively, and one in Lawrence, Lake, Codington, Yankton, Oglala Lakota, Todd, Roberts, Brookings, and Clay counties respectively.⁵ Following the mid-August openings nine of these counties had an R_t that remained above 1 for between 7-14 days and four had R_t values that fluctuated both below and above 1. However, it is important to note that many of these counties have a very small population with small daily incident case count, which resulted in wide CrIs for R_t estimates around this time period. This initial increases in R_t and case count might be related to the return of students to in-person learning. In most counties with colleges and universities the epidemiologic increases seen in mid-August transitioned into the overall state-level fall surge that peaked around November. It is also noteworthy that the Sturgis Motorcycle Rally was held in Meade County, South Dakota, August 7-16, 2020, which coincide with the school reopening on August 13, 2020. Therefore, it is difficult to attribute the increase in Rt to a single policy decision (schools reopening) or a single mass gathering (Sturgis Motorcycle Rally). Both could have been factors involved; please refer to the discussion in the Limitations section in the main text.

Idaho. Idaho schools opened on-site instruction for the fall 2020 semester with a framework on decision making based on four different levels of community transmission.⁶ The state is home for four public colleges and 4-year universities, four regional community colleges, and eight private colleges and universities. Of these, four located in Boise, two in Moscow, two in Nampa, two in Idaho Falls, and one in Rexburg, Caldwell, Meridian, Twin Falls, Pocatello, Lewiston, McCall, and Coeur d'Alene respectively.⁷ Following the mid-August openings, Boise County had an R_t fluctuated below and above 1 but mainly above 1; however, the daily case counts were very small, less than 15, therefore the CrI of R_t was wide. Boise County is a college town for four colleges and universities. These small daily case counts may relate to effective non-pharmaceutical interventions or issues of underreporting.


Montana. Montana was one of the 2 states that resumed in-person instructions in schools in the academic year 2019-2020, on May 7, 2020.⁸ As seen in Figure 3, R_t rose above one from early May and stayed high until mid-July which reflected the impact of this policy in the state, in contrast to most of the other states of the county. There are more than 20 colleges and universities in Montana, with approximately 40,000 students, in Gallatin county, Yellowstone county, Hill county, Missoula county, Silver Bow county, Beaverhead county, Dawson county, Flathead county, Custer county, Lewis and Clark county, Cascade county, and tribal colleges in the counties of Blaine, Glacier, Rosebud, Roosevelt, Big Horn, Lake and Hill.⁹ Fall semester 2020 started in the second half of August in almost all of these academic institutions with online and on campus courses, using new guidelines to improve safety of students. No significant change in R_t was observed in these counties after the Fall openings.

Wyoming. Wyoming K-12 schools, colleges and universities, and trade schools were reopened in limited on-site instructions on May 7, 2020, and later fully opened for face-to-face on-site instruction on June 10, 2020.¹⁰ The state is home for one 4-year university, eight public colleges, one tribal college, and two private colleges. Except for the University of Wyoming and Wyoming Technical Institute located in Laramie, the rest of the colleges scatter across different counties.¹¹ Following the reopening in June, Laramie County had an R_t spike of 4 and R_t fluctuated below and above 1 but mainly above 1 between June and December, and R_t dropped below 1 between December and January. The largest daily case count

was 246 on November 14, 2020. The trend of R_t and case count increase might be related to the school resumed face-to-face instruction in June and following fall semester.

Appendix E: Linear regression between log₁₀-transformed per capita cumulative case number (ccn) and log₁₀-transformed population size by county for North Dakota, South Dakota, Idaho, Montana, and Wyoming, on June 30th, August 31st, October 31st, and December 31st, 2020 (date of report) are presented in Figures S1-S5.

Figure S1. Linear regression between log₁₀-transformed per capita cumulative case number (ccn) and log₁₀-transformed population size by county for North Dakota (53 counties), on June 30th, August 31st, October 31st, and December 31st, 2020 (date of report). Counties that were outliers (defined as below 2.5 percentile or above 97.5 percentile of the distribution of the log₁₀-transformed per capita ccn) were marked in red.



North Dakota

Figure S2. Linear regression between log₁₀-transformed per capita cumulative case number (ccn) and log₁₀-transformed population size by county for South Dakota (66 counties), on June 30th, August 31st, October 31st, and December 31st, 2020 (date of report). Counties that were outliers (defined as below 2.5 percentile or above 97.5 percentile of the distribution of the log₁₀-transformed per capita ccn) were marked in red.




Figure S3. Linear regression between log₁₀-transformed per capita cumulative case number (ccn) and log₁₀-transformed population size by county for Idaho (44 counties), on June 30th, August 31st, October 31st, and December 31st, 2020 (date of report). Counties that were outliers (defined as below 2.5 percentile or above 97.5 percentile of the distribution of the log₁₀-transformed per capita ccn) were marked in red.




Figure S4. Linear regression between log₁₀-transformed per capita cumulative case number (ccn) and log₁₀-transformed population size by county for Montana (56 counties), on June 30th, August 31st, October 31st, and December 31st, 2020 (date of report). Counties that were outliers (defined as below 2.5 percentile or above 97.5 percentile of the distribution of the log₁₀-transformed per capita ccn) were marked in red.





Figure S5. Linear regression between log₁₀-transformed per capita cumulative case number (ccn) and log₁₀-transformed population size by county for Wyoming (23 counties), on June 30th, August 31st, October 31st, and December 31st, 2020 (date of report). Counties that were outliers (defined as below 2.5 percentile or above 97.5 percentile of the distribution of the log₁₀-transformed per capita ccn) were marked in red.



Supplemental Table 1. List of counties in North Dakota, South Dakota, Idaho, Montana, and Wyoming (organized by state).¹²

State	Counties
North Dakota (53 counties)	Adams, Barnes, Benson, Billings, Bottineau, Bowman, Burke, Burleigh, Cass, Cavalier, Dickey, Divide, Dunn, Eddy, Emmons, Foster, Golden Valley, Grand Forks, Grant, Griggs, Hettinger, Kidder, LaMoure, Logan, McHenry, McIntosh, McKenzie, McLean, Mercer, Morton, Mountrail, Nelson, Oliver, Pembina, Pierce, Ramsey, Ransom, Renville, Richland, Rolette, Sargent, Sheridan, Sioux, Slope, Stark, Steele. Stutsman, Towner, Traill, Walsh, Ward, Wells, Willams.
South Dakota (66 counties)	 Aurora, Beadle, Bennett, Bon Homme, Brookings, Brown, Brule, Buffalo, Butte, Campbell, Charles Mix, Clark, Clay, Codington, Corson, Custer, Davison, Day, Deuel, Dewey, Douglas, Edmunds, Fall River, Faulk, Grant, Gregory, Haakon, Hamlin, Hand, Hanson, Harding, Hughes, Hutchinson, Hyde, Jackson, Jerauld, Jones, Kingsbury, Lake, Lawrence, Lincoln, Lyman, McCook, McPherson, Marshall, Meade, Mellette, Miner, Minnehaha, Moody, Oglala Lokota, Pennington, Perkins, Potter, Roberts, Sanborn, Spink, Stanley, Sully, Todd, Tripp, Turner, Union, Walworth, Yankton, Ziebach.
Idaho (44 counties)	Ada, Adams, Bannock, Bear Lake, Benewah, Bingham, Blaine, Boise, Bonner, Bonneville, Boundary, Butte, Camas, Canyon, Caribou, Cassia, Clark, Clearwater, Custer, Elmore, Franklin, Fremont, Gem, Gooding, Idaho, Jefferson, Jerome, Kootenai, Latah, Lemhi, Lewis, Lincoln, Madison, Minidoka, Nez Perce, Oneida, Owyhee, Payette, Power, Shoshone, Teton, Twin Falls, Valley, Washington.
Montana (56 counties)	Beaverhead, Big Horn, Blaine, Broadwater, Carbon, Carter, Cascade, Chouteau, Custer, Daniels, Dawson, Dear Lodge, Fallon, Fergus, Flathead, Gallatin, Garfield, Glacier, Golden Valley, Granite, Hill, Jefferson, Judith Basin, Lake, Lewis and Clark, Liberty, Lincoln, McCone, Madison, Meagher, Mineral, Missoula, Musselshell, Park, Petroleum, Phillips, Pondera, Powder River, Powell, Prairie, Ravalli, Richland, Roosevelt, Rosebud, Sanders, Sheridan, Silver Bow, Stillwater, Sweet Grass, Teton, Toole, Treasure, Valley, Wheatland, Wibaux, Yellowstone.
Wyoming (23 counties)	Albany, Big Horn, Campbell, Carbon, Converse, Crook, Fremont, Goshen, Hot Springs, Johnson, Laramie, Lincoln, Natrona, Niobrara, Park, Platte, Sheridan, Sublette, Sweetwater, Teton, Uinta, Washakie, Weston.

Supplemental Table 2: Control measures implemented by state government agencies in North Dakota, South Dakota, Idaho, Montana, and Wyoming, USA

	Label	Date (mm/dd/yyyy)	Implemented measure(s)
North Dakota	А	03/16/2020	Executive Order 2020-04.1. The governor amended school closure order for certain schools and programs. ¹³
	В	03/30/2020	Executive Order 2020-10.1. The governor enabled remote participation for public meetings to promote physical distancing. He amended school facilities closure to accommodate child care needs and suspended deadlines for livestock auctions, pesticide applicators, and public libraries. ¹⁴
	С	05/26/2020	Testing Order Long Term Care. The State Health Officer Order for testing for 2019- nCoV/COVID-19 as a Disease Control Measures to prevent the spread 2019- nCoV/COVID-19. ¹⁵
	D	09/21/2020	Order #2020-02.4 Lifted Travel Quarantine Order. The State Health Officer Order lifted the 14-day quarantine order for those returning from international travel. ¹⁶
	E	11/13/2020	Order #2020-08 Mask Order. The mask order required face coverings in indoor businesses and indoor public settings, as well as in outdoor business and public settings when it was not possible to maintain physical distancing. ¹⁷ Executive Order 2020-43. The governor enacted mitigation measures for food service, events, sports, and extracurricular activities. ¹⁸
	F	12/21/2020	The governor amended the order to allow restaurants and bars to resume normal hours. ¹⁹
South Dakota	А	3/13/2020	State of emergency declared, executive state government workers given work from home order and restricted out of state travel (EO# 2020-05); K-12 schools closed for the rest of the year; higher education encouraged to close campus; and long-term care facilities implemented visitor restrictions. ²⁰
	В	4/6/2020	All residents shall review and practice CDC COVID-19 guidelines; all employers, local, and municipal government shall implement CDC COVID-19 guidelines to encourage telework, social distancing, and reduced numbers; all healthcare workers shall prepare for increased COVID-19 cases (EO# 2020-12) ²¹ ; stay-at-home order except for essential workers and essential outings for all Minnehaha and Lincoln "vulnerable individuals" (EO# 2020-13). ²²
	С	4/28/2020	All citizens shall implement "back to normal plan" leaving COVID-19 precautions up to each resident, employers, and schools; ended Minnehaha and Lincoln "vulnerable individuals" stay-at-home order; (EO# 2020-20). ²³
	D	8/13/2020	All K-12 schools were mandated by the South Dakota Governor to start in-person classes again. ²⁴
	Е	9/25/2020	Long-term care facilities began to relax visitor restrictions. ²⁴
Idaho	A	3/25/2020	Statewide stay-home order. This isolation order required Idaho residents to stay and work from home as much as possible while ensuring all essential services and business remain available. This isolation order did not prohibit outdoor activity such as walking, hiking running, and biking, but a safe distance of six (6) feet must be kept between those who do not live in the same household. ²⁵
	В	5/1/2020	Stay Healthy Order-Stage 1: Businesses and governmental agencies resumed operations at physical locations in the state of Idaho except for those businesses identified in this order. All businesses must adhere to the social distancing and sanitation requirement. Certain individuals entering Idaho required self-quarantine for 14 days. Gathering, both public and private, should be avoided. Non-essential travel should be avoided or minimized. ²⁶
	С	6/13/2020	Stay Healthy Guidelines-Stage 4: Gathering of any sizes were allowed but should adhere to the physical distancing and sanitation requirements. Non-essential travel could resume. ²⁷
	D	10/27/2020	Stay Health Order-Stage 3: Indoor and outdoor gathering, both public and private must adhere to the requirement identified in this order. Patrons of bars, nightclubs, and restaurants must remain seated. Face coverings were required at long-term care facilities. ²⁸

	E	11/14/2020	Stay Health Order-Stage 2: Gathering of more than 10 people, both public and private, were prohibited, including attendance at extracurricular activities such as sporting events. People at increased risk for severe illness living in the state of Idaho were strongly encouraged to stay home and limit their movement outside of their place of residence. ²⁹
Montana ³⁰	A	3/13/2020	State of Emergency declared and directive implementing Executive Orders 2-2020 and 3-2020 and providing for measures to combat the spread of COVID-19 Novel Coronavirus. Court closure.
	В	5/7/2020	School reopened.
	С	7/15/2020	Statewide mask requirement required individuals to wear masks inside certain businesses and at outdoor gatherings of greater than 50 people where social distancing was not possible.
	D	9/1/2020	Schools' Fall semester began.
	Е	11/18/2020	Additional mitigation measures. Gatherings were limited to 25 people when social distancing is not possible. Bars, restaurants, and casinos will have a 10 p.m. curfew every night and were limited to 50% capacity.
Wyoming	А	3/13/2020	Statewide Public Health Order #1. Closed all restaurants, bars, theaters, gymnasiums, childcare facilities, K-12 schools, colleges, universities, and trade schools, in the State of Wyoming, with certain exceptions. This Order was effective immediately and shall remain in effect until April 3, 2020. ³¹
	В	4/28/2020	Statewide Public Health Order, 3rd Continuation and Modification. This order authorized gymnasiums and childcare facilities to reopen under certain conditions. This order was effective on May 1, 2020 and shall remain in effect through May 15, 2020. Forbade gathering for 10 people or more. ^{32,33}
	С	6/10/2020	Statewide Public Health Order, 6th Continuation and Modification. This order authorized K-12 schools, colleges, universities, and trade schools to provide on-site instruction to students and allow others to use of their facilities under certain conditions. Forbade gathering for 50 people or more. ^{10,34}
	D	7/13/2020	Statewide Public Health Order, 8th Continuation and Modification. This order removed some conditions and restrictions that were applicable to restaurants. ^{35,36}
	E	10/14/2020	Statewide Public Health Order, 13th Continuation and Modification. This order required patrons at different tables (at restaurants/bars) to be seated at least 6 feet apart and did not apply to booths, and patrons could be in groups up to 8 instead of 6. This order also authorized groups of attendees seated or standing together during an event in groups of 8 instead of 6. Forbade gathering for 50 people or more. ^{37,38}
	F	12/7/2020	Statewide Public Health Order #4. Required face coverings in certain places, with exceptions. This order was effective on December 9, 2020, and shall remain in effect through January 8, 2021. ³⁹

Supplemental Table 3. The slope (and 95% Confidence Intervals) of the regression line between log₁₀-transformed per capita cumulative case count and log₁₀-transformed population size, by states, North Dakota, South Dakota, Idaho, Montana, and Wyoming, USA, on June 30, August 31, October 31, and December 31, 2020.

	June 30, 2020	August 31, 2020	October 31, 2020	December 31, 2020
North Dakota	0.2758	0.2171	0.0729	0.0986
	(0.0961, 0.4554)	(0.0846, 0.3496)	(0.0014, 0.1443)	(0.0366, 0.1605)
South Dakota	0.1603	0.2322	0.0052	0.0186
	(-0.0766, 0.3972)	(0.1018, 0.3627)	(-0.0794, 0.0898)	(-0.0402, 0.0774)
Idaho	0.2398	0.1857	0.0662	0.0798
	(-0.1067, 0.5863)	(0.0261, 0.3497)	(-0.0498, 0.1822)	(0.0150, 0.1446)
Montana	-0.1423	0.1741	0.0129	0.0359
	(-0.3975, 0.1130)	(-0.0028, 0.3510)	(-0.1170, 0.1428)	(-0.0397, 0.1116)
Wyoming	0.2765	0.3043	0.2210	0.1206
	(-0.1647, 0.7178)	(-0.0544, 0.6630)	(0.0659, 0.3761)	(0.0274, 0.2138)

Supplemental Table 4. The adjusted linear regression model between log₁₀-transformed per capita incident case count and log₁₀-transformed population size for Idaho, March 2020 -January 2021*

Variables	Parameter Estimate	Standard Error	t-Statistic	P-value
Population	0.106	0.030	3.606	0.0003
Below poverty estimate percentage	0.007	0.003	2.252	0.0244
No high school diploma percentage	0.005	0.006	0.827	0.4086
Age 65+ percentage	-0.014	0.005	-2.780	0.0055
Population with disability percentage	0.009	0.005	1.873	0.0613
Minority percentage	0.010	0.003	3.039	0.0024
Crowding percentage	-0.030	0.010	-2.921	0.0035
*Note: adjusted for weeks				

Supplemental Table 5. The adjusted linear regression model between log₁₀-transformed per capita incident case count and log₁₀-transformed population size for Montana, March 2020 - January 2021*

Variables	Parameter Estimate	Standard Error	t-Statistic	P-value
Population	0.0290	0.026	1.136	0.2561
Below poverty estimate percentage	-0.017	0.004	-4.401	0.0000
No high school diploma percentage	0.008	0.004	2.196	0.0282
Age 65+ percentage	-0.001	0.004	-0.304	0.7614
Population with disability percentage	-0.007	0.005	-1.373	0.1698
Minority percentage	0.008	0.002	4.703	0.0000
Crowding percentage	0.015	0.008	1.820	0.0689
*Note: adjusted for weeks				

Supplemental Table 6. The adjusted linear regression model between log₁₀-transformed per capita incident case count and log₁₀-transformed population size for North Dakota, March 2020 - January 2021*

Variables	Parameter Estimate	Standard Error	t-Statistic	P-value
Population	0.195	0.033	5.961	0.0000
Below poverty estimate percentage	-0.0002	0.004	-0.069	0.9454
No high school diploma percentage	0.007	0.005	1.608	0.1079
Age 65+ percentage	-0.003	0.005	-0.635	0.5258
Population with disability percentage	0.009	0.006	1.543	0.1230
Minority percentage	-0.001	0.002	-0.597	0.5507
Crowding percentage	0.017	0.011	1.449	0.1475
*Note: adjusted for weeks				

Supplemental Table 7. The adjusted linear regression model between log₁₀-transformed per capita incident case count and log₁₀-transformed population size for South Dakota, March 2020 - January 2021*

Variables	Parameter Estimate	Standard Error	t-Statistic	P-value
Population	0.127	0.030	4.278	0.0000
Below poverty estimate percentage	-0.008	0.002	-3.373	0.0008
No high school diploma percentage	0.015	0.005	3.167	0.0016
Age 65+ percentage	-0.009	0.004	-2.383	0.0172
Population with disability percentage	0.0004	0.005	0.079	0.9374
Minority percentage	0.004	0.001	2.657	0.0079
Crowding percentage	-0.010	0.004	-2.392	0.0168
*Note: adjusted for weeks				

Supplemental Table 8. The adjusted linear regression model between log₁₀-transformed per capita incident case count and log₁₀-transformed population size for Wyoming, March 2020 - January 2021*

530.0810060.006		0.0606 0.9145
	0.107	0.9145
		0.7110
0.012	-2.206	0.0277
00 0.007	0.012	0.9901
30 0.010	2.937	0.0034
12 0.005	2.562	0.0106
90 0.018	5.017	0.0000
)(0000.007300.010120.005	0000.0070.012300.0102.9371120.0052.562

References:

- 1. Smithfield Foods. Smithfield Foods To Close Sioux Falls, SD Plant Indefinitely Amid COVID-19. <u>https://www.smithfieldfoods.com/press-room/company-news/smithfield-</u> <u>foods-to-close-sioux-falls-sd-plant-indefinitely-amid-covid-19</u>. Published 2020. Accessed February 28, 2021, 2021.
- School responses in North Dakota to the coronavirus (COVID-19) pandemic during the 2020-2021 school year. <u>https://ballotpedia.org/School_responses_in_North_Dakota_to_the_coronavirus_(COVID</u> -19) pandemic during the 2020-2021 school year. Accessed March 1, 2021.
- 3. National Center for Education Statistics. College Navigator for North Dakota. <u>https://nces.ed.gov/collegenavigator/?s=ND&pg=1</u>. Accessed March 1, 2021.
- 4. Toman J, Mercer T. Public Universities Adjust Fall Semester Schedule <u>https://www.sdbor.edu/mediapubs/New%20Press%20Releases/060320FallAdjust.pdf</u>. Published 2020. Accessed February 28, 2021.
- South Dakota Board of Regents Public Universities & Special Schools. Higher Education Institutions Web site. <u>https://www.sdbor.edu/administrative-offices/infogovtrelations/Pages/South-Dakota-Higher-Education-Institutions.aspx</u>. Published 2021. Accessed February 28, 2021.
- 6. Idaho State Board of Education. Idaho Back to School Framework Published 2021. Accessed February 28, 2021.
- Idaho State Government. Colleges and Universities <u>https://www.idaho.gov/education/colleges-universities/</u>. Accessed February 28, 2021.
- School responses to the coronavirus (COVID-19) pandemic during the 2020-2021 academic year <u>https://ballotpedia.org/School_responses_to_the_coronavirus_(COVID-19)_pandemic_during_the_2020-2021_academic_year</u>. Published 2021. Accessed February 28, 2021.
- 9. Montana University System. Colleges and Universities <u>https://mus.edu/Universities/</u>. Accessed February 28, 2021.
- Wyoming State Health Officer. Statewide Public Health Order, 6th Continuation and Modification. . <u>https://drive.google.com/file/d/1sQJbSSvr0ZnRKuKQ0kNbqsyyhYe99OzG/view</u>. Published 2020. Accessed February 28, 2021.
- List of Colleges and Universities in Wyoming. <u>https://myschoolhelp.com/colleges-in-wyoming/</u>. Accessed February 28, 2021.
- 12. The United States Census Bureau. County population totals 2010-2019. United States Census Bureau. <u>https://www.census.gov/data/tables/time-series/demo/popest/2010s-counties-total.html</u>. Published 2019. Accessed January 11, 2021.
- 13. North Dakota State Government. Executive Order 2020-04.1. <u>https://www.governor.nd.gov/sites/www/files/documents/executive-orders/Executive%20Order%202020-04.1%20COVID-19.pdf</u>. Published 2020. Accessed February 28, 2021.
- 14. North Dakota State Government. Executive Order 2020-10.1. https://www.governor.nd.gov/sites/www/files/documents/executive-

orders/Executive%20Order%202020-10.1.pdf. Published 2020. Accessed February 28, 2021.

- 15. State of north Dakota Department of Health. Testing orders, Long Term Care. <u>https://www.health.nd.gov/sites/www/files/documents/Files/MSS/coronavirus/State%20</u> <u>Health%20Officer%20Orders/05-27-2020%20Testing%20Orders.pdf</u>. Published 2020. Accessed February 28, 2021.
- 16. State of North Dakota Department of Health. Lifted Travel Quarantine Order <u>https://www.health.nd.gov/sites/www/files/documents/Files/MSS/coronavirus/State%20</u> <u>Health%200fficer%200rders/2020_02.4_Lifted_Travel_Order.pdf</u>. Published 2020. Accessed February 28, 2021.
- 17. State of North Dakota Department of Health. Mask Order. <u>https://www.health.nd.gov/sites/www/files/documents/Files/MSS/coronavirus/State%20</u> <u>Health%20Officer%20Orders/2020-08_Mask_Order.pdf</u>. Published 2020. Accessed February 28, 2021.
- North Dakota State Government. Executive Order 2020-43. <u>https://www.health.nd.gov/sites/www/files/documents/Files/MSS/coronavirus/State%20</u> <u>Health%20Officer%20Orders/2020-08_Mask_Order.pdf</u>. Published 2020. Accessed February 28, 2021.
- 19. North Dakota State Government. Executive Order 2020-43.3. https://www.governor.nd.gov/sites/www/files/documents/Executive%20Order%202020-43.3.pdf. Published 2020. Accessed February 28, 2021.
- 20. State of South Dakota Office of the Governor. Executive Order 2020-05. https://covid.sd.gov/docs/2020-05.pdf. Published 2020. Accessed February 28, 2021.
- 21. State of South Dakota Office of the Governor. Executive Order 2020-12. https://covid.sd.gov/docs/2020-12.PDF. Published 2020. Accessed February 28, 2021.
- 22. State of South Dakota Office of the Governor. Executive Order 2020-13. https://covid.sd.gov/docs/2020-13.PDF. Published 2020. Accessed February 28, 2021.
- 23. State of South Dakota Office of the Governor. Executive Order 2020-20. https://covid.sd.gov/docs/2020-20.pdf. Published 2020. Accessed February 28, 2021.
- 24. Documenting South Dakota's path to recovery from the coronavirus (COVID-19) pandemic, 2020-2021. https://ballotpedia.org/Documenting_South_Dakota%27s_path_to_recovery_from_the_c_oronavirus_(COVID-19)_pandemic, 2020-2021. Published 2020. Accessed February 28, 2021.
- 25. Idaho Department of Health and Welfare. Statewide Stay Home Order. <u>https://coronavirus.idaho.gov/wp-content/uploads/2020/06/statewide-stay-home-order_032520.pdf</u>. Published 2020 Accessed February 28, 2021.
- 26. Idaho Department of Health and Welfare. Stay Healthy Order, Stage 1. <u>https://coronavirus.idaho.gov/wp-content/uploads/2020/06/stay-healthy-order-stage1.pdf</u>. Published 2020. Accessed February 28, 2021.
- 27. Idaho Department of Health and Welfare. Stay Healthy Guidelines, stage 4. <u>https://rebound.idaho.gov/wp-content/uploads/stage4-stay-healthy-guidelines.pdf</u>. Published 2020. Accessed February 28, 2021.
- 28. Idaho Department of Health and Welfare. Stay Healthy Order, Stage 3 Modified <u>https://coronavirus.idaho.gov/wp-content/uploads/2020/10/stay-healthy-order-stage3_2020oct27.pdf</u>. Published 2020. Accessed February 28, 2021.

- 29. Idaho Department of Health and Welfare. Stay Healthy Order, Stage 2 Modified <u>https://coronavirus.idaho.gov/wp-content/uploads/2020/11/stage-2-modified-order.pdf</u>. Published 2020. Accessed February 28, 2021.
- Documenting Montana's path to recovery from the coronavirus (COVID-19) pandemic, 2020-2021.
 <u>https://ballotpedia.org/Documenting_Montana%27s_path_to_recovery_from_the_corona</u> virus (COVID-19) pandemic, 2020-2021. Published 2020. Accessed February 28, 2021.
- 31. Wyoming State Health Officer. Statewide Public Health Order #1, Closures. <u>https://drive.google.com/file/d/1qZ4MrBFvBxAXnyi4-vNo6CHCfGgS1wgO/view</u>. Published 2020. Accessed February 28, 2021.
- 32. Wyoming State Health Officer. Statewide Public Health Order, 3rd Continuation and Modification. https://drive.google.com/file/d/1kDaNfimewmLISuGscROII93SRPCzDduX_/view

https://drive.google.com/file/d/1kDaNfjmewmUSuGscRQII93SRPCzDduX_/view. Published 2020. Accessed February 28, 2021.

- 33. Wyoming State Health Officer. Statewide Public Health Order, 3rd Continuation and Modification (Gathering). <u>https://drive.google.com/file/d/1BrGn4eoQCHG3-</u> <u>Prm87sqUOxG2Vp-whlM/view</u>. Published 2020. Accessed February 28, 2021.
- 34. Wyoming State Health Officer. Statewide Public Health Order, 6th Continuation and Modification (Gathering). <u>https://drive.google.com/file/d/1WBQpBzmWL3neZJKSdHlo-lin8C1msEqw/view</u>. Published 2020. Accessed February 28, 2021.
- 35. Wyoming State Health Officer. Statewide Public Health Order, 8th Continuation and Modification. https://drive.google.com/file/d/1rDsIe2HHxZYybtuXpABimw8rUFoSyOIg/view.

https://drive.google.com/file/d/IrDsle2HHxZYybtuXpABimw8rUFoSyOIq/vi Published 2020. Accessed February 28, 2021.

- 36. Wyoming State Health Officer. Statewide Public Health Order, 8th Continuation and Modification (Gathering) <u>https://drive.google.com/file/d/1LIm6_UudyJEbTK3NGL_x6KamMDzIZ4jl/view</u>. Published 2020. Accessed February 28, 2021.
- Wyoming State Health Officer. Statewide Public Health Order, 13th Continuation and Modification. <u>https://drive.google.com/file/d/1J6VGUYFN3-N4I8NNO8-</u> ci5 TwrheA11T/view. Published 2020. Accessed February 28, 2021.
- Wyoming State Health Officer. Statewide Public Health Order, 13th Continuation and Modification (Gathering). <u>https://drive.google.com/file/d/1D-fu6mv72ZXIrcHJTDlrM3jzNyceXUQV/view</u>. Published 2020. Accessed February 28, 2021.
- Wyoming State Health Officer. Statewide Public Health Order#4: Requiring Face Coverings in Certain Places, with Exceptions. <u>https://drive.google.com/file/d/1zDkPo6KLZ2HnJAZFYYQNH87wHS5jxCTq/view</u>. Published 2020. Accessed February 28, 2021.