What We Learned from COVID-19: From endotheliitis to treatment

Adem Dirican[1], Selin Ildir[2], Tugce Uzar[2], Irem Karaman[2], Sevket Ozkaya[3]

[1]MD, Pulmonologist, VM Samsun Medicalpark Hospital, Department of Pulmonary Medicine,

Samsun, Turkey.

[2]Medical Student/Intern, Bahcesehir University School of Medicine, Istanbul, Turkey.

[3]MD, Professor, Bahcesehir University Faculty of Medicine, Department of Pulmonary Medicine,

Istanbul, Turkey

Corresponding Author

Sevket Ozkaya, MD, Pulmonologist, Professor, Bahcesehir University, Faculty of Medicine,

Department of Pulmonary Medicine, Istanbul, Turkey.

E-mail: ozkayasevket@yahoo.com, mobile phone: +90 532 474 13 09

Word count: 5030

Number of figures: 1

Number of tables: 1

Abstract

Objective: COVID-19 may yield a variety of clinical pictures, differing from pneumonitis to Acute Respiratory Distress Syndrome (ARDS) along with vascular damage in the lung tissue, named as *endotheliitis*. To date, no specific treatment strategy was approved by any authority for the prevention or treatment of COVID-19 in terms of endotheliitis-related comorbidities. Here, we present our experience of COVID-19 by evaluating 11,190 COVID-19 patients with the manifestations of endotheliitis in skin, lung, and brain tissues according to the different phases of COVID-19.

Methods: After a retrospective examination, patients were divided into three groups according to their repercussions of vascular distress, which were represented by radiological, histopathological, and clinical findings. (Group A: no or mild pulmonary involvement, Group B: moderate pulmonary involvement with clinical risk of deterioration, Group C: severe pulmonary involvement and respiratory failure). We presented the characteristics and disease course of seven representative and complicated cases which represents the different phases of the disease, and discussed the treatment strategies in each group. The current pathophysiological mechanisms responsible from SARS-CoV-2 infection, COVID-19 related respiratory failure and current treatment strategies were reviewed and discussed in detail.

Results: Among 11.190 patients, 9294 patients met the criteria for Group A, and 1376 patients were presented to our clinics with Group B characteristics. Among these patients, 1896 individuals(Group B and Group C) were hospitalized. While 1220 inpatients were hospitalized within the first 10 days after the diagnosis, 676 of them were worsened and hospitalized 10 days after their diagnosis. Among hospitalized patients, 520 of them did not respond to group A and B treatments and developed hypoxemic respiratory failure (Group C) and 146 individuals needed ventilator support and were followed in the intensive care unit, and 43 (2.2%) patients died.

Conclusion: Distinctive manifestations in each COVID-19 patient, including non-respiratory conditions in the acute phase and the emerging risk of long-lasting complications, suggest that COVID-19 has an *endotheliitis*-centred thrombo-inflammatory pathophysiology. Endotheliitis can also explain the mechanism behind the respiratory failure in COVID-19, and the difference of COVID-19 related ARDS from ARDS seen in other critical conditions. In addition, use of early corticosteroid in patients with early symptoms and early tocilizumab in ICU helps to reduce mortality and progression of the disease. Endotheliitis-based pathophysiological mechanisms are known to be momentarily changing and difficut to manage due to their risk of sudden aggrevation. Hence, daily evaluation of clinical, laboratory and radiological findings of patients and deciding appropriate pathophysiological treatment would help to reduce the mortality rate of COVID-19.

Keywords: COVID-19, Acute Respiratory Distress Syndrome, Endotheliitis, Cytokine Storm, Plasma

Introduction

Novel coronavirus disease 2019 (COVID-19) has resulted in a dramatic pandemic crisis by causing mainly a respiratory disease that can rapidly progress to pneumonia and, in severe cases, to acute respiratory distress syndrome (ARDS) (1). Globally, as of July 15[th] of 2021, there have been 187,519,798 confirmed cases of COVID-19, including 4,049,372 deaths, reported to World Health Organization (WHO)(2). The heterogeneity of the disease serves a spectrum from asymptomatic cases to respiratory failure. With the experiences from 11.190 COVID-19 patients since March 2020, we observed that distinctive clinical, radiological and histopathological manifestations of each COVID-19 patient, including non-respiratory conditions in the acute phase and the emerging risk of long-lasting complications, suggests an *endotheliitis*-centred thrombo-inflammatory pathophysiology for COVID-19 (3). Therefore, understanding the pathophysiology of SARS-CoV-2 infection, and more significantly the host response against it, is a fundamental tool to develop a personalized treatment for the patient's need and momentary response. Accordingly, we adopted our treatment strategies depending on both the respiratory and vascular distresses observed in patients. In this paper, we aimed to present the clinical, radiological, and histopathologic manifestations of COVID-19 related endotheliitis, as well as present our treatment categories which were focused on preventing endotheliitis-related consequences in different phases of the disease.

Methods

Patient Analysis and Classification

All COVID-19 patients that were diagnosed and treated in Samsun VM Medicalpark Hospital, Turkey, between March 2020 and April 2021 were retrospectively evaluated. Patients who were suspected to be infected by SARS-CoV-2 were confirmed with clinic, laboratory (positive reverse-transcriptase–polymerase-chain-reaction assay of nasopharyngeal swabs or serological IgM/IgG rapid antibody test against SARS-CoV-2) and radiologic (consistent HRCT findings) results and included in the study. Demographic characteristics, presenting symptoms of the patients at the time of admission, radiological images, hospitalization status and the presence of need for respiratory support were retrieved from patient records at the time of admission. Confidentiality of the study participants' information was maintained throughout the study. The study was performed in accordance with the Helsinki Declaration and approval for this study procedure was obtained from the Istinye University Institutional Review Boards/ethical committees with respect to its scientific content.

Three Main Groups for Different Phases SARS-CoV-2 Infection

This heterogeneous population of patients were divided into three groups (A,B,C) according to their repercussions of vascular distress, which were represented by radiological, histopathological, and clinical findings. Division into three groups were made according to a combined criteria that was

adopted from WHO severity classification and the extent of endotheliitis, which were represented by
the clinical symptoms, baseline oxygenations status, radiologic findings (chest x-ray/CT findings), and
hemodynamic differences (4). Accordingly, three escalating phases of COVID-19 disease progression
with associated signs, symptoms, and potential phase-specific treatments were described as early
infection phase(Group A), progressive phase(Group B), and severe and dissipative phase(Group C).

The treatment modalities for each group were adopted from current local guidelines, as well as
interventions that were later standardized according to clinical observation of momentary changes by
two different pulmonologists (5-7).

Group A. Group A included patients who had any of the various signs and symptoms of COVID-19
(e.g., fever, cough, sore throat, headache, muscle pain, diarrhea, loss of taste and smell) but did not
have shortness of breath. Group A patients either presented no imaging findings of pneumonia, or they
presented with the typical radiological representation of minimal patchy, subpleural, peripheral,
perivascular ground-glass opacities (GGO) (Fig. 1-I and 1-II). The placement of GGOs in the early
presentation of the disease is compatible with the distribution of microvascular capillaries of lung.

If the pulmonary involvement was absent or mild-to-moderate and the patient was suitable for
ambulatory treatment, Favipiravir tablet (1600 mg BID for the first day, followed by 600 mg BID for
four days, making five days in total), Dexamethasone (0.1-0.2 mg/kg per day), Azithromycin tablet,
low-molecular-weight-heparin(LMWH) and acetylsalicylic acid(ASA) therapies were applied for one
week.

Group B. Patients who showed evidence of lower respiratory disease during clinical assessment
(respiratory symptoms) or imaging(findings of pneumonia) but an oxygen saturation(SpO2) ≥94% on
room air at sea level, were hospitalized for close follow-up. Patients with comorbidities or special
conditions (i.e. age>65, diabetes, cancer, obesity, cardiovascular disease, chronic lung disease, sickle
cell disease, chronic kidney disease, being pregnant, cigarette smoking, transplant or
immunosuppression recipient) were also hospitalized due to their high risk of severity[4]. Patients who
had moderate pulmonary involvement or unresponsive to the Group A treatment in terms of the
clinical symptoms, with no evident respiratory failure, but had been indicated for hospitalization were
classified as Group B. Group B corresponds to the progressive phase of the disease, which is
characterized by multifocal, bilaterally diffused GGOs with poorly circumscribed consolidations
scattered in peripheral zones of lungs, along with vascular and intra-lobular septal thickenings called
"crazy paving pattern" (Fig.1-IIIb) (7).

For this group, Favipiravir tablet (1600 mg BID for the first day, followed by 600 mg BID for four
days, making five days in total), Dexamethasone (0.1-0.2 mg/kg per day) or methylprednisolone (1-2
mg/kg per day), Azithromycin tablet or fluoroquinolone, LMWH, ASA and Famotidine tablet
therapies were prescribed. Although the treatment takes one week, we observed that this phase has the
peak stage in 10-13 days and may include potential secondary complications. Thus, patients require a
close follow-up by serial chest X-rays to establish a baseline to assess the improvement of aeration.

Group C. Patients with moderate to severe pulmonary involvement(lung infiltrates >50%.),
accompanied by respiratory failure(SpO2 <94% on room air at sea level and , a ratio of arterial partial
pressure of oxygen to fraction of inspired oxygen (PaO2/FiO2) <300 mm Hg, respiratory frequency
>30 breaths/min), or to the patients who were recalcitrant to standard therapy and had deteriorating
clinical status with laboratory (particularly increasing ferritin levels) and radiological findings were
classified as Group C. In addition, patients who showed rapid progession(>%50) on CT imaging and
who presented with respiratory failure or shock within 24-48 hours also included in this group. This
group represents the progressive, peak, dissipative and severe phases of the disease. The radiological
findings of Group C were microscopic lacerations and infiltrations in perialveolar vessels which were
radiologically appeared as developing pulmonary GGOs, consolidation and diffuse alveolar damage
(DAD), which may be accompanied by pneumothorax, pneumomediastinum or intracranial
hemorrhage at this stage(Fig. 1-IV-V-VI-VII).

For this severe group of patients, Favipiravir tablet (1600 mg BID for the first day, followed by 600
mg BID for 4 days, continued for 5 or 10 days in total), methylprednisolone (250-1000 mg/day for at
least three days), convalescent plasma(once in a day during the first and second days of radiological
detoriation, applied for maximum three days), tocilizumab(400 mg QD/daily, applied two times),
supported by Piperacilin/Tazobactam, LMWH and famotidine therapies were administered. The
effectiveness of this group of treatment was measured by overall mortality ratio.

Results
A total of 11.190 patients were evaluated. The mean age was 59.2 ±17.3 years and male to female
ratio was 5507/5683. Among 11.190 patients, 9294 (83,05%) patients met the criteria for Group A,
and 1376 (12.29) patients were presented to our clinics with Group B characteristics. Among these
patients, 1896 (16.9%) individuals(Group B and Group C) were hospitalized. While 1220 inpatients
were hospitalized within the first 10 days after the diagnosis, 676 of them were worsened and
hospitalized 10 days after their diagnosis. Among hospitalized patients, 520 (4.64%) of them did not
respond to group A and B treatments and developed hypoxemic respiratory failure (Group C). Among
patients who developed hypoxemic respiratory failure, 146 (7.7%) individuals needed ventilator
support (non-invasive/invasive mechanical ventilation) and were followed in the intensive care unit,

and 43 (2.2%) patients died. As a complication, among 11.190 patients, 30 patients developed either
spontaneous pneumothorax, pneumomediastinum or subcutaneous emphysema, and 4 patients died
with a 13.3% mortality rate.
188
Radiologic images and histopathologic samples of representative groups are presented in **Figure 1**
which consists of 7 representative and complicated cases with variable stages of COVID-19, togethet
with additional examples of findings of early and severe endotheliitis in brain, pulmonary and skin
tissues. Representative cases for early phase:Group A (Fig.1-I,II), progressive phase:Group B(Fig.1-
III), severe and dissipative phase:Group C(Fig.1-IV,V,VI,VII) were presented and cases were
presented in Figure 1.
195
Case 1(Figure 1-I) was a female patient who was in her 50's and admitted to the hospital with
complaints of cough and fever. She was diagnosed with COVID-19 due to PCR positivity. HRCT
images showed patchy, subpleural and peripheral perivascular ground-glass opacities, corresponding
the early phase of pneumonitis. She received Group A treatment. GGOs in transaxial images were
located in the subpleural area, where the microvascular *endoteheliitis* and endothelial destructions
mostly likely occur, due to the interaction with toxic plasma in the capillaries of pulmonary interstitial
space. She was completely recovered at 14^{th} days at control appointment, after 5 days of treatment
with favipiravir.
204
Case 2(Figure 1-II) was a male patient who was 47-52 years old and admitted to the hospital with
complaints of fever and chest pain. His COVID-19 PCR test was positive. In addition to the
characteristics of an early phase COVID-19 pneumonitis (end-capillary microangiopathy, explained in
Fig.1-I) his HRCT images showed perivascular consolidation. It appeared as the continuation of crazy-
paving pattern, which was demonstrated by the thickening in inter-lobular and intra-lobular septa.
Lung involvement was limited and monofocal. Hence, the patient received Group A treatment. He was
completely recovered at 14^{th} days at control appointment, after 5 days of treatment with favipiravir.
212
Case 3(Figure 1-III) was a male patient whose age ranged between 63-68 and admitted to the hospital
with complaints of fever and cough. His PCR test resulted positive for SARS-CoV-2. Transaxial
HRCT image in the first day of positivity shows bilateral and patchy nodulary GGOs as expected in
early phase **(IIIa)**. He received group B treatment and hospitalized for close follow-up. At the 5th day
of positivity, affected pulmonary areas were advanced into scattered consolidations (**IIIb**). This
appearance was noted as the progressive phase of pneumonitis and was considered as representation of
clinical deterioation clinically (i.e. dyspnea, respiratory failure). On the 15^{th} day of positivity,
fibroreticular consolidations were conspicuous (**IIIc**). The dissipative phase was the healing process,
characterized by the resolution in lung parenchyma and residual GGO, observed after 35 days of

positivity (**IIId**). Parenchymal bands, originated from previous fibroreticular proliferation, were also

visible (**IIId**).

If the patients have a tendency of severe phase and/or unresponsiveness to Group B therapy, **Group C**

treatment was used. Representative cases of this group are shown in sections **IV-V-VI-VII.** This group

of patients also underwent early and progressive phases of COVID-19 but they showed further

deterioration of lung involvement and endothelial damage, therefore, they received the last group of

treatment.

Case 4(Figure 1-IV) was a male patient whose age ranged between 53-58 and admitted to the hospital

with complaints of fever. His PCR test was positive results positive for SARS-CoV-2. Coronal,

sagittal and axial planes of HRCT images initially showed small GGO with a local subpleural sparing,

particularly around the microvascular area on the first day of positivity (**IVa**). In the 5^{th} day of

treatment, increased GGOs were visible in the progressive phase (**IVb**). After 18 days of positivity,

pneumomediastinum, Diffuse Alveolar Damage (DAD) and ARDS were seen in the severe phase

(**IVc**). Lastly, the dissipative phase was seen, after 30 days of positivity, with residual fibrotic

parenchyma (**IVd**).

Case 5(Figure 1-V) was a female patient whose aged ranged between 49-53 and admitted to the

hospital with complaints of fever and cough. Her COVID-19 PCR test was positive. Her first HRCT,

on the 2^{nd} day of positivity, showed COVID-19-related bilateral and multifocal nodular GGOs (**Va**).

After 10 days of positivity, DAD developed along with ARDS and pneumomediastinum characterized

with the severe phase (**Vb**). She recovered and discharged after 30 days of positivity with the

dissipative phase (**Vc**). The regression in pulmonary lesions was visible on the HRCT image two

months after diagnosis (**Vd).**

Case 6(Figure 1-VI) was a male patient whose age ranged between 50-55 and admitted to the

hospital with complaints of fever and cough. His COVID-19 PCR test was positive. His HRCT

showed moderate pneumonia in the early phase (**VIa**). After 12 days of positivity, the severe phase

develops with DAD and ARDS (**VIb**). In the 25^{th} day of positivity, recovery was observed in the

dissipative phase (**VIc**). The extent of improvement in pulmonary lesions can be noticed in **VId,**

which was 40 days after the diagnosis.

Case 7(Figure 1-VII) was a male patient whose age ranged between 48-53 and admitted to the

hospital with complaint of dyspnea. His COVID-19 PCR test was positive. His first HRCT showed

ARDS pattern with dense consolidations (**VIIa**). Pneumothorax developed after two weeks from

diagnosis (**VIIb**). In addition to the respiratory failure, hemorrhagical intracranial areas were seen in

the T2-weighted MRI. (**VIIc**). He was lost at the 20th day of positivity.

The demonstration of endotheliitis in the biopsy section of lung parenchyma (Fig.1-VIII), radiological

images of brain(Fig.1-IX) and lung(Fig.1-X,XI), and skin manifestation(Fig.1-XII) of endotheliitis are

also presented in Figure 1.

Discussion

Are Laboratory Findings' Reliable in Determining the COVID-19 Course & Prognosis?

Our classification of patient population and treatment groups was primarily based on the clinical and

radiological findings, rather than the laboratory findings. During a year of the pandemic, in our clinics,

we observed that even though the laboratory findings may show the degree of pathophysiology during

the disease course, it may mislead the clinician when it comes to clinical practice. While C-reactive

protein, ferritin, lactate dehydrogenase, erythrocyte sedimentation rate, IL-6 elevation and TNF-alpha

represent inflammatory changes; D-dimer, prolonged PTT and troponin elevation show the

predisposition to thrombosis and myocardial damage (8,9). Cheng et al. points out that severely

affected patients, along with the ones who could not survive the disease, are subject to significantly

higher ferritin levels in comparison to the non-severe and survivor groups of patients (10). However,

COVID-19 is an acute syndrome of different age and comorbidity groups. When the clinician fails to

detect the pre-infectious baseline values of the patient, overtreatment may worsen the situation**.** In our

study group, a patient with 250 mg/L ferritin received Group C treatment, while another with 2550

mg/L received Group A due to the difference in their previous normal values (ferritin increased 10 and

<1 times, respectively).   The change in the concentrations of inflammatory markers seem to be

significantly different in COVID-19 than in typical non-COVID-19-related ARDS, suggesting that

COVID-19 features its own unique, poorly understood, yet detrimental, inflammatory profile (11).

Therefore, we advise the categorization and treatment strategies to be selected by monitoring

radiological findings and the severity of clinical features (such as dyspnea and oxygen saturation).

Farsighted Evaluation: What are the Causes in Pathogenic Scene After Infection with SARS-

CoV-2?

Cytokine Storm and Emergence of Toxic Plasma

Pathophysiology of COVID-19 is closely related to cytokine storm, which arises from the consecutive

and intricate activation of numerous inflammatory cells that cause excessive and/or unregulated,

proinflammatory cytokine release (13). Cytokines force blood plasma to undergo a chemical

alteration, revealing toxic and irritant characteristics. Cytokine storm comprises the systemic

activation of unstimulated tissue cells, epithelial and endothelial cells in addition to hyperactivation of

hematopoietic cells, including B lymphocytes, natural killer (NK) cells, macrophages, dendritic cells,

neutrophils and monocytes which provoke the excessive release of proinflammatory cytokines (14).
This toxic setting not only causes inflammation but also a damage in various systemic tissues via the
signals of pro-apoptosis (15). Main clinical manifestations of cytokine storm appear as fever,
progressive dyspnea, tachypnea and elevated inflammatory markers such as IL-6, CRP and ferritin as
it is observed in COVID-19 patients (14,15). The uncontrolled production of pro-inflammatory factors
(IL-6, IL-8, IL-1β, and GM-CSF), and chemokines (CCL2, CCL3, CCL-5) together with reactive
oxygen species (ROS) cause ARDS which leads to pulmonary fibrosis and death. Abnormal nitric
oxide metabolism, upregulation of ROS and proteases, downregulation of endothelium-associated
antioxidant defense mechanisms, and induction of tissue factor altogether provide a basis for vascular
pathology in COVID-19 (3).

Additionally, the systemic inflammatory response against SARS-CoV-2 infection is supported by the
circulating mediators found in different organ systems, which are demonstrated in postmortem
histopathological analysis to indicate the organ-dependent cytokines (16,17). Fig.1-VIII represents the
areas of lung parenchyma with the mixt-type inflammatory-cell infiltration and exudative capillaritis
with thickened microvascular walls, in addition to the interstitial and intra-alveolar proliferation of
fibroblasts.

Vascular Effects: Endotheliitis, Thromboinflammation and Systemic Microangiopathy
Endotheliitis, hypercoagulability and thrombotic microangiopathy are namely the vascular hallmarks
of COVID‐19 (18,19). These vascular complications should be evaluated separately from ARDS. As
a matter of fact, the histopathologic changes observed in several tissue samples might be primarily the
result of C3-mediated pathways in thromboinflammation(18). Beigee et al. stated that vascular
widespread platelet–fibrin microthrombi was the main pathologic finding in the lung samples of
critical COVID-19 patients with severe hypoxemia and minor radiological abnormalities on imaging
(16). They also indicated that clinically not all patients with ARDS present DAD. However, the
presence of DAD with ARDS contributes worsening of clinical outcomes when compared with those
without DAD. Early and late endotheliitis lesions on pulmonary HRCT are seen in Fig.1-X-XI.

Varga et. al were first to demonstrate that irritant plasma, together with the SARS-CoV-2 infection,
cause '*endotheliitis*' in microvascular capillary endothelium, which is the primary pathology seen
under non-immune, corrosive and irritant conditions (20). Similarly, Zhang et al. demonstrated that
COVID-19 infection resembles more of the pathophysiology and phenotype of complement-mediated
thrombotic microangiopathies(TMA); rather than sepsis-induced coagulopathy or disseminated
intravascular coagulation (21). A common denominator of complement-mediated effects in TMAs is
angiocentric inflammation causing endothelial dysfunction; mononuclear and neutrophilic
inflammation of microvessels and as a result, microvascular thrombosis, which brings poor prognosis,

multiple organ dysfunction syndrome and ARDS(22). Microangiopathies in COVID-19 patients are characterized by anemia, increased lactate dehydrogenase, thrombocytopenia and organ damage (ex. skin lesions, neurological, renal and cardiac dysfunction) (22,23). In our cohort of patients, skin lesions were seen as COVID-19-associated papulovesicular exanthema scattered in trunk and mild pruritus (Fig.1-XII). Trellu et al. indicated that histopathological findings of papulovesicular eruption reveal the signs of *endotheliitis* and microthrombosis in the dermal vessels (24). Unfortunately, patients may die, not from respiratory failure, but due to the vascular coagulopathies (i.e.hemorrhage) in brain, kidneys and heart (Fig.1-IX).

Fundamentals Behind COVID-19 Related Respiratory Failure

Endothelial dysfunction plays a key role in understanding the multisystemic attack of SARS-CoV-2 infection. Microvascular capillary *endotheliitis* is the primary mechanism that cause clinical detoriation, particularly in those patients with advanced pulmonary involvement (25). The critical point in this manner is to differentiate non-immune *endotheliitis* from immune complex endotheliitis in lungs, and to consider it as the main pathology of COVID-19 since it is not directly mediated by the active antigen-antibody complexes or the virulence of SARS-CoV-2 itself (26,27).

Initially, the irritant action of plasma leads to the thickening of the vessel wall and deceleration in blood flow, which is responsible for the microthrombosis in capillary beds in lungs, which may lead to respiratory failure. Fox et al. demonstrated the microvascular thrombosis and hemorrhage in lungs, as a remarkable contributor of death, in the autopsies of COVID-19 non-survivors (28). They also proved that the cardiovascular damage is "non-immune" by demonstrating cardiac cell necrosis without lymphocytic myocarditis in deceased patients.

The concept of virus-induced pulmonary vasculitis is consistent with a substantial ventilation/perfusion mismatch in COVID-19 based on a right-to-left pulmonary shunt due to a vicious cycle beginning with an increase in respiratory effort and oxygen consumption in inflamed and hyperperfused lungs, failure of hypoxic vasoconstriction, and resulting fatal outcome (3). Therefore, the failure of simple ventilatory support in COVID-19 is commonly observed in patients who are unable to satisfy the oxygen demand due to reduced lung capacity (such as older patients and patients with obesity) and cardiovascular comorbidities.

In the clinical presentation, aggravation of dyspnea and hypoxemia symptoms were attributed to dysfunctional crosstalk between leukocytes and endothelial cells that manifest as vascular immunopathology predominantly confined to the lungs. Eventually, since microvascular walls are prone to damage, the destruction most conveniently occurs as *endotheliitis* at the site of pulmonary interstitial capillaries; with the help of lung elasticity and thin vascular walls, conveys into the

perivascular space(29). Remarkably, due to SARS-CoV-2's endotheliophilic nature, endothelial and

epithelial infections appear to be the predominating factors during the course of the disease (3). On the

other hand, alveolar-centered infection and the disruption of alveolar epithelial–endothelial barriers

contribute to the development of DAD and *pneumonitis,* which manifest as GGOs in alveolar spaces

(30,31) (Fig.1-III and VIII). The aforementioned endothelial damage may spread to different systems

and become lethal (Fig.1-VII).

Ekanem et al. stated that higher inflammatory markers (ferritin, CRP and fibrinogen), increased

fibrosis in HRCT images, and absence of receiving an interleukin-6 inhibitor or convalescent plasma

are associated with higher probability of severity and mortality via the spontaneous pneumothorax

(SPT) (32). They also suggested that there must be factors uniquely associated with COVID-19 that

contribute to the incidence of SPT, since half of the patients were not on a ventilator when the

pneumothorax was diagnosed. In Group C (Fig.1-IV-V-VI-VII), we also observed that endothelial

damage, along with thromboinflammation, brought increased incidence of pneumothorax secondary to

DAD in patients with ARDS. Here, among 11.190 patients, 30 patients developed either SPT,

pneumomediastinum or subcutaneous emphysema with a 13.3% mortality rate. The most important

reason behind these complications was most likely DAD, which stems from the high transpulmonary

pressure and alveolar wall vulnerability, with decreased compliance and increased frailty, resulting in

an air leakage into the chest compartments(33). SPT that is observed in severe COVID-19 patients, is

thought to be derived from reduced alveolar vessel caliber due to the virus-induced cytolysis,

mononuclear immunological response to injury and the small vessel thrombosis at the site of

perialveolar area, which should be differentiated from iatrogenic pneumothorax related to mechanical

ventilation (34).

Prevention of Aforementioned Microvascular Pathology and Mortality

Planning an effective therapy for COVID-19 infection is a complex process. According to Mastellos et

al. broader pathogenic involvement of C3-mediated pathways in thromboinflammation supports the

utilization of complement inhibitors in COVID-19, which result in diminished hyper-inflammation

and marked lung function improvement(35). Teuwen et al. suggested that normalization of vascular

walls through metabolic interventions might be considered as an additional potential target for the

therapy(36). Therefore, until a specific antiviral is discovered against SARS-CoV-2, convalescent

plasma therapy and immunomodulators play a significant role to control the consequences of SARS-

CoV-2 infection (i.e. cytokine storm), to reduce inflammatory cell infiltration in lungs and to prevent

fatal course in severe patients by reducing the likelihood of the Systemic Inflammatory Response

Syndrome (37).

The latest COVID-19 treatment guideline published by the Turkish Ministry of Health included the
favipiravir, ASA, famotidine, LMWH, dexamethasone , tocilizumab, which were shown to be safe and
in vitro effective against SARS-CoV-2 (5,6). Based on indirect evidence from different clinical trials,
convalescent plasma therapy and pulse steroid therapy were also suggested for severe patients (38).
The required interventions were made according to clinical severity of patients, vitals, time from
symptom onset and radiological images.

Convalescent plasma therapy plays a critical role in neutralizing the plasma and diminishing its
corrosiveness. Convalescent plasma not only demonstrates an antibody response but also denotes
immunomodulatory, anti-cytokine and proinflammatory effects, which appear as key factors to
minimize disease severity and mortality in COVID-19 cases(38). Gomez-Pastora et al. explain this
correlation by the phenomena of pro-inflammatory and anti-inflammatory cytokine activation, as a
result of macrophage associated hyperferritinemia(39). They stated that ferritin plays an active role as
a pathogenic mediator in COVID-19 and the therapeutic use of plasma is beneficial to reduce ferritin
and cytokine levels in the body. Our experience with convalescent plasma showed rapid and positive
results against the symptoms of dyspnea, hypoxemia, fever and radiologically seen infiltrations, which
was demonstrated **in Case 4-7.**

Antiviral drugs are being used to decrease the viral load. Correspondingly, Favipiravir was the drug of
choice that is recommended by Turkish Republic of Health Ministry guidelines (5,6). However, it
should be noted that antiviral therapy fails to prevent pulmonary involvement, which is the result of
inflammatory process rather than the effect of SARS-CoV-2 infection itself (40).

Systemic corticosteroid drugs (dexamethasone and methylprednisolone) are the only effective
therapeutic agents to repair non-immune capillary microvascular *endotheliitis*, hence, advised to be
used even in the presence of minimal ground-glass opacities(41). In our clinical experience, to benefit
the best of steroids, steroids should be utilized in the early phase, rather than the progressive phase
(approximately the second week of infection). Minimally distributed GGOs may easily progress to
severe ARDS, in the absence of steroid treatment (**Fig.1-VIIa-VIb**). In a multi-centered study
conducted by RECOVERY Collaborative Group, mortality ratio in patients, who receive oxygen
support and dexamethasone, found to be lower than the control group, especially if the patients are
receiving invasive mechanical ventilation (29.3% vs. 41.4%; rate ratio, 0.64; 95% CI, 0.51 to 0.81)
(42). Although the role of corticosteroids in COVID-19 has been well-recognized in the therapeutic
algorithm, the right timing, dosage and duration of corticosteroid use is still unknown. Pinna et al.
suggested that although the early use of corticosteroids might facilitate the viral replication in the
upper airways, late administration fails to prevent the alveolar damage(43). It has been suggested that
early initiation of low-dose methylprednisolone therapy (with up to 40mg/day for less than 10 days)

provides clinical, radiological improvement in patients with active disease state of COVID-19, by
reducing the immune cascade and progression into cytokine without any adverse effect (44-46).

Immunomodulatory therapies help to diminish the cytokine response of the body. Tocilizumab
therapy, a monoclonal IL-6 antagonist, reduced the likelihood of progression to the composite
outcome of mechanical ventilation or death (47). Capra et al. have demonstrated that tocilizumab can
be used as an immunomodulatory drug of choice in case of the severe COVID-19 to reduce mortality,
diminish oxygen intake and treat lung opacities as well (48). Findings by Gupta et al. also suppoerted
the early use of(within first 2 days of ICU) tocilizumab to reduce the in-hospital mortality among
critically ill patients with COVID-19 (49).

LMWH and ASA are well-known to prevent the formation of microvascular thrombosis and cure
hypoxemia, thus can be used as supportive treatment against cardiovascular complications of COVID-
19 (50). However, before the use of ASA, patient history should be questioned in terms of renal
failure, gastrointestinal system disease, cardiovascular disease. Local current guidelines recommended
the use of NSAIDs, especially in the first 5-10 days; and steroid treatment to be started in the early
period in patients who worsen after theoretical viral clearance is completed in the first 5-10 days (5).

Limitation
We acknowledge that there are confounding factors related to management of COVID-19 patients due
to lack of a standardized guideline for treatment of each and every patient. However, while the in-
hospital mortality was reported to be up to 25% in different prospective trials, we believe that our
standardized treatment approach, which has a result of 2.2% in-hospital mortality, represent the
success of the personalized management of each COVID-19 patient in a single center (50). As the
COVID-19 pandemic continues, our current strategy represents a snapshot that would most probably
change drastically over time.

As a limitation, the design of the current study does not allow for further analysis of the COVID-19
patients in terms of the change in their laboratory, clinical and radiological parameters and
investigation of the effects of different SARS-CoV-2 variants(such as Alpha, B.1.1.7, Beta B.1.351, or
Delta B.1.617.2) in the clinical presentation of the patients. The observational methodology suggested
may be a source of bias that could lead to wrong conclusions on the effectiveness of treatments and the
clinical representation of the underlying pathophysiology. Provided treatment strategies are
implemented to all patients regardless of their SARS-CoV-2 variant types. No corruptions from
Turkey's Ministry of Health guidelines are enforced to the patients in terms of new experimental drugs
and/or biological agents.

We would like to declare that, our study did not interfere any patient's right to recieve treatment by
adressing a control group in a pandemic situation. Here, we also did not aim to demonstrate the
effectiveness of the therapy in the means of laboratory data, however we aimed to present the
representations of the required interventions in the each group of patients, by preventing the
progression of endotheliitis. Therefore, we used the current literature to support our clinical
observation and own perspective. In this end of the study, our aim was to declare our own point of
view by using our clinical experience, clinical representation of endotheliitis and the current literature.
We would like to approach to the vascular distress phenomenon as a clinical parameter that can be
used practically in the clinics, both by recognizing from radiological images and clinical findings of
the patients

Conclusion

SARS-CoV-2 infection is a multisystemic disease which courses rapidly with respiratory failure and
complications secondary to vascular alterations (i.e. microvascular thrombosis, *endotheliitis*, and
cytokine induced plasma toxicity). Early detection of radiological detoriation before laboratory
findings, via monitoring chest X-rays daily, and planning personalized treatments constitute a crucial
and life-saving maneuver in the treatment of COVID-19. Our group suggests that an important key to
success relies on how closely the clinicians follow patients from diagnosis to treatment, including the
whole course of the disease from outpatient clinic to ICU, in order to differentiate instant clinical
deviations from previous general status

Distinctive manifestations in each COVID-19 patient, including non-respiratory conditions in the
acute phase and the emerging risk of long-lasting complications, suggest that COVID-19 has an
endotheliitis-centred   thrombo-inflammatory   pathophysiology.   Potential   pathophysiological
mechanisms   contributing   to   endotheliitis   includes   cytokine   storm   and   toxic   plasma,
thromboinflammation and systemic microangiopathy. Endotheliitis can also explain the mechanism
behind the respiratory failure in COVID-19, and the difference of COVID-19 related ARDS from
ARDS seen in other critical conditions. In our observations, utilization of early dexamethasone in
Group A prevented the progression of the COVID-19 into more severe form. In addition, use of early
steroid in Group A and early tocilizumab in group C helps to reduce mortality and progression of the
disease. Endotheliitis-based pathophysiological mechanisms are known to be momentarily changing
and difficut to manage due to their risk of sudden aggrevation. Hence, daily evaluation of patients and
deciding appropriate pathophysiological treatment for the momentary changes in clinical, laboratory
and radiological findings would help to reduce the mortality rate of this novel virus. The collaboration
of scientists and clinicians around the world is required in order to develop novel prognostic
biomarkers and establish precise predictive thresholds for known biomarkers to foresee the severity

for COVID-19 pneumonitis that is characterized by vasculopathies and wide range of immune

derangements.

Disclosure

The authors have no conflicts of interest to declare.

Ethics Statement

The patients signed written informed consents to be able to provide data in this study. The study was

performed in accordance with the Helsinki Declaration and approval for this study procedure was

obtained from the Istinye University Institutional Review Boards/ethical committees with respect to its

scientific content.

Authors' Contribution

AD and SO diagnosed, treated the patients and designed the analysis, SI, TU and IK collected the data,

analysis tools and wrote the paper.

Data Availability

The research article data used to support the findings of this study are available from the

corresponding author upon request.

References

1. Li X, Ma X. Acute respiratory failure in COVID-19: is it "typical" ARDS? Critical Care.
2020;24(1):1-5.

2. "Coronavirus Update (Live)-Worldmeter"(retrieved May 11,2021). www.worldometers.info

3. Osuchowski MF, Winkler MS, Skirecki T, Cajander S, Shankar-Hari M, Lachmann G, et al.
The COVID-19 puzzle: deciphering pathophysiology and phenotypes of a new disease entity.
The Lancet Respiratory Medicine. 2021.

4. Zu ZY, Jiang MD, Xu PP, Chen W, Ni QQ, Lu GM, et al. Coronavirus Disease 2019
(COVID-19): A Perspective from China. Radiology. 2020;296(2):E15-E25.

5. Republic of Turkey Ministry of Health. COVID-19 Interim Guidance (T.C. Sağlık Bakanlığı
COVID-19 (SARS-CoV-2 Enfeksiyonu) Rehberi) 2020. Available from:
https://covid19bilgi.saglik.gov.tr/depo/rehberler/COVID-19_Rehberi.pdf?type=file.

6.   Köktürk N, İtil BO, Altınışık G, Adıgüzel N, Akgün M, Akyıldız L, et al. COVID-19
Pandemic and the Global Perspective of Turkish Thoracic Society. Turkish thoracic journal.
2020;21(6):419-32.

7.   COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment
Guidelines. 2021. National Institutes of Health. Accessed 2021 Mar 28.
https://www.covid19treatmentguidelines.nih.gov/

8.   Merad M, Martin JC. Pathological inflammation in patients with COVID-19: a key role for
monocytes and macrophages. Nature Reviews Immunology. 2020;20(6):355-62.

9.   Grasselli G, Tonetti T, Protti A, Langer T, Girardis M, Bellani G, et al. Pathophysiology of
COVID-19-associated acute respiratory distress syndrome: a multicentre prospective
observational study. The lancet Respiratory medicine. 2020;8(12):1201-8.

10. Cheng L, Li H, Li L, Liu C, Yan S, Chen H, et al. Ferritin in the coronavirus disease 2019
(COVID□19): A systematic review and meta□analysis. Journal of clinical laboratory
analysis. 2020;34(10):e23618.

11. Sinha P, Calfee CS, Cherian S, Brealey D, Cutler S, King C, et al. Prevalence of phenotypes
of acute respiratory distress syndrome in critically ill patients with COVID-19: a prospective
observational study. The Lancet Respiratory Medicine. 2020;8(12):1209-18.

12. Ufuk F, Savaş R. Chest CT features of the novel coronavirus disease (COVID-19). Turkish
journal of medical sciences. 2020;50(4):664-78.
13. Coperchini F, Chiovato L, Croce L, Magri F, Rotondi M. The cytokine storm in COVID-19:
An overview of the involvement of the chemokine/chemokine-receptor system. Cytokine &
growth factor reviews. 2020;53:25-32.
14. Ye Q, Wang B, Mao J. The pathogenesis and treatment of the Cytokine Storm in COVID-19.
Journal of infection. 2020;80(6):607-13.

15. Becker RC. COVID-19-associated vasculitis and vasculopathy. Springer; 2020.

16. Beigee FS, Toutkaboni MP, Khalili N, Nadji SA, Dorudinia A, Rezaei M, et al. Diffuse
alveolar damage and thrombotic microangiopathy are the main histopathological findings in
lung tissue biopsy samples of COVID-19 patients. Pathology-Research and Practice.
2020;216(10):153228.

17. Elsoukkary SS, Mostyka M, Dillard A, Berman DR, Ma LX, Chadburn A, et al. Autopsy
findings in 32 patients with COVID-19: a single-institution experience. Pathobiology.
2021;88(1):55-67.

18. Mosleh W, Chen K, Pfau SE, Vashist A. Endotheliitis and endothelial dysfunction in patients
with COVID-19: its role in thrombosis and adverse outcomes. Multidisciplinary Digital
Publishing Institute; 2020.

19. Calabretta E, Moraleda JM, Iacobelli M, Jara R, Vlodavsky I, O'Gorman P, et al.
COVID□19□induced endotheliitis: emerging evidence and possible therapeutic strategies.
British Journal of Haematology. 2021;193(1):43.

20. Varga Z, Flammer AJ, Steiger P, Haberecker M, Andermatt R, Zinkernagel AS, et al.
Endothelial cell infection and endotheliitis in COVID-19. The Lancet. 2020;395(10234):1417-
8.

21. Zhang Y, Xiao M, Zhang S, Xia P, Cao W, Jiang W, et al. Coagulopathy and antiphospholipid
antibodies in patients with Covid-19. New England Journal of Medicine. 2020;382(17):e38.

22. Wang X, Sahu KK, Cerny J. Coagulopathy, endothelial dysfunction, thrombotic
microangiopathy and complement activation: potential role of complement system inhibition
in COVID-19. Journal of thrombosis and thrombolysis. 2021;51(3):657-62.

23. Thakur V, Ratho RK, Kumar P, Bhatia SK, Bora I, Mohi GK, et al. Multi-organ involvement
in COVID-19: beyond pulmonary manifestations. Journal of clinical medicine.
2021;10(3):446.

24. Trellu LT, Kaya G, Alberto C, Calame A, McKee T, Calmy A. Clinicopathologic aspects of a
papulovesicular eruption in a patient with COVID-19. JAMA dermatology. 2020;156(8):922-
4.

25. Rovas A, Osiaevi I, Buscher K, Sackarnd J, Tepasse P-R, Fobker M, et al. Microvascular
dysfunction in COVID-19: the MYSTIC study. Angiogenesis. 2021;24(1):145-57.

26. Azkur AK, Akdis M, Azkur D, Sokolowska M, van de Veen W, Brüggen MC, et al. Immune
response to SARS□CoV□2 and mechanisms of immunopathological changes in COVID□19.
Allergy. 2020;75(7):1564-81.

27. Gavriilaki E, Anyfanti P, Gavriilaki M, Lazaridis A, Douma S, Gkaliagkousi E. Endothelial
Dysfunction in COVID-19: Lessons Learned from Coronaviruses. Current hypertension
reports. 2020;22(9):1-12.

28. Fox SE, Akmatbekov A, Harbert JL, Li G, Brown JQ, Vander Heide RS. Pulmonary and
cardiac pathology in African American patients with COVID-19: an autopsy series from New
Orleans. The Lancet Respiratory Medicine. 2020;8(7):681-6.

29. Arachchillage DJ, Stacey A, Akor F, Scotz M, Laffan M. Thrombolysis restores perfusion in
COVID☐19 hypoxia. British journal of haematology. 2020;190(5):e270-e4.

30. Doglioni C, Ravaglia C, Chilosi M, Rossi G, Dubini A, Pedica F, et al. Covid-19 interstitial
pneumonia: histological and immunohistochemical features on cryobiopsies. Respiration.
2021;100(5):369-79.

31. Dumoulin DW, Gietema HA, Paats MS, Hendriks LE, Cornelissen R. Differentiation of
COVID-19 pneumonitis and ICI induced pneumonitis. Frontiers in oncology. 2020;10.

32. Ekanem E, Podder S, Donthi N, Bakhshi H, Stodghill J, Khandhar S, et al. Spontaneous
pneumothorax: An emerging complication of COVID-19 pneumonia. Heart & Lung.
2021;50(3):437-40.

33. Elhakim TS, Abdul HS, Romero CP, Rodriguez-Fuentes Y. Spontaneous
pneumomediastinum, pneumothorax and subcutaneous emphysema in COVID-19 pneumonia:
a rare case and literature review. BMJ Case Reports CP. 2020;13(12):e239489.

34. Zantah M, Castillo ED, Townsend R, Dikengil F, Criner GJ. Pneumothorax in COVID-19
disease-incidence and clinical characteristics. Respiratory Research. 2020;21(1):1-9.

35. Mastellos DC, da Silva BGP, Fonseca BA, Fonseca NP, Auxiliadora-Martins M, Mastaglio S,
et al. Complement C3 vs C5 inhibition in severe COVID-19: Early clinical findings reveal
differential biological efficacy. Clinical Immunology. 2020;220:108598.

36. Teuwen L-A, Geldhof V, Pasut A, Carmeliet P. COVID-19: the vasculature unleashed. Nature
Reviews Immunology. 2020;20(7):389-91.

37. Sun X, Wang T, Cai D, Hu Z, Liao H, Zhi L, et al. Cytokine storm intervention in the early
stages of COVID-19 pneumonia. Cytokine & growth factor reviews. 2020;53:38-42.

38. Altuntas F, Ata N, Yigenoglu TN, Bascı S, Dal MS, Korkmaz S, et al. Convalescent plasma
therapy in patients with COVID-19. Transfusion and Apheresis Science. 2021;60(1):102955.

39. Gómez-Pastora J, Weigand M, Kim J, Wu X, Strayer J, Palmer AF, et al. Hyperferritinemia in
critically ill COVID-19 patients–Is ferritin the product of inflammation or a pathogenic
mediator? Clinica Chimica Acta; International Journal of Clinical Chemistry. 2020.

40. Solaymani-Dodaran M, Ghanei M, Bagheri M, Qazvini A, Vahedi E, Saadat SH, et al. Safety
and efficacy of Favipiravir in moderate to severe SARS-CoV-2 pneumonia. International
immunopharmacology. 2021;95:107522.

41. Raju R, Prajith V, Biatris PS. Therapeutic role of corticosteroids in COVID-19: a systematic
review of registered clinical trials. Future Journal of Pharmaceutical Sciences. 2021;7(1):1-18.

42. Group RC. Dexamethasone in hospitalized patients with Covid-19. New England Journal of
Medicine. 2021;384(8):693-704.

43. Pinna SM, Scabini S, Corcione S, Lupia T, De Rosa FG. COVID-19 pneumonia: do not leave
the corticosteroids behind! Future Microbiology. 2021;16(5):317-22.

44. Erçen Diken , Yıldırım F, Yıldız Guˍlhan P, Özkaya Ş, Şimşek M, Yuˍcel C et al.
Corticosteroid use in COVID-19 pneumonia. Tuberk Toraks 2021;69(2):217-226.

45. Chrousos GP, Meduri GU. Critical COVID-19 disease, homeostasis, and the "surprise" of
effective glucocorticoid therapy. Clinical Immunology (Orlando, Fla.). 2020;219:108550.

46. Wang Y, Jiang W, He Q, Wang C, Wang B, Zhou P, et al. A retrospective cohort study of
methylprednisolone therapy in severe patients with COVID-19 pneumonia. Signal
transduction and targeted therapy. 2020;5(1):1-3.

47. Salama C, Han J, Yau L, Reiss WG, Kramer B, Neidhart JD, et al. Tocilizumab in patients
hospitalized with Covid-19 pneumonia. New England Journal of Medicine. 2021;384(1):20-
30.

48. Capra R, De Rossi N, Mattioli F, Romanelli G, Scarpazza C, Sormani MP, et al. Impact of low
dose tocilizumab on mortality rate in patients with COVID-19 related pneumonia. European
journal of internal medicine. 2020;76:31-5.

49. Gupta S, Wang W, Hayek SS, Chan L, Mathews KS, Melamed ML, et al. Association
between early treatment with tocilizumab and mortality among critically ill patients with
COVID-19. JAMA internal medicine. 2021;181(1):41-51.

50. Bikdeli B, Madhavan MV, Jimenez D, Chuich T, Dreyfus I, Driggin E, et al. COVID-19 and
thrombotic or thromboembolic disease: implications for prevention, antithrombotic therapy,
and follow-up: JACC state-of-the-art review. Journal of the American college of cardiology.
2020;75(23):2950-73.

51. Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L, et al. Features of 20
133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation
Protocol: prospective observational cohort study. bmj. 2020;369.

665
666
667

668   **Table 1. The Characteristics of Patients with COVID-19**

Characteristics	n(%)
Age (mean ±SD)	59.2 ±17.3
Gender	
Male/Female	5507/5683
Symptoms	
Asymptomatic	2460(21.9%)
Cough	6870(54.5%)
Pain	5370(47.9%)
Dyspnea	4140(36.9%)
Fever	3540(31.6%)
Pneumonitis	5061 (45.2%)
Patient groups	
Group A	9294 (83,05%)
Group B	1376(12.29%)
Group C	520(4.64%)
Hospitalized patients	1896(16.9%)
in first 10 days	1220(64.3%)
after 10 days	676 (35.7%)
Respiratory Failure	520 (27.4%)

NIMV + IMV**	*146(7.7%)*
Mortality	*43(2.2%)*

Figure 1 consists of 7 cases with variable stages of COVID-19 and examples of early and severe

endotheliitis in brain, pulmonary and skin tissues. Histopathological and radiological findings of

pneumonitis changed depending on the phase of the disease (early, progressive, severe and dissipative,

respectively), which leads to a divergence in treatment groups.

Examples of patients who were convenient for **Group A** *treatment are depicted in sections* **I-II**.

Figure 1-I. Case 1. HRCT images showed patchy, subpleural and peripheral perivascular ground-

glass opacities, corresponding the early phase of pneumonitis. She received Group A treatment. GGOs

in transaxial images were located in the subpleural area, where the microvascular *endoteheliitis* and

endothelial destructions mostly likely occur, due to the interaction with toxic plasma in the capillaries

of pulmonary interstitial space.

Figure 1-II. Case 2. HRCT images showed characteristics of an early phase COVID-19 pneumonitis

(end-capillary microangiopathy, explained in Fig.1-I) and perivascular consolidation. It appeared as

the continuation of crazy-paving pattern, which was demonstrated by the thickening in inter-lobular

and intra-lobular septa. Lung involvement was limited and monofocal. Hence, the patient received

Group A treatment.

An example of a patient who was convenient for **Group B** *treatment is depicted in section* **III**.

Figure 1-III. Case 3. Transaxial HRCT image in the first day of positivity shows bilateral and patchy

nodulary GGOs as expected in early phase **(IIIa)**. At the 5th day of positivity, affected pulmonary

areas were advanced into scattered consolidations **(IIIb)**. This appearance was noted as the

progressive phase of pneumonitis and was considered as representation of clinical deterioation

clinically (i.e. dyspnea, respiratory failure). On the 15[th] day of positivity, fibroreticular consolidations

were conspicuous (**IIIc**). The dissipative phase was the healing process, characterized by the

resolution in lung parenchyma and residual GGO, observed after 35 days of positivity (**IIId**).

Parenchymal bands, originated from previous fibroreticular proliferation, were also visible (**IIId**).

*If the patients have a tendency of severe phase and/or unresponsiveness to Group B therapy, **Group C***

*treatment was used. Representative cases of this group are shown in sections **IV-V-VI-VII**. This group*

of patients also undergo early and progressive phases of COVID-19 but they also showed further

deterioration of lung involvement and endothelial damage, therefore, they received the last group of

*treatment. For the description of early and progressive phases, sections **I, II** and can be referred.*

Figure 1-IV. Case 4. Coronal, sagittal and axial planes of HRCT images initially showed small GGO

with a local subpleural sparing, particularly around the microvascular area on the first day of positivity

(**IVa**). In the 5[th] day of treatment, increased GGOs were visible in the progressive phase (**IVb**). After

18 days of positivity, pneumomediastinum, Diffuse Alveolar Damage (DAD) and ARDS were seen in

the severe phase (**IVc**). Lastly, the dissipative phase was seen, after 30 days of positivity, with residual

fibrotic parenchyma (**IVd**).

Figure 1-V. **Case 5.** The first HRCT image, on the 2[nd] day of positivity, showed COVID-19-related

bilateral and multifocal nodular GGOs (**Va**). After 10 days of positivity, DAD developed along with

ARDS and pneumomediastinum characterized with the severe phase (**Vb**). She recovered and

discharged after 30 days of positivity with the dissipative phase (**Vc**). The regression in pulmonary

lesions was visible on the HRCT image two months after diagnosis (**Vd).**

Figure 1-VI. Case 6. HRCT image showed moderate pneumonia in the early phase (**VIa**). After 12

days of positivity, the severe phase develops with DAD and ARDS (**VIb**). In the 25[th] day of

positivity, recovery was observed in the dissipative phase (**VIc**). The extent of improvement in

pulmonary lesions can be noticed in **VId,** which was 40 days after the diagnosis.

Figure 1-VII. Case 7. First HRCT showed ARDS pattern with dense consolidations (**VIIa**).

Pneumothorax developed after two weeks from diagnosis (**VIIb**). In addition to the respiratory failure,

hemorrhagical intracranial areas were seen in the T2-weighted MRI. (**VIIc**). He was lost at the 20[th]

day of positivity.

*Representative histopathological and radiological images of endotheliitis **is seen in section VIII, IX,***

X, XI.

Figure 1-VIII Haematoxylin&Eosin-stained sections from representative areas of lung parenchyma
were seen with the mixt-type inflammatory-cell infiltration of lung tissue and exudative capillaritis
with thickened microvascular walls. Interstitial and intra-alveolar proliferation of fibroblasts are noted.
742
743   **Figure 1-IX. Diffusion MRI of** head image of COVID-19 positive patient showing characteristic
744   COVID-19 related *endotheliitis* causing characteristic lesions and intracranial hemorrhage.
745
746   **Figure 1-X** Early endotheliitis.
747   **Figure 1-XI** Late endotheliitis.
748   **Figure 1-XII** Papulovesicular eruptions as the skin manifestations of COVID-19 related endotheliitis.
749
750