Sex-specific epidemiological and clinical characteristics of Covid-19 patients in the southeast region of Bangladesh

Md. Aminul Islam[a,1], Abdullah Al Marzan[b], Md. Sydul Islam[c,1], Saima Sultana[d], Md. Iftakhar Parvej[e], Mohammad Salim Hossain[f], Mohammad Tohid Amin[f], Farzana Ehetasum Hossain[a], Md Abdul Barek[f], Md. Shafiul Hossen[f], Md. Shariful Islam[a], Foysal Hossen[a], Newaz Mohammed Bahadur[c], Md. Shahadat Hossain[g], Md. Didar-ul Alam[h], Firoz Ahmed[a,1,*]

[a]Department of Microbiology, Noakhali Science and Technology University, Noakhali-3814,Bangladesh

[b]Department of Biochemistry and Molecular Biology, Shahjalal University of Science and Technology, Sylhet-3114, Bangladesh

[c]Department of Applied Chemistry and Chemical Engineering, Noakhali Science and Technology University, Noakhali-3814, Bangladesh

[d]Quality Assurance & Management, Marie Stopes Bangladesh, Dhaka-1207, Bangladesh

[e]Department of Statistics, Noakhali Science and Technology University, Noakhali-3814, Bangladesh

[f]Department of Pharmacy, Noakhali Science and Technology University, Noakhali-3814, Bangladesh

[g]Department of Biotechnology and Genetic Engineering, Noakhali Science and Technology University, Noakhali-3814, Bangladesh

[h]Professor and Honorable Vice-Chancellor, Noakhali Science and Technology University, Noakhali-3814, Bangladesh

[1]These authors contributed equally to this work as first authors.

*Corresponding Author: Dr. Firoz Ahmed
                       Professor and Chairman & Focal Point
                       COVID-19 Diagnostic Lab, Department of Microbiology, Noakhali
                       Science and Technology University, Noakhali-3814, Bangladesh
                       Email: firoz@nstu.edu.bd; firoz19701016@gmail.com
                       Mobile: +8801713195658

Abstract

Purpose: The present study aimed to compare and analyze the sex-specific epidemiological,

clinical characteristics, comorbidities, and other information of confirmed COVID-19 patients

from the southeast region in Bangladesh for the first time.

Methods: 385 lab-confirmed cases were studied out of a total of 2471 tested samples between

June 5 and September 10, 2020. RT-PCR was used for COVID-19 identification and SPSS

(version 25) for statistical data analysis.

Results: We found that male patients were roughly affected compared to females patients (male

74.30% vs. female 25.7%) with an average age of 34.86 ± 15.442 years, and B (+ve) blood group

has been identified as a high-risk factor for COVID-19 infection. Workplace, local market, and

bank were signified as sex-specific risk zone ($p < 0.001$). Pre-existing medical conditions such as

diabetes, hypertension, cardiovascular and respiratory diseases were identified among the

patients. Less than half of the confirmed COVID-19 cases in the southeast region were

asymptomatic (37.73%) and more prevalent among females than males (male vs. female: 36.84%

vs. 40.51%, $p = 0.001$).

Conclusions: The findings may help health authorities and the government to take necessary

steps for identification and isolation, treatment, prevention, and control of this global pandemic.

Keywords: COVID-19, Coronavirus disease, Epidemiological, Clinical features, Asymptomatic,

Comorbidities

1 Introduction

COVID-19 (Coronavirus disease-2019) was first identified in December 2019 in a person at a seafood market in Wuhan city, in Hubei province, China (Chan et al., 2020). It is caused by an unknown pathogenic virus, and somedays later, scientists isolated this new virus from the lower respiratory tract of patients. This virus is a new type of Beta coronavirus, corresponds to a comparatively noted Coronaviridae family, and is identical to Bat SARS-type coronavirus (having 88% identity) and have identity nucleotide with the actual severe acute respiratory syndrome (SARS) (80%) and Middle East respiratory syndrome coronavirus (MERS-CoV) (about 50%) epidemic virus (Zheng, 2020). The pathogen of this disease has been confirmed by molecular methods as a new coronavirus and was primarily named a 2019 novel coronavirus (2019-nCoV). World Health Organization (WHO) declared the COVID-19 outbreak as the sixth public health emergency of international concern on 30 January 2020 and declared a fresh name, called Corona Virus Disease (COVID-19) on February 11, 2020, for the pandemic disease: Corona Virus Disease (COVID-19) (Lai et al., 2020).

On the grounds of taxonomy, phylogeny, and developed practice, the International Committee on Taxonomy of Viruses (ICTV) announced the official name of this coronavirus as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) on 12 February 2020 (Lai et al., 2020). Since the disease's occurrence, it is become a global threat and spreading quickly to provinces in China and more than 210 countries, including Bangladesh, worldwide (Worldometers, n.d.). By January 2, 2020, 41 lab-confirmed COVID-19 patients had been detected from hospitals and less than 50% of them had implicit illness including high blood pressure, hyperglycemia, and cardiovascular diseases (C. Huang et al., 2020). As of 22 January 2020, there were 571 registered

cases of 2019-new coronaviruses (COVID-19) in 25 provinces in China and 17 deaths until 22

January 2020 (Lu, 2020). Now, 4634 patients died up to 2 July 2020 in China (Worldometers,

n.d.).

COVID-19 infects more than ten million patients and about 516558 (8%) died on 2 July 2020

worldwide. USA has accounted for the highest number of COVID-19 patients(2758044),

followed by Brazil (1426913), Russia (654405), and the UK (313483) on 2 July 2020. Every

country of South Asia is affected by this zoonotic virus. About 604808 cases in India have been

identified, one of the top five affected countries (Worldometers, n.d.). Bangladesh announced the

first confirmed coronavirus cases on 8 March 2020, and this number increased day by day

(Anwar et al., 2020). It was reported that about 512,496 cases were identified, 7,531 patients

died, and 456,070 patients are recovered by 30 Dec 2020 (Worldometers, n.d.). It is now

spreading all over the country and the highest affected area was observed in the Dhaka division.

Based on district-wise case distribution, around 5546 and 2309 patients were identified and

5058 and 2232 patients were recovered in Noakhali and Lakshmipur, respectively, on 31 Dec

2020 (Civil Surgeon Office, Noakhali, and Laxmipur, 2020, Corona Information, n.d.). It

diffuses via people-to-people transmission, for instance, sneezing and coughing while some

research indicates the transmission chances even amongst the asymptomatic (Ningthoujam,

2020). It can also spread through stool, sweat, urine, and respiratory secretions.

The general incubation period of this virus is 1 to 14 days (Baud et al., 2020), however, a longer

latent period has been reported in some cases, which makes it more dangerous as the host may

transmit the disease to others unknowingly (Kong, 2020). The most frequently occurring

symptoms are fever, cough, sore throat, tiredness, ache, and headache, etc. Still, severe illnesses

including- diarrhea, pneumonia, pulmonary edema, acute respiratory distress syndrome, multiple

organ failure are also found among patients (Chen et al., 2020; Y. Meng et al., 2020). Both males

and females are affected by COVID-19, but several epidemiological and clinical studies suggest

that males are more progressive to infection than females (Barek et al., 2020; Y. Meng et al.,

2020). Many studies have already found that single or multiple comorbidities like high blood

pressure, cardiovascular disease, high blood pressure, liver problem, diabetes, lung disease, and

kidney disease patients are more affected by this virus (Barek et al., 2020; Emami et al., 2020).

Researchers are trying to develop medicine and vaccine to combat COVID-19 and upcoming

SARS and MERS viruses (Amanat and Krammer, 2020). Now, public concern and

consciousness are the main prohibitions for the COVID-19 outbreak.

At present, epidemiological, clinical characteristics, the pervasiveness of comorbidities, other

information is most important to take necessary steps for treatment, prevention, and control of

the current outbreak. There have no published demographic and clinical characteristics study

data regarding COVID-19 patients of Bangladesh. Considering the above-mentioned facts, we

analyzed 385 laboratory-confirmed COVID-19 positive patients' demographic data, clinical

manifestations, medical history, blood group, travel history, previous vaccination, and other data

from the southeast region in Bangladesh for the very first time which help to proper decisions to

reduce the risk of infection.

2 Methods

2.1 Study design and participants

We selected only laboratory-confirmed Covid-19 cases between 5 June 2020 to 10 September 2020 who were tested at Noakhali Science and Technology University COVID-19 testing laboratory, Noakhali, Bangladesh approved by the government (The Daily Star, 2020). The specimen was collected by nasopharyngeal swab, and reverse transcriptase-polymerase chain reaction (RT-PCR) assay was used for Covid-19 confirmation. Three hundred eighty-five patients (aged above 5 years) were identified as positive from the southeast region, Bangladesh (Noakhali and Lakshmipur) in where 286 patients were male, and 99 were female. Pregnant women, children, adults, and older patients were included in this study. To prevent the chance of bias, participants were selected and listed with the help of the Noakhali Science and Technology University Covid-19 testing laboratory database.

2.2 Data collection

The questionnaire was adopted from formerly published studies, and our research team added, modified, and developed some questions. There were various sections in the survey including demographic information, clinical characteristics, pre-existing medical condition, blood group, earlier visited information, vaccination, etc. The investigators were told that their participation was anonymous and entirely voluntary, and there was no reward for taking part. All the participants willingly participated in this work by giving written permission. They were invited to complete the questionnaire. We were present on hand to answer questions or clarify any doubts that they might have. All filled questionnaires were collected by us one by one.

2.3 Definition

Patients with SARS-CoV-2 infection have no clinical signs and symptoms were defined as

asymptomatic cases, and symptomatic patients were defined with SARS-CoV-2 infection

presenting with clinical characteristics from medical interviews. Comorbidities were ascertained

from physician documentation.

2.4 Specimen collection and RT-PCR assay for SARS-CoV-2

We detected the presence of SARS-CoV-2 by the qPCR method with the novel coronavirus

(2019-nCoV) nucleic acid diagnostic Kit (PCR-fluorescence probing). The test utilizes the novel

coronavirus (2019-nCoV) ORF 1ab and the specific conserved sequence of coding nucleocapsid

protein N gene as the target regions to detect sample RNA through fluorescence signal changes.

These diagnostic criteria were based on the recommendations by the Institute of Epidemiology,

Disease Control and Research (IEDCR), Bangladesh. Test facility and quality reconfirmed by

IEDCR.

2.5 Statistical Analysis

Descriptive statistics were used to summarize the data, continuous variables were expressed as

mean ±SD, and categorical variables were summarized as counts and percentages. There are

several kinds of models for analyzing the relationship between the data sets. As our data are

parametric, we used Pearson's Correlation Coefficient (2-tailed). All statistical analysis was

conducted using the IBM Statistical Package for the Social Sciences (SPSS), version 25.0

software. $P < 0.05$ was considered statistically significant. Correlation between Sex and many

other factors i.e. BMI, age, comorbidities, etc. had been analyzed and determined.

3 Results

The demographic characteristics of all the confirmed COVID-19 cases (N=385) were shown in **Table 1**. Of them, 74.3% of cases were male and 25.7% were female with an average age of 34.86 ± 15.442 years. The highest number of patients was in the age range of 21-50 years (71.91%). However, around 15% of cases were found older than them. As for Body Mass Index (BMI), 23.80% of cases were overweight, and 8.10% were obese, while only 3.40% were underweight. A (+ve), B (+ve), and O (+ve) blood groups were the most identified blood group with a percentage of 25.46%, 32.52%, and 18.40%, respectively. In the meanwhile, only a few percent of patient cases were observed with O (-ve) (3.07%), B (-ve) (3.68%), and A (-ve) (3.37%) blood group. Of the male with a positive blood group, the most common blood group was B (+ve) (35.22%), followed by A (+ve) (22.76%) and AB (+ve) (16.46%). In the female positive blood group, the most typical blood group was still B (+ve) and O (+ve) (24.05%) in each case, followed by A (+ve) (21.05%) and AB+ (15.79%) (**Figure 1**).

About 28.50% of cases passed the higher secondary certificate (HSC) level, while only 9.40% continued their education until the secondary school certificate (SSC) level. Graduated (25.50%) and post graduated (11.75%) cases were also observed in our study. Notably, most of the COVID-19 patients were businessmen (18.80%), students (15.43%), civil servants (14.09%), and health workers (8.05%). In male cases, 23.04% of cases were businessmen, followed by 17.83% civil servants and 8.26% health workers. However, a two-thirds portion of the female cases (61.76%) were housewives, followed by 20.59% of students (**Figure 2**). All of the confirmed cases ensured that they did not travel to any other countries within the nearest time before being infected. However, most of them visited their workplace (65.95%), local market (45.39%), and

health care center (13.80%) within the earlier time. About 19.63% and 18.09% of cases visited

the bank and relative house, respectively. Our study also explored that only 15.50% of infected

persons had clear contact with confirmed cases as far as they know. Additionally, diabetes

(5.19%), hypertension (5.97%), asthma (7.53%), allergy (0.52%) cancer (0.78%), pregnancy

(2.34%) and others (1.29%) were the most common comorbidities identified among the

confirmed cases. Most of the patients were vaccinated with BCG (86.2%), Polio (87.9%),

Mumps (54%), Tetanus (82.5%). However, only a few of them were vaccinated with Hepatitis B

(14.8%), Chickenpox (15.1%), and Measles (3%).

Analysis of demographic characteristics based on gender-specific is explored in **Table 2**. This

analysis showed that BMI, and earlier visited places showed significant ($p < 0.05$) differences

between male and female patients. The highest percent of both males (74.24%) and females

(68.1%) were in the age range of 21-50 years. About 88.66% male cases and 92.40% female

cases had positive Rh factor ($p = 0.115$). Additionally, earlier visited place such as workplace (p

< 0.001), bank ($p < 0.001$), local market ($p < 0.001$), and social gathering ($p < 0.001$) were also

reported as the significant risk factors for sex-specific COVID-19 infection. Among all

comorbidities, our study found significant differences between males and females in the case of

only chickenpox ($p = 0.014$).

Signs and symptoms identified during testing among COVID-19 confirmed cases are tabulated in

Table 3. Our study also confirmed the sex-specific clinical characteristics of COVID-19. This

present study found that less than half of the confirmed COVID-19 cases (37.73%) were

symptomatic, and this asymptomatic characteristic was more prevalent among females (40.51%)

than that of males (36.84%) patients ($p = 0.001$). Conversely, mortality cases were found higher

for males than females with no significant correlation ($p = 0.724$). Of the patients showing

symptoms, the most frequent symptom was fever (29.45%), followed by cough (34.05%) and

headache (13.50%), with no significant difference between genders. Fever and cough were

observed mainly among females (55.32% and 44.87%) than those of male patients (44.87% and

53.85%) cases. Also, 8.28% of cases and 13.50% of cases experienced sore throat and headache.

Though very few female cases showed fever ($p = 0.240$); however, about 37.82% of male cases

experienced sore throat in comparison to only 25.53% of female cases ($p = 0.204$).

4 Discussion

The new beta coronavirus is the third type of zoonotic coronavirus, the seventh human

coronavirus, and has similarities with before coronavirus (Zhou et al., 2020; Zhu et al., 2020).

This contagious virus shows the same receptor binding domain with SARS coronavirus (SARS-

CoV) and MERS-CoV (Lu et al., 2020). The number of COVID-19 cases continuously rising

worldwide, but no specific treatment has been confirmed to be fully effective against COVID-19

(irrespective of different strains). Therefore, it is crucial to identify the clinical demographical

characteristics, clinical manifestation, comorbidities, and other information of COVID-19

patients. And are more important to help in early detection and isolation of infected individuals,

and also minimize the spread of the disease, severity, and death rate. To our best knowledge, this

is the first study undertaken to investigate the demographic, clinical characteristics,

comorbidities, and excrescent information of the confirmed COVID-19 patients from the

southeast region in Bangladesh.

We analyzed 200 positive samples from Noakhali and 185 positive samples from Lakshmipur,

the southeast region of Bangladesh. In our study, we observed that the number of male patients

was largely (74.3%), almost three times higher than female patients (25.7%). Several studies

conducted in China and other countries corroborated a higher number of infected males than that

of females (Novel, 2020; Z. Wang et al., 2020). Meanwhile, a research study by Zhou *et al.*

suggests that the expression of ACE2, the receptor for COVID-19, is higher in males than that in

females which may be the reason behind the higher proportion of COVID-19 infected males

(Zhou et al., 2020). It is also thought that as females have more X chromosomes and sex

hormones, these factors play an important role in innate and adaptive immunity (Jaillon et al.,

2019). On the other hand, Females are less associated with a bad lifestyle than males. Similarly,

several studies found that males are more infected than females during MERS-CoV and SARS-

CoV pandemic (Badawi and Ryoo, 2016; Channappanavar et al., 2017). Additionally, the

average age of all cases was 34.86 ± 15.442 years old and nearly similar to several studies

(Azlan et al., 2020; Q. Huang et al., 2020; Kim et al., 2020). It might be due to the outgoing of

middle-aged people for their respective work or for their family need as they have to meet all

family and society demands. However, our study indicated that patients at a wide age range can

be infected by SARS-CoV-2. This present study also demonstrated a slightly greater mean BMI

value than those found in the Hubei (Yang et al., 2020) and Jiangsu (R. Huang et al., 2020)

province of China. Notably, more than three fourth of patients involved in our study had a

positive blood group including A (+ve) (25.46%), B (+ve) (32.52%), and O (+ve) (18.40%)

similar to Li *et al.* and Zietz *et al.* studies (Li et al., 2020; Zietz et al., 2020). According to one

study, anti-A antibodies inhibit S protein/angiotensin-converting enzyme 2-dependent adhesion

of these cells to an angiotensin-converting enzyme 2 expressing cell line (Guillon et al., 2008).

Besides, the majority of infected cases were business professionals (18.80%), followed by

students (15.43%), civil servants (14.09%), and health workers (8.05%), which is more in this

study (Azlan et al., 2020; L. Wang et al., 2020). Consistent with the previous study conducted in

Wuhan (D. Wang et al., 2020), this present study also explored males are mainly visited in the

workplace (82.97%, $p < 0.001$) and local market (58.08%, $p < 0.001$). Conversely, males need

to attend their daily work for livelihood and go to the local market for their daily needs. Besides,

both males (12.66%) and females (16.49%) are early visited in the health care center. Noticeably,

no patients had traveled to any other country within the nearest time before getting infected, and

only 15% of cases reported the known contact with confirmed cases. Of the confirmed cases, the

other frequently observed comorbidities were hypertension (male vs. female: 7.29% vs. 2.06%, p

$= 0.060$), diabetes mellitus (male vs. female: 4.86% vs. 6.18%, $p = 0.612$), asthma (male vs.

female: 8.33% vs. 5.15%, $p = 0.306$), cancer (male vs. female: 0.01% vs. 1.00%, $p = 0.745$) and

pregnancy (female: 5.26%, $p < 0.001$). However, they were observed with a lower percentage

than the foregoing observation in China and Italy (Colaneri et al., 2020; R. Huang et al., 2020; H.

Meng et al., 2020). Maximum patients take necessary vaccines such as Polio (87.9%), BCG

(86.2%), Tetanus (82.5%), Mumps (54%) were taken in life. About 37.73% of cases were

asymptomatic confirmed COVID-19 cases in the southeast region ($p = 0.001$). Only 5% of

asymptomatic cases were observed in Beijing (Tian et al., 2020), and this asymptomatic

characteristic was more prevalent among females (40.51%) than that of male (36.84%) patients

($p = 0.001$), which is similar to our findings. We recorded all the identified signs and symptoms

of the symptomatic cases. Out of these, the most dominant symptom was cough (34.05%),

followed by fever (29.45%). Other symptoms of fatigue (21.78%), headache (13.50%), sore

throat (8.28%), ache (12.58%), and diarrhea (3.07%) were also found among patients. In the

meanwhile, several studies conducted in Wuhan (Y. Meng et al., 2020), Shanghai (Yu et al.,

2020), Beijing (Niu et al., 2020), Hubei (Yang et al., 2020), Jiangsu (R. Huang et al., 2020), and

United States (Aggarwal et al., 2020) found that fever and cough as dominant clinical symptoms.

However, 29.45% of our study's symptomatic patients showed fever that was comparatively

higher than that in South Korea (11.6%) (Kim et al., 2020)**.**

There have no published studies on sex-specific demographic, clinical characteristics, and

comorbidity of COVID-19 infected patients, Bangladesh. This is the first cross-sectional study

and the most comprehensive assessment and robust evidence to date of patients' clinical

characteristics and comorbidities with COVID-19. Albeit this is a novel study, several limitations

might be noted in the present study. Firstly, the present study was performed only in a single

institution obtaining data via face-to-face information when patients came for coronavirus

testing. Hence, the represented data does not give the whole scenario of all COVID-19 patients

of the country. Secondly, the limited number of study populations, especially for female patients.

Finally, we did not include any data from hospitalized patients and laboratory outputs. However,

our study analyzed the demographic and clinical characteristics of COVID-19 patents that aid in

identifying possible risk factors and reducing the risk of COVID-19 susceptibility to control this

outbreak.

5 Conclusions

Nowadays, COVID-19 is a contagious disease and led to a major global health concern. In the

present study, we found that males were more affected than female patients, and middle-aged

people were mainly affected in both sexes. Fever, cough, and tiredness were the common

symptoms found among patients, and several comorbidities (diabetes, hypertension, and asthma)

were present among patients. Maximum COVID-19 patents were B (+ve), and businessmen were

mostly affected by this infectious virus. We also observed that male patients were mainly

affected by the coronavirus may due to the earlier visit to the workplace, local-market, and bank.

This study raises the knowledge about COVID-19 and helps to take effective and proper

decisions to identify risk factors to control this pandemic.

Funding disclosure

This work was supported by the Research Cell, Noakhali Science and Technology University,

Bangladesh.

Ethical approval

The study protocol was approved by the institutional ethical clearance committee of Noakhali

Science and Technology University, Bangladesh (Approval ID: 44-2020).

Acknowledgments

We acknowledge the sincere help extended by several COVID-19 Diagnostic Lab Volunteers to

name a few are: Md. Sahedul Islam, Md. Amzad Hossain, Mahmudul Islam Rakib, Golam

Shamdani, Amor Chandra Nath, and Md. Julker Nyne, of Noakhali Science and Technology

University, Bangladesh.

References

Aggarwal S, Garcia-Telles N, Aggarwal G, Lavie C, Lippi G, Henry BM. Clinical features,

laboratory characteristics, and outcomes of patients hospitalized with coronavirus disease 2019

(COVID-19): Early report from the United States. Diagnosis 2020;7:91–6.

14

Amanat F, Krammer F. SARS-CoV-2 Vaccines: Status Report. Immunity 2020;52:583–9.

https://doi.org/10.1016/j.immuni.2020.03.007.

Anwar S, Nasrullah M, Hosen MJ. COVID-19 and Bangladesh: Challenges and How to Address

Them. Front Public Heal 2020;8:154. https://doi.org/10.3389/fpubh.2020.00154.

Azlan AA, Hamzah MR, Sern TJ, Ayub SH, Mohamad E. Public knowledge, attitudes and

practices towards COVID-19: A cross-sectional study in Malaysia. PLoS One 2020;15.

https://doi.org/10.1371/journal.pone.0233668.

Badawi A, Ryoo SG. Prevalence of comorbidities in the Middle East respiratory syndrome

coronavirus (MERS-CoV): a systematic review and meta-analysis. Int J Infect Dis 2016;49:129–

33.

Barek MA, Aziz MA, Islam MS. Impact of age, sex, comorbidities and clinical symptoms on the

severity of COVID-19 cases: A meta-analysis with 55 studies and 10014 cases. Heliyon

2020;6:e05684. https://doi.org/10.1016/j.heliyon.2020.e05684.

Baud D, Qi X, Nielsen-Saines K, Musso D, Pomar L, Favre G. Real estimates of mortality

following COVID-19 infection. Lancet Infect Dis 2020.

Chan JF-W, Yuan S, Kok K-H, To KK-W, Chu H, Yang J, et al. A familial cluster of pneumonia

associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of

a family cluster. Lancet 2020;395:514–23.

Channappanavar R, Fett C, Mack M, Ten Eyck PP, Meyerholz DK, Perlman S. Sex-based

differences in susceptibility to severe acute respiratory syndrome coronavirus infection. J

Immunol 2017;198:4046–53.

Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical

characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive

study. Lancet 2020;395:507–13.

Civil Surgeon Office, Noakhali, and Laxmipur. 2020.

Colaneri M, Sacchi P, Zuccaro V, Biscarini S, Sachs M, Roda S, et al. Clinical characteristics of

coronavirus disease (COVID-19) early findings from a teaching hospital in Pavia, North Italy, 21

to 28 February 2020. Eurosurveillance 2020;25. https://doi.org/10.2807/1560-

7917.ES.2020.25.16.2000460.

Corona Information. Corona News n.d. https://corona.gov.bd/corona-news.

Emami A, Javanmardi F, Pirbonyeh N, Akbari A. Prevalence of Underlying Diseases in

Hospitalized Patients with COVID-19: a Systematic Review and Meta-Analysis. Arch Acad

Emerg Med 2020;8:e35. https://doi.org/10.22037/aaem.v8i1.600.

Guillon P, Clément M, Sébille V, Rivain JG, Chou CF, Ruvoën-Clouet N, et al. Inhibition of the

interaction between the SARS-CoV Spike protein and its cellular receptor by anti-histo-blood

group antibodies. Glycobiology 2008;18:1085–93. https://doi.org/10.1093/glycob/cwn093.

Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with

2019 novel coronavirus in Wuhan, China. Lancet 2020;395:497–506.

https://doi.org/10.1016/S0140-6736(20)30183-5.

Huang Q, Deng X, Li Y, Sun X, Chen Q, Xie M, et al. Clinical characteristics and drug therapies

in patients with the common-type coronavirus disease 2019 in Hunan, China. Int J Clin Pharm

2020:1.

Huang R, Zhu L, Xue L, Liu L, Yan X, Wang J, et al. Clinical findings of patients with

coronavirus disease 2019 in Jiangsu province, China: A retrospective, multi-center study. PLoS

Negl Trop Dis 2020;14:e0008280.

Jaillon S, Berthenet K, Garlanda C. Sexual dimorphism in innate immunity. Clin Rev Allergy

Immunol 2019:1–14.

Kim GU, Kim MJ, Ra SH, Lee J, Bae S, Jung J, et al. Clinical characteristics of asymptomatic

and symptomatic patients with mild COVID-19. Clin Microbiol Infect 2020;26:948.e1-948.e3.

https://doi.org/10.1016/j.cmi.2020.04.040.

Kong T. Longer incubation period of coronavirus disease 2019 (COVID-19) in older adults.

AGING Med 2020;3:102–9. https://doi.org/10.1002/agm2.12114.

Lai C-C, Shih T-P, Ko W-C, Tang H-J, Hsueh P-R. Severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): the epidemic and the

challenges. Int J Antimicrob Agents 2020:105924.

Li J, Wang X, Chen J, Cai Y, Deng A, Yang M. Association between ABO blood groups and

risk of SARS-CoV-2 pneumonia. Br J Haematol 2020;190:24–7.

https://doi.org/10.1111/bjh.16797.

Lu H. Drug treatment options for the 2019-new coronavirus (2019-nCoV). Biosci Trends

2020;14:69–71.

Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and epidemiology of

2019 novel coronavirus: implications for virus origins and receptor binding. Lancet

2020;395:565–74.

Meng H, Xiong R, He R, Lin W, Hao B, Zhang L, et al. CT imaging and clinical course of

asymptomatic cases with COVID-19 pneumonia at admission in Wuhan, China. J Infect

2020;81:e33–9. https://doi.org/10.1016/j.jinf.2020.04.004.

Meng Y, Wu P, Lu W, Liu K, Ma K, Huang L, et al. Sex-specific clinical characteristics and

prognosis of coronavirus disease-19 infection in Wuhan, China: A retrospective study of 168

severe patients. PLoS Pathog 2020;16:e1008520.

Ningthoujam R. COVID 19 can spread through breathing, talking, study estimates. Curr Med

Res Pract 2020;10:132–3. https://doi.org/10.1016/j.cmrp.2020.05.003.

Niu S, Tian S, Lou J, Kang X, Zhang L, Lian H, et al. Clinical characteristics of older patients

infected with COVID-19: A descriptive study. Arch Gerontol Geriatr 2020;89.

https://doi.org/10.1016/j.archger.2020.104058.

Novel CPERE. The epidemiological characteristics of an outbreak of 2019 novel coronavirus

diseases (COVID-19) in China. Zhonghua Liu Xing Bing Xue Za Zhi= Zhonghua

Liuxingbingxue Zazhi 2020;41:145.

The Daily Star. Coronavirus Test Lab Ina Noakhali Sci Technol Univ 2020.

Tian S, Hu N, Lou J, Chen K, Kang X, Xiang Z, et al. Characteristics of COVID-19 infection in

Beijing. J Infect 2020;80:401–6. https://doi.org/10.1016/j.jinf.2020.02.018.

Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized

patients with 2019 novel coronavirus–infected pneumonia in Wuhan, China. Jama

2020;323:1061–9.

Wang L, Duan Y, Zhang W, Liang J, Xu J, Zhang Y, et al. Epidemiologic and clinical

characteristics of 26 cases of COVID-19 arising from patient-to-patient transmission in

Liaocheng, China. Clin Epidemiol 2020;12:387.

Wang Z, Chen X, Lu Y, Chen F, Zhang W. Clinical characteristics and therapeutic procedure for

four cases with 2019 novel coronavirus pneumonia receiving combined Chinese and Western

medicine treatment. Biosci Trends 2020.

Worldometers. Confirmed Cases Deaths by Country, Territ or Convey n.d.

https://www.worldometers.info/coronavirus/.

Yang K, Sheng Y, Huang C, Jin Y, Xiong N, Jiang K, et al. Clinical characteristics, outcomes,

and risk factors for mortality in patients with cancer and COVID-19 in Hubei, China: a

multicentre, retrospective, cohort study. Lancet Oncol 2020.

Yu X, Sun X, Cui P, Pan H, Lin S, Han R, et al. Epidemiological and clinical characteristics of

333 confirmed cases with coronavirus disease 2019 in Shanghai, China. Transbound Emerg Dis

2020.

Zheng J. SARS-CoV-2: an emerging coronavirus that causes a global threat. Int J Biol Sci

2020;16:1678.

Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak

associated with a new coronavirus of probable bat origin. Nature 2020;579:270–3.

Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients with

pneumonia in China, 2019. N Engl J Med 2020.

Zietz M, Zucker J, Tatonetti NP. Associations between blood type and COVID-19 infection,

intubation, and death. Nat Commun 2020;11:1–6.

Tables

408 **Table 1: Demographic characteristics of COVID-19 cases from the southeast region in**

409 **Bangladesh**

Characteristics	Response	Patients
Sex (N = 385)	Male	286 (74.3%)
	Female	99 (25.7%)
Age (years) (N=381)	Mean ± SD	34.86 ± 15.442
Age group	0-20	50(13.12%)
	21-50	274 (71.91%)
	>51	57 (14.96%)
BMI (N= 298)	Mean ± SD	24.1012 ± 4.117
BMI range	<18.5 (Underweight)	10 (3.40%)
	18.5-24.9 (Normal)	193 (64.7%)
	25-29.9 (Overweight)	71 (23.80%)
	>30 (Obese)	24 (8.10%)
Blood group (N=326)		
	A (+ve)	83(25.46%)
	B (+ve)	106 (32.52%)
	AB (+ve)	43 (13.19%)
	O (+ve)	60 (18.40%)
	A (-ve)	11 (3.37%)
	B (-ve)	12 (3.68%)
	AB (-ve)	1 (0.31%)
	O (-ve)	10 (3.07%)
Educational status (N= 298)		
	Masters	35 (11.75%)
	Graduate	76 (25.50%)
	Degree	8 (2.68%)
	HSC	85 (28.50%)
	SSC	28 (9.40%)
	Below SSC	46 (15.44%)
	No education	20 (6.73%)
Occupation (N=298)		
	Civil servant	42 (14.09%)
	Health worker (doctor, nurse, sample collector)	24 [7,8,9] (8.05%) [2.68%,2.35%,3.02%]
	Engineer	1 (0.33%)
	Banker	8 (2.68%)
	Private job	43 (14.43%)
	Businessmen	56 (18.80%)
	Homemaker	42 (14.09%)

	Student	46 (15.43%)
	Others	36 (12.08%)
Traveling to other countries (N=298)		
	Yes	6 (2.01%)
	No	292 (97.99%)
Earlier visited places (N=326)		
	Workplace	215 (65.95%)
	Bank	64 (19.63%)
	Local market	148 (45.39%)
	Relatives house	59 (18.09%)
	Social gathering	36 (11.04%)
	Health care center	45 (13.80%)
Comorbidities (N= 385)		
	Diabetes	20 (5.19%)
	Hypertension	23 (5.97%)
	Asthma	29(7.53%)
	Allergy	2(0.52%)
	Cancer	3(0.78%)
	Pregnancy	9 (2.34%)
	Others	5 (1.29%)
	No risk factors	296 (76.35%)
Vaccine in life (N=298)		
	BCG	257 (86.2%)
	Polio	262 (87.9%)
	Hepatitis B	44 (14.8%)
	Mumps	161 (54.00%)
	Tetanus	246 (82.5%)
	Chickenpox	45 (15.1%)
	Measles	9 (3.00%)
Death Case		
Age (years) (N=130)	Mean ± SD	62.56 ± 14.252
Age group	0-20	1 (0.77%)
	21-50	23 (17.70%)
	>51	106 (81.54%)

412 **Table 2: Sex-specific analysis of demographic characteristics of confirmed COVID-19 cases**

Category	Male	Female	*p*-value
Age group (Years) (M=284, F= 97)			
0-20	30 (10.56%)	20 (20.6%)	
21-50	208 (73.24%)	66 (68.1%)	0.275
>51	46 (16.20%)	11 (11.3%)	
BMI range (M=230, F= 68)			
<18.5 (Underweight)	6 (2.61%)	4 (5.9%)	
18.5-24.9 (Normal)	152 (66.09%)	41 (60.29%)	< 0.001**
25-29.9 (Overweight)	52 (22.60%)	19 (27.94%)	
>30 (Obese)	17 (8.70%)	4 (5.88%)	
Rh factor (M=247, F=79)			
Positive	219 (88.66%)	73 (92.40%)	
Negative	28 (11.34%)	6 (7.60%)	0.115
Traveling to other countries (M= 230, F= 68)			
Yes	5 (2.17%)	1 (1.47%)	0.769
No	225 (97.83%)	67 (98.53%)	
Earlier visited places (M=229, F=97)			
Workplace	190 (82.97%)	25 (25.77%)	< 0.001**
Bank	55 (24.01%)	9 (9.28%)	< 0.001**
Local market	133 (58.08%)	15 (15.46%)	< 0.001**
Relatives house	38 (16.60%)	21 (21.65%)	0.404
Social gathering	34 (14.85%)	2 (2.06%)	< 0.001**
Health care center	29 (12.66%)	16 (16.49%)	0.400
Comorbidities (M= 288, F=97)			
Diabetes	14 (4.86%)	6 (6.18%)	0.612

Hypertension	21 (7.29%)	2 (2.06%)	0.060
Asthma	24 (8.33%)	5 (5.15%)	0.306
Cancer	2 (0.01%)	1 (1.00%)	0.745
Pregnancy	---	9 (9.27%)	< 0.001**
Others	3 (1.04%)	2 (2.06%)	0.444
No risk factors	224 (77.78%)	72 (74.22%)	---
Vaccine in life (M=230, F=68)			
BCG	81 (100.00%)	55 (80.9%)	0.084
Polio	81 (100.00%)	59 (86.8%)	0.519
Hepatitis B	12 (14.81%)	14 (20.6%)	0.159
Mumps	77 (95.06%)	40 (58.8%)	0.551
Tetanus	75 (92.59%)	54 (79.4%)	0.318
Chickenpox	1 (1.23%)	17 (25.00%)	0.014*
Measles	7 (3.04%)	2 (2.94%)	0.928
Diphtheria	196 (85.22%)	55(80.9%)	0.269

Table 3: Sex-specific signs and symptoms of COVID-19 patients identified during testing

Items	Total (%)	Male (%)	Female (%)	*p*-value
Asymptomatic	123 (37.73%)	91 (36.84%)	32 (40.51%)	0.001**
Symptomatic	203 (62.27%)	156 (63.16%)	47 (59.49%)	
Symptoms **(M=156, F=47)**				
Fever	96 (29.45%)	70 (44.87%)	26 (55.32%)	0.240
Cough	111 (34.05%)	84 (53.85%)	27 (57.45%)	0.666
Sore throat	27 (8.28%)	22 (14.10%)	5 (10.64%)	0.542
Headache	44 (13.50%)	34 (21.79%)	10 (21.27%)	0.817
Tiredness	71 (21.78%)	59 (37.82%)	12 (25.53%)	0.204
Ache	41 (12.58%)	28 (17.95%)	13 (27.66%)	0.080
Diarrhea	10 (3.07%)	7 (4.48%)	3 (6.38%)	0.601
Death significance (M=108, F=22)				
Age (years)				
0-20	1 (0.77%)	1 (0.93%)	0 (0.00%)	
21-50	23 (17.70%)	17 (15.70%)	6 (27.27%)	0.724
>50	1 (0.77%)	90 (83.33%)	16 (72.72%)	

*. *Correlation is significant at the 0.05 level (2-tailed)*

**. *Correlation is significant at the 0.01 level (2-tailed)*

419   **Figures**

422   **Figure 1.** Percentage of sex-specific blood group of COVID-19 infected patients

Figure 2. Percentage of sex-specific occupations of COVID-19 patients