Cecile F. Frampas[1], Katie Longman[1], Matt P. Spick[1], Holly M. Lewis[1], Catia D. S. Costa[2], Alex Stewart[3], Deborah Dunn-Walters[3], Danni Greener[4], George E. Evetts[4], Debra Skene[3], Drupad Trivedi[5], Andrew R. Pitt[5], Katherine Hollywood[5], Perdita Barran[5] and Melanie J. Bailey[1,2]

[1] Faculty of Engineering and Physical Sciences, University of Surrey, Guildford, GU2 7XH, UK

[2] Surrey Ion Beam Centre, University of Surrey, Guildford, GU2 7XH, UK

[3] Faculty of Health and Medical Sciences, University of Surrey, Guildford, GU2 7XH, UK

[4] Frimley Park Hospital, Frimley Health NHS Trust, GU16 7UJ, UK

[5] Manchester Institute of Biotechnology, University of Manchester, M1 7DN, United Kingdom

Corresponding Author: Dr Melanie Bailey, tel +44 (0)1483 682593, m.bailey@surrey.ac.uk, Faculty of Engineering and Physical Sciences, University of Surrey, Guildford, GU2 7XH, UK

Preprint location: MedRxiv

CF, KL and MS contributed equally to this paper.

Keywords: Saliva, Metabolomics, COVID-19, Prognostic Test, Liquid Chromatography Mass Spectrometry

23 **Abbreviations**

COVID-19	Coronavirus disease 19
CRP	C-reactive protein
HTN	Hypertension
IHD	Ischemic heart disease
KEGG	Kyoto Encyclopedia of Genes and Genomes
LC	Liquid chromatography
LC-MS	Liquid chromatography mass spectrometry
LOOCV	Leave-one-out cross validation
MS	Mass spectrometry
MS/MS or MS^2	Tandem mass spectrometry
NPA	Negative percent agreement
PCA	Principal components analysis
PCR	Polymerase chain reaction
PLS-DA	Partial least squares-discriminant analysis
PPA	Positive percent agreement
QC	Quality control
RT-PCR	Reverse transcription polymerase chain reaction
SARS-CoV-2	Severe acute respiratory syndrome coronavirus 2
T2DM	Type 2 diabetes mellitus
VIP	Variable importance in projection

ABSTRACT

Background

The COVID-19 pandemic is likely to represent an ongoing global health issue given

the potential for vaccine escape and the low likelihood of eliminating all reservoirs of

the disease. Whilst diagnostic testing has progressed at pace, there is an unmet

clinical need to develop tests that are prognostic, to triage the high volumes of patients

arriving in hospital settings. Recent research has shown that serum metabolomics has

potential for prognosis of disease progression. [1] In a hospital setting, collection of

saliva samples is more convenient for both staff and patients, and therefore offers an

alternative sampling matrix to serum. We demonstrate here for the first time that saliva

metabolomics can reveal COVID-19 severity.

Methods

88 saliva samples were collected from hospitalised patients with clinical suspicion of

COVID-19, alongside clinical metadata. COVID-19 diagnosis was confirmed using RT-

PCR testing. COVID severity was classified using clinical descriptors first proposed by

SR Knight et al. Metabolites were extracted from saliva samples and analysed using

liquid chromatography mass spectrometry.

Results

In this work, positive percent agreement of 1.00 between a PLS-DA metabolomics

model and the clinical diagnosis of COVID severity was achieved. The negative

percent agreement with the clinical severity diagnosis was also 1.00, for overall

percent agreement of 1.00.

Conclusions

This research demonstrates that liquid chromatography-mass spectrometry can

identify salivary biomarkers capable of separating high severity COVID-19 patients

from low severity COVID-19 patients in a small cohort study.

1. Introduction

The SARS-CoV-2 pandemic has caused a sustained threat to global health since the discovery of the virus in 2019. [2] Whilst great strides have been made in both treatment and vaccination development, [3,4] the disease has inflicted multiple waves of infection throughout the world during 2020 and into 2021. [5,6] COVID-19 has higher fatality rates than seasonal influenza, [7] and in addition, new variants are constantly evolving with the potential for either reduced vaccine effectiveness or altered lethality. [8] As a consequence, there is a continuing need both for better understanding of the impact of COVID-19 on the host metabolism as well as for prognostic tests that can be used to triage the high volumes of patients arriving in hospital settings.

Nasopharyngeal swabs followed by polymerase chain reaction (PCR) have been adopted worldwide for SARS-CoV-2 detection. However, supply chains for swabs rapidly collapsed amongst exponential increases in demand for testing, highlighting the urgency for alternative sample types and testing approaches. Furthermore, whilst PCR tests are easily deployable and highly selective for the virus, these approaches yield no prognostic information and cannot easily be delivered for rapid turnaround at the point of care, for example during a hospital admissions process. In contrast, tests based on mass spectrometry can be provided in minutes, with mass spectrometry instrumentation typically available in hospital pathology laboratories. Prognostic tests, whilst challenging due to the varied phenotypes that may present themselves, [9] could be used to manage demand for hospitalisation and treatment, especially should vaccine escape lead to future waves of COVID-19 infection.

Metabolic biomarkers in serum have been identified that carry prognostic information, [10,11] but sampling blood is invasive. Our experience in collecting and analysing patient

samples is that saliva samples are significantly easier to collect and handle than blood.

Blood collection requires trained phlebotomists, causes discomfort to patients and

must be spun soon after collection to preserve the metabolome. In contrast, a saliva

sample can be donated quickly and painlessly by a patient. Saliva is itself a carrier of

the coronavirus, [12] and has been proposed as a gold standard for SARS-CoV-2

detection. [13,14] It additionally offers information via its own characteristic metabolites.

[15] To date, saliva as a biofluid for metabolism analysis has been used for breast,

pancreatic and also oral cancers. [16,17] Here we explore the potential of saliva

metabolomics to distinguish between severe and mild COVID-19 infection, with a view

to providing a prognostic test that can be used to triage hospital patients, for example

to identify patients who would benefit from immunomodulating drugs such as

Tocilizumab. [18]

This work took place as part of the wider efforts of the COVID-19 International Mass

Spectrometry (MS) Coalition. [19,20] This consortium aims to provide molecular level

information on SARS-CoV-2 in infected humans, in order to better understand,

diagnose and treat cases of COVID-19 infection. Data related to this work will be

stored and fully accessible on the MS Coalition open repository. The website URL is

https://covid19-msc.org/

2. Materials and Methods

2.1 Participant recruitment and ethics

Ethical approval for this project (IRAS project ID 155921) was obtained via the NHS

Health Research Authority (REC reference: 14/LO/1221). 88 participants were

recruited at NHS Frimley NHS Foundation Trust hospitals by researchers from the

University of Surrey. Participants were identified by clinical staff to ensure that they

had the capacity to consent to the study, and were asked to sign an Informed Consent Form; those that did not have this capacity or who did not sign the form were not sampled. Consenting participants were categorised by the hospital as either "query COVID" (meaning there was clinical suspicion of COVID-19 infection) or "COVID positive" (meaning that a positive COVID test result had been recorded during their admission). All participants were provided with a Patient Information Sheet explaining the goals of the study.

Inclusion for participants was determined by reverse transcription polymerase chain reaction (RT-PCR) results; participants with an inconclusive RT-PCR test (clinically positive only and/or inconclusive test result, n=6) or where the time lag between initial RT-PCR test and sampling exceeded fourteen days were excluded (n=7). These additional exclusion criteria reduced the participant population from 88 to 75.

2.2 Sample collection, extraction and instrumental analysis

Patients were sampled immediately upon recruitment to the study in two waves, one between May and August 2020 and the second between October and November 2020. The range in time between symptom onset and saliva sampling ranged from 1 day to > 1 month, an inevitable consequence of collecting samples in a pandemic situation. Each participant provided a sample of saliva by spitting directly into a falcon tube which was placed on ice immediately after collection. Samples were transferred on ice from the hospital to the University of Surrey by courier within 4 hours of collection, to minimise changes to salivary metabolites. [21] Once received at University of Surrey, the samples were stored at minus 80 °C until analysis.

Alongside saliva collection, metadata for all participants was also collected covering *inter alia* sex, age, comorbidities (based on whether the participant was receiving

treatment), the results and dates of COVID PCR tests, bilateral chest X-Ray changes,

smoking status, drug regimen, and whether and when the participant presented with

clinical symptoms of COVID-19. Values for lymphocytes, CRP and eosinophils were

also taken; values obtained within five days of the saliva sampling were recorded.

Each participant was attributed a "severity score" in relation to their fitness

observations at the time of hospital admission using the metadata collected. This score

used the "mortality scoring" approach of SR Knight et al. [8] adapted to disregard age,

sex at birth and comorbidities, and ranged from 0 to 6; patients scoring 0 to 3 were

attributed low severity and patients scoring 4 to 6 were attributed high severity.

Sample preparation and processing followed the guidelines set out by the COVID-19

Mass Spectrometry Coalition. [22] Saliva samples were separated into aliquots: 50 μL

of saliva was added to 200 μL of ice-cold isopropanol to precipitate protein, and this

also had the advantage of deactivating the virus to allow transfer into a lower biological

safety level laboratory. The samples were agitated for one hour, sonicated three times

for 30 seconds, with resting on ice for 30 seconds between each sonication. Each

sample was then left to stand on ice for 30 minutes then centrifuged for 10 minutes at

10 000 g. The supernatant was removed and the precipitated protein pellet reserved

for future analysis. The supernatant then underwent centrifugal filtration (0.22 μm

cellulose acetate) for five minutes at 10,000 g, and the filtered supernatant was then

dried under nitrogen and stored at minus 80 °C.

Samples were reconstituted on the day of analysis in 100 μL water:methanol (95:5)

with 0.1% formic acid by volume. 10 μL of each sample was set aside for combination

into a pooled QC. The samples were analysed over a period of eleven days. Each day

consisted of a run incorporating blank injections (n=2), field blank injections (n=3),

pooled QC injections (n=6, 3 at the start and finish), as well as QCs to measure

instrumental and extraction variation (n=7 and 3 respectively), and 10 participant

samples, randomised for positive/negative, with 3 repeat analyses for each.

2.3 Materials and chemicals

The materials and solvents utilised in this study were as follows: 2 mL microcentrifuge

tubes (Eppendorf, UK), 0.22 µm cellulose acetate sterile Spin-X centrifuge tube filters

(Corning incorporated, USA), 200 µL micropipette tips (Starlab, UK) and Qsert^{TM} clear

glass insert LC vials (Supelco, UK). LC-MS grade 2-propanol was used as an

inactivation solvent. Optima^{TM} LC-MS grade methanol and water were used as

reconstitution solvents and mobile phases. LC-MS grade formic acid was added to the

mobile phase solvents at 0.1% (v/v). Solvents were purchased from Fisher Scientific,

UK.

2.4 Instrumentation and operating conditions

Analysis of samples was carried out using a UltiMate 3000 UHPLC equipped with a

binary solvent manager, column compartment and autosampler, coupled to a Q

Exactive™ Plus Hybrid Quadrupole-Orbitrap™ mass spectrometer (Thermo Fisher

Scientific, UK) at the University of Surrey's Ion Beam Centre. Chromatographic

separation was performed on a Waters ACQUITY UPLC BEH C18 column (1.7 µm,

2.1 mm x 100 mm) operated at 55 ºC with a flow rate of 0.3 ml min^{-1}.

Mobile phase A was water: methanol (v/v 95:5) with 0.1% formic acid, whilst mobile

phase B was methanol:water (v/v, 95:5) with 0.1% formic acid (v/v). An injection

volume of 5 µL was used. The initial solvent mixture was 2% B for one minute,

increasing to 98% B over 16 minutes and held at this level for four minutes. The

gradient was finally reduced back to 2% B and held for two minutes to allow for column

equilibration. Analysis on the Q-Exactive Plus mass spectrometer was performed with

a scan range of *m/z* 100 to 1 000, and 5 ppm mass accuracy. MS/MS validation of

features was carried out on Pooled QC samples using data dependent acquisition

mode and normalised collision energies of 30 and 35 (arbitrary units). Operating

conditions are summarised in Table S1 (Supplementary Material).

177   2.5 Data processing

LC-MS outputs (.raw files) were pre-processed for alignment and peak identification

using Compound Discoverer version 3.1 and Freestyle 1.6 (Thermo Fisher Scientific,

UK). Peak picking was set to a mass tolerance ±5 ppm, and alignment to a retention

time window of 120 seconds. Missing values were imputed using a K-nearest

neighbour approach. [23] Features identified by mass spectrometry were initially

annotated using accurate mass match with reference to external databases (KEGG,

Human Metabolome Database, DrugBank, LipidMaps and BioCyc), and then

validation was performed using data dependent MS/MS analysis. This process yielded

an initial peak:area matrix with 10,700 discrete features. Two criteria were used for

inclusion in the final analysis: only those features with identities validated by MS/MS

were used, reducing the number of features to 1,874, and 1,514 features that were

present in less than 30% of participant samples were excluded. This left 360 features

that were used in the analysis. Normalisation was performed using EigenMS in

NOREVA for each dataset analysed, [24,25] i.e. independently for the diagnostic

population (COVID-19 positive versus negative) and prognostic population (COVID-

19 positive: high severity versus low severity).

194   2.6 Statistical Analysis

PCA analyses were conducted in SIMCA (Sartorius Stedim Biotech, France). PLS-DA

and additional machine learning was conducted in R Studio Version 1.3.959 and

MetaboAnalyst. [26,27] Leave-one-out cross-validation was used for model validation test

accuracy, sensitivity and specificity; variable importance in projection (VIP) scores

were used to assess feature significance alongside p-values and effect sizes (fold

count). Batch effects were assessed by PCA analysis of both collection batches

(waves one and two) and also instrument and extraction batching by day (Figures S1

and S2, Supplementary Material), showing no clustering by batches. KEGG pathway

analysis was performed using MetaboAnalyst.

In prognostic analysis, given the lack of a "gold standard" reference test for whether

COVID-19 is likely to be high severity or low severity (as this depends on clinical

judgement), positive percent agreement (PPA) between the generated model and a

high severity clinical diagnosis was used in preference to sensitivity, which measures

the detection of positive instances of a disease relative to a ground truth value.

Similarly, negative percent agreement (NPA) between the model and a high severity

clinical diagnosis was used in preference to specificity, which measures the absence

of a disease relative to a ground truth value. In diagnostic analysis, given that RT-PCR

tests were available to establish a ground truth, sensitivity and specificity values were

calculated alongside diagnostic accuracy.

3. RESULTS

3.1 Population metadata overview

The study population analysed in this work included 75 participants, comprising 47

participants presenting with a positive COVID-19 RT-PCR test and 28 participants

presenting without. Of the positive participants, 10 were classed as presenting with

high severity COVID-19, 34 were classed as presenting with low severity COVID-19,

and 3 lacked sufficient clinical information for severity scoring. A summary of the

metadata is shown in Table 1.

Table 1: Summary of clinical characteristics by participant cohort

Parameters	Covid-19 Low severity	Covid-19 High severity	p-value High vs Low Severity	Covid-19 Negative	p-value Pos vs Neg
n	34	10		28	
Age (mean, standard deviation; years)	59.7 ± 18.4	63.1 ± 12.7	0.61	61.6 ± 21.9	0.74
Male / Female (n)	16 / 18	8 / 2	0.083	16 / 12	0.26
Treated for Hypertension (n)	6	6	.041	12	0.21
Treated for High Cholesterol (n)	2	0	1.00	6	.05
Treated for Type 2 Diabetes Mellitus (n)	5	3	0.39	10	0.29
Treated for Ischemic Heart Disease (n)	1	2	0.149	7	0.09
Current Smoker (n)	1	0	1.00	0	NA
Ex-Smoker (n)	12	5	0.71	8	0.46
Medical Acute Dependency admission (n)	10	6	0.26	4	0.06
Intensive Care Unit admission (n)	0	0	N/A	0	NA
Survived Admission (n)	34	8	0.048	27	1.00
Lymphocytes (mean, standard deviation; cells / µL)	0.8 ± 0.5	0.9 ± 0.7	0.77	1.0 ± 0.5	0.302
C-Reactive Protein (mean, standard deviation; mg / L)	115.4 ± 84.9	170.0 ± 83.4	0.075	127.3 ± 104.7	0.80
Eosinophils (mean, standard deviation; 100 / µL)	0.1 ± 0.1	0.0 ± 0.0	0.018	0.3 ± 0.4	0.002
Bilateral Chest X-Ray changes (n)	15	8	0.26	3	0.0009
Continuous Positive Airway Pressure (n)	1	1	0.442	3	0.36
O_2 required (n)	9	4	0.69	8	1.00

In this study all participants were recruited in a hospital setting with at least potential

suspicion of COVID-19 infection; controls were age matched and had similar profiles

in terms of gender, oxygen requirements and survival rates. Significantly more COVID-

19 positive patients had bilateral chest X-ray changes (p-value 0.0009) and higher

levels of eosinophils (p-value 0.002), in agreement with literature observations, [10] but

not for C-reactive protein (CRP, p-value 0.80). Type 2 diabetes mellitus (T2DM) was

more prevalent in the COVID-19 negative population than the positive population,

being observed in 36% of COVID-19 negatives versus 30% of high severity COVID-

19 patients and 15% of low severity COVID-19 patients, and a similar observation of greater comorbidity being seen in the negative population was also true for ischemic heart disease (IHD) and hypertension (HTN). The greater preponderance of underlying comorbidities within the negative population represents a confounding factor.

Within the COVID-19 positive cohort, comorbidities were again age matched, but the high severity grouping had more males (80% male for high severity versus 47% for low severity) and had a statistically significant difference in proportion presenting with hypertension (p-value 0.04) and a statistically significant decrease in eosinophil levels (p-value 0.02). CRP was increased by a 1.5x fold count in high severity participants versus low (p-value 0.08). There was no statistically significant increase in CRP for low severity versus COVID-19 negative participants, but this may reflect changes to the inflammatory response caused by interventions reducing CRP levels in cases of mild COVID-19.

3.2 Overview of features identified by Liquid Chromatography Mass Spectrometry (LC-MS)

360 features with MS/MS validation were identified as being present in 30% or more of participant samples. Of these 360 features, 37 were identified as related to medical interventions or food and were excluded, leaving 323 for statistical analysis. Of the 323, 38 were annotated by *m/z* value, 171 were annotated by formula (elemental composition), and 114 were annotated as metabolites.

3.3 Analysis of cohorts by multivariate techniques

Initially separation of COVID-19 positive versus negative participants was tested, as well as separation of COVID-19 high severity and low severity. As shown in Figure 1A,

separation for diagnostic purposes showed no clear separation by visual inspection

and delivered R2Y of 0.78 and Q2Y of 0.18. Leave-one-out cross-validation (LOOCV)

provided sensitivity of 0.74 (95% confidence interval of 0.60 - 0.86) and specificity of

0.75 (0.55 - 0.89), which was considered insufficient to justify further investigation. The

most significantly dysregulated identified metabolites (measured by p-value) between

positive and negative COVID-19 status are listed in table S2 (Supplementary Material).

Figure 1B shows separation for COVID-19 high severity participants versus low

severity participants. The optimal separation was found using 5 components. Using

leave-one-out cross validation, PPA for COVID-19 high severity was 1.00 (95%

confidence interval of 0.69 - 1.00) and NPA was 1.00 (0.90 - 1.00), for overall percent

agreement with the clinical diagnosis of 1.00 (0.92 - 1.00).

Figure 1: Saliva metabolomics analysis for COVID-19 diagnosis and prognosis via

LC-MS, showing:

A   PLS-DA plot for 75 participants, COVID-19 positive / negative

B   PLS-DA plot for 44 participants, high severity / low severity

C LOOCV confusion matrix, COVID-19 positive / negative

D LOOCV confusion matrix, high severity / low severity

A volcano plot is shown in Figure 2. The most significantly dysregulated identified

metabolites (measured by p-value) are shown as boxplots in Figure 3 below and a

complete list of metabolites showing statistically significant differences between high

and low COVID severity populations is shown in table S3 (Supplementary Material).

Amino acids are highlighted as this class of metabolites was the most dysregulated

between high and low severity of the identified features.

Figure 2: Volcano plot of statistical significance versus effect size for MS/MS validated

features separating participants presenting with high severity versus low severity

COVID-19

Figure 3:   Boxplots of features down regulated with Covid-19 severity

[corresponding p-values: 0.066 ; 0.004 ; 0.051]

MS/MS spectra for the significant features presented in Figure 3 are additionally

shown in Figures S3 to S5 (Supplementary Material).

The normalised prognostic dataset was also processed for pathway analysis to

explore changes in metabolic pathways relating to COVID-19 (Figure S6). No

pathways met the criteria for both meaningful impact and statistical significance,

possibly due to the number features identified in each pathway being notably smaller

than typically achieved in serum or plasma, consistent with saliva being a filtrate and

in general featuring lower metabolic concentrations. [28]

3.4 Validation set

Whilst no fully independent prognostic validation set was available, it was decided to

project the PLS-DA model obtained for high severity versus low severity participants

on to COVID-19 negative participants. Given that these participants should not show

features associated with high severity COVID-19, this was considered to offer

additional information. The confusion matrix for the results of the projection is shown

in Table 2 below.

Table 2: Confusion matrix for PLS-DA model projected on to COVID-19 negative

participants

	COVID-19 negative participants
PLS-DA model result: High Severity	1
PLS-DA model result: Low Severity	27

4. Discussion

Whilst age and recruitment venue were well matched (all participants were recruited

in a hospital setting including controls), a number of variables within the metadata

illustrate the natural difficulties in experimental design experienced during a pandemic.

Age ranges of participants were large, a wide range of comorbidities were present,

and the time between symptom onset and saliva sampling ranged from 1 day to > 1

month. Participant recruitment of the most severely affected was limited by ethics

approval only covering patients who could give informed consent, thereby precluding

the participation of patients with the highest COVID severity. Furthermore, given the

small n in this pilot study, precision was necessarily low and confidence intervals wide.

In this study, saliva samples were provided under conditions that could be practically

achieved in a hospital pandemic setting, albeit this meant no scope for abstinence

from food and / or drink before saliva sampling, and no prior rinsing of the mouth,

leading to potential confounding factors. Separation of COVID-19 positive COVID-19 versus negative participants was limited, possibly due to COVID-19 negative participants also being hospitalised and in poor health, perhaps having similar inflammatory responses to some COVID-19 positive participants. In spite of this, superior differentiation by multivariate analysis was achieved in relation to COVID-19 severity. PLS-DA showed separation of High Severity COVID-19 positive participants from Low Severity COVID-19 positive participants, with PPA and NPA of 100% by LOOCV. Furthermore, whilst not a true independent validation set, projecting the PLS-DA model on to COVID-19 negative participants, i.e. the controls, showed that the model classified them with 97% consistency as "low risk", i.e. that the features associated with high severity were present neither in low severity nor in COVID-19 negative participants.

A number of identified metabolites showed statistically significant differences between the high and low severity participants. Both valine (p-value 0.02, fold-count 0.48) and leucine (p-value 0.04, fold-count 0.67) showed statistically significant changes between high and low severity. As shown in Figure 3, amino acids constituted the class of metabolites seeing the most change between high and low severity, similar to literature observations of changes in either amino acids or ratios of amino acids, albeit specific amino acids commonly cited in the literature (for example kynurenine, arginine or ratios thereof) [1,29,30] did not feature in the saliva analysis presented here. It should be noted, however, that the correlation of metabolites between saliva and blood has previously been found to be weak or in some cases non-existent, [31,32] and the same may be true for the saliva and blood of individuals testing positive for COVID-19. Direct analysis of paired blood samples would be required to draw any definitive conclusion on differential dysregulation of metabolites between serum and saliva.

5. Conclusion

In this work a number of features have been identified for the first time that may differentiate the saliva of those presenting with high severity and low severity COVID-19. We believe that saliva has potential to add to understanding of the progression and severity of COVID-19. In addition, saliva may be collected less invasively than other biofluids, and mass spectrometry techniques have the advantage of being often located within hospitals, making MS-based techniques useful in a clinical setting. Consequently, we view saliva as a worthy biofluid for consideration for prognostic testing.

Data sharing statement

Participant metadata data with identifiers, alongside peak:area matrices used in this work will be made available on the Mass Spectrometry Coalition website upon publication of this study. The analytical protocols used as well as mass spectrometry .raw files, sample and participant data will be openly available for all researchers to access. The website URL is https://covid19-msc.org/

Declaration of Competing Interest

The authors have no competing interest to declare.

Author's contributions

CF and KL collected and extracted all samples used in this work, were responsible for LC-MS method development, data processing and statistical analysis in SIMCA. MS conducted machine learning and pathway analyses and additionally drafted the manuscript. CC and HL supported mass spectrometry method development and advised on analytical methods. AS and DDW obtained ethical approval and biobanking

of samples. GE and DG facilitated access to participants and collected participant

metadata. DS, DT, PB, KH and AP supported experimental and statistical design and

also provided editorial comment on the draft manuscript. MB oversaw all aspects of

this work, including obtaining funding for the study, clinical access, experimental

design, analysis and was responsible for supervision of the research team.

Funding

The authors would like to acknowledge funding from the EPSRC Impact Acceleration

Account for sample collection and processing, as well as EPSRC Fellowship Funding

EP/R031118/1, the University of Surrey and BBSRC BB/T002212/1. Mass

Spectrometry was funded under EP/P001440/1.

Acknowledgements

The authors acknowledge Samiksha Ghimire from Groningen Medical School for

translation of participant information sheets and consent forms into Nepalese, as well

as Kyle Saunders of the University of Surrey for access to batch controls. The authors

are additionally grateful to Thanuja Weerasinge (Jay), Manjula Meda, Chris Orchard

and Joanne Zamani of Frimley Park NHS Foundation Trust for their help with ethics

approvals and access to hospital patients.

Supplementary Material

Table S1: Operating conditions of the mass spectrometer used in this research.

Parameter	Operating condition
Spray voltage	3.5 kV
Capillary temperature	275 °C
S-lens RF level	50
Sheath gas flow rate	40
Aux gas flow rate	0
Scan range	100 *m/z* to 1 000 *m/z*
Resolution	70 000
Polarity	Positive
AGC target	10^6
Maximum inject time	200
MS/MS Parameter	**Operating condition**
Mode	Full Scan MS / dd-MS^2 with inclusion lists
Resolution	35 000 at m/z 200
Loop Count	6
Intensity threshold	$5 * 10^4$
Collision energy	30 and 35
Dynamic Exclusion	5 seconds

Figure S1: Principal Component Analysis of each patient sample (circles) and batch

QC's (squares), coloured according to extraction batch, showing no significant

clustering of patient samples according to extraction batch (square: QC ; circle:

patients)

398 **Figure S2:** Principal Component Analysis of each patient sample and run QC,

399 showing low levels of QC variation according to position in the run sequence

400 **Table S2:** Features distinctive between COVID-19 positive and negative

Feature	Fold Count	p-value
C25 H36 N6 O6	0.52	0.0121
DL-Phenylalanine	1.39	0.0164
C7 H10 N6 O2	1.35	0.0179
2-linoleoyl-sn-glycero-3-phosphoethanolamine	1.85	0.0180
C39 H71 N2 O16 P3	0.81	0.0226
DC2810000	1.60	0.0231
179.8965	2.09	0.0251
1-Hexadecanoylpyrrolidine	1.84	0.0287
Taurine	1.44	0.0292
2707	1.19	0.0363
148.0048	0.70	0.0379
C47 H84 N9 O13 P3	0.73	0.0415
C22 H36 N8 O9	1.39	0.0439

Feature	Fold Count	p-value
3721	1.70	0.0443
butyl acrylate	1.62	0.0470
C2 H3 N3 O2 P2	1.25	0.0501

402 **Table S3:** Features distinctive between COVID-19 high severity and low severity

Feature	Fold Count	p-value
C44 H74 N8 O16	0.50	0.0000
C39 H67 N7 O15	0.55	0.0003
C29 H56 N7 O11 P	0.57	0.0013
C42 H72 N8 O17	0.54	0.0038
C49 H81 N9 O17	0.61	0.0148
C37 H67 N6 O12 P3	0.51	0.0160
C47 H84 N9 O13 P3	0.66	0.0178
Valine	0.48	0.0198
C6 H13 N2 O6 P	1.92	0.0237
C9 H21 N O6	2.17	0.0241
C17 H22 N4 O8	1.98	0.0247
C8 H12 O5 P2	2.34	0.0254
C2 H8 O5 P2	1.41	0.0258
C60 H86 N10 O9 P2	0.65	0.0270
C32 H48 N8 O9	0.47	0.0289
Lys-phe	0.48	0.0367
N-{3-[(4-Acetamidobutyl)amino]propyl}acetamide	0.59	0.0382
Leucine	0.67	0.0405
C42 H71 N7 O15	0.44	0.0453
273.86553	1.29	0.0475

Figure S3: MS/MS spectra of proline

Figure S4: MS/MS spectra of valine

Figure S5: MS/MS spectra of phenylalanine

Figure S6: Pathways analysis of metabolites : high severity versus low severity

References

1 Danlos F-X, Grajeda-Iglesias C, Durand S, *et al.* Metabolomic analyses of

COVID-19 patients unravel stage-dependent and prognostic biomarkers. *Cell*

Death Dis 2021; **12**: 258.

2 WHO. Novel Coronavirus – China. 2020. https://www.who.int/csr/don/12-

january-2020-novel-coronavirus-china/en/ (accessed July 27, 2020).

3 Africa S. Comment Oxford – AstraZeneca COVID-19 vaccine efficacy. 2021;

397: 174–6.

4 The RECOVERY Collaborative Group. Dexamethasone in Hospitalized

Patients with Covid-19 — Preliminary Report. *N Engl J Med* 2020.

DOI:10.1056/nejmoa2021436.

5   Cacciapaglia G. Second wave COVID - 19 pandemics in Europe : a temporal

playbook. *Sci Rep* 2020; : 1–8.

6   Lai JW, Cheong KH. Superposition of COVID-19 waves , anticipating a

sustained wave , and lessons for the future. 2020; : 1–12.

7   Piroth L, Cottenet J, Mariet A-S, *et al.* Comparison of the characteristics,

morbidity, and mortality of COVID-19 and seasonal influenza: a nationwide,

population-based retrospective cohort study. *Lancet Respir Med* 2020; **2600**:

1–9.

8   WHO. SARS-CoV-2 Variant – United Kingdom of Great Britain and Northern

Ireland. 2020. https://www.who.int/csr/don/21-december-2020-sars-cov2-

variant-united-kingdom/en/ (accessed Jan 25, 2021).

9   Darmon M, Dumas G. Anticipating outcomes for patients with COVID-19 and

identifying prognosis patterns. *Lancet Infect Dis* 2021; **21**: 744–5.

10   Knight SR, Ho A, Pius R, *et al.* Risk stratification of patients admitted to

hospital with covid-19 using the ISARIC WHO Clinical Characterisation

Protocol: development and validation of the 4C Mortality Score. *BMJ* 2020;

370: m3339.

11   Shen B, Yi X, Sun Y, *et al.* Article Proteomic and Metabolomic

Characterization of COVID-19 Patient Sera ll Article Proteomic and

Metabolomic Characterization of COVID-19 Patient Sera. 2020; : 59–72.

12   To KK, Tsang OT, Yip CC, *et al.* Consistent Detection of 2019 Novel

Coronavirus in Saliva. 2020; **71**: 841–3.

13   Costa G, Pereira CM, Kelly T, Valente AP. Saliva NMR-Based Metabolomics

in the War Against COVID-19. 2020. DOI:10.1021/acs.analchem.0c04679.

14   Tan SH, Allicock O, Armstrong-Hough M, Wyllie AL. Saliva as a gold-standard

sample for SARS-CoV-2 detection. *Lancet Respir Med* 2021.

DOI:10.1016/S2213-2600(21)00178-8.

15   Mandel ID. The Functions of Saliva. *J Dent Res* 1987; **66**: 623–7.

16   Sugimoto M, Wong DT, Hirayama A, Soga T, Tomita M. Capillary

electrophoresis mass spectrometry-based saliva metabolomics identified oral,

breast and pancreatic cancer-specific profiles. *Metabolomics* 2010; **6**: 78–95.

17   Assad DX, Acevedo AC, Cançado E, *et al.* Using an Untargeted Metabolomics

Approach to Identify Salivary Metabolites in Women with Breast Cancer. 2020;

: 1–13.

18   Janik E, Niemcewicz M, Podogrocki M, Saluk-Bijak J, Bijak M. Existing Drugs

Considered as Promising in COVID-19 Therapy. *Int J Mol Sci* 2021; **22**: 5434.

19   Struwe W, Emmott E, Bailey M, *et al.* The COVID-19 MS Coalition—

accelerating diagnostics, prognostics, and treatment. *Lancet* 2020; **395**: 1761–

2.

20   Spick M, Longman K, Frampas C, *et al.* Changes to the sebum lipidome upon

COVID-19 infection observed via rapid sampling from the skin.

EClinicalMedicine 2021; : 100786.

21   Duarte D, Castro B, Pereira JL, Marques JF, Lu A, Gil AM. Evaluation of Saliva

Stability for NMR Metabolomics : Collection and Handling Protocols. ; : 1–15.

22   COVID-19 Mass Spectrometry Coalition. COVID-19 metabolomics and

lipidomics protocol. 2020. https://covid19-msc.org/metabolomics-and-

lipidomics-protocol/ (accessed Jan 25, 2021).

23   Beretta L, Santaniello A. Nearest neighbor imputation algorithms: A critical

evaluation. *BMC Med Inform Decis Mak* 2016; **16**. DOI:10.1186/s12911-016-

0318-z.

24   Karpievitch Y V, Taverner T, Adkins JN, *et al.* Normalization of peak intensities

in bottom-up MS-based proteomics using singular value decomposition.

Bioinformatics 2009; **25**: 2573–80.

25   Li B, Tang J, Yang Q, *et al.* NOREVA: Normalization and evaluation of MS-

based metabolomics data. *Nucleic Acids Res* 2017; **45**: W162–70.

26   R Core Team. R: A Language and Environment for Statistical Computing.

2020. https://www.r-project.org/.

27   Chong J, Wishart DS, Xia J. Using MetaboAnalyst 4.0 for Comprehensive and

Integrative Metabolomics Data Analysis. *Curr Protoc Bioinforma* 2019; **68**: e86.

28   Rossini EL, Milani MI, Lima LS, Pezza HR. Paper microfluidic device using

carbon dots to detect glucose and lactate in saliva samples. *Spectrochim Acta*

Part A Mol Biomol Spectrosc 2021; **248**: 119285.

29   Thomas T, Stefanoni D, Reisz JA, *et al.* COVID-19 infection alters kynurenine

and fatty acid metabolism, correlating with IL-6 levels and renal status. *JCI*

Insight 2020; **5**. DOI:10.1172/jci.insight.140327.

30   Fraser DD, Slessarev M, Martin CM, *et al.* Metabolomics Profiling of Critically

Ill Coronavirus Disease 2019 Patients: Identification of Diagnostic and

Prognostic Biomarkers. *Crit Care Explor* 2020; **2**: e0272.

31   Do KT, Kastenmüller G, Mook-Kanamori DO, *et al.* Network-Based Approach

for Analyzing Intra- and Interfluid Metabolite Associations in Human Blood,

Urine, and Saliva. *J Proteome Res* 2015; **14**: 1183–94.

32   Bel'skaya L V, Sarf EA, Kosenok VK. Age and gender characteristics of the

biochemical composition of saliva: Correlations with the composition of blood

plasma. *J Oral Biol Craniofacial Res* 2020; **10**: 59–65.