Scoping Review and Interpretation of Myofascial Pain/Fibromyalgia Syndrome: An Attempt to Assemble a Medical Puzzle

Shiloh Plaut[1]

St. George's University of London

shilowhale@gmail.com

Key words: fascial armoring, myofascial pain syndrome, myofibroblast, tensegrity, fibromyalgia, global chronic compartment syndrome.

Shiloh Plaut- sole researcher and author of this manuscript including conception, organization, literature review, integration of information, findings analysis, manuscript preparation and writing.

Summary/Abstract

Introduction: Myofascial Pain Syndrome (MPS) is a common overlooked and underdiagnosed condition, and with significant burden. MPS is often dismissed by clinicians while patients remain in pain for years. MPS can evolve into fibromyalgia, however, effective treatments for both are lacking due to absence of a clear mechanism. Many studies focus on central sensitization. Therefore, the purpose of this scoping review is to systematically search cross-disciplinary empirical studies of MPS, focusing on mechanical aspects, and suggest an organic mechanism explaining how it might evolve into fibromyalgia. Hopefully, it will advance our understanding of this disease.

Methods: Systematically searched multiple phrases in MEDLINE, EMBASE, COCHRANE, PEDro and medrXiv, majority with no time limit. Inclusion/exclusion based on title and abstract, then full text inspection. Additional literature added on relevant side topics. Review follows PRISMA-ScR guidelines. PROSPERO yet to adapt registration for scoping reviews.

Findings: 734 records included. Fascia can adapt to various states by reversibly changing biomechanical and physical properties. Trigger points, tension, and pain are a hallmark of MPS. Myofibroblasts play a role in sustained myofascial tension. Tension can propagate in fascia, possibly supporting a tensegrity framework. Movement and mechanical interventions treat and prevent MPS, while living sedentarily predisposes to MPS and recurrence.

Interpretation: MPS can be seen as a pathological state of imbalance in a natural process; manifesting from the inherent properties of the fascia, triggered by a disrupted biomechanical interplay. MPS might evolve into fibromyalgia through deranged myofibroblasts in connective tissue, or "fascial armoring", which may explain pain, distribution of pain, decreased pressure/pain threshold, and other phenomena eg, occasional complete resolution after surgery. Movement is an underemployed requisite in modern lifestyle. Lifestyle is linked to pain and suffering. The mechanism of needling is suggested to be more mechanical than currently thought. MPS should no longer be overlooked but solidified in the domain of common modern medicine.

Funding: none

Scoping Review and Interpretation of Myofascial Pain/Fibromyalgia Syndrome: An Attempt to Assemble a Medical Puzzle

Introduction

Chronic pain is a major cause of morbidity and has a significant impact on quality of life.[1] Myofascial pain denotes a pain arising from muscle and fascia. Commonly known as "muscle knots" myofascial pain usually arises in trigger points (TrPs) or 'tender spots'.[2,3,4] TrPs are small and sensitive areas in a contracted muscle, that spontaneously or upon compression cause pain to a distant region, known as a referred pain zone.[3,4]

Myofascial pain syndrome (MPS) is an entity that still lacks a clear definition. Some define it as a regional pain disorder,[2] others define it on the basis of tenderness and associated painful spots.[3] Nonetheless, TrPs and myofascial pain are a hallmark.[2,4,5] A fundamental difficulty arises when there is no clear definition, epidemiology, pathophysiology, or diagnosis of MPS. Having no accepted definition or criteria, clearly raises issues for both diagnosis and potential studies.[6-8] The purpose of this paper is to systematically search the empirical studies and components of MPS in an effort to assemble them into a suggested organic mechanism, explaining its pathophysiology and how it may evolve into fibromyalgia.

"Harrison's principles of internal medicine" (Kasper et al. 2018)[3] characterizes MPS by multiple areas of musculoskeletal pain and tenderness associated with painful points. Pain is deep and aching. It may arise after trauma, overuse or sedentarism.[3] A

study found laborers who exercise heavily are less likely to develop manifestations of MPS than sedentary workers.[3] Palpation of a TrP can reproduce or accentuate the pain.[3,4,9] However, these findings are not unique to MPS: in a controlled study they were also present in "normal" subjects.[3] Literature estimates 45% of men and 54% of women in the general population have TrPs.[2,9] Estimates are 37 to 65 percent of the population have myofascial pain, which costs the united states $47 billion every year.[4,10-11] MPS is one the most frequently under-diagnosed, under-treated and misunderstood sources of the ubiquitous aches and pains of humankind.[4,9] However, even recently, MPS is considered by some to be fiction, or otherwise paired with psychosomatic disorders since it involves pain and has no clear pathophysiology.[11-13]

Although MPS and fibromyalgia are defined as separate entities, the two may co-exist or considerably overlap.[2,3] They are both characterized by painful points, silent routine laboratory investigations and no systemic inflammation.[2,3,4] In some patients, regionally localized MPS may seem to evolve into fibromyalgia.[3] The puzzle of the mechanism of MPS and how it may develop into fibromyalgia still eludes our understanding. The marked dissociation between the estimated prevalence and burden of MPS, and the length of text on it in common medical textbooks, reflects this lack of understanding.

Studies of fibromyalgia reveal odd findings eg, complete resolution after laparoscopic surgery, and strong overlap with other conditions.[3] Many theories exist for fibromyalgia,[2] the most accepted and studied seems to be central sensitization, but no single theory seems to explain a wide range of empirical evidence, and the pathophysiology and etiology are still not clear.[2] Treatments are insufficient,

meanwhile patients suffer. This scoping review focuses on the organic mechanical aspects of myofascial pain with the hope it might advance our understanding of MPS and fibromyalgia.

Methods

A systematic search was held for multiple combinations of keywords in multiple databases. Keywords included : "fascia tension pain", "trigger point satellite", "fascia stiffness pain", "Fascia Densification", "trigger point densification movement", "biotensegrity", for all fields with no time limit; and for "Risk factors myofascial pain syndrome" only for title/abstract; Systematic reviews, meta-analysis and randomized controlled trials were searched in title or abstract for "myofascial pain syndrome", "spinal mobilization sympathetic nervous system", "sympathetic activity induced by pain" since 2015. The databases used were National Library of Medicine PUBMED (MEDLINE), COCHRANE, EMBASE, PEDro and medRxiv. Web of science was used to find items through a cited reference search. All searches were done between September 2020 and April 2021. Only items published in English were included. This work follows PRISMA guidelines for scoping reviews. The systematic search yielded 979 items. After removal of duplicates (N=221), 758 items were screened by assessing both title and abstract. Items off topic, foreign language, and in journal ranked Q4 were excluded (N=364). For the remaining items (N=394) a full text determination of eligibility was then performed. Four items were excluded. Final number of items included in systematic search was N=390.

When side topics were encountered, literature was added to broaden scoping review via medical textbooks, www.uptodate.com and more studies from searching databases (n=344). These side topics were not searched systematically. An attempt was made to cite more than one source in such cases. Total items was 734. Figure 1 shows flowchart of the scoping review.

Figure 1: Flowchart of scoping review.

Results

A total of 734 items were included. Numbers of items by recurrent themes were: "MPS and trigger points" (n=91), "properties of fascia" (n=40), "anatomy and movement" (n=43), "nervous system" (n=30), "biotensegrity" (n=25), "myofascial pain treatment" (n=206), "myofibroblasts" (n=85), "fibromyalgia" (n=74), "reviews" (n=33), and "other" eg, somatic syndromes, stress, chronic pain, Dupuytren's disease, plantar fasciitis (n=107). The systematic search included 127 clinical trials,

33 systematic review, and 19 meta-analysis. The findings are presented with the purpose of understanding the elements relevant to MPS and fascia, in order to assemble them and discuss a suggested mechanism of MPS and fibromyalgia. A microscopic view will set the stage for a macroscopic understanding.

Properties of fascia

General

Fascia is an important yet often misunderstood tissue in medicine. Essentially, fascia is connective tissue composed of irregularly arranged collagen fibers, distinctly unlike the regularly arranged collagen fibers seen in tendons, ligaments or aponeurotic sheets.[14] When connective tissue collagen is pressed, it can chondrify.[15] Fascia pervasively extends from head to toe, it envelops and permeates muscles, bones, blood vessels, nerves, and viscera, creating various layers at different depths.[14,16] Fascia supports the human body in vital functions such as posture, movement, and homeostasis,[14,17,18] as well as containing various sensory receptors for proprioception, nociception, and even hormones.[18,19] Nociception is influenced by the state of the fascia.[18] This means pain can arise from changes in the connective tissue.[18,19] The interconnectedness of fascia and its ability to transmit force are at the base of its functions.[18] Fascia is continuous from the trunk across the upper and lower limbs and hence has the potential to influence range of motion.[18] Fascia seems to connect the distant hip and the ankles, not only anatomically, but also mechanically, supporting the concept of a myofascial continuity.[18]

Biophysical and mechanical functions

Fascia has properties that enable it to reversibly change its stiffness and decrease the forces of tension experienced by it. This is a mechanical 'stress relaxation' resulting from the viscoelasticity of fascia that expresses high or low stiffness, depending on the rate of loading. Experimental tests demonstrate a 90% of stress relaxation achieved in about four minutes, afterwards the stress relaxation curve is near linear horizontal.[20,21,22] Studies observe increase in stiffness when ligaments are successively stretched, ie, strains produced by successive and identical loads decrease. They recover to baseline after a rest period. Isometrically maintained stress resulted in gradual tightening of the tissue.[14,23]

Blood vessels and nerves are interspersed in fascia, therefore, involvement of these structures due to fascial changes is commonly seen.[14] It is noted a tender point in the gluteus medius can refer pain down the leg and mimic sciatica.[3] Often the patient is aware of numbness or paresthesia rather than pain.[9] Pain is the result of a microenvironment around the nerve composed of connective tissues (eg, deep fascia, epineurium). Tissue modifications can be translated into change in nerve mobility, with a consequent decrease in the independency of the nerve from its surroundings. This leads to entrapment and "internal stretch lesion".[24] It is suggested that nerve compression syndromes are inevitably associated with a fascial pathology.[25] Circulation and perfusion can also be compromised. For example, dysfunction of the posterior layer of thoracolumbar fascia has been reported as a "chronic compartment syndrome" of the paraspinal muscles.[26]

Hyaluronic acid

Hyaluronic acid (HA) has a key role for fascia. It is the major glycosaminoglycan of the extracellular matrix (ECM) and a major constituent of connective tissue.[27] Its concentration determines, together with temperature and other physical parameters, the density and properties of the matrix.[16,27-30] In fascia, HA is present within the sublayers and facilitates normal free sliding.[27] Research suggests mechanical therapy increases sliding of layers and decreases pain by transforming the fascia from its gel state to a more fluid solution state.[16,28] Infra-red spectroscopy of viscous HA indicates water molecules can be arranged tetrahedrally almost like as in ice.[29] HA interchain interactions are reversible, disaggregation occurring with an increase in pH and temperature.[29] A gel-like to fluid-like transition occurs at 35-40 °C. Solutions of HA can be highly viscous with non-Newtonian flow properties,[28] and HA's osmotic activity is relatively high; Its stiffness occurs in part due to its long chains forming an entangled large volume network.[29,30] Once viscosity increases, HA can no longer function as an effective lubricant; this increases resistance in layers with abnormal sliding.[27-29]

Fascia and movement

Fascia acts as a key player for generating movement. Although often a dropped prefix, fascia is part of the (fascio)musculoskeletal system.[16,17,25] It is suggested to be able to transmit tension and affect other muscles, reflecting the direction of force vectors, and play a role in properly coordinated movements of the body.[16,31] The anomalous viscosity of HA suggests that it should be an ideal biological lubricant, eg, by reducing the workload during rapid movements.

Sedentarism

Non-versatile movement patterns and sedentarism are an important lifestyle component for fascial changes and pain.[3] A meta-analysis found that low work task variation is a risk factor for non-specific neck pain.[32] Immobilization of a limb or body segment can lead to an increase in the concentration of HA within and between the fascial compartments, which can increase the fluid viscosity and be perceived as stiffness.[28] Studies of immobilization suggest an increase in HA concentration and shortening sarcomere length occur as early as one week after immobilization; After about 4 weeks the collagen fibrils become arranged anbormally.[33,34] This suggests subtle changes in the turnover of HA and in the properties of the extracellular matrix with immobility can lead to structural and functional changes in the (fascio)musculoskeletal system.[33,34] Regular physical activity is recommended for treatment of chronic pain and its effectiveness has been established in clinical trials for people with a variety of pain conditions.[5,35] Proper posture and resting positions are important in preventing muscle tension.[3] There is also a high recurrence rate of pain unless appropriate exercises are prescribed.[5]

Tensegrity

The fact that fascia can transmit tension to a distance is a basis for a "bio-tensegrity" framework.[36,37] Bio-tensegrity applies principles of tensegrity to our understanding of human movement. Tensegrity is an architectural principle. According to this, structures (or tensegrity systems) are stabilized by continuous tension with discontinuous compression.[37]

When the tension of fascia increases, the connective tissue can distribute the forces throughout the surrounding areas, propagating along the myofascial system.[14,16,18] A study by Stecco et al.[38] of upper limb fascia shows all the specimens had deep fascia forming a sheath covering the entire arm. Another study shows manual traction to the biceps femoris tendon results in displacement of the deep lamina of the thoracolumbar fascia up to level L5–S1.[14] Acute bout of stretching of the lower limbs increases the maximal range of motion of the distant upper limbs and vice versa.[18] An aching ankle has been suggested to cause urogenital and visceral symptoms, such as dyspareunia.[39] This can be explained by the muscular connections existing between the pelvis and the ankle, causing increased tone of pelvic musculature.[18,39] A patient can develop various symptoms in areas far from the original dysfunctional point.[16,18,40]

Acceptance of the concept that a fascial tensegrity system connects the whole body is not necessary for a scientific discussion of tension spreading to other areas of muscle groups or structures. Some studies indicate that a tensegrity model might not be a true representation of the whole fasciomusculoskeletal system, and some evidence supports tensional propagation only to certain areas.[41]

Myofascial Pain Syndrome

There are several phenomena that accompany MPS including: trigger points (TrPs), active loci, taut bands, satellites, and the local twitch response.

Trigger points - Changes in fascia are associated with pain in trigger points (TrPs) and tender spots. Most patients with active (ie, painful) TrPs have other areas showing

the same electrophysiologic findings that are not symptomatic, called latent TrPs. Latent TrPs seems to only cause pain upon palpation, whereas active TrPs cause pain and symptoms both at the site of palpation and referred elsewhere.[2,4,9,37] The arising of pain is attributed to those palpable areas in the connective tissue, which seem to activate nociception.[2,4]

Theories of the origin of TrPs emphasize the chronic contraction associated with them or their tendency of appearing at the muscle spindle.[2,9] Others suggest TrPs represent hyperactive end-plate regions.[12] More theories that either deny the existence of MPS or believe it represents a focal dystonia or are of central nervous system origin are also hypothesized.[2,12] Critics of MPS argue that the findings seen at TrPs are simply variants of a normal physiologic finding.[12] There is reason to suspect this is true: about 50% of adults have active or latent TrPs,[2,9] and infants have been observed with point tenderness of the rectus abdominis muscle and colic. These both were relieved by sweeping a stream of vapocoolant, which inactivates myofascial TrPs, over the muscle.[5]

TrPs are extremely common. Among 224 non-specific neck pain patients, TrPs were found in all of them.[9] In studies in pain clinics, 74-85% of those presented to a clinic had TrPs the primary cause of their pain.[9] TrPs are one of the most frequent causes of back pain and neck pain.[42] The severity of symptoms caused by TrPs ranges from the agonizing, incapacitating pain caused by extremely active TrPs to the painless restriction of movement and distorted posture caused by latent TrPs.[5] The influence of latent TrPs on physical function is commonly overlooked.[5] Patients who had other kinds of severe pain, such as that caused by a heart attack, broken bones, or renal

colic, say that the myofascial pain can be just as severe.[5] Myofascial TrPs are present in up to 85% of individuals with colorectal, urological and gynecological pelvic pain syndromes, and can be responsible for some, if not all, of the symptoms related to these syndromes.[43] It was suggested that TrPs or myofascial tension, not exerted by external forces, can apply enough force to cause various abnormalities, including compression of anatomical structures,[3,44] altered position of organs and prolapse,[45] reduced visceral organ motility,[46] chronic changes in bowel movements,[9] disruption of blood and lymph flow,[25,40] etc.[47] If TrPs do not specifically refer pain, these abnormal tissues may still play a role in the patient's pain and may be part of the patient's scoliotic pattern of body organization.[48]

Active loci - TrPs show electrical abnormalities called 'active loci'. This is a small region in a muscle that exhibits spontaneous electrical activity (SEA), often characterized as endplate noise measured on electromyography (EMG), leading to chronic contractions.[9] Although some literature does not regard TrPs as a discriminating factor for MPS, the 'Active Locus' seems to be a consensus.[7]

Taut Band - A taut band is thought to be composed of several TrPs and shows excessive endplate potential activity.[9] Sustained abnormal activation of acetylcholine (Ach) is hypothesized to create an "Energy Crisis" component.[9]

TrP Satellite - TrP satellites are another disputed aspect of MPS. Some literature states TrPs themselves can induce motor activity (referred spasm) in other muscles.[9] Illustrated by an example where pressure on a TrP in the right soleus muscle induced a strong spasm response in the right lumbar paraspinal muscles. Pressure applied to

a TrP in the long head of the triceps brachii muscle indued a strong motor unit response in the upper trapezius muscle only during the 20 seconds that pressure was being applied. This response failed to be reproduced following inactivation of the triceps TrP. It is known as a 'satellite' phenomenon.[9,44,49] Moreover, it seems TrPs can spread, since an active TrP in one muscle can induce an active satellite TrP in another muscle.[9]

Various hypotheses have been suggested to explain the mechanism of MPS, among them are trigger points, non-muscular sensitization of the nervi nervorum, central nervous system, and several other theories.[9,12,50] Some relate the muscle contractions in TrPs to a myotatic (stretch) reflex evoked by fascial tension.[9,12]

Myofibroblasts

Fascia itself is able to actively contract. Tensional alterations are caused by contractile cells, creating a fascial tonus that is independent of neurological modulation.[51] Myofibroblasts are present in some developing or normal adult tissues, altering tissue tension.[30,51] Although most tissues exist under a mechanical tension, the same is not necessarily true of their resident cells; These are protected from the relatively massive external loads by the mechanical properties of the surrounding matrix.[51] In engineering terms, this is called 'stress-shielding', occurring due to the matrix material that they deposit and remodel.[51]

Normal fibroblasts are highly sensitive to the physical stimuli. Tomasek et al. show changes in tissue rigidity, shear forces, and strains are mechanical cues sensed by fibroblasts to transduce into new phenotypes.[51] Thus, the trans-differentiation from

fibroblasts into myofibroblasts is influenced by mechanical stress. Under tension, fibroblasts will differentiate into proto-myofibroblasts, which contain cytoplasmic actin stress fibres that terminate in fibronexus adhesion complexes.[51,52] The adhesion complex bridges the myofibroblast's internal microfilaments with extracellular fibronectin domains. This serves as a contractile mechanism that enables these cells to generate force in the nearby ECM by traction. This contractile force is maintained over time and reinforced further collagen deposition.[51] Increased expression of alpha smooth muscle actin (α-SMA) is directly correlated with increased force generation by myofibroblasts in a positive feedback regulation.[51] A vicious cycle is suggested, in which tension facilitates TGF-β1 signaling, which induces α-SMA expression. In turn, this increases development of more tension, which upregulates TGF-β and α-SMA repeatedly.[51] In short, contracting cells generate the mechanical conditions that make them even more contractile in a detrimental loop.[51] In contrast to the reversible short-lived contraction of striated and smooth muscles, myofibroblast contractility is different: together with ECM synthesis and degradation, they lead to connective tissue remodeling, followed by irreversible and long contractures in a process that can span for a period of time.[51] It is thought that myofibroblasts use a lockstep or ratchet mechanism of cyclic and incremental contractile events via Rho-kinase system, aside the myosin light chain kinase (MLCK) system.[51] When viewed at time frames of seconds and minutes the forces of myofibroblast tissue contractions are not sufficient to exert a significant effect on mechanical joint stability. However, when viewed in a time frame of several minutes and longer, such cellular contractions can impact motoneuronal

coordination.[53] Over a time-window of days to months, this cellular activity can induce long-term and severe tissue contractures.[25,51]

It is also possible that myofibroblasts can couple their activity directly to other cells like myocytes via gap junctions and contract as a unite.[51] Interestingly, if the collagen lattice is released from its points of attachment such that stress in the matrix is lost, the cells rapidly undergo isotonic contraction and subsequently lose their stress fibres and fibronexus adhesion complexes.[51]

Treatments for MPS

Several pharmacological and non-pharmacological treatments are suggested in literature.

Non-pharmacological:[4,9,11,54-57] including manipulative therapy, vibration therapy, exercise, etc. Since it is shown that needle insertion reduces pain and SEA on EMG,[56,58] this suggests a therapeutic role for mechanical needling.

Pharmacological: [4,9-11,56,57] including non-steroidal anti-inflammatory drugs,[56] opioids,[56] topical creams,[56] and TrP injections.[55,56]

Discussion

Synthesis of the data gathered in this review leads the author to the following analysis and suggested mechanism:

There are three aspects of fascial induced pain to consider, from the standpoints of physics, biomechanics, and cellular biology. These seem relevant to fascial tension, TrPs, and MPS.

The standpoint of physics relates to energy. Thermodynamically, the human body can be seen as an open system of energy, and so can the fascial system. Although energy and entropy are modulated by various factors, by nature, entropy of this system will spontaneously tend to increase. To tense the fascia energy needs to be expended. Once fascia is tense the energy can remain as potential elastic energy or change due to other modifications, for example molecular rearrangement. HA has a molecular structure that enables it to have a high degree of entropy, like an entropic "sponge". When entropy increases it generates osmotic pressure and diffusion of fluids and other processes. Thus, when entropy of HA increases, osmotic pressure increases, and stress decays. If stimulus is not removed after this acute stressing, fascia will remain in the failure part of the stress relaxation curve and will be remodeled in this state. Although entropy goes together with osmosis and stress decay, each oscillation between different energy states will likely cause a loss of energy being a non-ideal system, for example due to friction. Fascia has hysteresis and is not a perfect "energetic spring" to oscillate between high-tension and low-tension states, ie, it dissipates energy.[20]

Secondly, from a biomechanical standpoint the review shows that (i) simple stress decay and recovery from it is a reversible process due to elasticity of fascia; (ii)

trigger points (TrPs) and myofibroblasts are found in normal individuals; TrPs are seen even in childhood and infancy, and in animals.[58] (iii) fascia continuously remodels itself and HA is continuously synthesized and excreted, (iv) a sedentary lifestyle and repetitive overuse of muscles, with low variability of movements, all lead to fascial changes/fibrosis and possibly MPS; (v) exercise and proper movement relieve and prevent MPS; (vi) MPS seems to recur unless appropriate exercises are prescribed.

These suggest an underlying ongoing interplay between movement and sedentarism. Since tensional changes are reversible in fascia of normal individuals and depend on the mechanical state of the human body, this indicates fascia is meant to withstand and continuously undergo dynamic changes according to varying dynamic mechanical states. Since immobility often leads to HA viscosity, collagen alterations, and pain, we can see MPS as a natural consequence of the sedentary lifestyle. Resulting from the evolutionary "price" biology paid in a trade-off with chemistry when trying to "come up" with a tissue as effective, pliable, compliant and strong as the fascia. This "bargain" served us well to function effectively as an organism dominated by continuous movement. It seems that by living sedentarily, the natural disadvantages of the properties of fascia (chondrifying, HA aggregation, failure to decay, and fibrosis) manifest pathologically. If indeed nature designed a way to relax fascia during movements and reverse these changes it might happen through (i) dissipation of fascial energy as mentioned above; (ii) mechanical effect of movement that ruptures fascial fibers, (a study shows forces exerted by weights as low as 2-6 kg are enough to rupture the fascia)[38] or shaking off fibers, which might explain why

extra-corporeal shock wave therapy is shown to be effective for MPS;[57] (iii) friction between sliding layers of fascia elevating the local tissue temperature, or "warming up" myofascial tissue. Literature suggests the three-dimensional supramolecular assembly of HA breaks down progressively at 35-40 degrees celcius.[29] The more viscous is HA, the more friction will act to counter viscosity;[28] (iv) increasing clearance of HA due to increased lymph flow. Studies suggest HA increases after immobilization and is perceived as stiffness, and that higher concentrations of HA lead to its more gel-like state;[29] (v) pandiculation; (vi) self-aware "palpation" as a part of one's lifestyle.

The third aspect to consider is a cellular one: induction of myofibroblast by tension. Chronic tension is exacerbated with more tension generated by myofibroblast smooth muscle actin fiber contractions. This makes it increasingly more difficult to maintain a relaxation. Tension is converted to HA aggregation and entropy so long as compensation along the stress relaxation curve can allow for it. Increase in HA concentration is expected to keep HA in the denser gel-like state,[29,59] which is perceived as stiffness.[28] If the values of tension reached at the stress relaxation curve plateau tend be above the value of the threshold for myofibroblast differentiation, it will be a major driver for MPS (values may vary between individuals). Biologically, the reinforcing of fascial cells with actin fibers may be the way of the body to express support of the repetitive muscular effort. Accumulation of many microscopic stretch lesions along a sensory or sympathetic nerve, due to extracellular fibers pulling from multiple directions, might cause peripheral nerves to be hyperirritable chronically.

These three dimensions seem to be relevant to fascial tension, TrPs and the etiology of MPS.

Pathophysiological considerations

Manifestation of MPS is seen in TrPs. Latent TrPs may be points that have previously nociceptively sensed tension (ie, active TrPs), for example from sedentary behavior or a 'satellite', and then were mechanically induced to stress shield themselves via myofibroblasts. The stress shielding does not necessarily eliminate tension (and pain) completely, thus preserving some mechanical drive for further proto/myofibroblast activity. Moreover, new myofibroblasts generate alpha smooth muscle actin (α-SMA) and contractions themselves.[51] When mechanical stimuli continue, fascia remains more towards the failure part of the relaxation curve, and cells have time for stress shielding and matrix remodeling. Then, latent TrPs (via myofibroblasts) start to exert their own tension. This reinforces the abnormality. However, that tension would be more diffuse as it spreads along an intricate web of fibers (ie, stress shield), inducing more tensions, more force gradients in fascia, more stretch lesions, and other painful TrPs further away. If mechanical tension is sufficiently reduced by shielding or by other means, myofibroblast will dedifferentiate or undergo apoptosis,[51] leaving behind a remodeled fascia. Multiple iterations of the contractile cycle result in incremental and irreversible tissue contracture.[60]

These dynamics may be the basis for the unexplained "metastasis" of TrPs and satellites that Quinter et al. (2015)[7] refer to. When sedentary behavior continues, it

feeds more tension down the cascade. The clinical manifestation would be determined not only by the amount of myofibroblasts, but by the interconnectedness of their adhesion complexes, gap junctions, and fiber-cellular networks as well. Studies suggest myo/fibroblasts can form a body wide cellular network and exhibit coordinated calcium oscillations,[62,63] and they can actively respond to mechanical stimuli.[61] Studies suggest myo/fibroblasts have stretch activated calcium channels, and that intracellular calcium and myofibroblast contractility is mechanistically linked.[60,61] Langevin et al.[62] state: "Our findings indicate that soft tissue fibroblasts form an extensively interconnected cellular network, suggesting they may have important and so far unsuspected integrative functions at the level of the whole body."

Activating a latent TrP is hypothesized to involve interference with the stress shield or stretch-activating myofibroblasts to rapidly contract, thus exposing the area to higher tension and initiate nociception. Eliminating a TrP seems to induce a local twitch response, which possibly signifies the neuron or muscle's calcium driven response to a recovery from an active locus or shielded tension. Alternatively, it may be on-off "flickering" of the myotatic reflex due to nearby myofibroblast rapid MLCK contractions and their waves of calcium flux. Active loci may be a multifactorial phenomenon. For example, stress shielding (with or without gap junctions) superimposed on sympathetic activity, ie, a myotatic reflex on top of sympathetic drive. It would seem plausible to expect one to "learn to relax" or "learn to cognitively override the sympathetic aspect by focusing attention", but not to willfully override a reflex response. The fact that muscle spindles are surrounded by

a capsule of connective tissue, an area that sometimes might be palpated as a nodule (ie, TrP) suggests that if there are microscopic tensions, they might possess a sensitivity to generate more tension at the spindle (via proto/myofibroblasts), thus locking the stretch reflex to a certain degree. Locking the reflex would "turn on" an active locus in the associated muscle. To complicate the above, if electrical coupling of this structure occurs with an endplate (or non-intrafuseal fibers), via myofibroblasts, it might perpetuate an "arrhythmic" picture on EMG, with background noise, and random spikes upon myofibroblast contractions. Mechanical disruption of this structure eg, manipulative therapy, should unlock this pathology, at least partially. This might be one of the reasons for different degrees of body relaxation and the inability of some to experience it.

Notwithstanding, other factors are not excluded. Factors associated with the central nervous system could also be implicated in the genesis of MPS, as well as genetics, environmental factors, chronic sympathetic "freeze" reaction to everyday psychological stress, etc. Based on this discussion, we might expect factors affecting fascial myofibroblasts to have potential to cause tension, pain, or even lead to MPS. For example, diet is believed to play a role in MPS,[9] and studies suggest myofibroblasts are promoted by certain diets and by pesticides/herbicides.[64-67] Similarly, the molecule CCN2/CTGF is suggested to play a role in development of pain due to overuse,[68] and CCN2/CTGF is shown to be important in myofibroblast α-SMA synthesis.[69] Applying this sort of logic, we may find more factors linking myofibroblasts to myofascial pain.[70-73]

It is unclear where the tipping point between a status quo of movement-sedentarism and a deranged myofibroblast activity is. It is likely multifactorial, but lifestyle seems to be a major factor. A mechanical stimulus causes differentiation to myofibroblast,[51,74] some studies suggest a threshold of ~20-24 kPa.[75,76] Therefore, mechanical forces might be one of other possible factors contributing to fascial sustained tension and myofibroblast induction. For example: sedentarism and muscle overuse,[3,9,75] infection or inflammation,[4,28,78] and trauma/fracture.[4,9,74,79,80] Since MPS is so prevalent, also common overlooked factors in a lifestyle that is not evolution-oriented may be suspected, eg, medications that are shown to both induce myofibroblasts and increase risk of chronic pain conditions.[72,73]

Symptomatologic considerations

TrPs and myofascial pain, particularly when severe, can cause various symptoms that are not limited to the fascial-muscular-skeletal system. It is suggested that the sympathetic nervous system (SNS) causes TrPs.[12] Reciprocally, pain and TrPs can activate the SNS.[2] This activation may actually be due to coactivation of larger diameter mechano-sensitive afferents rather than nociceptive afferents.[81] A recent meta-analysis finds osteopathic treatment exerts greater parasympathetic responses when stimulating the cervical and lumber regions, while greater sympathetic responses are found when stimulating the thoracic region.[82] If the SNS is activated by paraspinal mechanical stimuli,[81,82] then maybe it can be activated by muscular tension alone. If so, this may serve as a slippery slope for further TrPs, active loci and MPS. Moreover, it seems that when MPS is severe, a widespread pathology may manifest like a fasciomuscular armor under the skin (ie, "fascial armoring"). For

example, immunohistochemical examination of fascia from low back pain patients demonstrates a myofibroblast density comparable to that found in "frozen shoulders".[14]

People with MPS fail to relax muscles due to a mechanical reason; Myofibroblasts can synthesize α-SMA fibers, which would transform fascia into a muscular contractile tissue. It is not necessarily innervated and can remain contracted for years.[51,83] Since this tension is not innervated by voluntary control, and despite involvement of the SNS, seeing these patients as people who just "need to relax" as first line management is counterproductive. Failure to address the organic aspect of myofascial pain is expected to lead to a detrimental cycle of pain and over medicating (a cycle seen in studies addressing ineffective treatments)[11,84-86]

Finally, sedentary behavior and active lifestyle are widely emphasized in public health. However, much focus is dedicated to cardiovascular and metabolic aspects. This review sheds light on the link between lifestyle and pain and suffering. It is worth noting that the most common recommended treatment of MPS found in this literature review is mechanical, including needle insertion, movement (passive or active, breathwork and vibration), and massage/manipulative therapy. It helps treat MPS or reduce fascial related pain.[2,3,4,9,11,19 ,43,54-56]

Therapeutic considerations

Needling was found as a recurrent modality in literature. Three elements might explain how needling can help treat MPS; These are (I) tensegrity mindset (II) trigger points and satellites (III) myofibroblast generated tension and stress shielding.

Since needle insertion itself seems to relieve pain,[87] this may indicate needling

creates a focus for a mechanical tearing action on the fascia. Meaning, the tension

and shear stress inside fascia would pull on the weak point of fine needle insertion

over time, until the sum force vectors, or horizontal components, at that point is

eliminated or needle is removed. Cutting off tension and fascial fibers is expected to

cause cells to rapidly contract, and subsequently lose their stress fibres and adhesion

complexes.[51] If proto/myo/fibroblasts are present (as a node or a line), tearing fascia

would affect the tensegrity structure. Mechanical stimuli can induce myofibroblasts

stretch activated calcium channels, and calcium is coupled to contraction in these

cells.[60,61] After calcium influx, MLCK activity might lead to increased contractile

forces and tension acutely, if they do not lose their stress fibres and adhesion

complexes. If needles continue to tear the fibers, tension will decrease. Studies

observe increase or decrease in contractility and stiffness, and witness needle grasp,

depending on mechanical stimuli.[88-91] Fibroblasts create tension in the attached

matrix, and upon release of tension cells undergo contraction resulting in a rapid

contraction of the collagen lattice.[89] So, it may be that any "damage" by the needle

actually allows for cutting off tensions between tense nodes in the network of

(myofibroblast) TrPs and satellites. Mechanically inducing changes,[88,91-93] and

allowing for the returning of fascia to its place. If inserted deep enough, it might also

free edematous fluid, HA, or other factors trapped within the layers (eg, serotonin)[94].

This would gently change the dynamics of the structure and the entropy (hidden

tension) of the system. Studies indicate needling lowers myofascial stiffness as

measured by shear wave elastography,[95-97] suggesting a potential for a mechanical

"re-alignment" of the fasciomusculoskeletal system and actual tensional release.

The more stress-shielding is present at the locus of insertion, the longer it should take for relief to happen. Stretch lesions along sensory nerves should be relieved too. This treatment acts like breaking the connections in a geodesic dome of the tensegrity model and then allowing for new healthier connections to form by natural remodeling. (The following video observations may help grasp this model)[98-101] Eliminating one node is not enough in this framework, on the contrary: if done improperly, it can only add tensions to other nodes in the structure and exacerbate the general pathology. Starting the relaxation process away from the focus of pain makes sense in this framework. It slowly and progressively begins to relax the primary, secondary, tertiary etc. nodes in the satellite network, approaching the highest area of unshielded tension later and collapsing the tension in a concentric way. According to the biotensegrity model, the peripheries carry tension affecting the focus and are equally important. Noting a statement by Li et al.: "Currently available needle delivery systems deform and move soft tissue and organs" (Li et al. 2020),[102] thus, alignment of the system should be done properly. The question is: What is properly? What is proper needling? Where, in which order, depth and time, is best for which symptom? Studies suggest needling affects facia at the point of insertion and at a distance.[103-105] How fast should the fascia be released? Indeed, it would be beneficial if we could monitor the mechanical state of fascia by palpation. By releasing the tension properly, the tissues can return to their healthy alignment without other areas exerting pulling forces. Following the "damage" of needling, the fascia will regenerate via myo/fibroblasts matching the bio-mechanical state. Evidence suggests that fascia regenerates within approximately 3-24 months after fasciectomy/fasciotomy.[106,107] In fact, it is unlikely that tearing the fascia (if done

properly) will seriously compromise physiology as this would have manifested as a complication after every invasive surgical operation. New extracellular fiber deposition will be based on optimized structural conformation.[21] Some may say inserting several needles subcutaneously sounds like a surgical intervention of the fascia. A gross illustration of this framework is in Figure 2.

Examples of these principles may have transpired in several studies. Three findings are briefly presented as follows. A study of 37 patients who were all diagnosed with plantar fasciitis and all had been treated with corticosteroid injection into the calcaneal origin of the fascia.[108] All had a presumptive diagnosis of plantar fascia rupture. 30% described a sudden tearing in the heel, while the rest had a gradual change in symptoms. Most of the patients had relief of the original heel pain, but it had been replaced by a variety of new problems, including dorsal and lateral midfoot pain, swelling, foot weakness, metatarsal pain, and even metatarsal fracture. In all 37 patients there was a palpable diminution and footprints showing flattening of involved arch. MRI showed fascial attenuation. The study's author concluded that plantar fascia rupture had occurred. The majority had resolution of the new symptoms within 12 months. In the rest, symptoms remained.[108]

A second study encountered the same enigma: 68% of patients with plantar fasciitis had a sudden rupture associated with a corticosteroid injection. Although it relieved the original pain, these patients developed new problems: arch or midfoot strain, lateral plantar nerve dysfunction, hammertoe deformity, and stress fracture.[47]

These examples might suggest there is room for a more sophisticated strategy in treating the fascia instead of simply aiming and firing at the pain. Maybe this should be interpreted not as an adverse event, but as a testimony that the treatment modality undermines the tensegrity system. Conceivably, behind this is a release of fascia in an area of very high tension, causing too drastic of a change in the tensegrity structure, instead of approaching this tension gradually. A sudden change might shift the forces to other areas and exacerbate the imbalances,[109] perhaps sufficient to cause bone fracture. Moreover, it was found that steroid injection decreases fibromatosis, fibrosis, and myofibroblasts in adhesive capsulitis,[110] suggesting injection to the plantar fascia has potential to affect myofibroblast generated tensegrity forces.

The third finding that might be interpreted under these principles, is the phenomena of: "following a tooth extraction, a pain behind the ear and on the side of the face in the day or so prior to facial weakness often constitute the earliest symptom of Bell's palsy."[3] ("Harrison's principles of internal medicine" Kasper et al. 2018, p222-223). Guided with a tensegrity mindset, this may suggest a relationship in the "geodesic" face: a pain develops behind the ear possibly as a "reciprocal" point for tension in the jaw, which happens to be near the emergence of the facial and trigeminal nerves from the skull. This is a reasonably susceptible area with curved bones to serve as a focus for tension. Of course, it does not necessarily mean that releasing tension from around the mastoid process is the key to preventing the complication. Tensegrity domes are made of many nodes, and "to change a node is to change the dome". Therefore, pain behind the ear is first and foremost an indication of extreme tension,

and only then pain. A tension like every other myofascial tension, eg, plantar fasciitis. However, this pain is located in a dermatome of a cervical root: which is a distant nerve and is unlikely to be directly injured by tooth extraction. A sudden intervention in the face probably emulates the same action as a calcaneal injection. If high tension is present (for example a temporomandibular tension seen in up to 25% of the population)[2] it will have the same potential energy to damage structures in the face, as implied by a metatarsal fracture. In the face it seems to cause a palsy of a nerve, similar to the foot. The degree of damage is modulated by the degree of the hidden energy.[109] If correct, this means two possibilities present themselves: either this palsy is not a Bell's palsy, or a Bell's palsy is not idiopathic. In any case, it highlights the instrumental importance of gentle and progressive release of fascia in a proper way. Therefore, "not feeling much" may be a good thing when needling fascia. In fact, we can predict sometimes pain might worsen at first, reflecting a non-ideal modulation of force vectors. Less pain might also lead to better compliance from patients. Research has yet to establish empirically if fascial release prior to (or even after) tooth extraction, can affect the rates of this palsy. Modulation of tensegrity vectors may be a factor in cases, for example, where (i) treatment of Dupuytren's disease leads to Boutonniere deformity,[111] (ii) unexplained displacement of a mass occurs during surgery;[112] (iii) treating the elbow for tennis elbow requires a supposedly unrelated shoulder arthroscopic decompression,[113] (iv) simultaneous bilateral digital flexor rupture occur;[114] (v) spontaneous sequential compartment syndromes occur;[115] or (vi) spine surgery leads to compartment syndrome of the foot.[116] It seems plausible since fascia is a system. Fascia appears to connect the distant hip and ankle joints, not only anatomically, but also mechanically, supporting

the concept of myofascial connectivity.[18] The thoracolumbar fascia proceeds with

the gluteus maximus, involving the fascia of the thigh, the leg, and the plantar

fascia.[40] Also, fascia is continuous from the trunk across the upper and lower limbs,[18]

and manual therapy to the suboccipital region is suggested to transmit forces to the

hamstrings.[117]

In conclusion, under the framework of tensegrity, it might be logical to first needle

the dorsal foot and foreleg in plantar fasciitis, for example, assuming something

organic is pulling the plantar fascia and causing pain. Therefore, aiming to primarily

treat the underlying tensional abnormality of fascia, and through this improving the

symptoms of pain. Actually, it seems a TrP with severe localized pain may not be the

generator of tension because myofibroblasts stress shield their area from tension.

Tension is likely to focus more near hard and angled surfaces that have sharp force-

gradients; initially generating active TrPs (if a nerve is there to experience this) that

have yet to sufficiently stress-shield themselves with a fiber web. By this rationale

we may expect points to be concentrated more in areas of fascia adjacent to solid or

bony edges (spine, face, pelvis, chest, shoulders, knees, periosteum, perineurium,

etc.). Tension will be present,[118,119] but pain depends on the degree of sensory nerve

involvement.[120,121] This understanding may put MPS and fibromyalgia on the

spectrum of a common mechanism; whereby a more widespread pain and

tenderness appears, at least in part, perhaps due to an 'archipelagos' of stress-

shielded points and taut-bands that generate and mask extreme tension. If true, a

severe manifestation of this might be described by some as equivalent to a sort of

mild to moderate chronic compartment syndrome of the whole body. Which may

explain why "MPS" can "develop" into "fibromyalgia", why these orphan pain syndromes are epidemiologically associated with one another (as observed in population studies),[122,123] why significant overlap is seen between them (implying anatomical overlap of involved fascia), why fibromyalgia tender points seem to distribute more near bone protrusions, why a study finds fibromyalgia symptoms and neuropathic pain correlate with tender point count (Amris et al. 2010)[121], and why mechanically puncturing their fascia recurs in literature as an empirical treatment option for them. It can also help explain histological and clinical evidence indicating a picture of widespread non-severe chronic ischemia. Studies of fibromyalgia observe various organic abnormalities,[124-141] which might support this framework.

Considering fibromyalgia as an entity of compression

Studies show tissue stiffness is significantly increased around TrPs,[142] and blood vessels near TrPs have retrograde flow in diastole, "indicating a highly resistive vascular bed" (Sikdar et al. 2009)[138]. The above analogy might offer context to few findings that will be mentioned here briefly.

Various cardiovascular abnormalities in fibromyalgia suggest an underlying widespread organic pathology that impairs perfusion throughout the vascular system. Capillary microscopy studies of fibromyalgia show fewer capillaries in the nail fold and significantly more capillary dilatations and irregular formations than the healthy controls. The peripheral blood flow in fibromyalgia patients was much less than in healthy controls.[128] This suggests that functional disturbances are present in

fibromyalgia patients.[128] While another study shows a decrease in transcapillary

permeability in fibromyalgia patients,[129] the findings of Cho et al.[143] might provide

the most insight: fibromyalgia patients exhibited decreased endothelial-dependent

vasodilatation and endothelial-independent vasodilatation in arteries. These

parameters were in parallel with pain and fibromyalgia impact questionnaire

(FIQ). The study did not find strong evidence for a cardiovascular disease that could

explain this impairment. Moreover, the authors concluded this impairment may be

due to mechanical forces limiting the artery's ability to relax, but they do not

elaborate on the nature of these forces. Aortic distension seems to be decreased and

impairment correlates with pain and FIQ.[144,145] A chronic sympathetic stress

response should mainly affect arterioles rather than the aorta itself, and these

studies were adjusted for possible confounding factors (eg, cardiovascular diseases).

Similarly, a study of biopsies in fibromyalgia states: "Observations in 12 primary

fibromyalgia syndrome cases without obvious muscle trauma indicate definite but

nonspecific muscle changes which we suspect are secondary to chronic muscle

spasm and ischemia of unknown etiology."[124]

In other body structures, high rates of carpal tunnel syndrome were found in

fibromyalgia.[146] Nacir et al. found twenty percent of fibromyalgia patients have

carpal tunnel syndrome (some bilateral) while only three percent of controls were

affected.[147] A survey of 2,596 fibromyalgia patients finds patients ranked the

intensity of their pain lower than the intensity of morning stiffness.[148] Considering

central sensitization raises several questions: if central sensitization to pain is the

main underlying pathophysiology, why is morning stiffness so commonly reported

among fibromyalgia patients, and even ranked higher than pain?[149] Why are there no reports of an extreme sensation of vibration? Wouldn't some patients be sensitized to vibration due to a long hour daily commute to work? Is central sensitization random or is there a pattern? Does it prefer only certain neural tracts, like pressure and pain? If it is random, why does it seem to localize to the same points in different individuals? If it is not random, why can sensitivity appear in almost any area? Why are these areas not correlated with the homunculus, and what is special about the occiput, second costochondral junctions, knee pad, and two centimeters distal to the lateral epicondyle (Kasper et al. 2018; p2637)[3]? It is worth noting that Sugawara et al.[150] suggest mechanical compression of the dorsal root ganglion (DRG) by a mechanical stimulus lowers the threshold needed to evoke a response and causes action potentials to be fired.[150] Action potentials that may even persist after the removal of the stimulus and high mechanical sensitivity are suggested by this (in vitro) study. More in vivo and in vitro studies suggest the same.[151-153] It would be interesting if pulling could also lower firing thresholds.

Bizarrely, Wang et al. find a bidirectional association between fibromyalgia and gastroesophageal reflux disease (GERD).[154] How is central sensitization/somatization related to this bidirectional linkage? Should we consider GERD to contain a strong component of somatization that is overlooked, or should we give fibromyalgia more organic credit? This entity, if it is what it seems, would probably affect gastroesophageal pressures and relaxations, and theoretically, mechanically compress the paraspinal and intervertebral fascia and its contents eg, DRG and/or the sympathetic chain against a vertebral body. If compression has a similar effect on the sympathetic ganglion/trunk as it does on the DRG, it might add a component of sympathetic abnormalities. Indeed, studies suggest mechanical pressure on the sympathetic trunk due to osteophytes can cause various abnormalities.[155] Studies

suggest para-articular fibrosis in the spine can involve neural structures and cause tethering and irritation of nerves.[156] A compressive factor can explain decreased pressure/pain thresholds and explain why they happen to be strongly associated with autonomic abnormalities. It also implies these abnormalities can be illusive and reversible in certain conditions. By the same rationale we might expect higher incidence of obstructive sleep apnea even after correcting for body mass index.

More studies indicate fibromyalgia may include easy bruising,[149,157] hair-loss,[149] reduced skin innervation,[158] tingling, creeping or crawling sensations,[159] reduced fecundity or infertility,[160,161] urinary urgency,[159] changes in bowel habits,[3] decreased optic disk perfusion,[127] dry eyes, [159] functional voice disorders (including muscle tension dysphonia),[162] wheezing,[149], seizures,[149] impaired cognition,[149] and Raynaud's phenomenon.[163] Can central sensitization and/or somatization of pain, be the explanation of these manifestations? Do psychosocial theories or neuroplastic pain circuits explain these? No single theory seems to fully account for fibromyalgia. For instance, if it is sympathetic hyperactivity, why does it cause wheezing? If it is sympathetic-parasympathetic dysregulation, why is postural hypotension not in the diagnostic criteria, and why is it strongly associated with sicca or but lacks reports of chronic sialorrhea? If it is inflammatory or autoimmune, where are the inflammatory markers and why does it respond to tricyclic antidepressants? Moreover, why is it so common in individuals with acquired immunodeficiency syndrome,[164] and how is it that a study finds placebo is a better treatment for fibromyalgia than are steroids?[165]

If it is stress and low mood superimposed on childhood anxieties, how does it reduce skin innervation and why selective serotonin reuptake inhibitors (SSRIs) are not first line treatment? Central sensitization was discussed above, but still, why is exercise the cornerstone of treatment and what is behind the close association with obesity? The relatively modest efficacy of antipsychotics and/or melatonin and/or z-drugs implies it is probably neither an imaginary hallucination nor due to lack of sleep. Do neuroendocrine theories support the topography of the tender points? Of course, two parallel mechanisms might be at work and are not necessarily incompatible with each other. Gupta et al. 2007 conclude in their study: while a high tender point count is associated with the onset of new chronic widespread pain, a low pain

threshold at baseline is not. An increased sensitivity to pressure pain in subjects with chronic widespread pain is therefore likely to be a secondary phenomenon rather than the antecedent of pain.[166] With our new view of MPS it is logical, since tender point count would be a marker for a fascial tensional pathology, leading to widespread pain and a lower pain threshold later. If in the trunk, this constrictive pathology might be able to impair lung function and account for wheezing. A physiological basis for phenomena like seizures/pseudoseizures will be suggested when mentioning the myodural bridge later. Interestingly, studies suggest tricyclic antidepressants can downregulate myofibroblasts and ACTA2 gene expression,[167,168] while some SSRIs seem to enhance them,[169] like does gp120.[170,171]

A study states that in fibromyalgia a global pathology involving tryptophan might come at the expense of other serotonergic operations such as in the brain.[172] Studies suggest a mechanosensitive effect can trigger serotonin release from enterochromaffin and neuroendocrine cells,[173,174] and studies suggest serotonin is involved in the cellular signaling of connective tissue cells.[175-177] Studies of fibromyalgia patients indicate an elevated level of serotonin in myofascial tissue and platelets, while a decreased level of tryptophan and its derivatives, serotonin and kynurenine, is observed in the blood and/or brain.[94,178-184] A study finds fibromyalgia patients have decreased rate of transport of tryptophan across the blood brain barrier.[172]

Several studies point to a peculiar phenomenon. Saber et al.[185] witness high rates of complete resolution of fibromyalgia following laparoscopic Roux-en-Y surgery. By which mechanism does a relentless longstanding central sensitization completely resolve following abdominal surgery? Could this be a result of an overlooked tearing of the fascia? The authors of that study suggest resolution may be due to weight loss, increased physical activity, and lifestyle changes. However, this idea does not align with findings of Adkisson et al.[169]. They studied effect of parathyroidectomy on fibromyalgia among 76 patients diagnosed by specialist or primary care physician.

Findings show that 21 percent of fibromyalgia patients discontinue all fibromyalgia drug medications after parathyroidectomy, and 89 percent have relief of one or more fibromyalgia symptoms postoperatively. These changes occurring as early as one week after neck surgery.[186] Yet, 11% of patients had no improvement in symptoms at all. The authors explain the complete resolution in 21 percent by suggesting a misdiagnosis, ie, these patients actually did not have fibromyalgia in the first place. However, if this was true, we might expect these individuals to cluster more towards the group not diagnosed by a specialist. Surprisingly, 23 out of 76 were diagnosed by a rheumatologist, out of them 96% had symptom improvement, and 70% were able to discontinue all medications after parathyroidectomy. Uncannily, 88% of those who did not experience any improvement in fibromyalgia symptoms were diagnosed by a primary care physician ($p<0·05$). The statistics suggest some non-random phenomenon caused a bimodal-like distribution. In light of this study, it might be hypothesized fibromyalgia is related to a parathyroid cause. However, a similar enigma is found with irritable bowel syndrome (IBS), a functional disorder with no definite organic findings and closely associated (or overlaps) with fibromyalgia. Some suggest fibromyalgia and IBS are actually the same entity.[149] IBS is shown to be relieved below Rome II criteria in 80 percent of patients after laparoscopic fundoplication for GERD.[187] Looking for a common denominator one can hypothesize that relief is achieved due to intense pain exposure and general anesthesia that resets the brain circuits, somewhat similar to electroconvulsive therapy combining gate control. But it does not align with findings that tonsillectomy increases risk of IBS,[188] that umbilical hernia repair surgery seems to predispose to IBS,[189,] and that patients with fibromyalgia have a higher incidence of suffering IBS

after appendectomies.[190] Moreover, hysterectomy, with or without oophorectomy, seems to worsen fibromyalgia.[191]

Oddly, why does the location and type of surgery seem to regulate the positive or negative impact on these conditions? Indeed, it was suggested in this literature review that MPS/TrPs can cause chronic changes in bowel habits, and that changes in fascia can lead to reduced visceral organ movements. Should we suspect that in these associated functional disorders, the connecting motif is connective tissue? This may imply that the location of the surgery has different effects on fascia, even remotely, due to- and regulated by- tensegrity forces. The lateral raphe and common tendon of the transversus abdominis are suggested to form connections between, and transfer tensions between, the abdomen and the paraspinal muscles and sheath.[14,192] This can explain how laparoscopy might release a compression on the DRG. Interestingly, this would mean the clustering of patients experiencing benefit post-parathyroidectomy towards the group diagnosed by a rheumatologist, and those not experiencing benefit towards the group diagnosed by primary care physician, isn't due to misdiagnosis but precisely because rheumatologist really identify those with "fibromyalgia" statistically significantly more accurately.

Lastly, studies show obesity induced connective tissue fibrosis is dependent on mechanosensitive signaling.[193] Fibrosis, via myofibroblasts, alters subcutaneous tissue plasticity and increases connective tissue rigidity and stiffness.[193] Thus, obesity might be associated with fibromyalgia by way of mechanically compressing fascia with higher forces. A Study in vivo suggests applying tension on skin induces myofibroblasts.[74] Therefore, by this same rationale we might expect a sedentary

individual that tends to wear tight clothes and accessories that mechanically compress fascia to be at higher risk for this pathology over time. Studies suggest the association between the higher incidence of fibromyalgia and estrogen is unlikely.[185,194] Since estrogen is suggested to inhibit myofibroblast differentiation and is associated with lower fascial stiffness,[195-198] and since studies suggests hormonal contraceptives may actually reduce the risk of fibromyalgia in women,[199] this could be part of the explanation to why fibromyalgia is increased in certain females. Some suggest fibromyalgia may be linked to menopause.[200] Indeed, studies suggest applying a splint induces myofibroblasts,[80] and pony-tails/hijabs/braziers appear to cause myofascial pain.[201-205] This phenomenology looks to be a similar or the same as that of applying bandages to "rheumatism" patients (William Balfour 1815),[206] meaning: modulation of the tensegrity structure and/or induction of stress shielding but failing to resolve tension (eg, Mrs. M. Pages 179-180). Interestingly, experimental models of restraint stress apply mechanical restraint and/or limit movements. These models show mechanical and cold allodynia, depression and anxiety like behavior, gut dysmotility, and other abnormalities.[207-211] Out of curiosity, what is the pluripotent entity that restraint stress tries to model? Is it just emotional stress?

In any case, perhaps a treatment or prophylaxis that applies multiple needles in different areas of the tensegrity system, could be utilized as a global percutaneous needle fasciotomy. The theoretical end purpose this treatment would be to lower global fascial stiffness beneath the threshold necessary for myofibroblasts activity, and slowly experience a tensional collapse. Literature indicates fascia of patients

with fibromyalgia measure higher on shear wave elastography, and that needling

reduces pain.[93,95,212-218] Studies also show needling significantly lowers the shear

modulus of myofascial tissue as measured by elastography ($p < 0.01$),[93,95,97] causes

mechanical changes at the point of insertion and at a distance,[90,102,103] and improves

perfusion to the area.[219] A systematic review finds evidence that needling improves

pain and stiffness in people with fibromyalgia.[220] Also supporting this modality's

capability of improving perfusion, is a study showing infiltration of TrPs reduces

symptoms of intermittent claudication.[221] Symptomatically, a person constantly

exerting effort against such forces of a suffocating fascia would likely feel chronic

fatigue and soreness, whilst accumulating nociceptive substances in muscles in a

chronically tonic body, and widespread "energy crises". A reader adopting this

"biological ethos" should not be surprised, for example, by a study finding that

fibromyalgia patients have elevated serum tau and amyloid beta levels, since studies

suggest amyloid beta and the cytoskeletal protein tau and are affected by, or

compensate for, mechanical stress.[221-228]

At least we might want to exclude an organic fascial cause before perceiving these

cases as psycho-functional disorders, even if the mechanism is still unclear, and even

if it involves a degree of central sensitization. Is there enough empirical evidence to

recognize MPS and fibromyalgia as honorable organic diseases, like any other, rather

than psychosomatic? Medicine is used to entities that are either immune related,

infectious, malignant, endocrine, metabolic, genetic, cardiovascular, traumatic or

toxic, but not familiar with many mechanical ones. If medicine overlooked another

system causing pain besides fascia, like integumentary or skeletal, it would probably lead to overlying on psychology to explain it.

The enthusiast of the principle of Occam's razor (ie, minimize multiplication of entities) might consider seeing other orphan-of-a-mechanism somatic entities with a pathology known to be associated with pluripotent TrPs,[2,43,227-230] as "MPS"/"fibromyalgia" that is localized to certain anatomical areas. "Kelley's textbook of rheumatology" and others suggest functional somatic syndromes to be on a continuum of one entity with fibromyalgia,[149,162] In addition, MPS and fibromyalgia are suggested to be two sides of the same coin.[2,231-233] Etymology does not necessarily reflect pathophysiology, but term "fibromyalgia syndrome" basically carries the same meaning as "myofascial pain syndrome" but in Greek. The analysis based on this scoping review offers one mechanical mechanism as a possible common pathophysiology.

On a final note, based on this review, it seems that fascia is an intelligent and sophisticated tissue network, at least like any other tissue network we know. It is shown myo/fibroblasts have gap junctions that can couple and directly communicate with each other (Xu et al. 2021)[237] and with other types of cells (with heart,[238] hair follicles,[239] cancer cells,[240] etc.)[241-244] and these can contract as a unit or in patterns,[244,245] affecting the organ's functionality.[246-258] If this theoretical model of "fascial armoring" is verified empirically, it might be worth further exploring the possibility that they can couple in vivo (directly, or indirectly through myocytes/glial cells) to nerves. Studies suggest the meninges might be coupled to the cerebral cortex to form one large network.[259-262] Studies suggest the myodural bridge

connects the extracranial occipital fascia with the dura.[263] Myo/fibroblasts seem to

form a body-wide cellular network that can respond to mechanical cues and exhibit

spontaneous calcium oscillations and synchronized contractions,[245,264-267] and

fibroblast themselves can induce contracture through actin beta and gamma

pathways.[268]

We could consider the hypothetical possibility that this dynamic body-wide

mechanoelectrical network of cells and fibers has an overlooked interplay with the

network of the human nervous system. This bizarre idea might disguise some of the

pathophysiology of self-psychosomatics ("so everyone chronically pulls themselves

together. This feeling of tightness is the physical referent for the psychological image

of ourselves."[269] etc,)[269-273] and potentially help us understand why a remarkable 17

percent of primary care patients are diagnosed with somatic symptom disorder

(defined in studies as at least four or more unexplained symptoms)[274]. We know

biology does not separate itself into different medical specialties like we do

occupationally.

Lastly, if myofibroblast can engulf their surroundings,[275] and in certain conditions are

able to present antigens via MHC class II to T cells,[276-280] should we examine the

potential role of myo/fibroblast fascial networks (or differentiated/somatic mutated

clone populations of them) in immune related conditions? Empirics suggest a link

between fibromyalgia and other rheumatological disorders, and apparently, a study

of 20 patients with fibromyalgia shows the majority have anti smooth muscle

antibodies.[281,282]

On a spectrum of "normality" we may all have "MPS" to some extent, technically

speaking. It seems to be a question of how much we have and how much it affects our

life. A tradeoff is still being made, no longer by biology but by humans. Not regarding

the molecular design of fascia, but it is an ongoing tradeoff between not moving and moving. As it seems to be a natural condition worsened by an un-natural lifestyle, we may say living sedentarily misses our evolutionary target (ie, a "biological sin"). It is interesting a study by Younger et al. finds patients with myofascial pain have reduced grey matter in the hippocampus, anterior insula, and cingulate cortex, among several areas:[241] Forgetting our evolutionary purpose seems to impair our memory, while cognitive detachment from our body-awareness seems to impair our cognition, and because the brain's biogenic amine metabolites are abnormal, the above suggests a lifestyle with scarcity of movement will rarely make us blissful. Looking through the given titles of these entities, we might find what we most want to find where we least want to look. A body that cannot move with total freedom will hold a mind that is seldom totally free of dis-ease. This aligns with the notion that the body and the mind are accepted to be one being. Just as we can say music is a "collection of noises" or a "divine melody", so may we say MPS is "pulled muscles" or a "fascinating disorder of the fascia". Assembling these conclusions and concepts to creatively connect this medical puzzle consists the easier part of this not so easy effort. The major challenge now being the research needed to verify them and moving forward with evidence on this topic while moving ourselves…

Limitations:

The main limitation of this scoping review is that a sole researcher evaluated the studies. It is based on other empirical studies where some are in vitro studies, as mentioned in parts of the discussion. Being a scoping review, although a broad searching strategy was applied, not every topic was searched systematically. A systematic search was done for several main key phrases, in five databases. Systematically searching for relevant side topics (ie, topics not defined in the key phrases) was not feasible, unfortunately. For side topics, an attempt to find a reliable

source of information, like a textbook or study (or both), was the strategy to gather information. In such cases an attempt to cite more than one study was made. Worth noting, no on topic studies were found when searching "fibromyalgia myofibroblast" or "fibromyalgia alpha smooth muscle actin". The cytokine profile and the topic of vitamin D in fibromyalgia was not discussed even though a study finds vitamin D might play a role in myofibroblast attenuation or de-differentiation.

Figure 2: Illustration of needling in fascial tensegrity framework

a) Illustration to assist the reader in imagining a healthy "geodesic dome" connected by spandex sheets (fascia)
With permission from PACIFIC DOMES Inc. www.eventdome.wordpress.com/

b) Normal external strain applied will tense the fascia and distribute the tension through the network of nodes.
With permission from Dome guys international www.domeguys.com

c) Chronic over-strain causes pathological changes in fascia or changes of myofibroblast and smooth muscle fibers inside the fascia. The fascia fails to distribute forces properly and keep the integrity of the structure. Structures on the dome and inside the dome are affected.
With permission from PACIFIC DOMES Inc www.pacificdomes.com

d) After 're-setting' the system with needling or other techniques, regeneration and growth of new connections over time should be determined by natural forces and pullies.21 New areas in the fascia serve as points of focus for changes and realignment. Structures can relax and return to the minimum energy state. "Fascia is always subjected to remodeling pressures and responds to the local mechanical state. However, if spatial deposition of fibers is altered with respect to physiological conditions, the rebuilding will be pathological."21 Mobilization encourages correct healing in order to avoid formation of fibrosis.21

Taken by Jim Bourg, with permission from REUTERS News Agency.

Contributions: this work was done by the sole researcher and author of this

manuscript including conception, organization, literature review, integration of

information, findings analysis, manuscript preparation and writing.

Acknowledgements: Professor Pnina Ohanna Plaut for assisting with methodology

and editing. Dr. Bronya Gorney for assisting with editing. Carrie Rodomar, librarian of

University of Nicosia Medical school for assisting with methodology.

Conflict of interest: The author declares no conflict of interests.

Funding: none

References:

1.   WHO Scientific Group on the Burden of Musculoskeletal Conditions at the Start of the New Millennium. (2003 : Geneva, Switzerland). (2003). The burden of musculoskeletal conditions at the start of the new millenium : report of a WHO scientific group. Geneve : World Health Organization. https://apps.who.int/iris/handle/10665/42721. Accessed November 2020

2.   www.uptodate.com. search "myofascial pain syndrome", "pelvic pain", "tmj" accessed November 2020. "Trigger points" "Fibromyalgia", "differential diagnosis of fibromyalgia", accessed April 2021.

MPS is a regional pain disorder caused by the presence of trigger points within muscles or their fascia. "MPS may include other common regional pain disorders such as tension headaches, idiopathic low back and cervical strain disorders etc" Trigger points, the hallmark of myofascial pain… produce sensory, motor, autonomic symptoms (eg, coldness, sweating, tachycardia, coryza, lacrimation, and erythema) Some authors have concluded that FM and myofascial pain are "two sides of the same coin"

3.   Kasper DL, Fauci, A. S., Hauser, S. L., Longo, D. L. 1., Jameson, J. L., & Loscalzo, J. Harrison's Principles of Internal Medicine, 20e. New York; 2018. Pages 222-223, 2637-2639, 2644-2645

"Fibromyalgia can overlap in presentation with other chronic pain conditions… headaches, facial/jaw pain, regional myofascial pain particularly involving the neck or back, and arthritis. Visceral pain involving the gastrointestinal tract, bladder, and pelvic or perineal region is often present as well. Patients may or may not meet defined criteria for specific syndromes."

4.   Giamberardino MA, Affaitati G, Fabrizio A, Costantini R. Myofascial pain syndromes and their evaluation. *Best Pract Res Clin Rheumatol.* 2011; **25**(2):185-98.

5.   Siegfried Mense DGS, I. Jon Russell. Muscle Pain: Understanding Its Nature, Diagnosis, and Treatment. Philadelphia; 2001 Pages 1-10.

"Myofascial pain caused by trigger points…"

6.   Lucas N, Macaskill P, Irwig L, Moran R, Bogduk N. Reliability of physical examination for diagnosis of myofascial trigger points: a systematic review of the literature. *Clin J Pain* 2009; **25**(1): 80-9.

7.   Quintner JL, Bove GM, Cohen ML. A critical evaluation of the trigger point phenomenon. *Rheumatology (Oxford)* 2015; **54**(3): 392-9.

8.   Phan V, Shah J, Tandon H, et al. Myofascial Pain Syndrome: A Narrative Review Identifying Inconsistencies in Nomenclature. *PM R* 2020; **12**(9): 916-25.

9.   Simons DG, Janet G. Travell, and Lois S Simons. Travell & Simons' Myofascial Pain and Dysfunction: The Trigger Point Manual. 2nd ed. Baltimore; 1999. (Pages xi- 87, 940-955)

10.   Wheeler AH. Myofascial pain disorders: theory to therapy. *Drugs* 2004; **64**(1): 45-62.

11.   Fleckenstein J, Zaps D, Rüger LJ, et al. Discrepancy between prevalence and perceived effectiveness of treatment methods in myofascial pain syndrome: results of a cross-sectional, nationwide survey. *BMC Musculoskelet Disord* 2010; **11**: 32.

12.   Rivner MH. The neurophysiology of myofascial pain syndrome. *Curr Pain Headache Rep* 2001; **5**(5): 432-40.

13.   Panta P. The Possible Role of Meditation in Myofascial Pain Syndrome: A New Hypothesis. *Indian J Palliat Care* 2017; **23**(2): 180-7.

14.   Willard FH, Vleeming A, Schuenke MD, Danneels L, Schleip R. The thoracolumbar fascia: anatomy, function and clinical considerations. *J Anat* 2012; **221**(6): 507-36.

"It is an intriguing thought that some cases of LBP may be associated with a similar stiffening of the TLF, in which case such a condition could be described as 'frozen lumbars'"

15.   Atlas of Human Fascial Topography. Fascia Atlas Leipziger Universiätsverlag; 2018.

16.   Tozzi P. Selected fascial aspects of osteopathic practice. *J Bodyw Mov Ther* 2012; **16**(4): 503-19.

"Electron photo microscopic studies of the human fascia have shown smooth muscle cells embedded in the collagen fibres, including intrafascial capillaries and autonomic nerves." Autonomic nerve fibres may regulate a fascial pre-tension via the smooth muscle cells contractility, independently from muscular tonus.

17.   Barker PJ, Briggs CA. Attachments of the posterior layer of lumbar fascia. *Spine (Phila Pa 1976).* 1999; **24**(17):1757-64.

18.   Nordez A, Gross R, Andrade R, et al. Non-Muscular Structures Can Limit the Maximal Joint Range of Motion during Stretching. *Sports Med.* 2017; **47**(10):1925-1929.

"Fascia, like peripheral nerves, is continuous from the trunk across the upper and lower limbs and hence has the potential to influence range of motion." It has dense sensory innervations making it sensitive to stress/strain that can induce pain. It has been shown to be continuous structures that cross several joints. Unilateral stretching of one limb increased ROM of contralateral limb.

19.   Avila Gonzalez CA, Driscoll M, Schleip R, et al. Frontiers in fascia research. *J Bodyw Mov Ther* 2018; **22**(4): 873-80.

20.   Zeng YJ, Sun XP, Yang J, Wu WH, Xu XH, Yan YP. Mechanical properties of nasal fascia and periosteum. *Clin Biomech (Bristol, Avon).* 2003; **18**(8):760-4.

21.   Pavan PG, Stecco A, Stern R, Stecco C. Painful connections: densification versus fibrosis of fascia. *Curr Pain Headache Rep* 2014; **18**(8): 441.

22.   Nelson-Wong E, Glinka M, Noguchi M, Langevin H, Badger GJ, Callaghan JP. Acute Surgical Injury Alters the Tensile Properties of Thoracolumbar Fascia in a Porcine Model. *J Biomech Eng.* 2018; **140**(10):1010121–7.

23.   Yahia LH, Pigeon P, DesRosiers EA. Viscoelastic properties of the human lumbodorsal fascia. *J Biomed Eng.* 1993; **15**(5):425-9.

24.   Stecco A, Pirri C, Stecco C. Fascial entrapment neuropathy. *Clin Anat* 2019; **32**(7): 883-90.

25.   Klingler W, Velders M, Hoppe K, Pedro M, Schleip R. Clinical relevance of fascial tissue and dysfunctions. *Curr Pain Headache Rep* 2014; **18**(8): 439.

Nerve compression syndromes are always associated with a fascial pathology.

26.   Konno S, Kikuchi S, Nagaosa Y. The relationship between intramuscular pressure of the paraspinal muscles and low back pain. *Spine (Phila Pa 1976)*. 1994; **19**(19): 2186-9.

27.   McCombe D, Brown T, Slavin J, Morrison WA. The histochemical structure of the deep fascia and its structural response to surgery. *J Hand Surg Br* 2001; **26**(2): 89-97.

28.   Cowman MK, Schmidt TA, Raghavan P, Stecco A. Viscoelastic Properties of Hyaluronan in Physiological Conditions. *F1000Res* 2015; **4**: 622.

29.   Matteini P, Dei L, Carretti E, Volpi N, Goti A, Pini R. Structural behavior of highly concentrated hyaluronan. *Biomacromolecules* 2009; **10**(6): 1516-22.

30.   Young, Barbara (Pathologist). Wheater's Functional Histology : a Text and Colour Atlas 5th edition. Edinburgh ; New York :Churchill Livingstone, 2000. Page 67,69,101

31.   Vleeming A, Pool-Goudzwaard AL, Stoeckart R, van Wingerden JP, Snijders CJ. The posterior layer of the thoracolumbar fascia. Its function in load transfer from spine to legs. *Spine (Phila Pa 1976).* 1995; **20**(7):753-8.

32.   Jun D, Zoe M, Johnston V, O'Leary S. Physical risk factors for developing non-specific neck pain in office workers: a systematic review and meta-analysis. *Int Arch Occup Environ Health* 2017; **90**(5): 373-410.

33.   Okita M, Yoshimura T, Nakano J, Motomura M, Eguchi K. Effects of reduced joint mobility on sarcomere length, collagen fibril arrangement in the endomysium, and hyaluronan in rat soleus muscle. *J Muscle Res Cell Motil* 2004; **25**(2): 159-66.

34.   Williams PE, Goldspink G. Connective tissue changes in immobilised muscle. *J Anat* 1984; **138 (Pt 2)**: 343-50.

35.   Lima LV, Abner TSS, Sluka KA. Does exercise increase or decrease pain? Central mechanisms underlying these two phenomena. *J Physiol.* 2017; **595**(13):4141-4150.

36.   Caldeira P, Fonseca ST, Paulo A, Infante J, Araújo D. Linking Tensegrity to Sports Team Collective Behaviors: Towards the Group-Tensegrity Hypothesis. *Sports Med Open* 2020; **6**(1): 24.

Biotensegrity is a functional concept. Tensegrity properties enable the synergies underlying complex human movement in task- and context specific scenarios.

37.   Tadeo I, Berbegall AP, Escudero LM, Alvaro T, Noguera R. Biotensegrity of the extracellular matrix: physiology, dynamic mechanical balance, and implications in oncology and mechanotherapy. *Front Oncol* 2014; **4**: 39.

"The ECM is a dynamic and multifunctional regulator and has its own biotensegrity with Ret F and elastin fibers acting as tensional elements, and ground substance and Col F as compression-resistance elements."

38.   Stecco A, Macchi V, Stecco C, et al. Anatomical study of myofascial continuity in the anterior region of the upper limb. *J Bodyw Mov Ther* 2009; **13**(1): 53-62.

39.   Kotarinos R. Myofascial pelvic pain. *Curr Pain Headache Rep.* 2012;**16**(5):433-8.

40.   Bordoni B, Zanier E. Clinical and symptomatological reflections: the fascial system. *J Multidiscip Healthc* 2014; **7**: 401-11.

41.   Zhou JP, Yu JF, Feng YN, Liu CL, Su P, Shen SH, Zhang ZJ. Modulation in the elastic properties of gastrocnemius muscle heads in individuals with plantar fasciitis and its relationship with pain. *Sci Rep* 2020; **10**(1): 2770.

42.   Fricton JR, Kroening R, Haley D, Siegert R. Myofascial pain syndrome of the head and neck: a review of clinical characteristics of 164 patients. *Oral Surg Oral Med Oral Pathol* 1985; **60**(6): 615-23.

43.   Moldwin RM, Fariello JY. Myofascial trigger points of the pelvic floor: associations with urological pain syndromes and treatment strategies including injection therapy. *Curr Urol Rep* 2013; **14**(5): 409-17.

44.   Siddiq MAB, Clegg D, Hasan SA, Rasker JJ. Extra-spinal sciatica and sciatica mimics: a scoping review. *Korean J Pain* 2020; **33**(4): 305-17.

45.   Crowle A, Harley C. Development of a biotensegrity focused therapy for the treatment of pelvic organ prolapse: A retrospective case series. *J Bodyw Mov Ther* 2020; **24**(1): 115-25.

46.   Tozzi P, Bongiorno D, Vitturini C. Low back pain and kidney mobility: local osteopathic fascial manipulation decreases pain perception and improves renal mobility. *J Bodyw Mov Ther* 2012; **16**(3): 381-91.

47.   Acevedo JI, Beskin JL. Complications of plantar fascia rupture associated with corticosteroid injection. *Foot Ankle Int* 1998; **19**(2): 91-7.

48.   Whyte Ferguson L. Adolescent idiopathic scoliosis: The Tethered Spine III: Is fascial spiral the key? *J Bodyw Mov Ther.* 2017; **21**(4):948-971.

49.   Gerwin RD. Classification, epidemiology, and natural history of myofascial pain syndrome. *Curr Pain Headache Rep* 2001; **5**(5): 412-20.

50.   Quintner JL, Cohen ML. Referred pain of peripheral nerve origin: an alternative to the "myofascial pain" construct. *Clin J Pain* 1994; **10**(3): 243-51.

51.   Tomasek JJ, Gabbiani G, Hinz B, Chaponnier C, Brown RA. Myofibroblasts and mechano-regulation of connective tissue remodelling. *Nat Rev Mol Cell Biol* 2002; **3**(5): 349-63.

Expression of α-SMA is directly correlated with increased force generation. Tissue contraction depends on collagen remodelling, a process that is dominated by ECM reorganization under the mechanical control of myofibroblast contraction. Calcium dependent MLCK would be used to generate rapid contraction, whereas Rho kinase is

important in maintaining sustained contraction. Misconception underlies the idea that cell cooperativity is needed to produce shortening in a particular plane. The force that is generated in tissues could be non-directional, as the pre-existing matrix shape and properties will largely dictate the resultant vectors.

52.   Stempien-Otero A, Kim DH, Davis J. Molecular networks underlying myofibroblast fate and fibrosis. *J Mol Cell Cardiol* 2016; **97**: 153-61.

Fate-mapping studies suggest myofibroblasts revert back to their original cell fate, but with a modified genetic signature permissive of accelerated conversion to a fibrotic myofibroblast upon additional injury

53.   Schleip R, Klingler W. Active contractile properties of fascia. *Clin Anat* 2019; **32**(7): 891-5.

54.   Dommerholt J, Thorp JN, Chou LW, Hooks T. A critical overview of the current myofascial pain literature - January 2020. *J Bodyw Mov Ther* 2020; **24**(2): 213-24.

55.   Dommerholt J, Hooks T, Chou LW, Finnegan M. A critical overview of the current myofascial pain literature - January 2018. *J Bodyw Mov Ther.* 2018; **22**(1):184-191.

56.   Shah JP, Thaker N, Heimur J, Aredo JV, Sikdar S, Gerber L. Myofascial Trigger Points Then and Now: A Historical and Scientific Perspective. *PM R.* 2015 Jul; **7**(7):746-761.

57.   Urits I, Charipova K, Gress K, et al. Treatment and management of myofascial pain syndrome. *Best Pract Res Clin Anaesthesiol.* 2020; **34**(3):427-448.

58.   Chen JT, Chung KC, Hou CR, Kuan TS, Chen SM, Hong CZ. Inhibitory effect of dry needling on the spontaneous electrical activity recorded from myofascial trigger spots of rabbit skeletal muscle. *Am J Phys Med Rehabil* 2001; **80**(10): 729-35.

59.   Hughes EJ, McDermott K, Funk MF. Evaluation of hyaluronan content in areas of densification compared to adjacent areas of fascia. *J Bodyw Mov Ther* 2019; **23**(2): 324-8.

60.   Castella LF, Buscemi L, Godbout C, Meister JJ, Hinz B. A new lock-step mechanism of matrix remodelling based on subcellular contractile events. *J Cell Sci.* 2010; **123**(Pt 10):1751-60.

Suggest calcium oscillations and myofibroblast contractile activity are mechanistically linked.

61.   Arora PD, Bibby KJ, McCulloch CA. Slow oscillations of free intracellular calcium ion concentration in human fibroblasts responding to mechanical stretch. *J Cell Physiol* 1994; **161**(2): 187-200.

1% stretch induce no calcium response. 2.8% stretch stimulated initial calcium transient and the subsequent generation of calcium oscillations of up to 2,000 sec.

62.   Langevin HM, Cornbrooks CJ, Taatjes DJ. Fibroblasts form a body-wide cellular network. *Histochem Cell Biol.* 2004; **122**(1):7-15.

63.   Lembong J, Sabass B, Sun B, Rogers ME, Stone HA. Mechanics regulates ATP-stimulated collective calcium response in fibroblast cells. *J R Soc Interface.* 2015; **12**(108):20150140.

Cells cultured on softer substrates show higher percentages of oscillating cells than cells cultured on a stiff substrate.

64.   Wolfson B, Zhang Y, Gernapudi R, et al. A High-Fat Diet Promotes Mammary Gland Myofibroblast Differentiation through MicroRNA 140 Downregulation. *Mol Cell Biol.* 2017; **37**(4):e00461-16.

65.   Witt W, Buttner P, Jannasch A, Matschke K, Waldow T. Reversal of myofibroblastic activation by polyunsaturated fatty acids in valvular interstitial cells from aortic valves. Role of RhoA/G-actin/MRTF signalling. *J Mol Cell Cardiol* 2014; **74**: 127-38.

66.   Yu Q, Wang T, Zhou X, et al. Wld(S) reduces paraquat-induced cytotoxicity via SIRT1 in non-neuronal cells by attenuating the depletion of NAD. *PLoS One* 2011; **6**(7): e21770.

67.   Tai W, Deng S, Wu W, et al. Rapamycin attenuates the paraquat-induced pulmonary fibrosis through activating Nrf2 pathway. *J Cell Physiol* 2020; **235**(2): 1759-68.

68.   Barbe MF, Hilliard BA, Amin M, et al. Blocking CTGF/CCN2 reverses neural fibrosis and sensorimotor declines in a rat model of overuse-induced median mononeuropathy. *J Orthop Res.* 2020; **38**(11):2396-2408.

69.   Chen Y, Abraham DJ, Shi-Wen X, et al. CCN2 (connective tissue growth factor) promotes fibroblast adhesion to fibronectin. *Mol Biol Cell.* 2004; **15**(12):5635-46.

70.   Simms RW, Zerbini CA, Ferrante N, Anthony J, Felson DT, Craven DE. Fibromyalgia syndrome in patients infected with human immunodeficiency virus. The Boston City Hospital Clinical AIDS Team. *Am J Med* 1992; **92**(4): 368-74.

71.   Marts LT, Guidot DM, Sueblinvong V. HIV-1 Protein gp120 Induces Mouse Lung Fibroblast-to-Myofibroblast Transdifferentiation via CXCR4 Activation. *Am J Med Sci* 2019; **357**(6): 483-91.

72.   Chen TC, Chang SW, Wang TY. Moxifloxacin modifies corneal fibroblast-to-myofibroblast differentiation. *Br J Pharmacol* 2013; **168**(6): 1341-54.

73.   Ganjizadeh-Zavareh S, Sodhi M, Spangehl T, Carleton B, Etminan M. Oral fluoroquinolones and risk of fibromyalgia. *Br J Clin Pharmacol* 2019; **85**(1): 236-9.

74.   Squier CA. The effect of stretching on formation of myofibroblasts in mouse skin. *Cell Tissue Res* 1981; **220**(2): 325-35.

75.   Goffin JM, Pittet P, Csucs G, Lussi JW, Meister JJ, Hinz B. Focal adhesion size controls tension-dependent recruitment of alpha-smooth muscle actin to stress fibers. *J Cell Biol* 2006; **172**(2): 259-68.

Hence, α-SMA localization in stress fibers in vivo and in vitro only occurs at a critical substrate stiffness of approximately 20 kPa

76.   Guvendiren M, Perepelyuk M, Wells RG, Burdick JA. Hydrogels with differential and patterned mechanics to study stiffness-mediated myofibroblastic differentiation of hepatic stellate cells. *J Mech Behav Biomed Mater* 2014; **38**: 198-208.

77.   Kaergaard A, Andersen JH. Musculoskeletal disorders of the neck and shoulders in female sewing machine operators: prevalence, incidence, and prognosis. *Occup Environ Med* 2000; **57**(8): 528-34.

78.   Vulfsons S, Minerbi A. The Case for Comorbid Myofascial Pain-A Qualitative Review. *Int J Environ Res Public Health* 2020; **17**(14).

79.   Kalichman L, Lachman H, Freilich N. Long-term impact of ankle sprains on postural control and fascial densification. *J Bodyw Mov Ther* 2016; **20**(4): 914-9.

80.   Hinz B, Mastrangelo D, Iselin CE, Chaponnier C, Gabbiani G. Mechanical tension controls granulation tissue contractile activity and myofibroblast differentiation. *Am J Pathol* 2001; **159**(3): 1009-20.

Myofibroblast features, such as alpha-SMA appeared earlier in splinted than in unsplinted wounds.

81.   Burton AR, Brown R, Macefield VG. Selective activation of muscle and skin nociceptors does not trigger exaggerated sympathetic responses in spinal-injured subjects. *Spinal Cord.* 2008; **46**(10):660-5.

82.   Amoroso Borges BL, Bortolazzo GL, Neto HP. Effects of spinal manipulation and myofascial techniques on heart rate variability: A systematic review. *J Bodyw Mov Ther* 2018; **22**(1): 203-8.

83.   Hinz B, Phan SH, Thannickal VJ, et al. Recent developments in myofibroblast biology: paradigms for connective tissue remodeling. *Am J Pathol* 2012; **180**(4): 1340-55.

In contrast to the reversible and comparably short-lived contraction of striated and smooth muscles, myofibroblast contractile activity, together with ECM synthesis and

degradation, leads to connective tissue remodeling, followed by irreversible and long

contractures in a process that can span weeks, months, or even years

84.   Goldenberg DL, Clauw DJ, Palmer RE, Clair AG. Opioid Use in Fibromyalgia: A

Cautionary Tale. *Mayo Clin Proc.* 2016; **91**(5):640-8.

85.   Stournaras D, Petrovic K. Fibromyalgia treatment and management considerations

for nurse practitioners. *J Am Assoc Nurse Pract* 2019; **31**(6): 371-7.

86.   Rico-Villademoros F, Postigo-Martin P, Garcia-Leiva JM, Ordoñez-Carrasco JL,

Calandre EP. Patterns of pharmacologic and non-pharmacologic treatment, treatment

satisfaction and perceived tolerability in patients with fibromyalgia: a patients' survey. *Clin*

Exp Rheumatol 2020; **38 Suppl 123**(1): 72-8.

87.   Frost FA, Jessen B, Siggaard-Andersen J. A control, double-blind comparison of

mepivacaine injection versus saline injection for myofascial pain. *Lancet* 1980; **1**(8167): 499-

500.

88.   Langevin HM, Churchill DL, Cipolla MJ. Mechanical signaling through connective

tissue: a mechanism for the therapeutic effect of acupuncture. *FASEB J.* 2001;**15**(12):2275-

82.

Needle grasp: feels pulling and increased resistance to further movement of the inserted

needle. They hypothesize that needle grasp is due to mechanical coupling between the

needle and connective tissue with winding of tissue around the needle. Also, needle

pressures transmit a mechanical signal to connective tissue cells via mechanotransduction.

89.   Tomasek JJ, Haaksma CJ, Eddy RJ, Vaughan MB. Fibroblast contraction occurs on

release of tension in attached collagen lattices: dependency on an organized actin

cytoskeleton and serum. *Anat Rec* 1992; **232**(3): 359-68.

Fibroblasts can develop tension in an attached collagen lattice and that upon release of

tension the fibroblasts undergo contraction resulting in a rapid contraction of the collagen

lattice.

90.   Langevin HM, Churchill DL, Fox JR, Badger GJ, Garra BS, Krag MH. Biomechanical

response to acupuncture needling in humans. *J Appl Physiol (1985)* 2001; **91**(6): 2471-8.

"During acupuncture treatments, acupuncture needles are manipulated to elicit the

characteristic "de qi" reaction. De qi has a biomechanical component, needle grasp, which

we have quantified by measuring the force… We found 167 and 52% increases in pullout

force with UNI and BI, respectively, compared with NO"

91.   Leow MQH, Cui SL, Cao T, Tay SC, Ooi CC. Measuring the effects of acupuncture on

muscle stiffness with shear wave elastography. *Acupunct Med* 2017; **35**(4): 304-6.

Immediately after needle insertion and 5min later, muscle stiffness appeared to increase

in both subjects. After 5 minutes, muscle stiffness appeared to decline and plateau at a

level that remained relatively consistent throughout time needle was retained.

92.   Kural MH, Billiar KL. Myofibroblast persistence with real-time changes in boundary

stiffness. *Acta Biomater* 2016; **32**: 223-30.

"when tissues were entirely released from one post to minimize the ability of the cells to

generate tension, the proportion of apoptotic cells increased more than five-fold, and α-

SMA staining was abrogated in the central region. Together, these data indicate that an

increase in boundary stiffness, even if modest and temporary, can have lasting effects on

myofibroblast persistence in tissues, and that a large decrease in the ability of the cells to

generate tension is required to trigger dedifferentiation and apoptosis."

93.   Muller CE, Aranha MF, Gaviao MB. Two-dimensional ultrasound and ultrasound

elastography imaging of trigger points in women with myofascial pain syndrome treated by

acupuncture and electroacupuncture: a double-blinded randomized controlled pilot study.

Ultrason Imaging 2015; **37**(2): 152-67.

Compare changes in elastography between verum and sham acupuncture.

94.   Ernberg M, Hedenberg-Magnusson B, Alstergren P, Kopp S. The level of serotonin in the superficial masseter muscle in relation to local pain and allodynia. *Life Sci* 1999; **65**(3): 313-25.

95.   Maher RM, Hayes DM, Shinohara M. Quantification of dry needling and posture effects on myofascial trigger points using ultrasound shear-wave elastography. *Arch Phys Med Rehabil.* 2013;**94**(11):2146-50.

96.   Luan S, Zhu ZM, Ruan JL, et al. Randomized Trial on Comparison of the Efficacy of Extracorporeal Shock Wave Therapy and Dry Needling in Myofascial Trigger Points. *Am J Phys Med Rehabil* 2019; **98**(8): 677-84.

97.   Turo D, Otto P, Hossain M, et al. Novel Use of Ultrasound Elastography to Quantify Muscle Tissue Changes After Dry Needling of Myofascial Trigger Points in Patients With Chronic Myofascial Pain. *J Ultrasound Med.* 2015;**34**(12):2149-61.

98.   https://www.youtube.com/watch?v=uzy8-wQzQMY&t=3s&ab_channel=RonelleWood)

Microscopic view of the dynamic fiber network. Force vectors travel according to the complex architecture.

99.

https://www.youtube.com/watch?v=cqHtEtZ9mRQ&t=177s&ab_channel=MedFreelancers

2:59-3:09 top left – fascia tears without diathermy touching it under low forces, or manipulation in areas adjacent to it.

100.   https://www.youtube.com/watch?v=3u0hjs53IMU&ab_channel=JeffreyChua

Needles vibrate when inserted. Some seem to be moved by horizontal force vectors. Patterns of vibration change as a function of time. It seems some groups of needles vibrate together. Overall a complex dynamic behavior is observed.

101.   https://www.youtube.com/watch?v=eW0lvOVKDxE&ab_channel=UKyOrtho

102.   Li ADR, Putra KB, Chen L, Montgomery JS, Shih A. Mosquito proboscis-inspired needle insertion to reduce tissue deformation and organ displacement. Sci Rep. 2020; **10**(1):12248.

Currently available needle delivery systems deform and move soft tissue and organs. These data support connective tissue winding as the mechanism responsible for the increase in pullout force induced by needle rotation

103.   Fox JR, Gray W, Koptiuch C, Badger GJ, Langevin HM. Anisotropic tissue motion induced by acupuncture needling along intermuscular connective tissue planes. *J Altern Complement Med* 2014; **20**(4): 290-4.

104.   Langevin HM, Konofagou EE, Badger GJ, Churchill DL, Fox JR, Ophir J, Garra BS. Tissue displacements during acupuncture using ultrasound elastography techniques. *Ultrasound Med Biol.* 2004;**30**(9):1173-83.

105.   Langevin HM, Churchill DL, Wu J, Badger GJ, Yandow JA, Fox JR, Krag MH. Evidence of connective tissue involvement in acupuncture. *FASEB J.* 2002; **16**(8):872-4.

106.   Moog P, Buchner L, Cerny MK, Schmauss D, Megerle K, Erne H. Analysis of recurrence and complications after percutaneous needle fasciotomy in Dupuytren's disease. *Arch Orthop Trauma Surg* 2019; **139**(10): 1471-7.

107.   Soreide E, Murad MH, Denbeigh JM, et al. Treatment of Dupuytren's contracture: a systematic review. *Bone Joint J* 2018; **100-B**(9): 1138-45.

108.   Sellman JR. Plantar fascia rupture associated with corticosteroid injection. *Foot Ankle Int* 1994; **15**(7): 376-81.

109.   Wu L. Nonlinear finite element analysis for musculoskeletal biomechanics of medial and lateral plantar longitudinal arch of Virtual Chinese Human after plantar ligamentous structure failures. *Clin Biomech* (Bristol, Avon). 2007; **22**(2):221-9.

110.   Hettrich CM, DiCarlo EF, Faryniarz D, Vadasdi KB, Williams R, Hannafin JA. The effect of myofibroblasts and corticosteroid injections in adhesive capsulitis. J Shoulder Elbow Surg. 2016; **25**(8):1274-9.

111.   Sanjuan-Cervero R, Carrera-Hueso FJ, Vazquez-Ferreiro P, Gomez-Herrero D. Adverse Effects of Collagenase in the Treatment of Dupuytren Disease: A Systematic Review. *BioDrugs* 2017; **31**(2): 105-15.

112.   Lewkonia P, Medlicott SA, Hildebrand KA. Intramuscular myxoid lipoma in the proximal forearm presenting as an olecranon mass with superficial radial nerve palsy: a case report. *J Med Case Rep*. 2011; **5**:321.

113.   Mishra AK, Skrepnik NV, Edwards SG, et al. Efficacy of platelet-rich plasma for chronic tennis elbow: a double-blind, prospective, multicenter, randomized controlled trial of 230 patients. *Am J Sports Med* 2014; **42**(2): 463-71.

114.   Gottlieb NL, Riskin WG. Complications of local corticosteroid injections. *JAMA* 1980; **243**(15): 1547-8.

115.   Abisi S, Yong YP, Beech A, Oluwole A, Tennant W. Spontaneous sequential compartment syndrome of the lower limbs. *Vasc Endovascular Surg* 2013; **47**(7): 566-8.

116.   Stotts AK, Carroll KL, Schafer PG, Santora SD, Branigan TD. Medial compartment syndrome of the foot: an unusual complication of spine surgery. *Spine (Phila Pa 1976)* 2003; **28**(6): E118-20.

117.   Joshi DG, Balthillaya G, Prabhu A. Effect of remote myofascial release on hamstring flexibility in asymptomatic individuals - A randomized clinical trial. *J Bodyw Mov Ther*. 2018; **22**(3):832-837.

Improvement in hamstring flexibility via remote MFR can be explained by the fascially mediated transfer of forces through the SBL. Evidence points towards the transmission of forces through intramuscular and intermuscular connective tissue pathways.

118.   Soiné JR, Hersch N, Dreissen G, Hampe N, Hoffmann B, Merkel R, Schwarz US. Measuring cellular traction forces on non-planar substrates. *Interface Focus* 2016; **6**(5): 20160024.

119.   Jones C, Ehrlich HP. Fibroblast expression of alpha-smooth muscle actin, alpha2beta1 integrin and alphavbeta3 integrin: influence of surface rigidity. *Exp Mol Pathol* 2011; **91**(1): 394-9.

120.   Bordoni B, Bordoni G. Reflections on osteopathic fascia treatment in the peripheral nervous system. *J Pain Res* 2015; **8**: 735-40.

Impeded sliding of a nerve between its various layers and tissues that cross its fascial system will generate local and dysesthetic pain.

121.   Amris K, Jespersen A, Bliddal H. Self-reported somatosensory symptoms of neuropathic pain in fibromyalgia and chronic widespread pain correlate with tender point count and pressure-pain thresholds. *Pain* 2010; **151**(3): 664-9.

122.   Petersen MW, Schroder A, Jorgensen T, et al. Irritable bowel, chronic widespread pain, chronic fatigue and related syndromes are prevalent and highly overlapping in the general population: DanFunD. *Sci Rep* 2020; **10**(1): 3273.

123.   Creed F. A review of the incidence and risk factors for fibromyalgia and chronic widespread pain in population-based studies. *Pain* 2020; **161**(6): 1169-76.

124.   Kalyan-Raman UP, Kalyan-Raman K, Yunus MB, Masi AT. Muscle pathology in primary fibromyalgia syndrome: a light microscopic, histochemical and ultrastructural study. *J Rheumatol* 1984; **11**(6): 808-13.

125.   Sprott H, Salemi S, Gay RE, et al. Increased DNA fragmentation and ultrastructural changes in fibromyalgic muscle fibres. *Ann Rheum Dis* 2004; **63**(3): 245-51.

126.   Elvin A, Siösteen AK, Nilsson A, Kosek E. Decreased muscle blood flow in fibromyalgia patients during standardised muscle exercise: a contrast media enhanced colour Doppler study. *Eur J Pain.* 2006;**10**(2):137-44.

127.   Zdebik N, Zdebik A, Boguslawska J, Przezdziecka-Dolyk J, Turno-Krecicka A. Fibromyalgia syndrome and the eye-A review. *Surv Ophthalmol* 2021; **66**(1): 132-7.

128.   Morf S, Amann-Vesti B, Forster A, et al. Microcirculation abnormalities in patients with fibromyalgia - measured by capillary microscopy and laser fluxmetry. *Arthritis Res Ther* 2005; **7**(2): R209-16.

129.   Grassi W, Core P, Carlino G, Salaffi F, Cervini C. Capillary permeability in fibromyalgia. *J Rheumatol* 1994; **21**(7): 1328-31.

130.   Lambova SN, Muller-Ladner U. Capillaroscopic Findings in Primary Fibromyalgia. *Curr Rheumatol Rev* 2018; **14**(1): 36-41.

131.   Scolnik M, Vasta B, Hart DJ, Shipley JA, McHugh NJ, Pauling JD. Symptoms of Raynaud's phenomenon (RP) in fibromyalgia syndrome are similar to those reported in primary RP despite differences in objective assessment of digital microvascular function and morphology. *Rheumatol Int*. 2016; **36**(10):1371-7.

132.   Holman AJ. Positional cervical spinal cord compression and fibromyalgia: a novel comorbidity with important diagnostic and treatment implications. J Pain. 2008 Jul;9(7):613-22.

133.   Strobel ES, Krapf M, Suckfull M, Bruckle W, Fleckenstein W, Muller W. Tissue oxygen measurement and 31P magnetic resonance spectroscopy in patients with muscle tension and fibromyalgia. *Rheumatol Int* 1997; **16**(5): 175-80.

134.   Bengtsson A, Henriksson KG, Larsson J. Muscle biopsy in primary fibromyalgia. Light-microscopical and histochemical findings. *Scand J Rheumatol.* 1986;**15**(1):1-6.

135.   Ruggiero L, Manganelli F, Santoro L. Muscle pain syndromes and fibromyalgia: the role of muscle biopsy. *Curr Opin Support Palliat Care* 2018; **12**(3): 382-7.

136.   Bengtsson A, Henriksson KG, Larsson J. Reduced high-energy phosphate levels in the painful muscles of patients with primary fibromyalgia. *Arthritis Rheum* 1986; **29**(7): 817-21.

137.   Ciregia F, Giacomelli C, Giusti L, et al. Putative salivary biomarkers useful to differentiate patients with fibromyalgia. *J Proteomics* 2019; **190**: 44-54.

138.   Sikdar S, Shah JP, Gebreab T, Yen RH, Gilliams E, Danoff J, Gerber LH. Novel applications of ultrasound technology to visualize and characterize myofascial trigger points and surrounding soft tissue. *Arch Phys Med Rehabil* 2009; **90**(11): 1829-38.

Small arteries (or enlarged arterioles) near A-MTrPs showed retrograde flow in diastole, indicating a highly resistive vascular bed.

139.   Erkmen Uyar M, Sezer S, Bal Z, et al. Fibromyalgia and its clinical relevance in renal transplant recipients. *Transplant Proc* 2015; **47**(4): 1105-9.

140.   Maekawa K, Clark GT, Kuboki T. Intramuscular hypoperfusion, adrenergic receptors, and chronic muscle pain. *J Pain* 2002; **3**(4): 251-60.

141.   Choi DH, Kim HS. Quantitative analysis of nailfold capillary morphology in patients with fibromyalgia. *Korean J Intern Med* 2015; **30**(4): 531-7.

142.   Grabowski PJ, Slane LC, Thelen DG, Obermire T, Lee KS. Evidence of Generalized Muscle Stiffness in the Presence of Latent Trigger Points Within Infraspinatus. *Arch Phys Med Rehabil.* 2018;**99**(11):2257-2262.

143.   Cho KI, Lee JH, Kim SM, Lee HG, Kim TI. Assessment of endothelial function in patients with fibromyalgia--cardiac ultrasound study. *Clin Rheumatol.* 2011; **30**(5):647-54.

144.   Gunturk EE, Ekiz T, Akkaya H. Evaluation of carotid-femoral pulse wave velocity, aortic stiffness index, and aortic distensibility in patients with fibromyalgia. *Clin Rheumatol* 2019; **38**(6): 1731-5.

145.   Triantafyllias K, Stortz M, de Blasi M, Leistner C, Weinmann-Menke J, Schwarting A. Increased aortic stiffness in patients with fibromyalgia: results of a prospective study on carotid-femoral pulse wave velocity. *Clin Exp Rheumatol.* 2019; **37** Suppl 116(1):114-115.

146.   Perez-Ruiz F, Calabozo M, Alonso-Ruiz A, Herrero A, Ruiz-Lucea E, Otermin I. High prevalence of undetected carpal tunnel syndrome in patients with fibromyalgia syndrome. *J Rheumatol.* 1995; **22**(3):501-4.

147.   Nacir B, Genc H, Duyur Cakit B, Karagoz A, Erdem HR. Evaluation of upper extremity nerve conduction velocities and the relationship between fibromyalgia and carpal tunnel syndrome. *Arch Med Res.* 2012; **43**(5):369-74.

148.   Bennett RM, Jones J, Turk DC, Russell IJ, Matallana L. An internet survey of 2,596 people with fibromyalgia. *BMC Musculoskelet Disord.* 2007; **8**:27.

149.   Firestein GS, Kelley WN. Kelley's textbook of rheumatology. 9th ed. Philadelphia, PA: Elsevier/Saunders; 2013. Pages 733-750

"…chronic fatigue syndrome, irritable bowel syndrome, headache syndromes…these syndromes have been called functional somatic syndromes (FSS)….Fibromyalgia lies at the end of a continuum of polysymptomatic distress rather than being a discrete disorder."

150.   Sugawara O, Atsuta Y, Iwahara T, Muramoto T, Watakabe M, Takemitsu Y. The effects of mechanical compression and hypoxia on nerve root and dorsal root ganglia. An analysis of ectopic firing using an in vitro model. *Spine (Phila Pa 1976).* 1996; **21**(18):2089-94.

151.   Gotoh H, Takahashi A. Mechanical stimuli induce intracellular calcium response in a subpopulation of cultured rat sensory neurons. *Neuroscience.* 1999; **92**(4):1323-9.

152.   Song XJ, Hu SJ, Greenquist KW, Zhang JM, LaMotte RH. Mechanical and thermal hyperalgesia and ectopic neuronal discharge after chronic compression of dorsal root ganglia. *J Neurophysiol.* 1999; **82**(6):3347-58.

153.   Gladman SJ, Ward RE, Michael-Titus AT, Knight MM, Priestley JV. The effect of mechanical strain or hypoxia on cell death in subpopulations of rat dorsal root ganglion neurons in vitro. *Neuroscience.* 2010; **171**(2):577-87.

154.   Wang JC, Sung FC, Men M, Wang KA, Lin CL, Kao CH. Bidirectional association between fibromyalgia and gastroesophageal reflux disease: two population-based retrospective cohort analysis. *Pain.* 2017; **158**(10):1971-1978

155.   Nathan H. Osteophytes of the spine compressing the sympathetic trunk and splanchnic nerves in the thorax. *Spine (Phila Pa 1976).* 1987; **12**(6):527-32

156.   Weinberg H, Nathan H, Magora F, Robin GC, Aviad I. Arthritis of the first costovertebral joint as a cause of thoracic outlet syndrome. *Clin Orthop Relat Res.* 1972; **86**:159-63.

157.   Wolfe F, Rasker JJ, Häuser W. Hearing loss in fibromyalgia? Somatic sensory and non-sensory symptoms in patients with fibromyalgia and other rheumatic disorders. *Clin Exp Rheumatol.* 2012; **30**(6 Suppl 74):88-93

158.   Evdokimov D, Frank J, Klitsch A, et al. Reduction of skin innervation is associated with a severe fibromyalgia phenotype. *Ann Neurol.* 2019; **86**(4):504-516.

159.   https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-fibromyalgia-in-adults?search=fibrommyalgia&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1 By Don L Goldenberg, accessed March 2021.

160.   Raphael KG, Marbach JJ. Comorbid fibromyalgia accounts for reduced fecundity in women with myofascial face pain. *Clin J Pain* 2000; **16**(1): 29-36.

161.   Mancuso AC, Summers KM, Mengeling MA, Torner JC, Ryan GL, Sadler AG. Infertility and Health-Related Quality of Life in United States Women Veterans. *J Womens Health (Larchmt)* 2020; **29**(3): 412-9.

162.   Piersiala K, Akst LM, Hillel AT, Best SR. Chronic Pain Syndromes and Their Laryngeal Manifestations. *JAMA Otolaryngol Head Neck Surg.* 2020; **146**(6):543-549.

163.   Vaeroy H, Helle R, Forre O, Kass E, Terenius L. Elevated CSF levels of substance P and high incidence of Raynaud phenomenon in patients with fibromyalgia: new features for diagnosis. *Pain* 1988; **32**(1): 21-6.

164.   Simms RW, Zerbini CA, Ferrante N, Anthony J, Felson DT, Craven DE. Fibromyalgia syndrome in patients infected with human immunodeficiency virus. The Boston City Hospital Clinical AIDS Team. Am J Med. 1992 Apr;92(4):368-74.

165.   Clark S, Tindall E, Bennett RM. A double blind crossover trial of prednisone versus placebo in the treatment of fibrositis. J Rheumatol. 1985 Oct;12(5):980-3. PMID: 3910836.

166.   Gupta A, McBeth J, Macfarlane GJ, Morriss R, Dickens C, Ray D, Chiu YH, Silman AJ. Pressure pain thresholds and tender point counts as predictors of new chronic widespread pain in somatising subjects. *Ann Rheum Dis.* 2007; **66**(4):517-21.

167.   Chen JY, Newcomb B, Zhou C, et al. Tricyclic Antidepressants Promote Ceramide Accumulation to Regulate Collagen Production in Human Hepatic Stellate Cells. *Sci Rep.* 2017; **7**:44867

168.   Morrissey JJ. Pleiotropic effects of amitriptyline ameliorate renal fibrosis. *Kidney Int.* 2009; **75**(6):583-4

169.   Olianas MC, Dedoni S, Onali P. Antidepressants induce profibrotic responses via the lysophosphatidic acid receptor LPA1. Eur J Pharmacol. 2020 Apr 15;873:172963.

170.   Bruno R, Galastri S, Sacchi P, Cima S, Caligiuri A, DeFranco R, Milani S, Gessani S, Fantuzzi L, Liotta F, Frosali F, Antonucci G, Pinzani M, Marra F. gp120 modulates the biology of human hepatic stellate cells: a link between HIV infection and liver fibrogenesis. Gut. 2010 Apr;59(4):513-20.

171.   Marts LT, Guidot DM, Sueblinvong V. HIV-1 Protein gp120 Induces Mouse Lung Fibroblast-to-Myofibroblast Transdifferentiation via CXCR4 Activation. Am J Med Sci. 2019 Jun;357(6):483-491.

172.   Russell IJ, Vaeroy H, Javors M, Nyberg F. Cerebrospinal fluid biogenic amine metabolites in fibromyalgia/fibrositis syndrome and rheumatoid arthritis. *Arthritis Rheum* 1992; **35**(5): 550-6.

173.   Alcaino C, Knutson KR, Treichel AJ, Yildiz G, Strege PR, Linden DR, Li JH, Leiter AB, Szurszewski JH, Farrugia G, Beyder A. A population of gut epithelial enterochromaffin cells is mechanosensitive and requires Piezo2 to convert force into serotonin release. Proc Natl Acad Sci U S A. 2018 Aug 7;115(32):E7632-E7641.

174.   Pan J, Copland I, Post M, Yeger H, Cutz E. Mechanical stretch-induced serotonin release from pulmonary neuroendocrine cells: implications for lung development. Am J Physiol Lung Cell Mol Physiol. 2006 Jan;290(1):L185-93.

175.   Dolivo DM, Larson SA, Dominko T. Tryptophan metabolites kynurenine and serotonin regulate fibroblast activation and fibrosis. *Cell Mol Life Sci* 2018; **75**(20): 3663-81.

176.   Stunes AK, Reseland JE, Hauso O, et al. Adipocytes express a functional system for serotonin synthesis, reuptake and receptor activation. *Diabetes Obes Metab* 2011; **13**(6): 551-8.

177.   Lofdahl A, Rydell-Tormanen K, Muller C, et al. 5-HT2B receptor antagonists attenuate myofibroblast differentiation and subsequent fibrotic responses in vitro and in vivo. *Physiol Rep* 2016; **4**(15).

178.   Legangneux E, Mora JJ, Spreux-Varoquaux O, Thorin I, Herrou M, Alvado G, Gomeni C. Cerebrospinal fluid biogenic amine metabolites, plasma-rich platelet serotonin and [3H]imipramine reuptake in the primary fibromyalgia syndrome. Rheumatology (Oxford). 2001 Mar;40(3):290-6.

179.   Al-Nimer MSM, Mohammad TAM, Alsakeni RA. Serum levels of serotonin as a biomarker of newly diagnosed fibromyalgia in women: Its relation to the platelet indices. J Res Med Sci. 2018 Aug 23;23:71.

180. Amin OA, Abouzeid SM, Ali SA, Amin BA, Alswat KA. Clinical association of vitamin D and serotonin levels among patients with fibromyalgia syndrome. Neuropsychiatr Dis Treat. 2019 May 27;15:1421-1426.

181. Barjandi G, Louca Jounger S, Löfgren M, Bileviciute-Ljungar I, Kosek E, Ernberg M. Plasma tryptophan and kynurenine in females with temporomandibular disorders and fibromyalgia-An exploratory pilot study. J Oral Rehabil. 2020 Feb;47(2):150-157.

182. Russell IJ, Vipraio GA, Morgan WW, Bowden CL. Is there a metabolic basis for the fibrositis syndrome? Am J Med. 1986 Sep 29;81(3A):50-4

183. Yunus MB, Dailey JW, Aldag JC, Masi AT, Jobe PC. Plasma tryptophan and other amino acids in primary fibromyalgia: a controlled study. *J Rheumatol* 1992; **19**(1): 90-4.

184. Welsch P, Uceyler N, Klose P, Walitt B, Hauser W. Serotonin and noradrenaline reuptake inhibitors (SNRIs) for fibromyalgia. Cochrane Database Syst Rev 2018; 2: CD010292.

185. Saber AA, Boros MJ, Mancl T, Elgamal MH, Song S, Wisadrattanapong T. The effect of laparoscopic Roux-en-Y gastric bypass on fibromyalgia. *Obes Surg* 2008; **18**(6): 652-5.

186. Adkisson CD, Yip L, Armstrong MJ, Stang MT, Carty SE, McCoy KL. Fibromyalgia symptoms and medication requirements respond to parathyroidectomy. *Surgery.* 2014; **156**(6):1614-20; discussion 1620-1.

187. Raftopoulos Y, Papasavas P, Landreneau R, et al. Clinical outcome of laparoscopic antireflux surgery for patients with irritable bowel syndrome. *Surg Endosc.* 2004; **18**(4):655-9.

188. Wu MC, Ma KS, Wang YH, Wei JC. Impact of tonsillectomy on irritable bowel syndrome: A nationwide population-based cohort study. *PLoS One.* 2020; **15**(9):e0238242.

189.   Rosen JM, Adams PN, Saps M. Umbilical hernia repair increases the rate of functional gastrointestinal disorders in children. *J Pediatr.* 2013; **163**(4):1065-8.

190.   Yang CY, Wu MC, Lin MC, Wei JC. Risk of irritable bowel syndrome in patients who underwent appendectomy: A nationwide population-based cohort study. *EClinicalMedicine.* 2020; **23**:100383.

191.   Vincent A, Whipple MO, Luedtke CA, et al. Pain and other symptom severity in women with fibromyalgia and a previous hysterectomy. *J Pain Res.* 2011; **4**:325-9.

192.   Vleeming A, Schuenke MD, Danneels L, Willard FH. The functional coupling of the deep abdominal and paraspinal muscles: the effects of simulated paraspinal muscle contraction on force transfer to the middle and posterior layer of the thoracolumbar fascia. *J Anat.* 2014; **225**(4):447-62.

193.   Marcelin G, Silveira ALM, Martins LB, Ferreira AV, Clément K. Deciphering the cellular interplays underlying obesity-induced adipose tissue fibrosis. J Clin Invest. 2019; **129**(10):4032-4040.

Those effects appear to be dependent on mechanosensitive signaling involving integrin and YAP/TEAD pathways. Fibrosis alters adipose tissue plasticity. Obesity increases collagen deposition and myofibrocyte distribution in adipose tissue. Fibrosis can be observed around blood vessels and crown like structures with inflammatory cells.

194.   Okifuji A, Turk DC. Sex hormones and pain in regularly menstruating women with fibromyalgia syndrome. *J Pain* 2006; **7**(11): 851-9.

195.   Wu M, Han M, Li J, et al. 17beta-estradiol inhibits angiotensin II-induced cardiac myofibroblast differentiation. *Eur J Pharmacol.* 2009; **616**(1-3):155-9.

196.   Ilg MM, Stafford SJ, Mateus M, et al. Phosphodiesterase Type 5 Inhibitors and Selective Estrogen Receptor Modulators Can Prevent But Not Reverse Myofibroblast Transformation in Peyronie's Disease. *J Sex Med.* 2020; **17**(10):1848-1864.

197. Siqueira OH, Herani Filho B, Paula RE, et al. Tamoxifen decreases the myofibroblast count in the healing bile duct tissue of pigs. *Clinics (Sao Paulo).* 2013; **68**(1):101-6.

198. Jiang HS, Zhu LL, Zhang Z, Chen H, Chen Y, Dai YT. Estradiol attenuates the TGF-β1-induced conversion of primary TAFs into myofibroblasts and inhibits collagen production and myofibroblast contraction by modulating the Smad and Rho/ROCK signaling pathways. *Int J Mol Med.* 2015; **36**(3):801-7.

199. Tu CH, Lin CL, Yang ST, Shen WC, Chen YH. Hormonal Contraceptive Treatment May Reduce the Risk of Fibromyalgia in Women with Dysmenorrhea: A Cohort Study. *J Pers Med.* 2020; **10**(4):280.

200. Blümel JE, Palacios S, Legorreta D, Vallejo MS, Sarra S. Is fibromyalgia part of the climacteric syndrome? *Maturitas.* 2012; **73**(2):87-93.

201. Lim EC, Seet RC, Lee KH, Wilder-Smith EP, Chuah BY, Ong BK. Headaches and the N95 face-mask amongst healthcare providers. *Acta Neurol Scand.* 2006; **113**(3):199-202.

Headaches have been reported with the donning of objects with straps, such as helmets, goggles and brassieres. The etiopathogenesis of mask headaches may be related to mechanical factors.

202. Ramirez-Moreno JM, Ceberino D, Gonzalez Plata A, et al. Mask-associated 'de novo' headache in healthcare workers during the COVID-19 pandemic. *Occup Environ Med.* 2020; oemed-2020-106956.

"Another phenomenon probably related to the physiopathology of headache after PPE use is the external compression that it generates. In most cases there is a temporal relationship between the use of devices and the headache, as well as the topographical location of the headache."

203. Blau JN. Ponytail headache: a pure extracranial headache. *Headache* 2004; **44**(5): 411-3.

Ponytail headache is common. Its varied distribution is explicable by local cutaneous nerve stimulation, as well as traction on extracranial muscles and their fascial coverings, and not by neural referral. Suggest that ponytail headaches arise from two sources: cutaneous nerve endings and from fascia.

204.   Ansari HN, Solomon GD. Hijab (headscarf) headache. *Headache* 2015; **55**(3): 437-8.

 The headache is believed to be purely extracranial coming from pericranial muscle fascia and traction of tendon.

205.   Lane A. Bra strap headache. *Med J Aust.* 1983; **1**(4):155.

206.   Balfour W. Observations on the Pathology and Cure of Rheumatism. Edinb Med Surg J. 1815 Apr 1;11(42):168-187. PMID: 30329434; PMCID: PMC5743373.

207.   Bardin L, Malfetes N, Newman-Tancredi A, Depoortere R. Chronic restraint stress induces mechanical and cold allodynia, and enhances inflammatory pain in rat: Relevance to human stress-associated painful pathologies. *Behav Brain Res* 2009; **205**(2): 360-6.

208.   Chiba S, Numakawa T, Ninomiya M, Richards MC, Wakabayashi C, Kunugi H. Chronic restraint stress causes anxiety- and depression-like behaviors, downregulates glucocorticoid receptor expression, and attenuates glutamate release induced by brain-derived neurotrophic factor in the prefrontal cortex. *Prog Neuropsychopharmacol Biol Psychiatry* 2012; **39**(1): 112-9.

209.   Farhin S, Wong A, Delungahawatta T, et al. Restraint stress induced gut dysmotility is diminished by a milk oligosaccharide (2'-fucosyllactose) in vitro. *PLoS One* 2019; **14**(4): e0215151.

210.   Sugama S, Sekiyama K, Kodama T, et al. Chronic restraint stress triggers dopaminergic and noradrenergic neurodegeneration: Possible role of chronic stress in the onset of Parkinson's disease. *Brain Behav Immun* 2016; **51**: 39-46.

211. Spyrka J, Hess G. Repeated Neck Restraint Stress Bidirectionally Modulates Excitatory Transmission in the Dentate Gyrus and Performance in a Hippocampus-dependent Memory Task. *Neuroscience* 2018; **379**: 32-44.

212. Hu HT, Gao H, Ma RJ, Zhao XF, Tian HF, Li L. Is dry needling effective for low back pain?: A systematic review and PRISMA-compliant meta-analysis. *Medicine (Baltimore).* 2018; **97**(26):e11225.

213. Karayol KC, Karayol SS. A comparison of visual analog scale and shear-wave ultrasound elastography data in fibromyalgia patients and the normal population. *J Phys Ther Sci* 2021; **33**(1): 40-4.

214. Taş S, Korkusuz F, Erden Z. Neck Muscle Stiffness in Participants With and Without Chronic Neck Pain: A Shear-Wave Elastography Study. J Manipulative *Physiol Ther.* 2018; **41**(7):580-588.

215. Giamberardino MA, Affaitati G, Fabrizio A, Costantini R. Effects of treatment of myofascial trigger points on the pain of fibromyalgia. *Curr Pain Headache Rep.* 2011; **15**(5):393-9.

216. Deare JC, Zheng Z, Xue CC, Liu JP, Shang J, Scott SW, Littlejohn G. Acupuncture for treating fibromyalgia. *Cochrane Database Syst Rev.* 2013; **2013**(5):CD007070.

217. Zhang XC, Chen H, Xu WT, Song YY, Gu YH, Ni GX. Acupuncture therapy for fibromyalgia: a systematic review and meta-analysis of randomized controlled trials. J Pain Res. 2019; **12**:527-542.

218. Sánchez-Infante J, Bravo-Sánchez A, Jiménez F, Abián-Vicén J. Effects of Dry Needling on Muscle Stiffness in Latent Myofascial Trigger Points: A Randomized Controlled Trial. J Pain. 2021: **S1526-5900**(21)00024-9

219. Dong Z, Shun-Yue L, Shu-You W, Hui-Min M. Evaluation of influence of acupuncture and electro-acupuncture for blood perfusion of stomach by laser Doppler blood perfusion imaging. *Evid Based Complement Alternat Med.* 2011; **2011**:969231.

220.   Deare JC, Zheng Z, Xue CC, Liu JP, Shang J, Scott SW, Littlejohn G. Acupuncture for treating fibromyalgia. *Cochrane Database Syst Rev.* 2013; **2013**(5):CD007070.

221.   Dorigo B, Bartoli V, Grisillo D, Beconi D. Fibrositic myofascial pain in intermittent claudication. Effect of anesthetic block of trigger points on exercise tolerance. *Pain.* 1979; **6**(2):183-190.

222.   Drabik D, Chodaczek G, Kraszewski S. Effect of Amyloid-β Monomers on Lipid Membrane Mechanical Parameters-Potential Implications for Mechanically Driven Neurodegeneration in Alzheimer's Disease. Int J Mol Sci. 2020 Dec 22;22(1):18.

223.   Levy Nogueira M, Hamraz M, Abolhassani M, Bigan E, Lafitte O, Steyaert JM, Dubois B, Schwartz L. Mechanical stress increases brain amyloid β, tau, and α-synuclein concentrations in wild-type mice. Alzheimers Dement. 2018 Apr;14(4):444-453.

224.   Kellermayer MS, Grama L, Karsai A, Nagy A, Kahn A, Datki ZL, Penke B. Reversible mechanical unzipping of amyloid beta-fibrils. J Biol Chem. 2005 Mar 4;280(9):8464-70.

225.   Bigot M, Chauveau F, Amaz C, Sinkus R, Beuf O, Lambert SA. The apparent mechanical effect of isolated amyloid-β and α-synuclein aggregates revealed by multi-frequency MRE. NMR Biomed. 2020 Jan;33(1):e4174.

226.   Chiasseu M, Cueva Vargas JL, Destroismaisons L, Vande Velde C, Leclerc N, Di Polo A. Tau Accumulation, Altered Phosphorylation, and Missorting Promote Neurodegeneration in Glaucoma. J Neurosci. 2016 May 25;36(21):5785-98.

227.   Zhang L, Fu J, Cheng XH, Tang L. Tau protein function: The mechanical exploration of axonal transport disorder caused by persistent pressure in dorsal root ganglia. Mol Genet Genomic Med. 2019 Apr;7(4):e00580.

228.   Xu Z, Paparcone R, Buehler MJ. Alzheimer's abeta(1-40) amyloid fibrils feature size-dependent mechanical properties. Biophys J. 2010 May 19;98(10):2053-62.

229.   Trumbore CN. Shear-induced amyloid formation of IDPs in the brain. Prog Mol Biol Transl Sci. 2019;166:225-309.

230.   Lenaerts ME, Gill PS. At the crossroads between tension-type headache and fibromyalgia. *Curr Pain Headache Rep.* 2006; **10**(6):463-6.

131.   Conti PC, Costa YM, Goncalves DA, Svensson P. Headaches and myofascial temporomandibular disorders: overlapping entities, separate managements? *J Oral Rehabil* 2016; **43**(9): 702-15.

232.   Chen CK, Nizar AJ. Myofascial pain syndrome in chronic back pain patients. *Korean J Pain* 2011; **24**(2): 100-4.

233.   Aggarwal VR, Macfarlane GJ, McBeth J. A high tender point count is associated with the presence of multiple idiopathic pain disorders: results from a population study. *Eur J Pain.* 2012; **16**(8):1195-203.

234.   Bourgaize S, Janjua I, Murnaghan K, Mior S, Srbely J, Newton G. Fibromyalgia and myofascial pain syndrome: Two sides of the same coin? A scoping review to determine the lexicon of the current diagnostic criteria. *Musculoskeletal Care* 2019; **17**(1): 3-12.

235.   Alonso-Blanco C, Fernández-de-las-Peñas C, Morales-Cabezas M, Zarco-Moreno P, Ge HY, Florez-García M. Multiple active myofascial trigger points reproduce the overall spontaneous pain pattern in women with fibromyalgia and are related to widespread mechanical hypersensitivity. *Clin J Pain.* 2011; **27**(5):405-13.

In fibromyalgia significant positive correlation found between number of active TrPs and spontaneous pain intensity.

236.   Ge HY. Prevalence of myofascial trigger points in fibromyalgia: the overlap of two common problems. *Curr Pain Headache Rep.* 2010; **14**(5):339-45.

The considerable overlap of MTrPs and fibromyalgia in pain characteristics and pathophysiology suggests that fibromyalgia pain is largely due to trigger points.

237.   Xu Y, Hu J, Yilmaz DE, Bachmann S. Connexin43 is differentially distributed within renal vasculature and mediates profibrotic differentiation in medullary fibroblasts. Am J Physiol Renal Physiol. 2021 Jan 1;320(1):F17-F30.

238.   Miragoli M, Gaudesius G, Rohr S. Electrotonic modulation of cardiac impulse conduction by myofibroblasts. Circ Res. 2006 Mar 31;98(6):801-10.

239.   Iguchi M, Hara M, Manome H, Kobayasi H, Tagami H, Aiba S. Communication network in the follicular papilla and connective tissue sheath through gap junctions in human hair follicles. Exp Dermatol. 2003 Jun;12(3):283-8.

These findings suggest that the FP and the CTS form a communicating network through gap junctions, which may play a role in controlling the dynamic structural changes of hair follicles during hair cycling.

240.   Luo M, Luo Y, Mao N, Huang G, Teng C, Wang H, Wu J, Liao X, Yang J. Cancer-Associated Fibroblasts Accelerate Malignant Progression of Non-Small Cell Lung Cancer via Connexin 43-Formed Unidirectional Gap Junctional Intercellular Communication. Cell Physiol Biochem. 2018;51(1):315-336.

241.   Younger JW, Shen YF, Goddard G, Mackey SC. Chronic myofascial temporomandibular pain is associated with neural abnormalities in the trigeminal and limbic systems. Pain. 2010 May;149(2):222-228.

242.   Gabbiani G, Chaponnier C, Hüttner I. Cytoplasmic filaments and gap junctions in epithelial cells and myofibroblasts during wound healing. J Cell Biol. 1978 Mar;76(3):561-8.

243.   Spanakis SG, Petridou S, Masur SK. Functional gap junctions in corneal fibroblasts and myofibroblasts. Invest Ophthalmol Vis Sci. 1998 Jul;39(8):1320-8

244.   Woodward TL, Sia MA, Blaschuk OW, Turner JD, Laird DW. Deficient epithelial-fibroblast heterocellular gap junction communication can be overcome by co-culture with an

intermediate cell type but not by E-cadherin transgene expression. J Cell Sci. 1998 Dec;111 (

Pt 23):3529-39.

245.   Neuhaus J, Gonsior A, Cheng S, Stolzenburg JU, Berger FP. Mechanosensitivity Is a

Characteristic Feature of Cultured Suburothelial Interstitial Cells of the Human Bladder. Int J

Mol Sci. 2020 Jul 31;21(15):5474.

246.   Lembong J, Sabass B, Sun B, Rogers ME, Stone HA. Mechanics regulates ATP-

stimulated collective calcium response in fibroblast cells. J R Soc Interface. 2015 Jul

6;12(108):20150140.

Moreover, the frequency of calcium oscillations is independent of the rigidity of the

substrate but depends on ATP concentration.

247.   Ikeda Y, Fry C, Hayashi F, Stolz D, Griffiths D, Kanai A. Role of gap junctions in

spontaneous activity of the rat bladder. Am J Physiol Renal Physiol. 2007 Oct;293(4):F1018-

25.

In the bladders of adults, spontaneous activity was initiated at multiple sites, resulting in a

lack of coordination. These data suggest that coordinated spontaneous activity requires

gap junction upregulation in urothelial cells and lamina propria myofibroblasts

248.   Neuhaus J, Pfeiffer F, Wolburg H, Horn LC, Dorschner W. Alterations in connexin

expression in the bladder of patients with urge symptoms. BJU Int. 2005 Sep;96(4):670-6.

249.   Paw M, Borek I, Wnuk D, Ryszawy D, Piwowarczyk K, Kmiotek K, Wójcik-Pszczoła KA,

Pierzchalska M, Madeja Z, Sanak M, Błyszczuk P, Michalik M, Czyż J. Connexin43 Controls the

Myofibroblastic Differentiation of Bronchial Fibroblasts from Patients with Asthma. Am J

Respir Cell Mol Biol. 2017 Jul;57(1):100-110.

250.   Zlochiver S, Muñoz V, Vikstrom KL, Taffet SM, Berenfeld O, Jalife J. Electrotonic

myofibroblast-to-myocyte coupling increases propensity to reentrant arrhythmias in two-

dimensional cardiac monolayers. Biophys J. 2008 Nov 1;95(9):4469-80

251.   Askar SF, Bingen BO, Swildens J, Ypey DL, van der Laarse A, Atsma DE, Zeppenfeld K, Schalij MJ, de Vries AA, Pijnappels DA. Connexin43 silencing in myofibroblasts prevents arrhythmias in myocardial cultures: role of maximal diastolic potential. Cardiovasc Res. 2012 Mar 1;93(3):434-44.

252.   Nagaraju CK, Dries E, Gilbert G, Abdesselem M, Wang N, Amoni M, Driesen RB, Sipido KR. Myofibroblast modulation of cardiac myocyte structure and function. Sci Rep. 2019 Jun 20;9(1):8879.

253.   Daniel EE, Yazbi AE, Mannarino M, Galante G, Boddy G, Livergant J, Oskouei TE. Do gap junctions play a role in nerve transmissions as well as pacing in mouse intestine? Am J Physiol Gastrointest Liver Physiol. 2007 Mar;292(3):G734-45.

Target of innervation of these nerves does not appear to be the ICC-DMP, but rather fibroblast-like cells, which are SK3, CD34 positive, c-*kit* negative. They, too, are connected to outer CM cells by gap junctions… **Thus these gap junctions may play another role in neurotransmission, acting to ensure proximity of either ICC or fibroblast to outer CM cells and/or transferring a stretch or other physical or chemical effect from one cell to the other.**

254.   Mewe M, Bauer CK, Schwarz JR, Middendorff R. Mechanisms regulating spontaneous contractions in the bovine epididymal duct. Biol Reprod. 2006 Oct;75(4):651-9.

255.   Grand T, Salvarani N, Jousset F, Rohr S. Aggravation of cardiac myofibroblast arrhythmogeneicity by mechanical stress. *Cardiovasc Res* 2014; **104**(3): 489-500.

256.   Nguyen TP, Xie Y, Garfinkel A, Qu Z, Weiss JN. Arrhythmogenic consequences of myofibroblast-myocyte coupling. *Cardiovasc Res* 2012; **93**(2): 242-51.

257.   Ubogu EE. The molecular and biophysical characterization of the human blood-nerve barrier: current concepts. *J Vasc Res* 2013; **50**(4): 289-303.

258.   Squecco R, Chellini F, Idrizaj E, et al. Platelet-Rich Plasma Modulates Gap Junction Functionality and Connexin 43 and 26 Expression During TGF-beta1-Induced Fibroblast to Myofibroblast Transition: Clues for Counteracting Fibrosis. *Cells* 2020; **9**(5).

259.   Mercier F, Hatton GI. Connexin 26 and basic fibroblast growth factor are expressed primarily in the subpial and subependymal layers in adult brain parenchyma: roles in stem cell proliferation and morphological plasticity? J Comp Neurol. 2001 Feb 26;431(1):88-104.

"A continuum of Cx26-immunoreactive puncta was observed throughout the three meningeal layers, the perineurium of cranial nerves, and meningeal projections into the brain… found a close correlation between the distribution of Cx26 and vimentin immunoreactivities in the meninges and their projections into the brain. Previously showed vimentin and S100beta immunoreactivities through a network of meningeal fibroblasts in the three layers of meninges, perivascular cells, and ependymocytes and in a population of astrocytes. Considering that connexin immunoreactivity may reflect the presence of functional gap junctions, the present results are consistent with our hypothesis that all of these various cell types may communicate in a cooperative network"

260.   Grafstein B, Liu S, Cotrina ML, Goldman SA, Nedergaard M. Meningeal cells can communicate with astrocytes by calcium signaling. Ann Neurol. 2000 Jan;47(1):18-25.

261.   Mercier F, Hatton GI. Immunocytochemical basis for a meningeo-glial network. J Comp Neurol. 2000 May 15;420(4):445-65.

262.   Grafstein B, Liu S, Cotrina ML, Goldman SA, Nedergaard M. Meningeal cells can communicate with astrocytes by calcium signaling. Ann Neurol. 2000 Jan;47(1):18-25.

263.   Zheng N, Yuan XY, Chi YY, Liu P, Wang B, Sui JY, Han SH, Yu SB, Sui HJ. The universal existence of myodural bridge in mammals: an indication of a necessary function. Sci Rep. 2017 Aug 15;7(1):8248

264.   Langevin HM, Cornbrooks CJ, Taatjes DJ. Fibroblasts form a body-wide cellular network. Histochem Cell Biol. 2004 Jul;122(1):7-15.

265.   Follonier L, Schaub S, Meister JJ, Hinz B. Myofibroblast communication is controlled by intercellular mechanical coupling. J Cell Sci. 2008 Oct 15;121(Pt 20):3305-16.

…contraction is transmitted via adherens junctions and leads to the opening of mechanosensitive ion channels in adjacent cells. The resulting Ca2+ influx induces a contraction that can feed back on the first cell and/or stimulate other contacting cells.

266.   Camelliti P, Green CR, LeGrice I, Kohl P. Fibroblast network in rabbit sinoatrial node: structural and functional identification of homogeneous and heterogeneous cell coupling. Circ Res. 2004 Apr 2;94(6):828-35.

267.   Murata K, Hirata A, Ohta K, Enaida H, Nakamura KI. Morphometric analysis in mouse scleral fibroblasts using focused ion beam/scanning electron microscopy. Sci Rep. 2019 Apr 19;9(1):6329.

The cells frequently showed a cellular junction with the neighboring cells and formed cellular networks.

268.   Mechanisms by which fibroblasts promote wound contraction may include activation of contractile signaling pathways, which promote interaction between non-muscle myosin II and ACTB and ACTG1.

269.   https://www.youtube.com/watch?v=yCi-UQEVHKA&ab_channel=TheSpiritualLibrary

06:10 "What is it, then, that we feel? … it's a chronic habitual sense of muscular strain… So everyone chronically pulls themselves together. If you get a person to just lie on the floor and relax, there is the floor under you as firm as can be holding you up. Nevertheless, you will detect the person is making all sorts of tensions… this feeling of tightness is the physical referent for the psychological image of ourselves." A.W. Watts

270.   https://en.wikipedia.org/wiki/Wilhelm_Reich

"…he massaged his disrobed patients to dissolve their muscular armour"

271.

https://www.reichandlowentherapy.org/Content/Energy_and_Movement/armor.html

"Armor can be palpated as muscle tension. It can also be observed as impairment in movement. Patterns of armor do not strictly follow the pathways of voluntary motor nerves but rather show up as bands or segments."

272.   Alexander Lowen, Leslie Lowen. The Way to Vibrant Health: A Manual of Bioenergetic Exercises. New York; Harper & Row 1977.

Page 5-6 "Every stress produces a state of tension in the body. Chronic tensions, however, persist after the provoking stress has been removed as an unconscious bodily attitude or muscular set. Such chronic muscular tensions disturb emotional health by decreasing an individual's energy, restricting his motility and limiting his self expression… every contracted muscle is blocking some movement… A healthy body is in a constant state of vibration."

273.   https://www.eckharttolle.com/power-of-now-excerpt/

Page 4: "Then I felt drawn into what seemed like a vortex of energy. It was a slow movement at first and then accelerated… my body started to shake. I could feel myself being sucked into a void. It felt as if the void was inside myself rather than outside. Suddenly, there was no more fear, and I let myself fall into that void."

274.   https://www.uptodate.com/contents/somatic-symptom-disorder-epidemiology-and-clinical-presentation?search=somatic%20symptom%20disorder&source=search_result&selectedTitle=1~28&usage_type=default&display_rank=1#H3058265287

By James L. Levenson, Accessed January 2021 "Defined somatization as four or more unexplained physical symptoms in men and six or more unexplained physical symptoms in women; the median point prevalence of somatization was 17 percent"

275.   Nakaya M, Watari K, Tajima M, Nakaya T, Matsuda S, Ohara H, Nishihara H, Yamaguchi H, Hashimoto A, Nishida M, Nagasaka A, Horii Y, Ono H, Iribe G, Inoue R, Tsuda M, Inoue K, Tanaka A, Kuroda M, Nagata S, Kurose H. Cardiac myofibroblast engulfment of dead cells facilitates recovery after myocardial infarction. J Clin Invest. 2017 Jan 3;127(1):383-401.

276.   Saada JI, Pinchuk IV, Barrera CA, Adegboyega PA, Suarez G, Mifflin RC, Di Mari JF, Reyes VE, Powell DW. Subepithelial myofibroblasts are novel nonprofessional APCs in the human colonic mucosa. J Immunol. 2006 Nov 1;177(9):5968-79.

277.   Elyada E, Bolisetty M, Laise P, Flynn WF, Courtois ET, Burkhart RA, Teinor JA, Belleau P, Biffi G, Lucito MS, Sivajothi S, Armstrong TD, Engle DD, Yu KH, Hao Y, Wolfgang CL, Park Y, Preall J, Jaffee EM, Califano A, Robson P, Tuveson DA. Cross-Species Single-Cell Analysis of Pancreatic Ductal Adenocarcinoma Reveals Antigen-Presenting Cancer-Associated Fibroblasts. Cancer Discov. 2019 Aug;9(8):1102-1123.

278.   Fang M, Xia J, Wu X, Kong H, Wang H, Xie W, Xu Y. Adenosine signaling inhibits CIITA-mediated MHC class II transactivation in lung fibroblast cells. Eur J Immunol. 2013 Aug;43(8):2162-73.

279.   Sebastian A, Hum NR, Martin KA, Gilmore SF, Peran I, Byers SW, Wheeler EK, Coleman MA, Loots GG. Single-Cell Transcriptomic Analysis of Tumor-Derived Fibroblasts and Normal Tissue-Resident Fibroblasts Reveals Fibroblast Heterogeneity in Breast Cancer. Cancers (Basel). 2020 May 21;12(5):1307.

280.   Boots AM, Wimmers-Bertens AJ, Rijnders AW. Antigen-presenting capacity of rheumatoid synovial fibroblasts. Immunology. 1994 Jun;82(2):268-74. PMID: 7927499; PMCID: PMC1414827.

281.   Applbaum E, Lichtbroun A. Novel Sjögren's autoantibodies found in fibromyalgia patients with sicca and/or xerostomia. Autoimmun Rev. 2019 Feb;18(2):199-202.

282.   Jacobsen S, Høyer-Madsen M, Danneskiold-Samsøe B, Wiik A. Screening for autoantibodies in patients with primary fibromyalgia syndrome and a matched control group. APMIS. 1990 Jul;98(7):655-8.

283.   Ibrahim MM, Chen L, Bond JE, Medina MA, Ren L, Kokosis G, Selim AM, Levinson H. Myofibroblasts contribute to but are not necessary for wound contraction. Lab Invest. 2015 Dec;95(12):1429-38.