

CONSEQUENCES FOR UNATTACHED PATIENTS

1

1 **Title:** “What do you mean I can’t have a doctor? This is Canada!” – A Qualitative Study of the
2 Myriad Consequences for Unattached Patients Awaiting Primary Care Attachment

3
4 **Authors:** Emily Gard Marshall ^a, Sara Wuite ^b, Beverley Lawson ^a, Melissa K. Andrew ^c, Lynn
5 Edwards ^d, Adrian MacKenzie ^e, Ana Correa Woodrow ^f, Sarah Peddle ^f

6
7 **Affiliations:** ^a Department of Family Medicine, Primary Care Research Unit, Dalhousie
8 University, Halifax, Canada
9 ^b Public Health, Nova Scotia Health, Halifax, Canada
10 ^c Department of Medicine (Geriatrics), Dalhousie University, Halifax, Canada
11 ^d Primary Health Care & Chronic Disease Management, Nova Scotia Health, Halifax,
12 Canada
13 ^e Research and Innovation, Nova Scotia Health, Halifax, Canada
14 ^f Patient Partner, Halifax, Canada

15
16 **Corresponding Author:** Dr. E.G. Marshall
17 Associate Professor, Department of Family Medicine, Primary Care
18 Research Unit
19 Dalhousie University
20 1465 Brenton St., Suite 402
21 Halifax, NS
22 B3J 3T4
23 Canada
24 Emily.Marshall@dal.ca
25

26

27

28

29

30

31

32

33

34

35

36

37 **Abstract:**

38 **Background:** Patient access to primary healthcare (PHC) is the foundation of a strong healthcare
39 system and healthy populations. Attachment to a regular PHC provider, a key to healthcare
40 access, has seen a decline in some jurisdictions. This study explored the consequences of
41 unattachment from a patient perspective, an under-studied phenomenon to date.

42 **Methods:** A realist-informed qualitative study was conducted with unattached patients in Nova
43 Scotia, Canada. Semi-structured interviews with nine participants were conducted and
44 transcribed for analysis. The framework method was used to carry out analysis, which was
45 guided by Donabedian's model of assessing healthcare access and quality.

46 **Results:** Five key findings were noted in this study: 1) Participants experienced a range of
47 consequences from not having a regular PHC provider. Participants used creative strategies to 2)
48 attempt to gain attachment to a regular PHC provider, and, to 3) address their health needs in the
49 absence of a regular PHC provider. 4) Participants experienced negative feelings about
50 themselves and the healthcare system, and 5) stress related to the consequences and added work
51 of being unattached and lost care.

52 **Conclusions:** Unattached patients experienced a burden of care related to lost care and managing
53 their own health and related information, due to the download of medical record management
54 and system navigation to them. These findings may underestimate the consequences for further
55 at-risk populations who would not have been included in our recruitment. This may result in
56 poorer health outcomes, which could be mitigated by interventions at the structural level, such as
57 enhanced centralized waitlists to promote attachment. Such waitlists may benefit from a triage
58 approach to appropriately attach patients based on need.

59

60 **MeSH keywords:**

- 61 • Primary health care
- 62 • Primary Care
- 63 • Access
- 64 • Family practice
- 65 • Waiting lists
- 66 • Continuity of patient care

67

68

69 **Background**

70 Patient access to primary healthcare (PHC) is the cornerstone of a strong healthcare care
71 system; patient attachment to a regular PHC provider (a family physician or nurse practitioner)
72 helps prevent illness and death (1). At the health system level, it is associated with better
73 population health outcomes, lower overall costs of care, and reduced health disparities across
74 socioeconomic status (2). In Canada, primary care is the first line of contact with the healthcare
75 system, through which patients access diagnostics and further specialist care. For patients,
76 attachment is related to more timely treatment (3), better preventive care and chronic disease
77 management (4,5), lower likelihood of unmet health needs (6,7), and ultimately, better health
78 outcomes (1,2). However, some populations face challenges to accessing care and experience
79 lower attachment rates (7), such as recent immigrants (8), LGBTQ+ populations (9), and those
80 experiencing homelessness (10).

81 Historically, population-level attachment to a PHC provider has been high in Canada.
82 The percentage of Canadians with a regular PHC provider grew significantly from 83.2% in
83 2015 to 85.5% in 2019 (11). However, in Nova Scotia, a province with an aging population and
84 high burden of chronic illness, attachment instead dropped significantly from 88.7% to 85.6%
85 throughout that same time period (11). Among those Nova Scotians without a PHC provider,
86 41% reported that they previously had a provider who had left or retired (11). Because of this,
87 there has been widespread public scrutiny regarding the lack of PHC providers in the province
88 (12), an issue which has grown in the wake of the COVID-19 pandemic (13). Provincial
89 governments across Canada are responsible for healthcare operations and delivery, and recognize
90 the importance of PHC attachment. As of May 2021 seven Canadian provinces, including Nova

91 Scotia, have implemented centralized waitlists for their unattached patients in an attempt to
92 streamline attachment processes and increase attachment rates (14).

93 As patients move through phases of becoming unattached and back to attachment, such as
94 while they are on a centralized waitlist, they may use creative strategies to access care and to find
95 a new PHC provider (15,16). However, little is known about what the consequences of
96 unattachment are from a patient perspective or about the strategies that unattached patients use to
97 meet their health needs and access care. The purpose of this study is to inform health system
98 strategies for mitigating negative consequences of unattachment and for increasing attachment
99 by understanding patient-defined impacts of unattachment and the strategies they use to access
100 PHC. The study was guided by the following research questions: a) What are the lived
101 experiences of unattached patients in relation to their health, healthcare needs, and attempts to
102 find a regular PHC provider; and b) How are unattached patients' experiences related to the
103 social determinants of health and social vulnerability?

104 **Methods**

105 We conducted a qualitative study informed by a realist orientation, which focuses on the
106 experiences and realities of participants (17). This methodological approach is appropriate as the
107 topic of unattachment in PHC is novel in the literature, with only two qualitative studies in
108 chronically-ill subpopulations found to date (15,16). The perspectives of unattached patients,
109 especially in this geographic region, are unknown and require an exploratory approach. Our
110 analysis was informed by Donabedian's model of assessing healthcare access and quality,
111 focused on structure, process, and outcomes (18), and operationalized using the framework
112 method of qualitative analysis as outlined below (19).

CONSEQUENCES FOR UNATTACHED PATIENTS

5

113 Our study was approved by the Nova Scotia Health Authority Research Ethics Board
114 (File #1022763). We recruited unattached patients using invitational letters to patients on the
115 Nova Scotia Need a family Practice Registry. To maintain the confidentiality of people on the
116 registry, letters were sent by the registry custodians on behalf of the study team. We also
117 advertised on social media and local online marketplaces. For all recruitment methods, potential
118 participants were instructed to contact the study team if they were interested in participating. To
119 be considered for inclusion, potential participants had to be over 18, speak English, reside in
120 Nova Scotia, and not have a regular family physician or nurse practitioner. After receiving
121 expressions of interest via email from potential participants, we conducted pre-screening phone
122 calls and iteratively selected participants using purposive sampling based on geography, age,
123 gender, and healthcare needs (20,21). We continued to invite participants until saturation was
124 reached (22).

125 A trained qualitative researcher conducted face-to-face interviews in locations that were
126 convenient for each participant. The interview guide was developed by the study team, based on
127 unattached patient team member experience. Interviews lasted approximately one hour, were
128 digitally recorded, and transcribed verbatim.

129 Interview data were coded inductively and deductively. Initial transcript coding was
130 conducted by four team members, including the interviewer and principal investigator; final
131 coding of all transcripts was conducted by the interviewer. Initial coding categories were
132 developed based on the experience of unattached patient team members. Additional codes were
133 added inductively from the transcript data. The final coding and thematic framework structure
134 were developed iteratively and collaboratively among four team members, including an
135 unattached patient team member. All analysis was conducted in NVivo 11 (QSR International).

136 **Results**

137 We interviewed nine participants with varied socio-demographic representation (Table
 138 1).

139 Table 1: Socio-demographic characteristics of unattached patient study participants

	Number of Participants (%)
Gender	
Male	3 (33%)
Female	6 (66%)
Age (years)	
18-39	2 (22%)
30-49	1 (11%)
40-49	1 (11%)
50-59	1 (11%)
60-69	4 (44%)
Health Status (self-report)	
Very good	7 (77%)
Good	2 (22%)
Visible Minority	
Yes	2 (22%)
No	7 (77%)
Highest Education	
High School	1 (11%)
Trade Diploma	2 (22%)
University Certificate or College Diploma	3 (33%)
Bachelor's	3 (33%)
Employment Status	
Employed	5 (55%)
Self-Employed	1 (11%)
Other	3 (33%)
Household Income (\$CAD)	
10,000-19,000	1 (11%)
20,000-29,000	2 (22%)
30,000-39,000	0
40,000-49,000	0
50,000-59,000	0
61,000-69,000	0
70,000-79,000	2 (22%)
80,000-89,000	1 (11%)
90,000-99,000	1 (11%)
100,000-149,000	1 (11%)
150,000+	1 (11%)

CONSEQUENCES FOR UNATTACHED PATIENTS

7

140
141 Participants became unattached through both patient- and provider- initiated mechanisms. Most
142 commonly, participant moved either within the province or from outside of the province. In some
143 cases, a previous PHC provider retired or moved out of province. In one instance, a participant
144 had a previous PHC provider that may have been available to them after moving back to Nova
145 Scotia, but they were uncomfortable receiving care from them and so chose to look for a new
146 regular PHC provider. No participants in our sample were “fired”(23) from their most recent
147 PHC provider before participating in the study.

148 Guided by framework analysis (19), five main findings from interviews with unattached
149 patients were uncovered and are described in more detail below: 1) Unattached patients
150 experience a range of consequences from not having a regular PHC provider. Unattached
151 patients use creative strategies to 2) attempt to gain attachment to a regular PHC provider and to
152 3) address their health needs in the absence of a regular PHC provider. Unanimously, 4)
153 participants experienced stress related to the consequences and added work of being unattached
154 and 5) had negative feelings and sense of abandonment from not having a regular PHC provider.


155 Consequences of Unattachment

156 Participants experienced a range of consequences from being unattached related to the burden of
157 care, lost care, and health impacts (Figure 1).

158 Figure 1: Themes related to consequences of unattachment.

159

Consequences of Unattachment



160

161 *Care Burden*

162 All participants experienced an increased burden of managing their own healthcare in the
163 absence of a regular PHC provider. Participants experienced increased burden compared to when
164 they were attached for finding and managing information, managing their own medical history,
165 navigating health and other service systems, and in some cases, patient costs including time and
166 increased travel distances for care.

167 Participants look to their PHC providers as experts in medicine and in the healthcare
168 system. Without a regular PHC provider, several participants looked to other sources for
169 information that may have some healthcare expertise, such as the internet or family and friends.

CONSEQUENCES FOR UNATTACHED PATIENTS

9

170 Three participants expressed that they specifically felt the loss of an expert who could give
171 concrete answers and to help process information into something actionable.

172 *“Kind of that hub where, you know, all the information goes in and somebody*
173 *needs to kind of figure out what to do with it” – Participant 8*

174 All participants discussed the burden of having to manage their own medical record while
175 unattached. This manifested as needing to communicate their medical history to interim
176 providers, managing new healthcare information and instructions that would typically be held in
177 a medical record under the custodianship of a regular PHC provider, and extra work to track and
178 access their existing medical records from previous PHC providers. Participants expressed that
179 they felt stressed by these added responsibilities and worried that they may not be able to carry
180 them out as well as a regular PHC provider. One participant took detailed notes in order to
181 maintain their medical history while accessing walk-in clinics; another paid to have a print copy
182 of her whole medical history up until they became unattached.

183 *“...I did a lot of talking to bring the [walk-in clinic] doctor up to speed ...*
184 *what happens now if I got to somebody that I don't know and doesn't know me*
185 *or my history, and I have 15 minutes or whatever, and they're absolutely*
186 *overloaded with patients anyway?” – Participant 1*

187 *“Because when I got to the walk-in, and I tell them I had a surgery once but*
188 *they don't really know much about it, and I can't tell them a whole lot about it*
189 *... Like, I am not very good at, you know, explaining things when I am on the*
190 *spot.” - Participant 4*

CONSEQUENCES FOR UNATTACHED PATIENTS

10

191 At the intersection of the loss of a “home” for their medical records and an increased
192 burden of managing healthcare information, our study participants expressed the consequence of
193 having to navigate the healthcare system alone or with dis-jointed advice. Some participants
194 expressed confusion about how long is normal to wait for a follow-up from a referral or who is
195 even responsible for initiating that follow-up. Others felt challenged by having to navigate
196 different care providers and services without the central “hub” of a PHC provider to help
197 coordinate their healthcare around a particular health issue.

198 *“You’re just sort of roaming around a maze, you know, with drones at each*
199 *station.” –Participant 3*

200 Although healthcare in Canada, including PHC, walk-in clinics, and hospital care, is
201 funded through taxes and has no user access fees, some services, such as purchase of most
202 prescription drugs and services of some allied health professionals, incur fees unless covered by
203 an insurance plan. Participants in this study faced increased financial and time costs related to
204 not having a regular PHC provider. Most commonly, participants sought over the counter
205 medications, where if they had a regular PHC provider, they would have sought a prescription
206 (or other treatment avenue) that would then be covered through their medical insurance. Four
207 participants had costs related to accessing their medical records or sending medical information
208 to their insurance companies. Several participants also expressed costs related to travel, including
209 going to walk-in clinics or other healthcare services that are further than they would typically
210 travel; paying for taxis or in one case, an ambulance; and time away from work having to wait in
211 emergency rooms and walk-in clinics that do not offer pre-booked appointment times.

212

CONSEQUENCES FOR UNATTACHED PATIENTS

11

213 *Lost Care*

214 All patients had trouble accessing certain kinds of care and felt they “lost” some care due
215 to not having a regular PHC provider. There was a strong theme of the benefit of “being known”
216 to a regular PHC provider, which supported participant’s comfort in accessing care and
217 providers’ ability to provide continuous care. Without a regular PHC provider, participants did
218 not feel “known” by the interim providers they saw. This, grouped with the unavailability of
219 their whole medical history and bouncing between various interim care providers, resulted in
220 feeling that pieces of their care got lost.

221 *“Having to tell your story over and over and over again to ...and also I think*
222 *for a physician, they may be missing certain links. I think the impact has been*
223 *enormous, actually” – Participant 5*

224 *“The first time [with an emerging medical issue] I bounced from doctor to*
225 *doctor each visit ... History matters. It matters very much. Yeah, you know*
226 *your doctor and your doctor knows you” – Participant 2*

227 All participants faced challenges, sometimes insurmountable, in accessing either medical
228 follow up from a visit to an interim provider, prescriptions, or referrals to specialists. In
229 particular, getting prescriptions for controlled medications – such as those for ADHD, sleeping,
230 and other mental illnesses – posed serious problems at walk-in clinics, even if the participant had
231 been prescribed the medication they were seeking previously. Similarly, some participants had
232 trouble accessing referrals to specialists or for testing, and then if a referral was made, were
233 challenged by a lack of a “home” to manage follow-up, such as receiving test results and next
234 steps. For example, one patient struggled to have an issue diagnosed via walk-in clinics but

235 received one upon an emergency room visit for acute pain related to the issue. From the
236 emergency room, they were referred to a surgeon for treatment. The participant was unsure of
237 how long they should wait to hear from the surgeon; eventually, it became clear that the referral
238 either was not made or got lost. They then struggled to gain another referral from interim PHC
239 sources.

240 *“I mean, I didn’t know if I could have called the emergency room to ask who I*
241 *had been referred to. It didn’t seem like that was a place I could sort of go*
242 *back to to follow up. So, you just sort of feel a little bit sort of forgotten and*
243 *kind of lost in the system. Like I think I got lost in the system a couple of times,*
244 *and I didn’t really know how to follow up on it or just what to do next.” –*

245 *Participant 9*

246 *Health Impacts*

247 When asked directly if being unattached had direct impacts on their health, most
248 participants responded that it did not. However, when probed further about specific experiences,
249 health impacts emerged, including making lifestyle changes to improve their health, condition-
250 specific changes in health, incorrect, missed, or delayed diagnoses, self-diagnosis and
251 medication, and not seeking care.

252 Three participants made proactive lifestyle changes in an attempt to reduce their
253 healthcare needs and to manage existing symptoms. Lifestyle changes included diet, exercise,
254 being more cautious, and occupational health changes (e.g., shoe insoles for walking,
255 physiotherapy-inspired exercises for office work).

256 *“Even walking down stairs, like even when I’m moving around, I’m thinking*
257 *just take a second and think about what you’re doing...I’m trying to keep*
258 *myself from need to see [a doctor].” –Participant 3*

259 Six participants experienced some deterioration in their health or sustained symptoms,
260 related to a specific health issue that they each experienced, that they attributed to not having a
261 PHC provider.

262 *“...kind of waiting for these pain flare-ups to happen again because it would*
263 *sort of be unexpected flare-ups with terrible pain.” – Participant 9*

264 *“It’s just been kind of the same things having to deal with several years or*
265 *longer. For a long time at this point. So, it just kind of ends up becoming the*
266 *background noise of my life” – Participant 7*

267 Seven participants felt that they had diagnoses that were either incorrect, delayed, or
268 missed altogether due to the discontinuity of PHC and the absence of having a complete medical
269 record available for interim care without a regular PHC provider. For example, one participant
270 had a persistent issue for which they sought care at a walk-in clinic several times and then the
271 emergency room. They eventually had to call an ambulance and seek emergency care for the
272 issue. Another participant spent five months with debilitating symptoms before receiving a
273 diagnosis that was ultimately picked up with routine bloodwork, which had never been ordered
274 by the walk-in doctors she visited previously.

275 In the absence of a regular PHC provider, several participants reported self-diagnosing
276 and treating their health conditions, despite their perception that this was not “the right” thing to

CONSEQUENCES FOR UNATTACHED PATIENTS

14

277 do. Relatedly, some participants reported not seeking any care even when they felt it was
278 warranted and would have if they had a regular PHC provider. In both cases, participants felt
279 either the barriers of finding appropriate “professional” healthcare were insurmountable (e.g.,
280 waiting at a walk-in or in emergency), or, that their healthcare needs were not severe enough to
281 warrant discontinuous walk-in care or emergency care.

282 *“So, you don’t feel you should go in unless you’ve really got something serious*
283 *wrong with you. So yeah, you do self-medicate” – Participant 2*

284 *“And I guess the biggest difference is we sort of feel held back from dealing*
285 *with certain issues because it is not convenient to be able to access the care.*
286 *So maybe we would have gone to the doctor for less of a reason” – Participant*

287 6

288 Strategies for Care: Attachment Strategies

289 *Unattached Patients’ Registry*

290 Most participants were registered on the provincial unattached patient registry. Only one
291 of the participants attached through the registry at the time of their interview, and none of the
292 participants on the registry held much hope that they would find a provider through this strategy.

293 *“And I always kind of figured that I wouldn’t be on the list for very long ...*
294 *And so, I figured it was just a matter of time before they called me or contacted*
295 *me at all. But no, they never did. They still haven’t.” – Participant 7*

296 *“But we put ourselves on the list with the government for all the [expletive]*
297 *good that’s going to do” – Participant 3*

298 *Interim PHC Provider*

299 A common strategy was to gain or approximate attachment through an interim care
300 provider. Participants would either try to attach to the regular practice of the doctors that they
301 saw in walk-in clinics or rural emergency rooms, or they would attempt to return to the same
302 walk-in clinic doctor each time they visited. In one case, a walk-in doctor agreed to see a patient
303 continuously until a particular health issue was resolved.

304 *Cold-Calling PHC Practices*

305 Another common practice was to cold-call PHC offices to see if any of the providers
306 were taking new patients. Six of the participants tried this approach, but half have since stopped
307 due to feelings of discouragement. One participant had a dedicated approach of calling all the
308 practices in a geographical area that feels acceptable to them every two months. This method
309 elicited a lot of frustration among participants, but also empathy for the receptionists who receive
310 their calls, whom participants described as generally pleasant and sympathetic. No participants
311 found a new provider with this approach.

312 *“I’ve stopped [calling clinics] because it’s pointless. You know, I’m well*
313 *aware of how bad it is now. So, I’ve stopped trying” – Participant 6*

314 *Personal Networks*

315 Seven participants used their personal networks to try to find a new regular PHC
316 provider. Participants’ approaches ranged from asking friends and family if they knew of any
317 providers taking new patients, to requesting that they ask their providers directly (e.g., a mother
318 asked if her physician would take her unattached son), to asking directly among PHC providers
319 that they knew personally. There were mixed approaches to using social media – some

CONSEQUENCES FOR UNATTACHED PATIENTS

16

320 participants used all networks available to them, while some did not feel comfortable
321 “broadcasting” their healthcare issues in such a public way. Two participants indicated that they
322 used public and personal political avenues to try to inspire broader change or simply understand
323 the larger issue of unattachment to be better able to navigate the system to find a provider for
324 themselves.

325 *“You know, the only way you’re going to find a doctor in NS is if you know*
326 *somebody or if somebody happens to move into a new practice.” –Participant*

327 3

328 *Expanding Geographic Area*

329 Regardless of the active approach to finding a new PHC provider, four participants
330 expressed a willingness to attach to a provider in a geographic area outside of what they consider
331 convenient. Some participants expressed frustration that you could not indicate willingness to
332 travel on the unattached patients’ registry.

333 *Giving Up*

334 Finally, most participants simply gave up actively looking for a regular PHC provider.

335 *“...there is no doctor to be found. You know, we start paying attention to the*
336 *news and ... you know. And it was the start realization that, you know what,*
337 *it’s a systemic problem in this province” – Participant 6*

338 Accessing Care While Unattached

339 Similar to not immediately identifying health impacts related to being unattached,
340 participants frequently answered that they did not have any ongoing healthcare needs while
341 unattached. However, as participants’ narratives unfolded, it became clear that most had either an

CONSEQUENCES FOR UNATTACHED PATIENTS

17

342 emergent healthcare need or a controlled chronic condition during their unattachment. While
343 participants did not always identify a specific need in the time that they were unattached, most
344 expressed concern for the future, when things “get worse” or “come up.”

345 *Walk-in Clinics*

346 Walk-in clinics were the primary source of PHC for participants while unattached. Some
347 participants could access walk-in hours in the family practice clinic they used to attend, others
348 used dedicated walk-in clinics that provide no continuous care. While some were able to use the
349 same walk-in clinic repeatedly to approximate attachment, most found them to be frustrating due
350 to wait times, feeling rushed in the appointment, reluctance of walk-in doctors to order tests,
351 write referrals, or write certain prescriptions, having to be the reporter of their own medical
352 history, and the perceived incompetence of some walk-in physicians. While some participants
353 were thankful to have access to some form of care, most felt walk-ins were an inadequate
354 replacement for a regular PHC provider.

355 *“But having been to a couple of walk-in clinics for various things, like, you*
356 *know, a sinus infection for my daughter or strep throat for my son or whatever,*
357 *and having to deal with these people at these walk-in clinics was enough to*
358 *turn me off and say, you know what, it’s not worth dealing with.” – Participant*

359 6

360 *Emergency Care*

361 Overall, participants were reluctant to use emergency care unless they felt their issue was
362 truly an emergency. In some cases, participants did use emergency care when they felt it was

363 warranted, but that it could have been avoided if they had a regular PHC provider to address their
364 issue before it escalated.

365 *Specialists and Other Healthcare Professionals*

366 Participants used a variety of healthcare providers that were available to them. Two
367 participants had access to specialists for other ongoing health issues and had those specialists
368 address issues outside of their area of expertise that would typically be addressed by a PHC
369 provider. Pharmacists were accessed by seven participants in lieu of a PHC provider, primarily
370 for preventative health services like flu shots. Participants also access self-referral walk-in
371 clinics, methadone clinic physicians, chiropractors, and women's health clinics for issues that
372 would be appropriate for PHC.

373 Stress of Managing Unattachment and Concerns for Future Health

374 Overwhelmingly, due to the consequences and the increased work for accessing care, all
375 participants experienced stress related to being unattached. Most participants expressed concern
376 for the future, related to accessing care for either an exacerbation or recurrence of an existing
377 health condition or for a new issue, often related to aging. This stress was especially pronounced
378 among participants who had chronic health conditions that would benefit from a PHC provider's
379 involvement. Several participants also expressed frustration and worry about the perceived lack
380 of choice of providers – both at walk-in clinics and with a potential new regular PHC provider.

381 *“It probably wouldn't bother me a lot but I've been through cancer treatment.*

382 *So I'm in remission. But I have ongoing care from that experience. But what I*

383 *don't have is somebody [to care for me] if something else comes up even*

384 *related to my cancer.” - Participant 1*

385 *“I am fortunate right now that things are kind of stabilized and I am healthy.*

386 *But you know, if that changes, I’m not sure what I would do.” – Participant 9*

387 *“Health and stress go together in my life” – Participant 3*

388 Negative Feelings and Sense of Abandonment due to Unattachment

389 Finally, several participants expressed an unexpected feeling of abandonment or betrayal
390 by the healthcare system due to not having a regular PHC provider. Participants expressed
391 feeling demoralized and “like a beggar” in the process of seeking a new PHC provider. They also
392 expressed a desire to have access to a regular PHC provider and described the sense of comfort
393 and reassurance they would feel if they could become attached.

394 *“Yeah, you’re not expecting it. It’s kind of shocking. And then to realize there are no doctors.*

395 *What do you mean I can't have a doctor? This is Canada! We all have medical care, right? No,*

396 *we don’t. It was startling.” – Participant 2*

397 *“The biggest thing is not having that level of comfort. You know, that it’s not... It’s a bit of a*

398 *security blanket to know that you have a family doctor.” - Participant 6*

399 **Discussion**

400 Unattached patients experience a range of consequences related to not having a regular
401 PHC provider. They also use a variety of creative strategies to seek attachment and to address
402 their healthcare needs. Our study found that unattached patients experience a burden of care
403 related to managing their own medical records and worry about having to be the reporter of their
404 own medical history, especially when it is with an unfamiliar provider. This is consistent with
405 the findings of another qualitative study in British Columbia that found similar themes related to

406 medical records and a desire for a trusting relationship with a regular PHC provider (16).
407 Interestingly, patients were able to keenly identify the stress associated with not having a regular
408 PHC provider, but until prompted about specific experiences, were hesitant to express specific
409 health consequences or even healthcare needs. While participants were creative in their efforts to
410 obtain interim care, it was clear that many still had healthcare needs that went unmet. Indeed, not
411 having a regular PHC provider is a significant and strong predictor of having unmet healthcare
412 needs (7).

413 Another small study of unattached women with chronic illness identified similar
414 strategies to accessing interim care and attachment, including relying on specialists until their
415 healthcare needs exceeded the specialists' scope, and using walk-in clinics; this study also
416 identified some of the same frustrations with walk-in clinics, including wait times, inability to
417 book an appointment, and seeing different PHC providers each time (15). This study also found
418 similar strategies to gain attachment, including calling practices and a willingness to expand
419 geographical scope.

420 Building on these two qualitative studies on unattachment, our study adds a deeper
421 understanding of the impact of these added care responsibilities for a general population of
422 unattached patients, ranging from self-diagnosis and treatment through to giving up on both
423 seeking care and looking for a new PHC provider, improving health behaviours to avoid needing
424 care, and a negative feeling of abandonment from a health care system though thought they could
425 always rely on.

426 While we reached data saturation, our study is still limited by a small number of
427 volunteer participants within one province in a country with provincially delivered healthcare.
428 Our recruitment methods were also not designed to reach marginalized populations, members of

429 cultural and language minorities, socially isolated and/or vulnerable people, older adults, those
430 with limited access to technology, those with severe health conditions, or those with cognitive
431 impairment who are likely to experience amplified or different consequences of unattachment,
432 and for whom the compensating strategies to meet health care needs described by our
433 participants may not be feasible. However, recruited participants who were relatively empowered
434 and particularly motivated to share their experiences, our purposive sampling method allowed
435 for participant selection across several socio-demographic variables and unattachment
436 experiences, highlighting the pervasiveness of negative consequences of unattachment across
437 sociodemographic characteristics that are not limited to those with more challenging
438 circumstances.

439 **Conclusions**

440 Our study demonstrates the impact of unattachment and can be understood using the
441 Donabedian model of healthcare access of structure, process, and outcomes (18). While
442 unattached, much of the “process” of healthcare, including medical record management and
443 system navigation, is downloaded from PHC providers onto patients, which may contribute to
444 poorer health and social outcomes. Similarly, patients are responsible for seeking new
445 attachment via centralized wait lists and other means, as well as their care while unattached.
446 Patients may also have differential capacity to pursue attachment and may face discrimination in
447 that process (24). Hence, first-come-first-served centralized waitlists may be an insufficient
448 approach that has the potential to be unfair and result in inequitable care and re-attachment along
449 socio-demographic axes. As such, unattached patients may benefit from interventions at the
450 “structure” level, such as health human resource interventions to ensure an adequate PHC
451 workforce and patient registries/wait lists to facilitate patient attachment to PHC.

452 Given that patients with ongoing health needs cited greater stress related to not having a
453 PHC provider relative to patients without a regular PHC provider, our study suggests that patient
454 registries and waitlists may best support unattached patients by taking a triaged approach to
455 prioritize attachment for patients with greater needs, rather than using a purely first-come-first-
456 served, time-based algorithm. The more severe health impacts for participants that were
457 managing an acute healthcare need suggest that these people (and the health system as a whole)
458 may have benefited from more efficient attachment than those who did not have any severe
459 chronic or acute care concerns. Participants’ “concern for the future” when not perceiving an
460 immediate healthcare need and their reluctance to use emergency rooms unless absolutely
461 necessary suggests there is an openness among unattached patients to ranking need to shepherd
462 efficient and effective attachment. Triaged registries exist in British Columbia, Ontario, and
463 Quebec (14), where they have demonstrated effectiveness at prioritizing vulnerable patients (25),
464 with other structural factors promoting even stronger attachment for vulnerable patients (26).

465 Last, but not least, it is critical to consider how a potentially growing lack of confidence
466 in the health care system’s ability to provide access to having a regular primary care provider
467 may impact support for Canada’s decentralized, universal, publicly funded health system. Next
468 steps include sharing these findings with our provincial government, health authority, and patient
469 partner stakeholders who are active members of the study team. Future work could use survey
470 methods to observe the spread and scale of the concerns and consequences identified by our
471 research participants, and the potential impact of the COVID-19 pandemic on access to primary
472 care attachment.

473

474

475 **Declarations / Acknowledgements:**

476 ***Ethics approval and consent to participate***

477 This study was approved by the Nova Scotia Health Research Ethics Board (#1022374).

478 The informed consent discussion was conducted by a trained master-level interviewer, and
479 participants provided written informed consent.

480 ***Consent for publication***

481 Not applicable.

482 ***Availability of data and materials***

483 The data that supports the findings of this study are not publicly available due to the identifiable
484 nature of raw data (interview transcripts). Requests for additional information about study data or
485 results can be forwarded to the corresponding author (EGM).

486 ***Competing interests***

487 The authors declare that they have no competing interests.

488 ***Funding***

489 This study was funded by the Nova Scotia Health Research Fund (#1021744).

490 ***Authors' contributions***

491 EGM conceptualized the study, designed the methodology, established protocol, analyzed and
492 interpreted data, provided software, writing the manuscript, supervision of data collection and
493 analysis, funding acquisition. SW collected, analyzed and interpreted data, manuscript writing
494 and editing. BL supported funding application, study development, analysis review, writing-
495 reviewing, and editing. MKA supported funding application, study development, analysis
496 review, writing-reviewing, and editing. LE supported funding application, study development,
497 writing-reviewing, and editing. AM supported funding application, study development, analysis
498 review, writing-reviewing, and editing. ACW supported funding application, study development,

CONSEQUENCES FOR UNATTACHED PATIENTS

24

499 analysis review, writing-review, active Patient Partner for this study. SP supported funding
500 application, study development, analysis review, writing-review and editing, active Patient
501 Partner for this study.

502 *Acknowledgements*

503 We gratefully acknowledge our study participants who generously shared their time and lived
504 experiences as unattached patients.

505 References

- 506 1. Starfield B, Shi L, Macinko J. Contribution of primary care to health systems and health.
507 The milbank quarterly. 2005;83(3):457–502.
- 508 2. Starfield B, Shi L. The medical home, access to care, and insurance: a review of evidence.
509 Pediatrics. 2004;113(Supplement 4):1493–8.
- 510 3. DeVoe JE, Fryer GE, Phillips R, Green L. Receipt of preventive care among adults:
511 insurance status and usual source of care. American journal of public health.
512 2003;93(5):786–91.
- 513 4. McIsaac WJ, Fuller-Thomson E, Talbot Y. Does having regular care by a family physician
514 improve preventive care? Canadian Family Physician. 2001;47(1):70–6.
- 515 5. Xu KT. Usual source of care in preventive service use: a regular doctor versus a regular
516 site. Health services research. 2002;37(6):1509–29.
- 517 6. Statistics Canada. Unmet health care needs, 2014 [Internet]. Health Facts Sheets. 2016
518 [cited 2021 May 12]. Available from: [https://www150.statcan.gc.ca/n1/pub/82-625-](https://www150.statcan.gc.ca/n1/pub/82-625-x/2016001/article/14310-eng.htm)
519 [x/2016001/article/14310-eng.htm](https://www150.statcan.gc.ca/n1/pub/82-625-x/2016001/article/14310-eng.htm)
- 520 7. Awe OA, Okpalauwaekwe U, Lawal AK, Ilesanmi MM, Feng C, Farag M. Association
521 between patient attachment to a regular doctor and self-perceived unmet health care needs
522 in Canada: A population-based analysis of the 2013 to 2014 Canadian community health
523 surveys. The International journal of health planning and management. 2019;34(1):309–23.
- 524 8. Hay C, Pacey M, Bains N, Ardal S. Understanding the unattached population in Ontario:
525 evidence from the Primary Care Access Survey (PCAS). Healthcare Policy. 2010;6(2):33.
- 526 9. Gahagan J, Subirana-Malaret M. Improving pathways to primary health care among
527 LGBTQ populations and health care providers: key findings from Nova Scotia, Canada.
528 International Journal for Equity in Health. 2018 Jun 13;17(1):76.
- 529 10. Khandor E, Mason K, Chambers C, Rossiter K, Cowan L, Hwang SW. Access to primary
530 health care among homeless adults in Toronto, Canada: results from the Street Health
531 survey. Open Medicine. 2011;5(2):e94.
- 532 11. Statistics Canada. Primary health care providers, 2019 [Internet]. 2020 [cited 2021 May
533 13]. Available from: [https://www150.statcan.gc.ca/n1/pub/82-625-](https://www150.statcan.gc.ca/n1/pub/82-625-x/2020001/article/00004-eng.htm)
534 [x/2020001/article/00004-eng.htm](https://www150.statcan.gc.ca/n1/pub/82-625-x/2020001/article/00004-eng.htm)
- 535 12. Draus A. List of patients in need of a family doctor growing in N.S. rural areas [Internet].
536 Global News. [cited 2021 May 13]. Available from:
537 [https://globalnews.ca/news/6403926/list-of-patients-in-need-of-a-family-doctor-growing-](https://globalnews.ca/news/6403926/list-of-patients-in-need-of-a-family-doctor-growing-in-n-s-rural-areas/)
538 [in-n-s-rural-areas/](https://globalnews.ca/news/6403926/list-of-patients-in-need-of-a-family-doctor-growing-in-n-s-rural-areas/)

- 539 13. Grant T. In this pandemic year, the number of Nova Scotians waiting for a family doctor
540 has grown. *CBC*. 2020;6.
- 541 14. Breton M, Wong ST, Smithman MA, Kreindler S, Jbilou J, Sutherland J, et al. Centralized
542 waiting lists for unattached patients in primary care: learning from an intervention
543 implemented in seven Canadian provinces. *Healthcare policy*. 2018;13(4):65.
- 544 15. Randall E, Crooks VA, Goldsmith LJ. In search of attachment: a qualitative study of
545 chronically ill women transitioning between family physicians in rural Ontario, Canada.
546 *BMC family practice*. 2012;13(1):1–12.
- 547 16. Crooks VA, Agarwal G, Harrison A. Chronically ill Canadians’ experiences of being
548 unattached to a family doctor: a qualitative study of marginalized patients in British
549 Columbia. *BMC Family Practice*. 2012;13(1):69.
- 550 17. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative research in*
551 *psychology*. 2006;3(2):77–101.
- 552 18. Donabedian A. The quality of care. How can it be assessed? *JAMA*. 1988 Sep
553 23;260(12):1743–8.
- 554 19. Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework method for the
555 analysis of qualitative data in multi-disciplinary health research. *BMC Medical Research*
556 *Methodology*. 2013 Sep 18;13(1):117.
- 557 20. Patton MQ. *Qualitative research & evaluation methods: Integrating theory and practice*.
558 Sage publications; 2014.
- 559 21. Green J, Thorogood N. *Qualitative Methods for Health Research*. In: *Qualitative Methods*
560 *for Health Research*. 3rd ed. Los Angeles: Sage Publications, Inc; 2013. p. Pg. 360.
- 561 22. Mason M. Sample size and saturation in PhD studies using qualitative interviews. In:
562 *Forum qualitative Sozialforschung/Forum: qualitative social research*. 2010.
- 563 23. College of Physicians and Surgeons of Nova Scotia. *Professional Standard and Guidelines*
564 *for Ending the Physician-Patient Relationship* [Internet]. 2015. Available from:
565 <https://www.cpsbc.ca/files/pdf/PSG-Ending-the-Patient-Physician-Relationship.pdf>
- 566 24. Marshall EG, Burge F, Gibson RJ, Lawson B, O’Connell C. Accepting new patients who
567 require opioids into family practice: results from the MAAP-NS census survey study. *BMC*
568 *family practice*. 2019;20(1):1–8.
- 569 25. Breton M, Smithman MA, Brousselle A, Loignon C, Touati N, Dubois C-A, et al.
570 Assessing the performance of centralized waiting lists for patients without a regular family
571 physician using clinical-administrative data. *BMC family practice*. 2017;18(1):1–13.

CONSEQUENCES FOR UNATTACHED PATIENTS

27

- 572 26. Breton M, Smithman MA, Touati N, Boivin A, Loignon C, Dubois C-A, et al. Family
573 Physicians Attaching New Patients From Centralized Waiting Lists: A Cross-Sectional
574 Study. *Journal of primary care & community health*. 2018;9:2150132718795943.

575

576

577