Title

Long-distance migration was associated with increased prevalence of post-traumatic

stress disorder in Syrian refugees

Authors

Andreas Halgreen Eiset[1]*, ORCID: 0000-0002-5487-8359

Michaelangelo P. Aoun[2]

Monica Stougaard[1]

Annemarie Graa Gottlieb[1]

Ramzi S. Haddad[2]

Morten Frydenberg[3]

Wadih J. Naja[2]

[1]Department of Affective Disorders, Aarhus University Hospital–Psychiatry, Aarhus,

Denmark

[2] Faculty of Medical Sciences, Lebanese University, Beirut, Lebanon

[3]Consultant biostatistician, Olsted, Denmark

*Corresponding author

Abstract

Background: Refugees are forced migrants but there is a large variation in the distance that refugees cover and there is a knowledge gap on how this may affect refugees' health and health care needs. We investigated the association between long-distance migration and post-traumatic stress disorder (PTSD), a serious psychiatric disorder associated with deteriorating mental and somatic health. **Methods:** Included from 2016-2019 were 712 adult Syrian refugees and asylum seekers in Lebanon and Denmark arrived up to 12 months prior. PTSD was assessed using the Harvard Trauma Questionnaire and the estimate of association was obtained by multiply imputing missing data and adjusting for confounding by propensity score-weighting with covariates age, sex, socioeconomic status, trauma experience and WHO-5-score, reporting the bootstrap 95-percentile confidence interval (95% CI). Additionally, a number of sensitivity analyses were performed. **Results:** After multiply imputing missing data and propensity score-weighted adjustment for confounding, migration to Denmark instead of Lebanon was associated with an increase in PTSD prevalence of 9 percentage point (95% CI [-1; 19] percentage point). **Discussion:** We found that long-distance migration was associated with an increase in the prevalence of PTSD suggesting that long-distance migration may be a factor to consider when assessing refugees' and asylum seekers' health. To secure timely and appropriate health care for this vulnerable group we need to remedy the current knowledge gap regarding the health effects of migration. This study is the first step in doing so.

Introduction

Refugees suffer from high risk of post-traumatic stress disorder (PTSD) [1] and are exposed to many risk-factors for PTSD before, during and after migration [2–4]. Risk factors for PTSD in the country of origin and in the host country has been examined in detail; however, the contribution of the migration itself has not received much attention in refugee health research. PTSD is a disorder that may develop after a traumatic event and is associated with significant loss of functioning and increased morbidity and mortality [5,6]. Symptoms of PTSD include intrusion (e.g. "flash-backs" and nightmares), avoidance and arousal, characterized by persistent perception of threat which may result in sleep deprivation, irritability and inability to concentrate [7,8]. Risk factors for developing PTSD comprise multiple trauma, female sex, being spouseless, ethnicity, lower educational status, comorbid mental disorders and type of trauma (for example intentional trauma, sexual assault etc.) [7]. Additionally, gene-environment interactions may play a key role [9]. Dysfunction of the amygdala (emotional learning and memory modulation) and medial prefrontal cortex (executive function) are key elements in PTSD and the disease is associated with epigenetic changes, increased blood levels of norepinephrine and decreased blood levels of glucocorticoids [7]. The life-time prevalence of PTSD in the general population is estimated to be around 5% for males and 10% for females with considerable variation according to the study population and diagnostic tools utilized [7,10]. In comparison, a recent meta-analysis produced a summary prevalence estimate for PTSD of 31% across all refugees [1] with little difference between males (29%) and females (34%) though with considerable variation between the country of origin, host countries and diagnostic tools; however,

not taking migration history into account. The impact of migration and the role and

measurement of "distance" (geographical, cultural, linguistic etc.) has been discussed at

length for a number of outcomes [11,12] and long-distance migration has been

associated with poorer self-rated health in the general population [13], hypothesizing a

link with increasing difficulty in acculturation [14,15]. In the present study, we aimed at

taking an initial step in assessing the effect of migration on refugees' and asylum

seekers' health by comparing the prevalence of PTSD in newly arrived Syrian refugees

and asylum seekers in Lebanon and Denmark.

Methods

In a cross-sectional design with one-stage cluster randomized sampling between 2016

and 2019, newly arrived adult Syrian refugees were included in Lebanon and newly

arrived Syrian asylum seekers were included in Denmark and assessed for PTSD using

the Harvard Trauma Questionnaire (HTQ)-scale with cut-off score at 2.5. The

implemented inclusion criteria were: (a) adult ($\geq$18 years of age), (b) Syrian-born, (c)

had left Syria after the onset of the ongoing civil-war (after February 2011), (d) arrived

in Lebanon or Denmark less than 12 months prior to inclusion and (e) resident in the

Lebanese settlements or Danish asylum seeker centres at the time of inclusion.

Exclusion criteria were physical or mental illness that prevented participation. Variables

to be collected to address confounding were decided upon *a priori* after discussions

among the authors guided by graphical representations of our assumptions. See the

Supplement for a detailed description of the data collection instruments and the directed

acyclic graph illustrating our assumptions about the relation of the variables. Also the

study protocol is published and freely available [16].

Lebanon is a country of about 6 million people living in an area of nearly 10.000 km²—

comparable in size to the two major Danish island (Zealand and Funen). Since 2015

Lebanon has stopped accepting asylum applications from Syrians seeking refuge in

Lebanon. Nevertheless, being a close neighbour, in both geographical, linguistic and

cultural distance, the influx of refugees from the Syrian war has endured. The majority

live in structures ranging from regular (though dilapidated) structures to improvised

tents clustered in informal gatherings scattered across Lebanon [17]. We stratified

Lebanon into five regions based on geography, infrastructure and insights about local

concerns (political, religious etc.). In each region, we collaborated with the local health

service operators to draw up the sampling frame of gatherings and randomly decided on

the inclusion sites, where all individuals eligible for inclusion were invited to participate

(i.e. stratified cluster sampling [18,19]). See the Supplement for details on the sampling

frame. We obtained permission to include participants in all regions of Lebanon, that is,

we had access to include participants from areas that are usually outside the reach of

researchers due to "security considerations". Should a chosen site be deemed unsafe at

the scheduled time of inclusion we relocated to the closest setting deemed appropriate

within the same region.

Denmark is a country of about 6 million people living in an area of about 50.000 km². In

Denmark, all eligible residents at a random sample of asylum centres were invited. The

Danish asylum seeker centres are run by the Danish Immigration Services but

outsourced to different operators. Several types of asylum seeker centres exist with the

most common being the accommodation centre, to which the asylum seeker is allocated

by random shortly after arrival and applying for asylum in Denmark. Of five available

operators, three responded to the invitation, all accepting to participate in the study,

representing a total of 11 accommodation asylum centres. Data-collection teams of two

to four healthcare professionals were instructed during several pre-collection sessions in

the practicalities and ethical considerations involved in the task: this encompassed

obtaining written and informed consent, appropriate handling of confidential

information, collection of the questionnaire data, the mental health self-rating scales,

clinical measurements and the biological samples. At consent, the participant was

instructed to independently complete a background questionnaire and mental health

scales. The data collectors assisted those who could not read or write or had further

questions regarding the study. Afterwards, the participant was invited to a semi-

structured clinical interview and a short clinical exam of basic parameters (blood

pressure, height, weight). Participants could at any time opt out of the data collection; if

choosing to do so the participant was asked if she would care to provide a reason. Also,

individuals who were invited but refused to participate were asked for a short non-

structured interview including basic demographics and reason for refusal. The

procedure was piloted in its entirety in Denmark including 10 participants in an asylum

seeker centre, resulting in minor adjustments to the workflow. All relevant permissions

were obtained prior to inclusion of any participant in both Lebanon and Denmark.

In the following we give a brief summary of the applied statistical methodology. Further

details are given in the Supplement and a thorough discussion of the statistical

methodological considerations necessary to combine multiple imputation and propensity

score-weighted analysis is provided in Eiset and Frydenberg [20]. Missing data was

multiply imputed and confounding was address by propensity score-weighting including

age, sex, socioeconomic status, experienced trauma and WHO-5-score in the model.

The "standardised mortality ratio"-weights [21] was computed to estimate the weighted

prevalence difference of PTSD in the population that fled to Denmark instead of

Lebanon and presented with a 95-percentile confidence interval found by bootstrapping.

A number of sensitivity analyses were carried out to assess the robustness of the

estimated association between long-distance migration and PTSD. All data

management, analysis and plots were done in R [22]. Further details are given in the

Supplement as well as directly in the R code available together with the *a priori*

described analysis plan and a number of the exploratory plots from

https://github.com/eiset/ARCH.

Results

A total of 599 individuals were included in Lebanon and 113 were included in Denmark.

Table 1 gives a statistical summary of the inclusion process and non-participation. The

study populations differed on key variables such as sex (female: 73% in Lebanon, 47%

in Denmark) and experience of violence (24% in Lebanon, 39% in Denmark) and there

were missing values for all variables except long-distance migration. Figure 1 illustrates

the missing proportion of key variables stratified on exposure status and Table 2 gives a

detailed summary of the study population characteristics for both the observed

realization and the imputed data set including missing for each covariates in the

observed data.

The unadjusted prevalence of PTSD was higher in Denmark (60.2%) compared with

Lebanon (55.1%). The prevalence difference of PTSD increased from 5.1 percentage

point (95-percentile CI [-4.6; 15.0]) to 8.8 percentage point (95-percentile CI [-1.4; 18.6

percentage point]) after multiply imputing missing data and adjusting for confounding

by propensity score-weighting. All sensitivity analyses produced estimates in the same

direction and of the same magnitude, except when forcing all missing in the "Violence"

variable to "Yes" in the study population included in Denmark, thus, grossly violating

the missing-at-random assumption of multiple imputation. Table 3 presents the point

estimate for the propensity score-weighted analysis and each of the sensitivity analysis

accompanied by Figure 2 which includes different types of bootstrap confidence

intervals.

Discussion

This study aimed at estimating the association between long-distance migration and

prevalence of PTSD. The estimate of association corresponded to 87 additional cases of

PTSD for every 1000 Syrian refugees that migrated to Denmark instead of Lebanon.

The 95% confidence interval indicate that, under the model, the estimate may be as high

as 190 additional cases or as low as 4 cases less per 1000 refugees, thus, the estimate

did not reach statistical significance, which may be ascribed to the relatively low

number of participants. The targeted sample size was not reached neither in Lebanon

nor in Denmark due to structural and political obstacles during the data collection.

To the best of the authors' knowledge this is the first investigation of the association of

long-distance migration and PTSD in a refugee population. A recent study [23] of

Syrian refugees in Lebanon and Norway found a prevalence of PTSD (assessed as

HTQ-score $\geq$ 2.5) of 4% in Lebanon and 7% in Norway. The study conflated

heterogeneous groups for example asylum-seekers and refugees and participants with

short and long stay in the host country prior to inclusion in the study. The study found

that "Host country, Norway" was associated with an average increase in HTQ-score of

3% (95% CI [-5%; 12%]) when adjusting for age and sex; however, did not provide

estimates of how this translated into absolute number of participants with PTSD. With a

mean HTQ-score of 1.49 in the study population in Lebanon the number of additional

cases in Norway due to "host country" would be marginal. The underlying cause for the

observed increase in PTSD prevalence in Syrian refugees and asylum seekers after long-

distance migration has never been investigated. In our study, we approach this by

minimized the effect of *living* in the host country, thus, "long-distance migration"

represented the migration process itself (travelling through Europe). We found a high

prevalence of PTSD and an increased prevalence difference compared with the

aforementioned study. Some studies hypothesise that great distance between populations

is associated with difficulty acculturating which in turn is associated with worse health

outcomes, including mental health [13,15,24]. The presented study is a first step in

examining this in a refugee population.

The prevalence estimates of PTSD was high in both Lebanon and Denmark compared

with previous studies of Syrian refugees and across different study designs and utilized

instruments for assessing PTSD [1,25–27]. In the present study, we utilized the widely

used HTQ-score [28] which is considered valid across multiple settings and diagnostic

classification systems [29] although the specific cut-off has been debated [30]. The

sensitivity analysis showed little change in the estimate of association including when

changing the HTQ-score cut-off. An effort was made to make the data collection in

Lebanon and Denmark as comparable as possible; however, differences may exist

introducing information bias. For example, differential misclassification cannot be ruled

out, if, during data collection in Denmark, doubt remained among the participants

whether specific answers would affect their asylum application. This may explain (some

of) the difference in prevalence of PTSD in Lebanon and Denmark; however, judged

from extensive exploratory plotting of the covariates, there was no difference between

the exposure groups in the association of the reported HTQ-score and WHO-5 (general

mental well-being) or depression- or anxiety-scores, thus, the participants would have

had to be very consistent in their misclassification of own mental health. The proportion

that refused to answer, or did not know what to answer, in the question on

socioeconomic status was high and differed between the study population in Denmark

(22%) and in Lebanon (7%). After discussing this variable with Syrian natives, we

speculate that "socioeconomic status" was not defined in sufficient detail for all

participants to be able to reflect on the question; however, we do not have an

explanation for the difference in proportion of missing between Lebanon and Denmark.

We consider the risk of selection bias low given the randomized sampling design and

very high participation rates. It was, however, not possible to retrieve or create a precise

sampling frame in Lebanon and thus, it was based on best information available at the

time, knowing that refugees frequently relocate and there is little registration of

individuals outside the formalized refugee camps.

In the present study, the interpretation of the association between long-distance

migration and PTSD is of course limited by the cross-sectional study design. Most

importantly, the time-order may be reversed so that instead of long-distance migration

leading to higher risk of PTSD it may be that PTSD affect the "risk" of undertaking

long-distance migration. A positive association between long-distance migration and

PTSD was found, thus, a reversal of the time order would mean that individuals with

PTSD are more likely to undertake long-distance migration. Although this cannot be

entirely rejected, taking into account the classical symptoms of PTSD such as

difficulties in concentrating and planning, this interpretation of the result seems far less

likely than our stated hypothesis: that undertaking an, at times, hazardous travel and

living with uncertainty about the near future, increases the risk of PTSD.

We addressed confounding by propensity score-weighting and obtained balance on all

covariates judged from our pre-specified criteria. To account for confounding we

included covariates in the propensity score model according to what the authors believe

is the current best evidence [7]; however, this does not alleviate us of speculations of

entirely other covariates that may change the estimate. Furthermore, the collapsing of

two levels of the socioeconomic status covariate diminishes the information this

variable contains and thus may be a source for residual confounding in both the multiple

imputation and in the propensity score weighted analysis. The missing mechanism was

"everywhere missing-at-random" allowing for frequentist inference from the multiply

imputed data. In the Supplement, we provide examples of the discussion necessary for

each partly observed variable to deem the missingness mechanism "everywhere

missing-at-random".

Whether the results presented in this paper is specific to Syrian refugees and host

countries Lebanon and Denmark must be examined in future studies. We note that the

direction of the association was consistent across different thresholds for PTSD

(effectively lowering and raising the estimated prevalence for PTSD in the study

population), different propensity score models and when imposing violation of

underlying assumptions of multiple imputation; the strength was also relatively

consistent although, forcing all missing values in "experienced violence" in the exposed

study population produced a relatively low prevalence difference estimate. We

hypothesize that the association will be reproducible at least in comparable populations

and settings—for example newly arrived adult refugees from a Middle Eastern country

to a Western European country. Also, the association between long-distance migration

and other health outcomes, for example other mental health issues, is of great interest.

The presented study is an initial analysis of the association between long-distance

migration and refugee health and provides the foundation upon which future studies

could build.

Tables

Table 1: Summary of the inclusion in Denmark and Lebanon

	Denmark n(%)	Lebanon n(%)
Source population	613[a]	1,011,366[a]
Examined for eligibility	124	641
Excluded	3	11
Invited	121 (100%)	630 (100%)
Refused to participate		
Total	8 (7%)	31 (5%)
Sex, female	2	5[b]
Age[c]	All between 20 and 30 years old	All younger than 50 years old
Reasons[d]		
Mistrust	5	1
No time	2	4
No personal gain	0	2
No answer	1	24
Included (participation rate)	113 (93%)	599 (95%)

[a]The number given for the source population represents a maximum and include for example children. It was not possible to obtain a more precise estimate of the study population. [b]Sex was only recorded for 8 non-participants in Lebanon. [c]Few non-participants provided age (two in Denmark and two in Lebanon), thus a best guess is provided. [d]Reasons provided from the non-participants, aggregated into smallest possible number of strata.

Table 2: Comparison of summary statistics of key variables in observed (with missing) and imputed data set stratified on exposure group

	Observed data		Imputed data	
	Lebanon (n=599)	Denmark (n=113)	Lebanon (n=599)	Denmark (n=113)
Age - years old	35 (20)	30 (12.5)	35 (20)	30 (12.9)
Missing	n=58 (9.7%)	n=10 (8.9%)	NA	NA
Sex				
Female	72.7%	46.9%	72.7%	46.9%
Missing	n=38 (6.3%)	n=0	NA	NA
WHO-5-score	24 (32)	24 (48)	24 (32)	24 (48)
Missing	n=47 (7.9%)	n=4 (3.5%)	NA	NA
Experienced violence				
Yes	23.8%	39.5%	23.8%	39.7%
Missing	n=37 (6.2%)	n=37 (32.7%)	NA	NA
Socioeconomic status				
Below average	25.5%	24.7%	25.3%	23.2%
On average	63.4%	50.7%	63.4%	50.0%
Above average[a]	4.5%	2.7%	11.3%[a]	26.8%[a]
Don't know/refuse to answer[a]	6.7%	21.9%		
Missing	n=61 (10.2%)	n=40 (35.4%)		
HTQ score	2.56 (0.94)	2.62 (1)	2.56 (0.95)	2.62 (1)
Missing	n=33 (5.5%)	n=0	NA	NA
PTSD (HTQ score ≥2.5)				
Overall	55.1%	60.2%	54.9%	60.2%
Female	59.4%	66.0%	59.1%	66.0%
Male	44.7%	55.0%	43.7%	55.0%
Missing	n=33 (5.5%)	n=0	NA	NA

Categorical variables are represented as % of observations. Continuous variables are

represented as median (interquartile range). Missing is given as n (%) of all included

participants for the observed data set. [a]In the substantive model, and thus in the multiple

imputation, the strata "Above average" and "Don't know/refuse to answer" was

collapsed due to numerical problems in the multiple imputation.

Table 3: Estimated prevalence difference in crude, propensity score analysis and

sensitivity analysis

	Prevalence difference; point estimate [95% CI][a]
Multiply imputed and PS-weighted analysis	8.8 [-1.4; 18.6][b]
Complete case analyes	
Crude	5.1 [-4.6; 15.0][b]
PS-weighted	5.1 [-5.0; 16.1][b]
Sensitivity analysisc	
Alternative PS model	8.0 [-1.4; 17.8]
HTQ score threshold for PTSD: 2.3	10.6 [0.6; 18.3]
HTQ score threshold for PTSD: 2.7	7.2 [-3.1; 16.7]
Missing in "SES" in Lebanon are forced to "Don't know/refuse to answer"	9.5 [-0.1; 19.6]
Missing in "Violence" in Denmark are forced to "Yes"	4.2 [-5.4; 14.8]

272   Abbreviations: CI, confidence interval; HTQ, Harvard Trauma Questionnaire; PS,

273   propensity score; SES, socio-economic status. [a]All numbers are percentage point. The

274   95-percentile confidence interval is used for all estimates; [b]Based on 999 bootstrap

275   replications, otherwise based on 250 bootstrap replications. [c]Multiply imputed and PS-

276   weighted. Also see Figure 2.

Figures

Figure 1: Missing fraction plot.

Variables included in the propensity score and multiple imputation models for

estimating the association between long-distance migration and PTSD among Syrian

refugees in Lebanon and Denmark. Abbreviations: HTQ-IV, Harvard Trauma

Questionnaire part IV; WHO-5, WHO-5 Mental-health scale; Violence, exposure

(directly or indirectly) to violence; HSCL-anxeity, Hopkins Symptom Check List,

anxiety part; Bp.sys, systolic blood pressure.

Figure 2: Estimate with different bootstrap confidence interval types.

Abbreviations: BCa, bias-corrected and accelerated bootstrap CI; CI, confidence

interval; PS, propensity score; Cc, complete case; HTQ, Harvard Trauma Questionnaire

part IV; SES, socioeconomic status; Violence, exposure (directly or indirectly) to

violence. From the top: the propensity score-weighted ("substantive model") and the

two complete case (no imputation) estimates with 999 bootstrap replications. The five

sensitivity analysis are at the bottom, each with 250 bootstrap replications (and thus no

BCa 95% CI).

Author contributions

Conceptualization: AHE; Methodology: AHE, MF; Formal analysis and investigation: AHE, MF; Writing - original draft preparation: AHE; Writing - review and editing: AHE, MPA, MF, AGG, MS, WJN, RSH; Funding acquisition: AHE; Resources: AHE, MPA, WJN, RSH, AGG, MS; Supervision: MF, RSH, MS, AGG, WJN.

Acknowledgements

The authors would like to thank the following for their invaluable help in the data collection, in no particular order: Dr. Fayez Saadaldin, Margrethe Nielsen, Susanne Løgsted, Rebekka Pedersen, Sine Søe Romedahl, Hanne Jeppesen, Dr. Chahid Farah, Dr. Fadi Nader, Dr. Malak Rammal, Dr. Mayssaloun Khairallah, Dr. Sultana Baydoun, Dr. Tanios Dagher, Dr. Maibritt Provstgaard, Maja Nielsen, Dr. Arendse Lykke Loua, Dr. Cecilie Blenstrup Patsche, Dr. Bjørg Olsen, Dr. Camilla Kjersgaard, Hawa-Idil Harakow.

Data sharing

The individual-level data cannot be shared because it contains sensitive information and its distribution would violate the agreement under which the data was provided as well as Danish law. The computing code for all parts of the data cleaning, analysis, and plotting is publicly available from the first author's Github repository: https://github.com/eiset/ARCH. This repository also contains all the exploratory plots produced.

Ethics and research committee approval

All relevant permissions and approvals were obtained prior to inclusion of any participant in Lebanon and Denmark including from the Lebanese Ministry of Public Health (2018/4/38918), the Ethics Committee of Mount Lebanon Hospital (PSY-2018-005), the Ethics Committee of the Lebanese University (CUMEB/D 144/232018) and the Danish Data Protection Agency (J.nr. 2015-41-4500). The Danish National Committee on Health Research Ethics waived the right to assess the project (F 257/2015).

Declaration of interests

All authors declare no competing interests.

Funding

This research was supported by the Danish Institute in Damascus and The Graduate School of Health, Aarhus University, Denmark. The funding agencies had no role in the study design, data collection, data analysis, data interpretation, or writing of the report.

References

1. Blackmore R, Boyle JA, Fazel M, Ranasinha S, Gray KM, Fitzgerald G, et al. The prevalence of mental illness in refugees and asylum seekers: A systematic review and meta-analysis. PLOS Med. Public Library of Science; 2020;17:e1003337.

2. Gushulak BD, MacPherson DW. The basic principles of migration health: population mobility and gaps in disease prevalence. Emerg Themes Epidemiol. 2006;3:3–3.

3. Kristiansen M, Mygind A, Krasnik A. Health effects of migration. Dan Med Bull. 2007;54:46–7.

4. Spallek J, Zeeb H, Razum O. What do we have to know from migrants' past exposures to understand their health status? a life course approach. Emerg Themes Epidemiol. 2011;8:6.

5. Katrinli S, Stevens J, Wani AH, Lori A, Kilaru V, van Rooij SJH, et al. Evaluating the impact of trauma and PTSD on epigenetic prediction of lifespan and neural integrity. Neuropsychopharmacology. Nature Publishing Group; 2020;45:1609–16.

6. Scherrer JF, Salas J, Schnurr PP. Repeated Assessment of Posttraumatic Stress Disorder Severity and the Risk of Mortality. JAMA Netw Open. 2020;3:e1920493.

7. Shalev A, Liberzon I, Marmar C. Post-Traumatic Stress Disorder. N Engl J Med. 2017;376:2459–69.

8. Wisco BE, Marx BP, Miller MW, Wolf EJ, Krystal JH, Southwick SM, et al. A comparison of ICD-11 and DSM criteria for posttraumatic stress disorder in two national samples of U.S. military veterans. J Affect Disord. 2017;223:17–9.

9. Sareen J, Murray S, Michael F. Posttraumatic stress disorder in adults: Epidemiology, pathophysiology, clinical manifestations, course, assessment, and diagnosis [Internet]. UpToDate. [cited 2020 Oct 9]. Available from: https://www.uptodate.com

10. Keane TM, Marshall AD, Taft CT. Posttraumatic stress disorder: etiology, epidemiology, and treatment outcome. Annu Rev Clin Psychol. 2006;2:161–97.

11. Spolaore E, Wacziarg R. Ancestry, Language and Culture. Palgrave Handb Econ Lang. Palgrave Macmillan UK; 2016.

12. Wilding S, Martin D, Moon G. How far is a long distance? An assessment of the issue of scale in the relationship between limiting long-term illness and long-distance migration in England and Wales. Popul Space Place. 2018;24:e2090.

13. Detollenaere J, Baert S, Willems S. Association between cultural distance and migrant self-rated health. Eur J Health Econ. 2018;19:257–66.

14. Klein EM, Müller KW, Wölfling K, Dreier M, Ernst M, Beutel ME. The relationship between acculturation and mental health of 1st generation immigrant youth in a representative school survey: does gender matter? Child Adolesc Psychiatry Ment Health [Internet]. 2020 [cited 2021 Apr 30];14. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7368737/

15. Lara M, Gamboa C, Kahramanian MI, Morales LS, Hayes Bautista DE. Acculturation and Latino Health in the United States: A Review of the Literature and its Sociopolitical Context. Annu Rev Public Health. 2005;26:367–97.

16. Eiset AH, Aoun MP, Haddad RS, Naja WJ, Fuursted K, Nielsen HV, et al. Asylum seekers' and Refugees' Changing Health (ARCH) study protocol: an observational study in Lebanon and Denmark to assess health implications of long-distance migration on communicable and non-communicable diseases and mental health. BMJ Open. British Medical Journal Publishing Group; 2020;10:e034412.

17. United Nations Children's Fund (UNICEF), United Nations High Commissioner for Refugees (UNHCR) and the United Nations World Food Programme (WFP). Vulnerability Assessment for Syrian Refugees in Lebanon (VASyR-2018) [Internet]. 2018 Dec. Available from: https://www.unhcr.org/lb/wp-content/uploads/sites/16/2018/01/VASyR-2017.pdf

18. Sedgwick P. Stratified cluster sampling. BMJ. 2013;347:f7016.

19. Sharon L. Lohr. Sampling: Design and Analysis [Internet]. 2. Brooks/Cole, Cengage Learning; 2010 [cited 2019 Feb 27]. Available from: https://www.sharonlohr.com/sampling-design-and-analysis

20. Eiset AH, Frydenberg M. Considerations for using multiple imputation in propensity score-weighted analysis. medRxiv. Cold Spring Harbor Laboratory Press; 2021;2021.06.30.21259793.

21. Sato T, Matsuyama Y. Marginal structural models as a tool for standardization. Epidemiol Camb Mass. 2003;14:680–6.

22. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2020. Available from: http://www.R-project.org/

23. Strømme EM, Haj-Younes J, Hasha W, Fadnes LT, Kumar B, Igland J, et al. Health status and use of medication and their association with migration related exposures among Syrian refugees in Lebanon and Norway: a cross-sectional study. BMC Public Health. 2020;20:341.

24. Salant T, Lauderdale DS. Measuring culture: a critical review of acculturation and health in Asian immigrant populations. Soc Sci Med. 2003;57:71–90.

25. Hendrickx M, Woodward A, Fuhr DC, Sondorp E, Roberts B. The burden of mental disorders and access to mental health and psychosocial support services in Syria and among Syrian refugees in neighboring countries: a systematic review. J Public Health. Oxford Academic; 2020;42:e299–310.

26. Selmo P, Knaevelsrud C, Mohamad N, Rehm J. Prevalence and predictors of psychopathology in the war-afflicted Syrian population. Transcult Psychiatry [Internet]. SAGE PublicationsSage UK: London, England; 2020 [cited 2020 Nov 16]; Available from: https://journals.sagepub.com/doi/10.1177/1363461520937931?url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrossref.org&rfr_dat=cr_pub++0pubmed

27. Tinghög P, Malm A, Arwidson C, Sigvardsdotter E, Lundin A, Saboonchi F. Prevalence of mental ill health, traumas and postmigration stress among refugees from Syria resettled in Sweden after 2011: a population-based survey. BMJ Open [Internet]. 2017 [cited 2018 Oct 24];7. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5778338/

28. Sigvardsdotter E, Malm A, Tinghög P, Vaez M, Saboonchi F. Refugee trauma measurement: a review of existing checklists. Public Health Rev. 2016;37:10.

29. Tay AK, Mohsin M, Rees S, Steel Z, Tam N, Soares Z, et al. The factor structures and correlates of PTSD in post-conflict Timor-Leste: an analysis of the Harvard Trauma Questionnaire. BMC Psychiatry [Internet]. 2017 [cited 2020 Oct 9];17. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5441051/

30. Ichikawa M, Nakahara S, Wakai S. Cross-cultural use of the predetermined scale cutoff points in refugee mental health research. Soc Psychiatry Psychiatr Epidemiol. 2006;41:248–50.