Characterization of A Novel Autoimmune Encephalitis Associated Antibody against CRMP2

Kaibiao Xu*, Dongmei Wang*, Yan He*, Shengnan Wang*, Guanghui Liu, Yue Pan, Haishan Jiang, Yu Peng, Fenliang Xiao, Yihua Huang, Qiqi Wang, Yongming Wu[¶], Suyue Pan[¶] and Yafang Hu[¶]

Department of Neurology, Nanfang Hospital, Southern Medical University, Guangzhou, Guangdong, China.

* These authors contributed equally to this paper.

[¶] Corresponding authors:

Yafang Hu, Department of Neurology, Nanfang Hospital, 1838 Northern Guangzhou Avenue, Guangzhou 510515, China. Tel: +86 20 62786544; Fax: +86 20 62787664; E-mail: yafanghu@smu.edu.cn.

Suyue Pan, Department of Neurology, Nanfang Hospital, 1838 Northern Guangzhou Avenue, Guangzhou 510515, China. Tel: +86 20 62786544; Fax: +86 20 61641961; E-mail: pansuyue@126.com.

Yongming Wu, Department of Neurology, Nanfang Hospital, 1838 Northern Guangzhou Avenue, Guangzhou 510515, China. Tel: +86 20 62787664; Fax: +86 20 62787664; E-mail: 13711530187@139.com.

Subtitle: anti-CRMP2 Autoimmune Encephalitis

ABSTRACT

Objective

Autoimmune encephalitis (AE) is a large category disorder urging antibody characterization. The aim was to identify a novel AE related autoantibody targeting an intracellular synaptic protein.

Methods

Suspected AE patients had negative conventional antibodies screening but strong immunolabel signals on rat brain sections with the serum and cerebrospinal fluid (CSF) samples were considered burdening unkown antibody. Immunoprecipitation from the rat brain protein lysate followed by mass spectrometry analysis was used to identify the targeting antigen. Western blotting and/or cell-based assay (CBA) with antigen-overexpressing HEK293T cells were used for antibody specificity, epitope and IgG subtype determination. Patients with similar immunostaining pattern on rat brain sections were retrospectively screened for the antibody.

Results

The antibody against collapsin response mediator protein 2 (CRMP2), a synaptic protein involved in axon guidance, was identified in a patient with suspected AE. The patient samples reactivated with HEK293T cell overexpressed CRMP2, rather than CRMP1, 3, 4, and 5. The patient samples mainly stained neuronal cytoplasm of the cortex, hippocampus and cerebellum Purkinje cells. This

reactivity was eliminated by pre-immunoabsorption with CRMP2-overexpressing

HEK293T cells. CRMP2 truncation experiments indicated that 536 amino acids at

C-terminus was necessary for the epitope. Subtype analysis showed that

anti-CRMP2 antibody was IgG4. Moreover, screening from 19 suspected AE

patients led to identification of anti-CRMP2 antibody in another patient with a

diagnosis of encephalomyelitis. The two patients responded to immunotherapy.

Conclusions

This study discovered a novel anti-CRMP2 antibody associated with AE. Testing

of the antibody might be promising for AE diagnosis and treatment.

KEYWORDS: auto**immune encephalitis, autoimmune encephalomyelitis**,

CRMP2

INTRODUCTION

Autoimmune encephalitis (AE) is a large category of inflammatory disorders

mediated by immune responses against neuronal intracellular antigens, cell-

surface or synaptic antigens, in some cases, accompanied by neoplasia.

Intracellular antigens include Hu (anti-neuronal nuclear antibody type 1, ANNA1) ,

Ri (ANNA2), and Ma2 in the nucleus, Yo

(Purkinje cell cytoplasmic antibody type 1, PCA1), amphiphysin, glutamate

decarboxylase 65 kDa isoform (GAD65), and Kelch-like Protein 11 in the cytosol,

which cause T cell mediated immune response and some cases respond to

immunotherapy [1-4]. Most AE caused by autoantibodies against antigens on the

neuronal surface or synaptic proteins, such as N-methyl-D-aspartate receptor

(NMDAR), leucine-rich gliomainactivated 1 (LGI1), contactin-associated protein 2

(Caspr2), gamma-aminobutyric acid (GABA) receptors (A/B),

alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic acid (AMPA) receptors,

dipeptidyl-peptidase-like protein-6 (DPPX) , delta/notch-like epidermal growth

factor-related receptor (DNER), dopamine-2 receptor (D2R), metabotropic

glutamate receptor 5 (mGluR5), voltage-gated calcium channel salpha-2/delta

subunit ($Ca_V\alpha2\delta$), and glutamate kainate receptor subunit 2 (GluK2), respond well

to immunotherapy[1, 5-9]. Thus, intensive testing of these antibodies in suspected

AE patients plays an important role in guiding the diagnosis and treatment of the

disease. Despite this, the cause of many cases of AE remains unexplained

because of limited known antibodies[10-12]. Searching for additional antibodies has

been desperately needed.

Family of collapsin response mediator proteins (CRMPs) consists five

homologous members, CRMP1-5, which are cytosolic microtubule-associated

phosphoproteins and play important roles in dendrite and axonal guidance,

regulating migration and synaptic dynamics [13]. Structurally, CRMP1-4 share

69-76% amino acids (aa) identity, while they share less identity to CRMP5,

approximately 50% [14]. Antibodies against CRMPs except CRMP2 have been

reported in several encephalitis. Previously, eight cases of anti-CV2/CRMP5 sera

from patients with autoimmune myelopathy were detected anti-CRMP3 antibody

positive, and one of them also had anti-CRMP1 antibody, one had anti-CRMP1, 4

antibodies[14-15]. Antibodies to CRMP3-4 were found in a case with subacute limbic

encephalitis and thymoma[16].

CRMPs antibodies also have been found in other disorders. Anti-CRMP1/2

antibodies are found in autoimmune retinopathy patients[17]. Antibodies to

CRMP1/2 in maternal sera increased the occurrence of autism spectrum

disorders in children[18]. Presence of antibodies to CRMP2 and/or GFAP in the

acute phase of spinal cord injury increased the occurrence of neuropathic pain six

months later [19]. Here, we found CRMP2 as the specific autoantigen in a patient

with intracranial *Mycoplasma pneumoniae* (*M.P.*) infection and possible

secondary immune-mediated encephalitis.

MATERIALS AND METHODS

Patients and autoantibody testing

This study was approved by the Ethical Review Committee of Nanfang hospital,

the Southern Medical University, Guangzhou, China. All patients or their families

signed the informed consents.

Patients with suspected autoimmune encephalitis were hospitalized in our

Neurology department between January 1, 2018 and December 31, 2020.

Patients' sera or CSF samples were screened for known autoantibodies, including

anti-NMDAR, LGI1, Caspr2, $GABA_BR$, $AMPA_{A/B}R$, DPPX, DNER, D2R, mGluR5,

GAD65 and IgLON5 antibodies by cell-based assay (CBA), some patients tested

anti-Hu, Yo, Ri, Ma2, PCA2, amphiphysin, and CRMP5 antibodies by blot assay

according to reference[1]. The negative samples were further analyzed by

tissue-based assay (TBA) of immunostaining on rat brain sections. CBA negative

and TBA positive samples were considered with unknown antibodies and applied

to antibody discovery process.

Immunohistochemistry (IHC)

5µm thick frozen tissue sections from the brain of adult SD rat or C57BL/6 mouse

were prepared and performed IHC (or TBA) with CSF, sera, and secondary

antibodies as previously reported[20]. Dilutions were used as CSF, 1:1, sera, 1:200,

HRP-labeled goat anti-human IgG secondary antibody (bs-0297G-HRP, Bioss,

Beijing, China), 1:3000, DyLight 488-labeled goat anti-human IgG secondary

antibody (ab97003, Abcam, Cambridge, MA), 1:200. Immunostaining images

were photographed under DM3000 microscope (Leica, Wetzlar, Germany) or

LSM980 confocal microscope (Zeiss, Oberkochen, Germany). For the absorption

assay, the serum diluted 1:200 was pre-immunoabsorbed six times by incubated

with acetone fixed CRMP2-overexpressing HEK293T cells for 1 hour at 37□ as

reported[6].

Western blotting, immunoprecipitation (IP) and liquid chromatography

tandem mass spectrometry (LC-MS/MS)

Fresh whole brain protein lysate of adult SD rats and Western blotting were

conducted as reported [20]. The lysis buffer (20 mmol/l Tris-HCl, pH 7.4, 150 mmol/l,

1% TritonX-100, 2 mmol/l EDTA, 5% glyserol, 1 mmol/l PMSF) was used. The

protein lysate was mixed with the patient's CSF at 4□ overnight, followed by

adding protein A agarose beads and incubation for another 2 hours. The beads

were washed and collected by centrifugation, then boiled with SDS-PAGE sample

buffer. The antibody-protein complexes pulled down were separated by 10% SDS

gel electrophoresis, transferred to nitrocellulose membrane for Western blotting or

stained with Coomassie brilliant blue. Correlated band compared with Western

blotting results from the gels was cut and sent for protein identification using

LC-MS/MS analysis by Fitgene Biotech Ltd. (Guangzhou, China).

Cell culture and immunofluorescence (IF)

Primary cortical neurons were prepared from 18 to 19 days C57BL/6 mouse

fetuses and cultured for 14 days as reported[21]. Ice acetone fixed neurons were

blocked with 10% goat serum at 37°C for 30 minutes and incubated with patient'

serum (1:200), rabbit anti-MAP2 antibody (1:200, 8707T, Cell Signaling

Technology, Danvers, MA) or rabbit anti-CRMP2 antibody (1:200, ab129082,

Abcam) at 4°C overnight. DyLight 488-labeled goat anti-human IgG and Alexa

Fluor 594-labeled goat anti-rabbit IgG (ab150092, Abcam) secondary antibodies

(1:200) were used and fluorescence images were photographed under LSM980

confocal microscope (Zeiss).

HEK293T cells were transfected with plasmids

pcDNA3.1-CRMP1/2/3/4-Flag, pcDNA3.1-CRMP5-eGFP, pcDNA3.1-CRMP2 T1

(1-141 amino acids of isoform 1) / T2 (1-36 amino acids of isoform 2)-eGFP, and

pcDNA3.1-CRMP2 T3 (142-677 amino acids of isoform 1)-Flag separately for 48

hours. Protein coding sequences in the plasmids referred to mRNA of CRMP1:

NM_001014809.3, CRMP2 (isoform 1): NM_001197293.3, CRMP2 (isoform 2):

NM_001386.6, CRMP2 (isoform 3): NM_001244604.2, CRMP3: NM_006426.3,

CRMP4: NM_001197294.2, and CRMP5: NM_020134.4. Full length isoform 1 of

CRMP2 was used unless illustrated. Transfected cells were fixed with ice acetone

for 10 minutes, washed with PBST, and blocked with 10% goat serum, then

incubated with patient's serum or rabbit anti-CRMP2 antibody (1:200) overnight at

4°C. After wash, the appropriate secondary antibodies were incubated at room

temperature for 1 hour, including DyLight 488-labeled goat anti-human IgG,

DyLight 550-labeled goat anti-human IgG (ab96908, Abcam), FITC-labeled

anti-human IgG1/2/3/4 (F0767/F4516/F4641/F9890, Sigma-Aldrich) and/or Alexa

Fluor 594-labeled goat anti-rabbit IgG secondary antibodies (1:200). The

fluorescence images were taken by using LSM980 confocal microscope (Zeiss) or

IX73 inverted microscope (Olympus, Tokyo, Japan).

RESULTS

Identification of an autoantibody specific targeting CRMP2 in a patient with

suspected AE

The index patient (P1) was diagnosed as intracranial *M.P.* infection and possible

secondary immune-mediated encephalitis. The known autoantibodies tests were

negative. However, IHC of P1's serum showed positive staining across the brain

(Fig. 1A). Immunofluorescence assay revealed neuronal staining in cortex,

hippocampus, and especially Purkinje cells of cerebellum (Fig. 1B). Further

analysis of P1's serum showed reaction with the cytosol compartments of cultured

cortical neurons (Fig. 1C).

Western blotting after IP revealed a band around 70 kDa, which was pulled

down by P1's CSF from rat brain protein lysate (Fig. 1D). The correlated visible

band from the Coomassie brilliant blue staining gel (Fig. 1E) was elucidated and

CRMP2 was identified as the possible target antigen by mass spectrometry (Fig.

1F, Table S1 in the supplement).

Verification of the anti-CRMP2 antibody in suspected AE patients

Next, we confirmed that both P1's serum and commercial anti-CRMP2 antibody

detected a band near 70 kDa from the pulled-down protein complex by the P1's

serum from rat brain protein lysate (Fig. 2A). With the CRMP2-overexpressing

HEK293T cells, CRMP2 were pulled down by commercial anti-CRMP2 antibody

and were recognized by P1's serum (Fig. 2B). There are three isoforms of

CRMP2 , isoform 1 (677 amino acids, 74kDa), isoform 2 (572 amino acids,

62kDa), and isoform 3 (536 amino acids, 58kDa), attribute to three transcript

variants, and the observed molecular weight of CRMP2 bands in Western blotting

might vary or be higher because of post-translational modification. Here we used

the human full-length isoform 1 CRMP2 for further tests unless illustrated.

Anti-CRMP2 antibody and P1's serum immunostained the overexpressed

CRMP2 in HEK293T cells and had well co-localization (Fig. 2C). As expected,

immunostaining of cultured neurons with P1's serum also co-localized with

anti-CRMP2 antibody signals (Fig. 2D). When P1's serum was

pre-immunoabsorbed with the CRMP2-overexpressing HEK293T cells to rule out

anti-CRMP2 antibody, a negative immunoreaction was found on the mouse brain

section. It indicates that no other neuronal antibodies presented (Fig.1B).

Further, 19 suspected AE patients with negative known antibodies testing

but similar immunostaining on rat brain sections were screened for anti-CRMP2

antibody by the CBA method. As indicated in Fig. 2C, another patient, P2, was

found having immunoreaction with CRMP2 in the serum. Moreover, the

immunoreaction was disappeared in P2's serum collected when the patient

recovered completely and was reevaluated 1.5 years later. All the data verify that

anti-CRMP2 antibody present in certain AE patients.

The antibody is IgG4 and specific to the C-terminus of CRMP2

Since CRMP2 belongs to the highly conserved CRMPs family now with five

members (Fig. 3A), we further determined whether the antibodies in our patients

are specific to CRMP2. Expression plasmids were constructed to express

full-length CRMP1-5. P1' and P2' sera showed immunoreaction only in the

HEK293T cells expression of CRMP2, rather than other CRMPs (Fig. 3B).

To further determine which part of CRMP2 responsible for the

antigen-antibody reaction, three expression plasmids (T1-3) carrying truncated

CRMP2 were constructed as schematically indicated and applied to CBA assay

(Fig. 3C). T1 contained 141 amino acids N-terminus of isoform 1, T2, 36

N-terminus of isoform 2, and T3, 536 C-terminus which shared by three CRMP2

isoforms. The results indicate that the 536 aa C-terminus of CRMP2 is necessary

for the immunoreaction (Fig. 3D).

We also checked the IgG subtype of anti-CRMP2 antibody in the patients. As

shown in Fig. 3E, only IgG4 mediated anti-CRMP2 immunoreaction. Same results

obtained with P2's serum (data not shown). The results suggest that anti-CRMP2

IgG4 recognize the C-terminus of CRMP2 in our AE patients.

Clinical features of the two patients with anti-CRMP2 antibody

P1 patient, a young adult female, was admitted due to persistent dizziness, visual

rotation, recurrent nausea and vomiting and fever for 8 days, and unstable

walking and blurred speech for one day. Fever was mild and the highest body

temperature was 38.5□. Physical examinations revealed signs of cerebellum

including poetry-like language, nystagmus, and bilateral finger-nose test

instability. Slight neck stiffness was noticed.

CSF examination revealed a normal opening pressure of 148mmH$_2$O with

only slightly elevated white blood cell count of 38 cells/μl (90% monocytes).

Screenings of paraneoplastic antigens and serological autoantibodies including

anti-nuclear, anti-thyroid peroxidase and anti-thyroglobulin antibodies were all

negative. Serum anti-*M.P.* IgM tests were positive twice in one week after

admission, and weak positive two weeks later. Serum anti-Widal's O test was

positive with a titer of 1:80 on admission and conversed to be negative two weeks

later. Next generation sequencing (Vision Medicals, Guangzhou, China) of CSF

sample detected 64 unique reads against *M.P.*. CSF test was negative for AE

antibodies, but was positive by TBA.

During hospitalization, the patient's symptoms deteriorated with recurrent

nausea and vomiting, also her mental status was changed to be apathy. Physical

examination showed opsoclonus-myoclonus involving her head and both arms.

She was diagnosed as intracranial *M.P.* infection and possible secondary

immune-mediated encephalitis. Azithromycin, doxycycline, intravenous

methylprednisone (MP, 40mg/day) and intravenous immunoglobin (IVIG, 0.4

g/kg/d) were prescribed. Levodopa and clonazepam were also given to control

the myoclonus. With the treatment, the patient's symptoms improved with mild

dizziness and opsoclonus. Cranial MRI was repeated and the result was not

remarkable with mild white matter degeneration (Fig. 4A-F). She was discharged

to a local hospital for rehabilitation. During the follow-up study, she recovered

partially with dizziness and a modified Rankin Scale (mRS) score of 2 at 3

months.

P2 patient, a young adult female, previously healthy, was admitted due to

chest pain, right arm fatigue for 10 days and headache, nausea, vomiting,

numbness in both legs for 6 days. Physical examination was not remarkable

except for hyperalgesia in the left chest and abdomen, decreased needle tingling

on the back of the left hand, and numbness on the feet. Chest CT and

electrocardiogram (ECG) were not remarkable. Spine MRI showed long-segment

spinal cord lesions from medulla to C6 segment (Fig. 4J-L). Lumbar puncture

revealed an opening pressure of 185 mmH$_2$O, with moderately elevated WBC of

240 cells/µl (90% monocytes), decreased glucose of 2.64 mmol/l, normal chloride

level of 123 mmol/l, and elevated protein level of 0.80 g/l. CSF pathogen

screenings for tuberculosis (TB) culture, TB-SPOT, X-pert, bacteria and fungi

cultures were all negative. Antibodies including autoimmune encephalitis panels,

anti-aquaporin 4 (AQP4), anti-myelin oligodendrocyte glycoprotein (MOG) were

all negative. Cranial MRI with contrast revealed multiple patchy abnormal signals

in the white matter area of the bilateral cerebral hemispheres and brain stem (Fig.

4G-I). TBA was negative for CSF, and positive for serum.

Possible tuberculous encephalomyelitis and acute demyelinating

encephalomyelitis (ADEM) were considered. The patient was given anti-TB

regimen (Rifampicin 0.45 g qd, Isoniazid 0.3 g qd, Pyrazinamide 0.5 g tid,

Ethambutol 0.75 g qd) and intravenous MP pulse therapy (1 g/day for 5 days, and

tapered to oral 48 mg/d). With the treatment, the patient's chest pain and

numbness were improved gradually. Lumber puncture was repeated 20 days later

and revealed that the WBC decreased dramatically to 20 cells/µl, the glucose and

protein returned to normal. She was followed up regularly in our clinic and the

steroids was tapered. Anti-TB therapy was discontinued after 6 month's

treatment. Repeated cranial MRI revealed that the abnormal signals in the white

matter area of the bilateral cerebral hemispheres and brain stem decreased

dramatically (Fig. 4M-O). Cervical MRI revealed that the abnormal signals of her

spinal cord disappeared (Fig. 4P-R).

The clinical features of the two patients were summarized in Table 1.

DISCUSSION

In this study, we identified IgG4-autoantibody specifically against CRMP2 in the sera and CSF samples of two AE patients with acute disease onset. Two patients had various symptoms. P1 presented with fever, nausea, vomiting, apathy, opsoclonus-myoclonus of head and arm, cerebellar signs including dizziness, slurred speech, and unstable walking. P2 had headache, nausea, vomiting, chest pain, right arm fatigue numbness in both legs. Both patients showed increased number of cells in CSF, and P2 had low glucose and increased level of protein. P1's CSF had unique reads against *M.P.*. MRI images showed abnormal signals in the white matter in the two patients and long segment lesions on the spinal cord in P2. P1 was first diagnosed as *M.P.* infection. However, treatment with antibiotics only did not improve the symptoms. TBA immunostaining strongly indicated a secondary AE. Indeed, in additional of immunotherapy achieved good improvement at discharge and had 2 score of mRS at 3 months. P2 was diagnosis as possible TB & ADEM. 20 days treatment with anti-TB and immunotherapy reached good outcome for the patient at discharge and had 0 score of mRS at 3 months. Based on the study, we may correct the diagnosis as possible TB and autoimmune encephalomyelitis. No neoplasm was found in these patients currently and will be continuously follow up in the future. Our study provide evidence for the diagnosis of AE, expanding anti-CRMP2 antibody to the AE spectrum and suggesting to test anti-CRMP2 antibody in suspected AE patients.

CRMPs highly express during embryonic development and decreases

significantly after birth, while each CRMP has a different subcellular localization

[22]. In adult mouse brain, CRMP2 is expressed in the soma, axons, and dendrites

of neurons from the cortex, hippocampus, cerebellum, immature and mature

oligodendrocytes [23, 24]. It is thought that antibodies against intracellular antigen

mediate T cell immunoreaction and impair cell function in the related area, while

the antibodies themselves will not cause a pathogenic antibody-antigen

interaction [25-27]. Therefore, symptoms of the two patients are consistent with the

subcellular distribution of CRMP2 in the brain, or probably in the spinal region.

Indeed, sera /and CSF from the two patients with anti-CRMP2 antibody had

robust staining of the cortex, CA3 region of hippocampus and Purkinje cells of

cerebellum.

M.P. encephalitis is one of the most severe extra-respiratory complications,

which accounts for 5-30% of all of the reported encephalitis cases[28]. The latency

period between the onset of respiratory symptoms and the development of

neurological abnormalities varies. In some cases, there was not a previous,

clinically evident respiratory episode at all [29]. P1 had definite intracranial infection

of *M.P.* and early onset of encephalitis evidenced by the presence of IgM antibody

against mycoplasma in serum and presence of *M.P.* DNA in CSF. Presence of

antibody to CRMP2 suggests that P1 had an immediately immune-mediated CNS

inflammation, which was also evidenced by the solely treatment with antibiotics

deteriorated the patient's symptoms. However, additional of IVIG and MP

treatment improved her symptoms gradually. We considered that the symptoms

of the patient were more attributed to the secondary inflammation caused by the

M.P. infection. Anti-ganglioside and galactocerebroside antibodies have been

found in association with mycoplasma neurological disorders. Thus, these studies

and ours provide direct evidence that infection of *M.P.* cause secondary

immune-mediated inflammation [30].

Infections by viral and bacterial pathogens are suspected to initiate a broad

range of neurological inflammatory disorders particularly AE [31]. Herpes simplex

virus (HSV) has reported to trigger some types of AE frequently more than others,

including anti-NMDAR, $GABA_AR$, mGluR5, Neure-xin3a, and D2R antibodies.

Among of them, near 64% AE patients post-HSV had anti-NMDAR antibody in

their serum or CSF, while the remaining had antibodies unidentified. TB was

associated with a lot of autoimmune disorders[32, 33]. Whether TB is a trigger of

anti-CRMP2 encephalitis in P2 patient, more studies are warranted.

Each CRMP has distinct neuronal function. *Crmp2* gene knockout mouse

shows that lack of CRMP2 leads to defects in axon guidance, axon pruning in

hippocampus and visual cortex and altered dendritic spine remodeling[34]. The

C-terminal region of CRMP2 involves in the post-translational modifications

including phosphorylation, SUMOylation, oxidation, and O-GlcNAcylation. Several

critical phosphorylation sites locate in the C-terminus[35-37]. The phosphorylated

CRMP2 inhibits its interaction with tubulin heterodimers and augments interaction

with NaV1.7[38]. CRMP2 binds to CaV2.2 and increases Ca^{2+} level on the cell

surface, interacts with several receptors including NMDARs, Kainate, and NCX

(Na^+/Ca^{2+} exchanger 3) to regulate Ca^{2+} and Na^+ neuronal homeostasis[39-40].

Although immunoreaction of anti-CRMP2 in related neurons is supposed not a

direct causative reason, it might lead to dysregulation of other receptors. Epitope

studies of CRMP2 indicated that C-terminus is responsive for the antibody

binding, implying that antigen-antibody reaction may affect the function of

CRMP2. The pathogenic properties of CRMP2 antibody should be clarified by

more studies.

The present study consists small samples data from a single clinic center,

thus, the detail clinical features and information of immunotherapy of anti-CRMP2

AE will be obtained in a large cohort study in the future. Our study characterized

anti-CRMP2 antibody in two AE patients, who responded well to immunotherapy.

We suggest to test anti-CRMP2 antibody in suggested AE patients.

ARTICAL INFORMATION

Contributors

Y.F.H., S.Y.P., Y.M.W. and K.B.X. designed the study and drafted the manuscript;

D.M.W. and S.N.W. took care of the index patient and responsible for the clinical

data preparation; Y.H. and G.H.L. did most experiments; Y.P. provided cultured

cells and neurons; H.S.J. and Y.P. were responsible for the other anti-CRMP2

antibody positive patients; F.L.X. prepared mouse and rat brain sections; Y.H.H.

took confocal images; Q.Q.W. did Western blotting; all the authors reviewed the

manuscript.

Funding

This work was supported by the National Natural Science Foundation of China

(81771225 to Y.F.H., 82071484 to Y.M.W.), Guangdong Provincial Scientific and

Technologic Progression Fund (2016A020215182 to Y.F.H.), Natural Science

Foundation of Guangdong Province (2019A1515011760 to Y.M.W.), President

Foundation of Nanfang Hospital (2019B007 to D.M.W., 2020B008 to K.B.X.),

Medical Science and Technology Foundation of Guangdong Province (A2021151

to K.B.X.).

Competing interests

A patent (202110461611.6) is pending for the assay to detect the anti-CRMP2

antibody in autoimmune encephalitis (S.Y.P., Y.F.H., S.N.W., G.H.L., D.M.W.,

K.B.X., Y.M.W.). No other disclosures to report.

Patient consent for publication

The two patients with anti-CRMP2 antibody and clinical features presented signed

the consents for publication.

Ethics approval

This study was approved by the Ethical Review Committee of Nanfang hospital,

the Southern Medical University, Guangzhou, China. All patients or their families

signed the informed consents.

Data availability statement

Data are available upon reasonable request. Anonymised data used in the

present study may be available upon reasonable request to the corresponding

author Y.F.H..

REFERENCES

1.   Graus F, Titulaer MJ, Balu R, et al. A clinical approach to diagnosis of

autoimmune encephalitis. *Lancet Neurol*. 2016;15(4):391-404.

doi:10.1016/S1474-4422(15)00401-9.

2.   Furneaux HM, Dropcho EJ, Barbut D, et al. Characterization of a cDNA

encoding a 34-kDa Purkinje neuron protein recognized by sera from patients with

paraneoplastic cerebellar degeneration. *Proc Natl Acad Sci U S A*.

1989;86(8):2873-2877. doi:10.1073/pnas.86.8.2873.

3.   Dropcho EJ. Antiamphiphysin antibodies with small-cell lung carcinoma and

paraneoplastic encephalomyelitis. *Ann Neurol*. 1996;39(5):659-667.

doi:10.1002/ana.410390516.

4.   Mandel-Brehm C, Dubey D, Kryzer TJ, et al. Kelch-like Protein 11 Antibodies

in Seminoma-Associated Paraneoplastic Encephalitis. *N Engl J Med*.

2019;381(1):47-54. doi:10.1056/NEJMoa1816721.

5.   de Graaff E, Maat P, Hulsenboom E, et al. Identification of delta/notch-like

epidermal growth factor-related receptor as the Tr antigen in paraneoplastic

cerebellar degeneration. *Ann Neurol*. 2012;71(6):815-824.

doi:10.1002/ana.23550.

6.   Sabater L, Gaig C, Gelpi E, et al. A novel non-rapid-eye movement and

rapid-eye-movement parasomnia with sleep breathing disorder associated with

antibodies to IgLON5: a case series, characterisation of the antigen, and

post-mortem study [published correction appears in Lancet Neurol. 2015

Jan;14(1):28]. *Lancet Neurol.* 2014;13(6):575-586.

doi:10.1016/S1474-4422(14)70051-1.

7. Lee ST, Lee BJ, Bae JY, et al. CaV α2δ Autoimmune Encephalitis: A Novel

Antibody and its Characteristics. *Ann Neurol.* 2021;89(4):740-752.

doi:10.1002/ana.26017.

8. Landa J, Guasp M, Míguez-Cabello F, et al. Encephalitis with Autoantibodies

against the Glutamate Kainate Receptors GluK2. *Ann Neurol.*

2021;90(1):101-117. doi:10.1002/ana.26098.

9. Parratt KL, Allan M, Lewis SJ, Dalmau J, Halmagyi GM, Spies JM. Acute

psychiatric illness in a young woman: an unusual form of encephalitis. *Med J*

Aust. 2009;191(5):284-286. doi:10.5694/j.1326-5377.2009.tb02787.x.

10. Granerod J, Tam CC, Crowcroft NS, Davies NW, Borchert M, Thomas SL.

Challenge of the unknown. A systematic review of acute encephalitis in

non-outbreak situations. *Neurology.* 2010;75(10):924-932.

doi:10.1212/WNL.0b013e3181f11d65.

11. Bhagavati S. Autoimmune Disorders of the Nervous System:

Pathophysiology, Clinical Features, and Therapy. *Front Neurol.* 2021;12:664664.

Published 2021 Apr 14. doi:10.3389/fneur.2021.664664.

12. Goodfellow JA, Mackay GA. Autoimmune encephalitis. *J R Coll Physicians*

Edinb. 2019;49(4):287-294. doi:10.4997/JRCPE.2019.407.

13. Stratton H, Boinon L, Moutal A, Khanna R. Coordinating Synaptic Signaling

with CRMP2. *Int J Biochem Cell Biol*. 2020;124:105759.

doi:10.1016/j.biocel.2020.105759.

14. Keegan BM, Pittock SJ, Lennon VA. Autoimmune myelopathy associated with

collapsin response-mediator protein-5 immunoglobulin G. *Ann Neurol*.

2008;63(4):531-534. doi:10.1002/ana.21324.

15. Honnorat J, Byk T, Kusters I, et al. Ulip/CRMP proteins are recognized by

autoantibodies in paraneoplastic neurological syndromes. *Eur J Neurosci*.

1999;11(12):4226-4232. doi:10.1046/j.1460-9568.1999.00864.x.

16. Gonzalez-Quintela A, Alende R, Gude F, et al. Serum levels of

immunoglobulins (IgG, IgA, IgM) in a general adult population and their

relationship with alcohol consumption, smoking and common metabolic

abnormalities. *Clin Exp Immunol*. 2008;151(1):42-50.

doi:10.1111/j.1365-2249.2007.03545.x.

17. Adamus G, Bonnah R, Brown L, David L. Detection of autoantibodies against

heat shock proteins and collapsin response mediator proteins in autoimmune

retinopathy. *BMC Ophthalmol*. 2013;13:48. Published 2013 Sep 25.

doi:10.1186/1471-2415-13-48.

18. Braunschweig D, Krakowiak P, Duncanson P, et al. Autism-specific maternal

autoantibodies recognize critical proteins in developing brain. *Transl Psychiatry*.

2013;3(7):e277. Published 2013 Jul 9. doi:10.1038/tp.2013.50.

19. Hergenroeder GW, Redell JB, Choi HA, et al. Increased Levels of Circulating

Glial Fibrillary Acidic Protein and Collapsin Response Mediator Protein-2

Autoantibodies in the Acute Stage of Spinal Cord Injury Predict the Subsequent

Development of Neuropathic Pain. *J Neurotrauma*. 2018;35(21):2530-2539.

doi:10.1089/neu.2018.5675.

20. Song P, Guan Y, Chen X, et al. Frameshift mutation of *Timm8a1* gene in

mouse leads to an abnormal mitochondrial structure in the brain, correlating with

hearing and memory impairment [published online ahead of print, 2020 Aug 20]. *J

Med Genet*. 2020;jmedgenet-2020-106925.

doi:10.1136/jmedgenet-2020-106925.

21. Facci L, Skaper SD. Culture of rodent cortical and hippocampal

neurons. *Methods Mol Biol*. 2012;846:49-56. doi:10.1007/978-1-61779-536-7_5.

22. Moutal A, White KA, Chefdeville A, et al. Dysregulation of CRMP2

Post-Translational Modifications Drive Its Pathological Functions. *Mol Neurobiol*.

2019;56(10):6736-6755. doi:10.1007/s12035-019-1568-4.

23. Ricard D, Stankoff B, Bagnard D, et al. Differential expression of collapsin

response mediator proteins (CRMP/ULIP) in subsets of oligodendrocytes in the

postnatal rodent brain. *Mol Cell Neurosci*. 2000;16(4):324-337.

doi:10.1006/mcne.2000.0888.

24. Bretin S, Reibel S, Charrier E, et al. Differential expression of CRMP1,

CRMP2A, CRMP2B, and CRMP5 in axons or dendrites of distinct neurons in the

mouse brain. *J Comp Neurol*. 2005;486(1):1-17. doi:10.1002/cne.20465.

25. Voltz R, Dalmau J, Posner JB, Rosenfeld MR. T-cell receptor analysis in

anti-Hu associated paraneoplastic encephalomyelitis. *Neurology*.

1998;51(4):1146-1150. doi:10.1212/wnl.51.4.1146.

26. Galli J, Greenlee J. Paraneoplastic Diseases of the Central Nervous

System. *F1000Res*. 2020;9:F1000 Faculty Rev-167. Published 2020 Mar 6.

doi:10.12688/f1000research.21309.1.

27. Lancaster E, Dalmau J. Neuronal autoantigens--pathogenesis, associated

disorders and antibody testing. *Nat Rev Neurol*. 2012;8(7):380-390. Published

2012 Jun 19. doi:10.1038/nrneurol.2012.99.

28. Christie LJ, Honarmand S, Talkington DF, et al. Pediatric encephalitis: what is

the role of Mycoplasma pneumoniae?. *Pediatrics*. 2007;120(2):305-313.

doi:10.1542/peds.2007-0240.

29. Narita M. Classification of Extrapulmonary Manifestations Due to

Mycoplasma pneumoniae Infection on the Basis of Possible Pathogenesis. *Front*

Microbiol. 2016;7:23. Published 2016 Jan 28. doi:10.3389/fmicb.2016.00023.

30. Komatsu H, Kuroki S, Shimizu Y, Takada H, Takeuchi Y. Mycoplasma

pneumoniae meningoencephalitis and cerebellitis with antiganglioside

antibodies. *Pediatr Neurol*. 1998;18(2):160-164.

doi:10.1016/s0887-8994(97)00138-0.

31. Piquet AL, Clardy SL. Infection, Immunodeficiency, and Inflammatory

Diseases in Autoimmune Neurology. *Semin Neurol*. 2018;38(3):379-391.

doi:10.1055/s-0038-1660820.

32. Balbi GGM, Machado-Ribeiro F, Marques CDL, Signorelli F, Levy RA. The

interplay between tuberculosis and systemic lupus erythematosus. *Curr Opin*

Rheumatol. 2018;30(4):395-402. doi:10.1097/BOR.0000000000000493.

33. Zhang Y, Wang D, Yin Y, et al. Tuberculosis Infection in Chinese Patients with

Giant Cell Arteritis. *Sci Rep.* 2019;9(1):14364. Published 2019 Oct 7.

doi:10.1038/s41598-019-50892-9.

34. Ziak J, Weissova R, Jeřábková K, et al. CRMP2 mediates

Sema3F-dependent axon pruning and dendritic spine remodeling. *EMBO Rep.*

2020;21(3):e48512. doi:10.15252/embr.201948512.

35. Brittain JM, Pan R, You H, et al. Disruption of NMDAR-CRMP-2 signaling

protects against focal cerebral ischemic damage in the rat middle cerebral artery

occlusion model. *Channels (Austin).* 2012;6(1):52-59. doi:10.4161/chan.18919.

36. Yamashita N, Goshima Y. Collapsin response mediator proteins regulate

neuronal development and plasticity by switching their phosphorylation

status. *Mol Neurobiol.* 2012;45(2):234-246. doi:10.1007/s12035-012-8242-4.

37. Nagai J, Baba R, Ohshima T. CRMPs Function in Neurons and Glial Cells:

Potential Therapeutic Targets for Neurodegenerative Diseases and CNS

Injury. *Mol Neurobiol.* 2017;54(6):4243-4256. doi:10.1007/s12035-016-0005-1.

38. Dustrude ET, Moutal A, Yang X, Wang Y, Khanna M, Khanna R. Hierarchical

CRMP2 posttranslational modifications control NaV1.7 function. *Proc Natl Acad*

Sci U S A. 2016;113(52):E8443-E8452. doi:10.1073/pnas.1610531113.

39. Moutal A, Luo S, Largent-Milnes TM, Vanderah TW, Khanna R.

Cdk5-mediated CRMP2 phosphorylation is necessary and sufficient for peripheral

neuropathic pain. *Neurobiol Pain.* 2019;5:100022.

doi:10.1016/j.ynpai.2018.07.003.

40. Nakamura F, Ohshima T, Goshima Y. Collapsin Response Mediator Proteins:

Their Biological Functions and Pathophysiology in Neuronal Development and

Regeneration. *Front Cell Neurosci.* 2020;14:188. Published 2020 Jun 23.

doi:10.3389/fncel.2020.00188.

FIGURE LEGENDS

Figure 1 Identification of the autoantibody in a patient with an acute

encephalitis.

P1 patient was diagnosed as brain *M.P.* infection and possible secondary

immune-mediated acute encephalitis. The CSF sample from control patient with

Carpal Tunnel Syndrome (nCSF) and the serum from a health person (nSerum)

were negative controls. (A) IHC on mouse brain sections. Serum from P1 patient

had stained on cortex, hippocampus, and cerebellum. Cell nuclei were

counterstained blue with hematoxylin. (B) P1 patient's serum stained obviously in

cytosolic compartments of the neurons in cortex, CA3 area of hippocampus and

cerebellum on mouse brain sections (green). White arrows, possible

oligodendrocytes; arrow heads, Purkinje cells. CRMP2-expressing HEK293T cell

absorbed P1 patient's serum had no obvious staining. Cell nuclei were stained

blue with DAPI. (C) P1patient's serum (green) stained positively in soma and

dendrites of cultured mouse cortical neurons which were labeled with anti-MAP2

Ab (red). (D, E) IP of P1 CSF with rat brain protein lysate. 500 µl (lane 1) and 100

µl (lane 2) P1 CSF were applied to IP, respectively. 500 µl nCSF was used for

negative control. A positive band around 70 kDa was pulled down by P1 CSF but

not by nCSF. Similar protein size band (red arrow head) was observed on a

parallel gel with Coomassie brilliant blue staining (E). (F) Characterization of the

autoantigen by mass spectrometry. Suspected protein band in E was collected for

protein identification by LC-MS/MS. 17 identified peptides (red) matched to rat

DPYL2 (UniProtKB ID: P47942), an alias of CRMP2, with a score 226 and

sequence coverage 40%. CRMP2, collapsin response mediator protein 2; CSF,

cerebrospinal fluid; DAPI, 6-diamidino-2-phenylindole; DPYL2,

dihydropyrimidinase like 2; GL, granular layer; IB, immunoblotting; IHC,

immunohistochemistry; IP, immunoprecipitation; LC-MS/MS, liquid

chromatography tandem mass spectrometry; *M.P.*, *Mycoplasma pneumonia*;

MAP2, microtubule associated protein 2; ML, molecular layer; PCL, Purkinje cell

layer; SO, stratum oriens; SP, stratum pyramidale; SR, stratum radiatum. Scale

bars represent 50 µm in B and 20 µm in C.

Figure 2 Verification of anti-CRMP2 antibody in the suspected AE patients.

(A) IP with P1 patient's serum was performed with rat brain protein lysate.

Immunoblottings were done with P1 patient's serum (upper panel) and

commercial anti-CRMP2 antibody (lower panel). (B) IP with commercial

anti-CRMP2 antibody was performed with HEK293T cells overexpressed full

length human CRMP2 (isoform 1). Anti-CRMP2 antibody, P1 patient or negative

control sera were used for Western blotting, separately. (C) CRMP2-expressing

HEK293T cells were immunostained with patients' sera and anti-CRMP2

antibody. The sera from the first collection of P1 and P2 patients showed positive

staining and co-localized with CRMP2. Serum collected from P2 patient after

recovered 1.5 years showed negative staining. (D) Colocalization of P1 patient's

serum and anti-CRMP2 Ab immunostaining in cultured mouse cortical neurons.

CRMP2, collapsin response mediator protein 2; DAPI,

6-diamidino-2-phenylindole; IB, immunoblotting; nIgG, normal IgG; IP,

immunoprecipitation; WB, Western blotting. Scale bars represent 10 µm in C and 20 µm in D.

Figure 3 Specificity, epitope and IgG subtype determination of the antibody.

(A) Homology analysis showed that CRMP2 had an identity (red) of 76.224%, 76.573%, 75.524% and 50.348% with CRMP1, 3, 4, and 5, respectively. Blue, UniProtKB accession number. (B) CBA assay of the patients' sera for CRMPs. pcDNA3.1 based plasmids carrying CRMP1, 2, 3, 4 and CRMP5-eGFP were transfected into HEK293T cells separately for protein overexpression. Secondary antibodies anti-human IgG were labeled with green (CRMP1-4) or red (CRMP5) fluorescence. Cell nuclei were stained blue with DAPI. P1 (upper panels) and P2 (lower panels) patients' sera were only immunoreactive with CRMP2 among CRMP family members. (C) Schematic draws of the three CRMP2 isoforms and of the truncation strategy. The length or amino acid sites were indicated with numerals. Identical sequences between the isoforms were illustrated with same line colors. (D) pcDNA3.1 based plasmids carrying CRMP2 truncates T1, T2 or T3 illustrated in C were transfected into HEK293T cells separately for CBA assay. P1 (upper panels) and P2 (lower panels) patient's sera were reactive with T3 or 536 amino acids C-terminus of CRMP2 (red). (E) Immunostaining of CRMP2-expressing HEK293T cells with P1 patient serum and FITC conjugated anti-human IgG1-4 secondary antibodies. Positive stained in IgG4. CRMP, collapsin response mediator protein; DAPI, 6-diamidino-2-phenylindole; Iso., isoform. Scale bars represent 20 µm in B, D and E.

Figure 4 Magnetic resonance images of the patients.

(A-F) Cranial MRI of patient 1 was not remarkable with mild whiter matter

degeneration. (G-I) Cranial MRI of patient 2 showed multiple abnormal signals of

white matters in bilateral cerebral hemisphere and brainstem. (J-L) Spine MRI of

patient 2 showed long-segment spinal cord lesions from medulla to C6 segment.

After 3 months of treatment, (M-O) the cranial MRI of patient 2 showed that the

abnormal signals of the white matter and brainstem were reduced compared with

G-I; (P-R) the cervical MRI showed that the original abnormal signals of the spinal

cord had disappeared.

Table 1 Clinical Features of the Patients with anti-CRMP2 antibody.

Parameters	Patient 1	Patient 2
Gender	Female	Female
Age range	36-40	41-45
Symptoms	Dizziness, fever, slurred speech, visual rotation, unstable walking, opsoclonus-myoclonus	Headache, nausea and vomiting when raised his neck, left chest pain, right arm fatigue, numbness in both legs
CSF	WBC: 38 cells/μl (90% monocytes)	WBC: 240 cells/μl (90% monocytes), glucose: 2.64 mmol/l[a], protein: 0.80g/l
Autoimmune Abs	AE Abs[b] (CSF, serum), anti-CRMP5 and Homer3 Abs (serum), thyroid related Abs[c] (serum), remaining serum autoimmune Abs[d], all above (-)	AE Abs (CSF, serum) (-), anti-MOG and AQP4 Abs (CSF, serum) (-), anti-GFAP Ab (CSF) (-), thyroid related Abs (serum) (-), anti-histone Ab (serum) (+), remaining serum autoimmune Abs[e] (-)
TBA	Serum (+), CSF (+)	Serum (+), CSF (-)
Infection tests	Serum anti-*M.P.* IgM Ab (+), CSF NGS: *M.P.* (+)	T-spot (-), TB DNA (-), X-Pert (-), AFB (-)
MRI	Mild white matter degeneration	Multiple abnormal signals of white matters in bilateral cerebral hemisphere and brainstem, long-segment spinal cord lesions form medulla to C6 segment
Tumor	Tumor antigens/biomarkers[f] (-)	NA
Diagnosis	*M.P.* encephalitis and possible autoimmune mediated encephalitis	Tuberculous encephalomyelitis and ADEM
Immunotherapy	IVIG, MP	MP
Prognosis (3 months mRS)	2	0

Abbreviations: Abs, antibodies; ADEM, acute demyelinating encephalomyelitis; AE, autoimmune encephalitis; AFB, acid-fast bacillus test; CSF, cerebrospinal fluid; IVIG, intravenous immunoglobin; *M.P.*, *Mycoplasma pneumoniae*; MP, methylprednisolone; MRI, magnetic resonance imaging; mRS, modified Rankin Scale; NA, not available; NGS,next generation sequencing; TB, tubercle bacillus; TBA, tissue-based assay (rat brain sections); WBC, white blood cell.

[a] Normal range: 2.8-4.0 mmol/l.

[b] Autoimmune encephalitis antibodies: anti-NMDAR, LGI1, Caspr2, $GABA_BR$,

$AMPA_{A/B}R$, DPPX, DNER, D2R, mGluR5, GAD65, IgLON5 and GlyRα1

antibodies.

[c] Thyroid related antibodies: anti-thyrotropin receptor, thyroid peroxidase and

thyroglobulin antibodies.

[d] Autoantibodies tested for patient 1 serum: anti-Hu, Ri, Ma2, ANNA3, SOX1,

dsDNA, Sm, U1-RNP, Zic4, Yo, amphiphysin, CRMP5, PCA2, recoverin, Titin, Tr

(DNER) antibodies.

[e] Autoantibodies tested for patient 2 serum: anti-dsDNA, Jo-1, SSA/Ro52, SSB,

Sm, U1-RNP, PCNA, AMAM2, CENPB, PM-Scl, Scl-70, AnuA antibodies.

[f] Tumor antigens/biomarkers: CA-125, CYFRA21-1, SCC, CA242, CA-724,

CA-199, CEA, NSE.

A

P1 Serum

nSerum

B

P1 Serum/DAPI

Cortex | Hippocampus-CA3 | Cerebellum

SO
SP
SR

ML | PCL | GL

Absorbed P1 Serum/DAPI

Hippocampus-CA3 | Cerebellum

C

P1 Serum | MAP2 | DAPI | Merge | nSerum/MAP2/DAPI

D

IP   IP
P1 CSF   Input   nCSF

MW (kDa)
180 –
130 –
100 –
70 –
55 –
40 –
35 –
30 –
25 –

IB: P1 CSF

E

IP: P1 CSF

MW (kDa)
180 –
130 –
100 –
70 –
55 –
40 –
35 –
25 –
15 –

F

sp|P47942|DPYL2_RAT, Score: 226, Sequence coverage: 40%

```
1    MSYQGKKNIP RITSDRLLIK GGKIVNDDQS FYADIYMEDG LIKQIGENLI
51   VPGGVKTIEA HSRMVIPGGI DVHTRFQMPD QGMTSADDFF QGTKAALAGG
101  TTMIIDHVVP EPGTSLLAAF DQWREWADSK SCCDYSLHVD ITEWHKGIQE
151  EMEALVKDHG VNSFLVYMAF KDRFQLTDSQ IYEVLSVIRD IGAIAQVHAE
201  NGDIIAEEQQ RILDLGITGP EGHVLSRPEE VEAEAVNRSI TIANQTNCPL
251  YVTKVMSKSA AEVIAQARKK GTVVYGEPIT ASLGTDGSHY WSKNWAKAAA
301  FVTSPPLSPD PTTPDFLNSL LSCGDLQVTG SAHCTFNTAQ KAVGKDNFTL
351  IPEGTNGTEE RMSVIWDKAV VTGKMDENQF VAVTSTNAAK VFNLYPRKGR
401  ISVGSDADLV IWDPDSVKTI SAKTHNSALE YNIFEGMECR GSPLVVISQG
451  KIVLEDGTLH VTEGSGRYIP RKPFPDFVYK RIKARSRLAE LRGVPRGLYD
501  GPVCEVSVTP KTVTPASSAK TSPAKQQAPP VRNLHQSGFS LSGAQIDDNI
551  PRRTTQRIVA PPGGRANITS LG
```