Association between telecommuting environment and low back pain

among Japanese telecommuting workers: A cross-sectional study

Ryutaro Matsugaki[1], Ph.D., Keiji Muramatsu[1], M.D., Ph.D, Seiichiro Tateishi[2], M.D.,

Ph.D., Tomohisa Nagata[3], M.D., Ph.D., Mayumi Tsuji[4], M.D., Ph.D., Ayako Hino[5],

M.D., Ph.D., Kazunori Ikegami[6], M.D., Ph.D., Yoshihisa Fujino[7], M.D., M.P.H., Ph.D.,

and Shinya Matsuda[1], M.D., Ph.D., for the CORoNaWork Project

[1] Department of Preventive Medicine and Community Health, School of Medicine,

University of Occupational and Environmental Health, Japan

[2] Department of Occupational Medicine, School of Medicine, University of

Occupational and Environmental Health, Japan

[3] Department of Occupational Health Practice and Management, Institute of Industrial

Ecological Sciences, University of Occupational and Environmental Health, Japan

[4] Department of Environmental Health, School of Medicine, University of Occupational

and Environmental Health, Japan

[5] Department of Mental Health, Institute of Industrial Ecological Sciences, University of

Occupational and Environmental Health, Japan

[6] Department of Work Systems and Health, Institute of Industrial Ecological Sciences,

University of Occupational and Environmental Health, Japan

[7] Department of Environmental Epidemiology, Institute of Industrial Ecological

Sciences, University of Occupational and Environmental Health, Japan

Corresponding author:

Yoshihisa Fujino, M.D., M.P.H., Ph.D.

Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,

University of Occupational and Environmental Health, Japan

1-1, Iseigaoka, Yahatanishiku, Kitakyushu, 807-8555, Japan

Tel: +81-93-691-7401

Email: zenq@med.uoeh-u.ac.jp

Abstract

Objectives: We evaluated the relationship between telecommuting environment and

low back pain (LBP) among desk-based workers in Japan.

Methods: This cross-sectional study included 3,663 desk-based, telecommuting

workers. LBP was assessed using a 0–10 numerical rating scale. The telecommuting

environment was evaluated using subjective questions. Mixed-effects logistic regression

analysis was used.

Results: Mixed-effects logistic model results revealed that not having a place or room

to concentrate on work, desk not well-lit enough for work, not having enough space on

the desk to work, not having enough legroom, and not having comfortable temperature

and humidity conditions in the workspace were significantly associated with higher

odds of LBP.

Conclusions: Our findings suggest that telecommuting environment is associated with

the prevalence of LBP.

Keywords: Low Back Pain, Telecommuting, Workplace, COVID-19, Japan

Introduction

The coronavirus disease (COVID-19) pandemic led to rapid expansion of telecommuting in Japan. The Japanese government recommended telecommuting to prevent the spread of COVID-19 infection[1,2]. Although the telecommuting rate of a Japanese worker was 14.8% as of October 2019, it increased to 23.0% by November 2020[3]. In actuality, 85.0% of workers who were teleworking as of November 2020 did so to prevent the spread of COVID-19 infection[3]. Telecommuting is a way to continue business even in times of emergency, and it may become an even more common way of working.

With the rapid expansion of telecommuting due to the COVID-19 pandemic, management of the work environment during telecommuting is an emerging occupational health problem. As with the office environment, the recommended working environment for a telecommute worker is a private, quiet, and safe dedicated space, with adequate lighting, comfortable temperature and humidity, ergonomic chairs/desk[4–6]. However, it is difficult to manage the telecommute work environment because, in contrast to the office, the workstation at home is difficult to supervise by managers, and managers do not have the authority to direct the private environment of an individual. In addition, neither companies nor workers were sufficiently prepared to optimize telecommuting environment due to the unexpected occurrence of COVID-19 [4]. It has been reported that more than 50% of telecommuting workers in Japan do not have a desk/ chair and more than 70% do not have a private room or space for work[3].

Low back pain (LBP) is an important health problem associated with office work. The prevalence of LBP is 34-56% among office workers [7–10]. In office workers, gender, body mass index, sleep disturbance, and previous symptoms of LBP are known to be risk factors for LBP [8,10–13]. In terms of work-related factors, it is known that sitting

time and sitting posture at work are known to be risk factors for LBP in an office

worker[8,14,15]. At home, the work environment is less developed than in the office

environment; therefore, the work environment of home workers is assumed to be an

important risk factor for low back pain.

Previous studies conducted in the COVID-19 pandemic have suggested an

association between telecommuting and musculoskeletal pain[16]. However, the

relationship between work environment and LBP in telecommuting workers is unclear.

Thus, the purpose of this study was to clarify the relationship between the work

environment and LBP of telecommuting workers.

Methods

Study Design and Subjects

This cross-sectional internet-based monitoring survey was conducted from

December 22 through to December 26, 2020, when the third wave of COVID-19

infections began in Japan. The details of the survey protocol have been previously

reported.[17] The data was collected from the workers with employment contracts at the

time of the survey. Of the 33,302 workers participating in the survey, 27,036 were

surveyed, excluding those who gave fraudulent answers. Of these, 3,663 (2,093 males

and 1,570 females) who responded that they mainly performed desk work (e.g., office

work, computer work) and telecommuted at least once a week were included in the

present analysis.

This study conformed to the principles of the Declaration of Helsinki. In addition,

the study was approved by the ethics committee of the University of Occupational and

Environmental Health, Japan (reference No. R2-079). Informed consent was obtained

online from the participants through the website.

The assessment of LBP

We assessed the presence of LBP in the participants based on two questions. First,

we asked all subjects "Have you experienced stiff shoulders or LBP in the past two

weeks?" and asked them to answer, "yes" or "no." If the subject answered "yes" to that

question, the following questions were asked to assess the severity of LBP such as, "what

was your average level of LBP in the past 2 weeks? (Please rate your pain from 0 to 10,

where 0=no pain at all and 10=the most intense pain you have experienced)." The

numerical rating scale (NRS) was used for evaluating the pain severity. In this study, a

score of 3 or higher on the NRS was defined as LBP.

The assessment of the telecommuting environment

The telecommuting environment was assessed among the telecommuting workers

with the following questions: 1) "Do you have a place or room where you can concentrate

on your work?" ; 2) "Is your desk well-lit enough for you to work?" ; 3) "Do you have

enough space on your desk to work?"; 4) "Is there enough space to stretch the legs?"; 5)

"Are the degrees of temperature and humidity in the room where you work appropriate

for working comfortably?"; 6) "Do you use an office desk or chair? (Including children's

study desks)." The respondents answered either "yes" or "no."

The assessment of participants' characteristics and other covariates

The following items were examined for socioeconomic factors: age, sex, body

mass index (calculated by dividing the weight by height squared), educational

background (junior high school; high school; vocational school; junior college or

technical college; university; or graduate school) and equivalent income (household

income divided by the square root of the household size).

The following items were examined for lifestyle factors: smoking (currently

smoking), drinking (alcohol consumption on two or more days per week), physical

activity (perform equivalent physical activities for at least 1 h a day in daily life for more

than 2 days a week), and exercise habit (exercise for 30 minutes or more for more than

two days a week).

For mental health status was assessed using the following question: "During the

past 30 days, how many days did you experience poor mental health, including stress,

depression, emotional problems, etc.?"

The following items were examined for work-related factors: we examined the

following items: industry type, working time (hours per week), frequency of

telecommuting (one day per week; more than two days per week; and more than four days

per week), company size (total number of employees at the company where the

respondent is working), and working hours per week.

Statistical analysis

The number of days that the participant experienced poor mental health during the

past 30 days are expressed as continuous variables using mean and standard deviation.

Other variables are presented as categorical variables using numbers and percentages.

Mixed-effects logistic regression analysis was conducted with the presence of

LBP as the dependent variable, subjective evaluation of the telecommuting environment

as the independent variable, with the city of residence as the random effects. We used age,

sex, body mass index, lifestyle habits, the number of days of poor mental health,

equivalent income, educational background, working hours, company size, and industry

type as covariates to adjust for potential confounders.

All statistical analysis were performed with Stata software (Stata Statistical

Software: Release 16; StataCorp LLC, TX, USA). A p-value of < 0.05 was considered

statistically significant.

Results

Table 1 presents the characteristics of the participants. The prevalence rate of

LBP in the study participants was 49.9%. Compared to the participants without LBP, the

participants who experienced LBP were younger; middle-aged workers (20–49 years:

43.1% vs. 52.2%) and more likely to be females (37.6% vs. 48.2%).

[Insert Table 1. here]

The associations of LBP and telecommuting environments are shown in Table 2.

There was a significant association of LBP with the question "Do you have a place or

room where you can concentrate on your work? (No)" (odds ratio [OR]: 1.37, 95% CI:

1.16–1.63, p<0.001), "Is your desk well-lit enough for you to work? (No)" (OR: 1.47,

95%CI: 1.22–1.79, p<0.001), "Do you have enough space on your desk to work? (No)"

(OR: 1.22, 95%CI: 1.04–1.43, p=0.012), "Is there enough space to stretch the legs? (No)"

(OR: 1.32, 95%CI: 1.11–1.57, p=0.002), and "Are the degrees of temperature and

humidity in the room where you work appropriate for working comfortably? (No)" (OR:

1.37, 95%CI: 1.16–1.61, p<0.001). There was no significant association of LBP with the

question "Do you use an office desk or chair? (NO)" (OR: 1.04, 95%CI: 0.90–1.19,

p=0.633). Similar results were obtained for Model 1, which was adjusted only for sex and

age, and Model 2, which was adjusted for other potential confounders.

[Insert Table 1. here]

Discussion

This study showed that telecommuting environment was associated with LBP

in telecommuting workers during the COVID-19pandemic. Specifically, it was

suggested that insufficient desk and foot space, inadequate desk lighting, uncomfortable

room temperature and humidity, and lack of room/space for concentrating on work were

associated with the prevalence of LBP.

In this study, inadequate desk and foot space, and insufficient lighting were

associated with LBP. It is suggested that an awkward posture and sitting for long

durations are risk factors for LBP [8,18–20]. Having enough space on the desk and at the

feet is effective in maintaining good posture when working as well as changing the

posture and stretching as needed. Also, inadequate lighting at the desk may contribute to

awkward posture of workers when looking at documents or computer screens on their

desk. It is necessary to manage the work environment to avoid awkward postures and

prolonged maintenance of the same posture during work.

The present study also revealed that uncomfortable temperature and humidity

in the telecommuting space was associated with LBP. Since this study was conducted

during the winter season, it can be speculated that the results imply that cold

temperature and low humidity is associated with LBP. The relationship between cold

temperature and LBP has been shown in previous studies [15,21–23]; this is also the same

for telecommuting workers. Although the most suitable room temperature for the

prevention of LBP is not clear, we consider that room temperatures that are subjectively

cold should be avoided as they may increase the risk of musculoskeletal symptoms and

injuries.

In this study, using an office chair/desk was not associated with LBP. This

finding is supported by the findings of previous studies reporting no significant

association between LBP and use of chair/desk in office workers [11,24]. Alternately,

previous studies have reported that LBP is related to the characteristics of the chair,

such as with or without lumbar support and adjustable back support[8,18]. This may

suggest that it is not simply a matter of whether an office chair is used, but what

function and shape of the chair is used for prevention of LBP.

The lack of room or space to concentrate on work was associated with LBP. It

is suggested that the teleworker's workstation should be in a dedicated space that is

private, quiet, and secure, preferably away from the flow of activity in the home [4]. The

results of this study support that argument. Previous studies have reported that

psychological stress is associated with LBP in workers [25–27]. The lack of room or space

to concentrate on work may cause psychological stress to telecommuting workers. The

results of this study suggest that it is important for telecommuting workers to have a

space where they can devote themselves to their work as much as possible, even if they

work from home.

Based on these findings, we suggest that the work environment of

telecommuting workers may be associated with LBP. Therefore, employers need

educate telecommuting workers of the importance of an appropriate home working

environment. If it is difficult to prepare an appropriate telecommuting environment due

to household situations and family structure, it is necessary to consider the use of

co-working spaces, satellite offices, and spaces near the employee's residence for

telecommuting.

Moreover, this study has three limitations. Firstly, the specific details such as

previous history, duration of the symptoms, and diagnosis of LBP are unknown as only

the NRS was used for its evaluation. It is uncertain how these factors affect the

relationship between the telecommuting environment and low back pain. Secondly, the

home environment was also evaluated using subjective questions; hence, the validity of

the responses is unclear. However, an objective method to evaluate the work

environment at home has not been established at this time. Thirdly, there may be a

selection bias in this study. If it is possible to choose telecommuting, people with LBP

may be more likely to choose telecommuting. The effect of this bias on the results of

this study is uncertain.

Conclusion

The present study suggests that telecommuting environment is associated with

the prevalence of LBP among telecommuting workers in Japan and suggests that

employers may need to consider providing telecommuting workers with a working

space where they can concentrate on their work with appropriate room temperature and

humidity control to prevent LBP.

References

1.   Decisions by the Headquarters for Novel Coronavirus Disease Control. Basic

Policies for Novel Coronavirus Disease Control. Ministry of Health, Labour and

Welfare. Published 2020. Accessed June 9, 2021.

https://www.mhlw.go.jp/content/10200000/000603610.pdf

2.   Decisions by the Headquarters for Novel Coronavirus Disease Control. Basic

Policies for Novel Coronavirus Disease Control (Revised on April 7, 2020).

Ministry of Health, Labour and Welfare. Published 2020. Accessed June 9, 2021.

https://www.mhlw.go.jp/content/10900000/000620733.pdf

3.   Vital statistics of telework (in Japanese). Ministry of Land, Infrastructure,

Transport and Tourism. Published 2021. Accessed June 8, 2021.

https://www.mlit.go.jp/toshi/daisei/content/001392107.pdf

4.   Robertson MM, Mosier K. Work from home: Human factors/ergonomics

considerations for teleworking. International Labour Organization. Published

2020. Accessed May 31, 2021.

http://www.ilo.org/global/topics/safety-and-health-at-work/events-training/events

-meetings/world-day-safety-health-at-work/WCMS_742061/lang--en/index.htm

5.   Work environment maintenance for teleworking at home (in Japanese). Ministry

of Health, Labour and Welfare. Published 2020. Accessed June 9, 2021.

https://www.mhlw.go.jp/stf/newpage_01603.html

6.   Guidelines for the Promotion of Appropriate Introduction and Implementation of

Telework (in Japanese). Ministry of Health, Labour and Welfare. Published 2021.

Accessed June 9, 2021.

https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/koyou_roudou/roudoukijun/shi

goto/guideline.html

7.   Janwantanakul P, Pensri P, Jiamjarasrangsri V, Sinsongsook T. Prevalence of

self-reported musculoskeletal symptoms among office workers. *Occup Med*

(Chic Ill). 2008;58(6):436-438. doi:10.1093/occmed/kqn072

8.   Spyropoulos P, Papathanasiou G, Georgoudis G, Chronopoulos E, Koutis H,

Koumoutsou F. Prevalence of low back pain in greek public office workers. *Pain*

Physician. 2007;10(5):651-659. Accessed May 12, 2021.

https://pubmed.ncbi.nlm.nih.gov/17876361/

9.   Fanta M, Alagaw A, Kejela G, Tunje A. Low back pain and associated factors

among civil service sectors office workers in Southern Ethiopia. *Int J Occup Saf*

Heal. 2020;10(1):53-63. doi:10.3126/ijosh.v10i1.29883

10.   Kaliniene G, Ustinaviciene R, Skemiene L, Vaiciulis V, Vasilavicius P.

Associations between musculoskeletal pain and work-related factors among

public service sector computer workers in Kaunas County, Lithuania. *BMC*

Musculoskelet Disord. 2016;17(1):1-12. doi:10.1186/s12891-016-1281-7

11.   Juul-Kristensen B, Søgaard K, Strøyer J, Jensen C. Computer users' risk factors

for developing shoulder, elbow and back symptoms. *Scand J Work Environ Heal*.

2004;30(5):390-398. doi:10.5271/sjweh.827

12.   Janwantanakul P, Sitthipornvorakul E, Paksaichol A. Risk factors for the onset of

nonspecific low back pain in office workers: A systematic review of prospective

cohort studies. *J Manipulative Physiol Ther*. 2012;35(7):568-577.

doi:10.1016/j.jmpt.2012.07.008

13.   Shariat A, Cardoso JR, Cleland JA, et al. Prevalence rate of neck, shoulder and

lower back pain in association with age, body mass index and gender among

Malaysian office workers. *Work*. 2018;60(2):1-9. doi:10.3233/wor-2738

14.   Janwantanakul P, Pensri P, Moolkay P, Jiamjarasrangsi W. Development of a

risk score for low back pain in office workers -a cross-sectional study. *BMC*

Musculoskelet Disord. 2011;12(1):23. doi:10.1186/1471-2474-12-23

15.   Ye S, Jing Q, Wei C, Lu J. Risk factors of non-specific neck pain and low back

pain in computer-using office workers in China: A cross-sectional study. *BMJ*

Open. 2017;7(4). doi:10.1136/bmjopen-2016-014914

16.   Yoshimoto T, Fujii T, Oka H, Kasahara S, Kawamata K, Matsudaira K. Pain

Status and Its Association with Physical Activity, Psychological Stress, and

Telework among Japanese Workers with Pain during the COVID-19 Pandemic.

Int J Environ Res Public Health. 2021;18(11):5595. doi:10.3390/ijerph18115595

17.   Fujino Y, Ishimaru T, Eguchi H, et al. Protocol for a nationwide Internet-based

health survey in workers during the COVID-19 pandemic in 2020. *medRxiv*.

Published online 2021. doi:https://doi.org/10.1101/2021.02.02.21249309

18.   Janwantanakul P, Pensri P, Moolkay P, Jiamjarasrangsi W. *Development of a*

Risk Score for Low Back Pain in Office Workers - a Cross-Sectional Study. Vol

12.; 2011. doi:10.1186/1471-2474-12-23

19.   Liira JP, Shannon HS, Chambers LW, Haines TA. Long-term back problems and

physical work exposures in the 1990 Ontario health survey. *Am J Public Health*.

1996;86(3):382-387. doi:10.2105/AJPH.86.3.382

20.   Lis AM, Black KM, Korn H, Nordin M. Association between sitting and

occupational LBP. *Eur Spine J*. 2007;16(2):283-298.

doi:10.1007/s00586-006-0143-7

21.   Jin K, Sorock GS, Courtney T, et al. Risk factors for work-related low back pain

in the People's Republic of China. *Int J Occup Environ Health*. 2000;6(1):26-33.

doi:10.1179/oeh.2000.6.1.26

22.   Dovrat E, Katz-Leurer M. Cold exposure and low back pain in store workers in

313   Israel. *Am J Ind Med*. 2007;50(8):626-631. doi:10.1002/ajim.20488

314   23.   Skandfer M, Talykova L, Brenn T, Nilsson T, Vaktskjold A. Low back pain

315   among mineworkers in relation to driving, cold environment and ergonomics.

316   *Ergonomics*. 2014;57(10):1541-1548. doi:10.1080/00140139.2014.904005

317   24.   Celik S, Celik K, Dirimese E, Tasdemir N, Arik T, Büyükkara İb. Determination

318   of pain in musculoskeletal system reported by office workers and the pain risk

319   factors. *Int J Occup Med Environ Health*. 2018;31(1):91-111.

320   doi:10.13075/ijomeh.1896.00901

321   25.   Vinstrup J, Jakobsen MD, Andersen LL. Perceived Stress and Low-Back Pain

322   Among Healthcare Workers: A Multi-Center Prospective Cohort Study. *Front*

323   *Public Heal*. 2020;8. doi:10.3389/fpubh.2020.00297

324   26.   Tsuboi Y, Ueda Y, Naruse F, Ono R. The Association between Perceived Stress

325   and Low Back Pain among Eldercare Workers in Japan. *J Occup Environ Med*.

326   2017;59(8):765-767. doi:10.1097/JOM.0000000000001062

327   27.   Yoshimoto T, Oka H, Katsuhira J, et al. Prognostic psychosocial factors for

328   disabling low back pain in Japanese hospital workers. *PLoS One*. 2017;12(5).

329   doi:10.1371/journal.pone.0177908

Table 1. Characteristics of participants

		Low back pain	
		-	+
		n (%)	n (%)
		1,834	1,829
Age (y)	20-29	88 (4.8%)	82 (4.5%)
	30-39	241 (13.1%)	319 (17.4%)
	40-49	462 (25.2%)	554 (30.3%)
	50-59	716 (39.0%)	629 (34.4%)
	60-69	327 (17.8%)	245 (13.4%)
Sex	Men	1145 (62.4%)	948 (51.8%)
	Women	689 (37.6%)	881 (48.2%)
Body mass index	< 18.5	199 (10.9%)	240 (13.1%)
	18.5-25.0	1265 (69.0%)	1200 (65.6%)
	25.0-30.0	317 (17.3%)	312 (17.1%)
	> 30.0	53 (2.9%)	77 (4.2%)
Educational background	Junior high school	15 (0.8%)	14 (0.8%)
	High school	268 (14.6%)	262 (14.3%)
	Vocational school	160 (8.7%)	173 (9.5%)
	junior college/technical college	157 (8.6%)	189 (10.3%)
	University	1030 (56.2%)	1014 (55.4%)
	Graduate school	204 (11.1%)	177 (9.7%)
Equivalent income (10,000yen)	<260	517 (28.2%)	485 (26.5%)
	261-450	458 (25.0%)	507 (27.7%)
	451-600	374 (20.4%)	393 (21.5%)
	> 601	485 (26.4%)	444 (24.3%)
Lifestyle habit	Smoking (yes)	447 (24.4%)	460 (25.2%)
	Drinking ($\geq$2d/w)	823 (44.9%)	855 (46.7%)
	Physical activity ($\geq$2d/w)	697 (38.0%)	685 (37.5%)
	Exercise habit ($\geq$2d/w)	531 (29.0%)	523 (28.6%)
Days experienced poor mental health		2.3 (5.9)	5.1 (8.4)

during the past 30 days, mean (SD)

Industry type	Energy, materials, industrial machinery	71 (3.9%)	85 (4.6%)
	Food	20 (1.1%)	14 (0.8%)
	Beverages/Tobacco products	8 (0.4%)	6 (0.3%)
	Pharmaceuticals/Medical supplies	38 (2.1%)	32 (1.7%)
	Cosmetics/Toiletries/Sanitary products	13 (0.7%)	12 (0.7%)
	Fashion and Accessories	20 (1.1%)	19 (1.0%)
	Precision Machinery and Office supplies	43 (2.3%)	45 (2.5%)
	Home appliances/AV equipment	61 (3.3%)	57 (3.1%)
	Automobiles and Transportation equipment	72 (3.9%)	67 (3.7%)
	Household goods	3 (0.2%)	4 (0.2%)
	Hobby/Sporting goods	8 (0.4%)	5 (0.3%)
	Real estate and Housing equipment	105 (5.7%)	96 (5.2%)
	Information and Communication	318 (17.3%)	288 (15.7%)
	Distribution and Retail	72 (3.9%)	91 (5.0%)
	Finance/Insurance	87 (4.7%)	98 (5.4%)
	Transportation and Leisure	20 (1.1%)	22 (1.2%)
	Restaurant and Other services	44 (2.4%)	49 (2.7%)
	Public offices and Organizations	50 (2.7%)	56 (3.1%)
	Education, medical services, religion	71 (3.9%)	97 (5.3%)
	Mass media	29 (1.6%)	30 (1.6%)
	Market research	7 (0.4%)	8 (0.4%)
	Other	674 (36.8%)	648 (35.4%)
Working time (h/w)	< 40	1346 (73.4%)	1293 (70.7%)
	40-48	237 (12.9%)	245 (13.4%)
	49-60	189 (10.3%)	232 (12.7%)
	> 60	62 (3.4%)	59 (3.2%)
Frequency of telecommute	1 d/w	266 (14.5%)	297 (16.2%)
	2-3 d/w	495 (27.0%)	563 (30.8%)
	≥ 4d/w	1073 (58.5%)	969 (53.0%)
Company size (person)	-9	732 (39.9%)	614 (33.6%)
	10-99	202 (11.0%)	228 (12.5%)
	100-999	304 (16.6%)	370 (20.2%)
	>1000	596 (32.5%)	617 (33.7%)

SD: standard deviation

Table 2. The association between low back pain and telecommuting environment

	Total		Low back pain		Model 1†				Model 2‡			
	n	%	n	%	OR	95% CI		P	OR	95% CI		P
Environment of telecommute												
Do you have a place or room where you can concentrate on your work? (No)	847	23.1	496	58.6	1.51	1.29	1.77	<0.001	1.37	1.16	1.63	<0.001
Is your desk well-lit for you to work? (No)	596	16.3	361	60.6	1.65	1.37	1.98	<0.001	1.47	1.22	1.79	<0.001
Do you have enough space on your desk to work? (No)	1054	28.8	582	55.2	1.33	1.15	1.54	<0.001	1.22	1.04	1.43	0.012
Is there enough space to stretch the legs? (No)	784	21.4	450	57.4	1.46	1.24	1.72	<0.001	1.32	1.11	1.57	0.002
Are the degrees of temperature and humidity in the room where you work appropriate for working comfortably? (No)	921	25.1	526	57.1	1.44	1.24	1.69	<0.001	1.37	1.16	1.61	<0.001
Do you use an office desk or chair? (Including children's study desks) (No)	1791	48.9	931	52.0	1.10	0.97	1.26	0.149	1.04	0.90	1.19	0.633

†Model 1: Adjusted for age, sex

‡Model 2: Adjusted for age, sex, body mass index, lifestyle habit (smoking, drinking, physical activity, and exercise habit), days experienced poor mental health during the past 30 days, equivalent income, educational background, industry type, working time, frequency of telecommute, company size

CI: confidence interval, OR: Odds ratio