

Title: Tobacco and Nicotine Use in the United States Army Trainee Soldier's Environment

Authors: Adam Edward Lang PharmD¹, Aleksandra Yakhkind MD², Elizabeth Prom-Wormley PhD, MPH³

1. Department of Primary Care, McDonald Army Health Center, Fort Eustis, Virginia

2. Department of Neurology, University of Pennsylvania Health System, Philadelphia, Pennsylvania

3. Department of Family Medicine and Population Health, Virginia Commonwealth University, Richmond, Virginia

Correspondence Author: Adam Edward Lang | a.edward.lang@gmail.com

McDonald Army Health Center, 576 Jefferson Avenue, Fort Eustis, Virginia 23604

The views expressed in this publication are those of the author and do not necessarily reflect the official policy of the Department of Defense, Department of the Army, U.S. Army Medical Department, or the U.S. Government.

Abstract

Tobacco and nicotine use rates within the United States military population are high, and rates among young trainee soldiers are not improving over time. Trainee soldiers are heavily influenced by role models in their environment, and nicotine product use rates in this environment has not previously been evaluated. This study found that over 30% of non-trainee U.S. Army soldiers, known as cadre, within the training environment use nicotine products. For the young trainee soldier, this creates an initial Army experience that continues to foster a culture of tobacco and nicotine use. We urge changes in Army tobacco and nicotine policy to protect the vulnerable trainee soldier population.

Abbreviations: AIT (Advanced Individual Training), BCT (Basic Combat Training, US (United States))

Introduction

Rates of tobacco and nicotine product use in United States (US) Army trainees in 2019 were shown to be over 35%, which was significantly higher compared to those aged 18-24 in the general US population that same year (1, 2). One promising strategy to reduce the rates of tobacco and nicotine use in soldiers is the implementation of policies restricting their use. Specifically, the use of these products is not allowed in the first 10-week phase of Army training (basic combat training, BCT). Rates of nicotine product use in trainees who smoked prior to BCT decreased by 31.5% after this stage of training (1). Nicotine use is allowed in the second phase (advanced individual training, AIT). This shift in policy towards nicotine use has allowed trainees who used prior to BCT to reinitiate early in a cessation attempt. It also puts trainees who never used nicotine products in an environment where peer nicotine product use is allowed and accepted.

In 2020, a nicotine-free policy was implemented throughout an AIT brigade on Fort Eustis in Virginia, where training spans on average three to six months, in order to encourage nicotine abstinence and extend the trainee nicotine-free period. AIT trainees were subject to this policy, but cadre were not. Cadre include all non-trainee active duty service members such as drill sergeants, course instructors, and those in command positions. Cadre hold enormous influence over trainees through constant daily interactions. Consequently, AIT trainees may be exposed to inconsistent messaging regarding social norms related to tobacco and nicotine use in the Army. Our study sought to identify prevalence of tobacco and nicotine use among cadre to understand the potential degree of trainee exposure to tobacco and nicotine during AIT.

Methods

Data were collected using a confidential online survey between January 08-25, 2020, just prior to the implementation of the AIT nicotine-free policy discussed above. All cadre in an AIT training brigade on Fort Eustis (N = 757) received an email with an invitation to participate in the survey as well as contact information for the tobacco treatment clinic to assist those interested in receiving treatment for tobacco and nicotine dependence. Four hundred and thirteen (N = 413) individuals participated in the survey (54.6% response rate). The study was deemed exempt from review by the Clinical Investigation Department at Naval Medical Center Portsmouth. Statistical analyses were conducted using R, version 3.6.1 (3).

Results

One hundred and ninety (46.0%) cadre participants engaged in any lifetime nicotine use, 159 (38.4%) used prior to joining the military, and 126 (30.8%) were current nicotine users. Most current users (74.0%) have used nicotine for seven years or more and 38.6% had been using for 15 years or more. Further, 51.1% of current users engaged in nicotine use six or more times per day. The most common product used was smokeless tobacco (40.6%), followed by conventional cigarette use (36.6%). Of current users, 20.5% engaged in dual or polyproduct use. The most common combination of use was combustible cigarette and smokeless tobacco (45.8%), followed by combustible cigarette and electronic cigarette use (20.8%, Table 1).

At the time of the survey, 29.9% of current users were interested in quitting. The proportion of interest to quit increased to 45.7% when participants were asked whether they wanted to quit within a year.

Discussion

Cadre tobacco and nicotine use in the military training setting has not previously been evaluated.

The results in this study indicate that almost one-third of cadre are current users and that over half engage in a high frequency of nicotine use (6 or more times per day). Consequently, tobacco and nicotine use in cadre is widespread and is expected to be a factor in trainee tobacco and nicotine use. In prior studies, young Air Force trainees as well as civilian adolescents report their tobacco use to be heavily influenced by cadre or role model tobacco use, peer tobacco use, and the individual's perception of social norms (4, 5). Of concern in our study is the relatively small change in prevalence of cadre nicotine use from before they joined the military until the present time, along with the high percentage of cadre using for greater than 15 years. Given the interest in quitting among cadre, a focus on policy change and appropriate treatment of tobacco and nicotine dependence in soldiers of every age and rank must be a priority for the United States Army. Interventions targeting seasoned soldiers would prevent the downstream effect of influencing use in the trainee soldier. This influence, which likely results in de novo, reinitiated, continued, and multiproduct use, is a major part of the cycle contributing to systematic use within the Army. Interventions are essential to prevent poor health outcomes in military, Veterans Affairs, and civilian populations, and the degradation of US Army readiness. Study limitations include the potential for recall bias of nicotine use prior to joining the military, absence of objective measurements of nicotine use, self-selection bias, inability to compare demographics of those who responded to survey non-responders, an underrepresentation of women compared to the military average, lack of differentiation between non-nicotine and nicotine e-cigarettes on the survey, and a population of soldiers only within the aviation field.

Future AIT nicotine-free policies must include cadre in order to be maximally effective. The Army trainee population is highly vulnerable. Policies to protect them are realistic in the highly controlled military setting, and lack of action allows the harmful cycle of military tobacco and nicotine use to continue.

References

1. Lang AE, Yakhkind A, Lamb RW, Stack KM. Effect of a Basic Training Nicotine-Free Policy on Soldiers in the United States Army [published online ahead of print, 2021 Mar 26]. *CHEST*. 2021; S0012-3692(21)00564-X. doi: 10.1016/j.chest.2021.03.039
2. Cornelius, Monica E et al. “Tobacco Product Use Among Adults - United States, 2019.” *MMWR. Morbidity and mortality weekly report* vol. 69,46 1736-1742. 20 Nov. 2020, doi:10.15585/mmwr.mm6946a4
3. R Core Team (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>
4. Green KJ, Hunter CM, Bray RM, Pemberton M, Williams J. Peer and role model influences for cigarette smoking in a young adult military population. *Nicotine Tob Res*. 2008;10(10):1533-1541. doi:10.1080/14622200802398763
5. National Center for Chronic Disease Prevention and Health Promotion (US) Office on Smoking and Health. Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General. Atlanta (GA): Centers for Disease Control and Prevention (US); 2012. 2, The Health Consequences of Tobacco Use Among Young People. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK99242/>

Table 1. Summary Statistics

<i>Variable</i>	<i>N</i>	<i>%</i>
Gender		
Man	370	89.6
Woman	34	8.2
Other	9	2.2
Total	413	100.0
Age		
18-29	65	15.7
30-39	247	59.8
40 and over	101	24.5
Total	413	100.0
Any Lifetime Nicotine Use		
No	223	54.0
Yes	190	46.0
Total	413	100.0
Nicotine Use Prior to the Military		
No	254	61.6
Yes	159	38.4
Total	413	100.0
Current Nicotine Use		
No	286	69.2
Yes	127	30.8
Total	417	100.0
Length of Use (in Years)		
Less than 1	9	7.1
1-3	12	9.4
4-6	12	9.4
7-10	22	17.3
11-15	23	18.1
15 and over	49	38.6
Total	127	99.9
Frequency of Daily Use		
Less than 1 time/day	8	6.3
1-5 times/day	54	42.5
6-10 times/day	36	28.3
More than 10 times/day	29	22.8
Total	127	99.9
Poly-product Use		
1 product	101	79.5
2 products	24	18.9
3 or more products	2	1.6
Total	127	100.0
Single Product-Specific Use		
Electronic Cigarette	20	19.8
Conventional Cigarette	37	36.6
Cigar	3	3.0
Smokeless	41	40.6
Dissolvable	0	0.0
Pipe	0	0.0
Total	101	100.0
Dual Use		
Cigarette/ Smokeless	11	45.8
Cigarette/ Cigar	5	20.8
Cigarette/ Electronic Cigarette	5	20.8
Cigarette/Pipe	1	4.2
Cigar/ Pipe	1	4.2
Electronic Cigarette/ Dissolvable	1	4.2
Total	24	100.0
Current Desire to Quit		
No	89	70.1
Yes	38	29.9
Total	127	100.0
Desire to Quit within a Year		
No	69	54.3
Yes	58	45.7
Total	127	100.0