# Identifying Louisiana Communities at the Crossroads of Environmental and Social Vulnerability, COVID-19, and Asthma

Arundhati Bakshi, PhD*; Alicia Van Doren, MD; Colette Maser, MPHTM; Kathleen Aubin, MSPH; Collette Stewart, MSPH; Shannon Soileau, MS; Kate Friedman, MNS; Alexis Williams, MPH

Section of Environmental Epidemiology and Toxicology, Office of Public Health, Louisiana Department of Health, Baton Rouge, LA-70802

*corresponding author
- Address: 628 N. 4th St. #516-5, Baton Rouge, LA-70802
- Email: Arundhati.Bakshi@LA.gov
- Office: 225-342-3279

**Short Title:** Intersection of Social and Environmental Vulnerability, COVID-19, and Asthma in Louisiana

**Keywords:** asthma, COVID-19, coronavirus, chronic disease, environment, social vulnerability, environmental justice, population health, health disparities

**Abstract**

The COVID-19 pandemic has disproportionately affected the socially and environmentally vulnerable, including through indirect effects on other health conditions. Asthma is one such condition, which may be exacerbated by both prolonged adverse in-home exposures if quarantining in unhealthy homes and prolonged outdoor exposures if the ambient air quality is unhealthy or hazardous. As both are often the case in Environmental Justice (EJ) communities, here we have analyzed data at the census tract (CT) level for Louisiana to assess any correlation between social and environmental vulnerability, and health issues like COVID-19 and asthma. Higher Social Vulnerability Index (SVI), Particulate Matter less than 2.5 µm in diameter ($PM_{2.5}$) and Ozone levels were associated with higher rates of cumulative COVID-19 incidence at various time points during the pandemic, as well as higher average annual asthma hospitalization rates and estimated asthma prevalence. Further, cumulative COVID-19 incidence during the first three months of the pandemic was moderately correlated with both asthma hospitalizations and estimated prevalence, suggesting similar underlying factors may be affecting both conditions. Additionally, 137 CTs were identified where social and environmental vulnerabilities co-existed, of which 75 (55%) had high estimated prevalence of asthma. These areas are likely to benefit from asthma outreach that considers both social and environmental risk factors. Fifteen out of the 137 CTs (11%) not only had higher estimated prevalence of asthma but also a high burden of COVID-19. Further research in these areas may help to elucidate any common social determinants of health that underlie both asthma and COVID-19 burdens, as well as better clarify the possible role of the environment as related to the COVID-19 burden in Louisiana.

## Introduction

In 2020, the COVID-19 pandemic laid bare the major health impacts of social and environmental vulnerability, by taking a heavy toll on populations suffering such vulnerabilities. It highlighted how a novel virus does not affect everyone equally; it exploits underlying health conditions that are more prevalent in vulnerable populations [1, 2]. According to the US Centers for Disease Control and Prevention (CDC), asthma is one such respiratory health condition that might be a risk factor for severe illness with COVID-19. It is not only more prevalent among African-Americans and people living in poverty (a population disproportionately affected by COVID-19 [3, 4]), but asthma is also heavily influenced by environmental factors [5-7]. As such, it is particularly important to tackle this issue in Environmental Justice (EJ) communities, which lack equitable access to protection from environmental and health hazards, and where socioeconomic, demographic and environmental stressors overlap to create environmental health disparities. For example, many people living in EJ communities do not have access to affordable healthy housing, and thus may be exposed daily to asthma triggers in the home [8]. At a time when people are spending more time in their homes than ever, many asthmatics living in unhealthy homes may experience worsening of their symptoms. People living in communities that have outdoor air concerns may experience prolonged adverse outdoor exposures to known asthma triggers, such as ozone and Particulate Matter less than 2.5 µm in diameter ($PM_{2.5}$), if they spend more time socializing outdoors in keeping with COVID-19 safety protocols. These, in turn, could translate into poor outcomes if they contracted COVID-19.

The need to tackle this vicious cycle was recently highlighted by a study that showed respiratory health risks from hazardous air pollutants and high incidence of COVID-19 converged in several counties across the southeastern United States [9]. These areas tended to have higher non-

Hispanic black populations, socioeconomically disadvantaged persons, people with disabilities

and people without health insurance. This county-level study served as an important starting

point for further exploring the impact of COVID-19 in EJ communities at a finer geographic

resolution on a state-by-state basis [9]. Since Louisiana (LA) was one of the states where a

majority of counties (called 'parishes' in LA) seemed to fall under the "high-high" category for

both respiratory hazard risk and COVID-19 risk, here we have undertaken a census tract (CT)-

level analysis to assess any correlation between social and environmental vulnerability, COVID-

19, and asthma. We have further identified several EJ communities with a high burden of asthma

and COVID-19 where mitigation measures may be targeted, as well as where further studies may

be conducted to elucidate the environment's role on COVID-19 outcomes in Louisiana.

**Methods**

<u>Indicators of Social Vulnerability</u>

To assess social vulnerability, we used the 2018 Social Vulnerable Index (SVI) derived by the

CDC Agency for Toxic Substances and Disease Registry (ATSDR) [10]. Along with the Overall

SVI, we considered the four specific SVI themes outlined by CDC/ATSDR – socioeconomic

status (Theme 1); household composition and disability (Theme 2); minority status and language

(Theme 3); and housing type and transportation (Theme 4). We also applied the US Census

Bureau's 2018 American Community Survey (ACS) data for percent population without health

insurance, below poverty level, unemployed, without a high school diploma, disabled, minority

(all persons except non-Hispanic white), age 65+, age 17 and under, and population (age 5+)

who speak English "less than well". We also took into account household level information on

those without access to a vehicle, single parent households and households with more people

than rooms (crowded homes). Detailed information on each of these variable is available from

CDC/ATSDR

(https://www.atsdr.cdc.gov/placeandhealth/svi/documentation/SVI_documentation_2018.html).

All datasets for social vulnerability **(Table 1)** were considered at the census tract level, and the

SVI scores classified into five categories based on the CT's ranking (percentile) against all CTs

in the state: 0-0.20; 0.21-0.40; 0.41-0.60; 0.61-0.80; 0.81-1.00. All mapping and spatial analyses

were performed in ESRI ArcGIS 10.8.1.

**Table 1. Summary of datasets assessed in this study.** All datasets were mapped at the census

tract level.

| Topic | Indicator (Measure) | Data Source | Data Year(s) |
|---|---|---|---|
| Social Vulnerability | Social Vulnerability Index and related indicators from ACS 2018 | CDC/ATSDR | 2018 |
| Environmental Vulnerability | NATA Respiratory Hazard Index | EPA | 2014 |
| | $PM_{2.5}$ level (average annual concentration in $\mu g/m^3$) | EPA | 2016 |
| | Ozone level (summer seasonal average of daily maximum 8-hour concentration in air in parts per billion) | EPA | 2016 |
| | Indoor mold concerns reported to IEQES program (average annual number of calls) | LDH | 2017-2019 |
| COVID-19 | Cumulative COVID-19 incidence rate at 3-, 6-, 9- and 12-month time points | LDH | Mar 2020- Mar 2021 |
| Asthma | Asthma Hospitalization (average annual crude rate, where asthma was a primary diagnosis among hospitalization cases) | LDH | 2017-2019 |
| | Estimated Asthma Prevalence (average annual crude rate, where asthma was any diagnosis among hospitalization cases) | LDH | 2017-2019 |

In terms of environmental datasets (outdoor air), we downloaded the following from US

Environmental Protection Agency's (EPA's) EJSCREEN tool version 2020

(https://www.epa.gov/ejscreen): National Air Toxics Assessment (NATA) Respiratory Hazard

Index (RHI), $PM_{2.5}$ levels and Ozone levels. The EJSCREEN technical documentation (available

from: https://www.epa.gov/sites/production/files/2021-

04/documents/ejscreen_technical_document.pdf) provides detailed definitions for each of these

measures. Briefly: The RHI is the ratio of the exposure concentration of Hazardous Air

Pollutants (HAPs) to a health-based reference concentration. The most recent year of RHI data

available (2014) take into account 181 HAPs (including diesel particulate matter), for which both

emissions and health-related toxicity data were available. In order to derive the RHI, US EPA

uses several modeling techniques to estimate both the concentration of the HAPs as well as the

risk of exposure and downstream respiratory health risks. Given that the susceptibility to air

pollution cannot be reliably estimated, these models take into consideration differences in

emissions and meteorology (which affects the concentration of pollutants in ambient air) and the

differences in the location of individuals (which affects exposure) among CTs. Thus, applying

these models then allow us to identify geographic patterns and assess relative risks at the CT-

level, with the caveat that all calculations become more uncertain with smaller geographies.

Complete details on RHI modeling and calculation methods are available from the 2014 NATA

Technical Support Document [11]. $PM_{2.5}$ levels refer to average annual concentrations in $\mu g/m^3$,

as calculated by US EPA using a fusion of modeled and monitored data (for methodological

details on air quality modeling, see [12]). Ozone levels were similarly calculated using a

combination of modeled and monitored data by the EPA [12], using the summer seasonal

average of daily maximum 8-hour concentration in air in parts per billion. The most recent data

available for both $PM_{2.5}$ and Ozone levels (2016) was used for this study.

Finally, to account for not only outdoor but also Indoor Environmental Quality (IEQ) concerns,

we surveyed internal data from the Louisiana Department of Health's (LDH's) Indoor

Environmental Quality Education Service (IEQES). Since 2000, LDH's IEQES has been

providing guidance to Louisiana homeowners and renters on a variety of IEQ concerns reported

to the program. The majority of these calls (~75%) are related to mold in residential settings;

however, the IEQES program provides guidance on all types of IEQ issues (including odors,

pests, radon, chemical exposures, etc.) in both private and public buildings (such as homes,

schools, places of worship, etc.). For this study, we have aggregated the three most recent years

of data available through IEQES (2017-2019) and calculated the average annual number of mold

concerns reported to the program as an indicator of IEQ concerns in an area. Mold concerns were

chosen for this study over all IEQ concerns since they comprised the majority (86%) of the calls

received. Further, mold concerns were most often associated with respiratory complaints (the

health focus of this study), as reported by the callers; of the 785 calls that reported respiratory

complaints, 708 (90%) were associated with mold. Mold exposure is also recognized as a

possible environmental trigger for asthma [13].

All environmental datasets (**Table 1**) were considered at the census tract level. For all datasets,

we used EJSCREEN's visualization method and mapped the RHI, $PM_{2.5}$, Ozone and IEQ

concerns as a percentile score relative to all CTs in the state (ranked using the

PERCENTRANK.INC function in MS Excel 2016). A choropleth map was then created using

progressively darker colors for five categories: 0-20th percentile; 21st-40th percentile; 41st-60th

percentile; 61st-80th percentile and 81st-100th percentile.

<u>Health Indicators (COVID-19 and Asthma)</u>

Total number of COVID-19 cases by CT were downloaded from LDH's COVID-19 dashboard

(https://ldh.la.gov/coronavirus/) and cumulative incidence rates calculated at four time points (3,

6, 9 and 12 months into the pandemic in LA) by dividing the total number of cases at each time

point by the ACS 2018 population of the CT. Of the 1145 CTs in LA, data were available for

approximately 1034 CTs at the 3-month time point, 1038 at the 6-month time point, and 1039 at

the 9- and 12-month time points. Case counts are not provided for CTs with a 2018 population of

less than 800 to safeguard privacy.

Since asthma surveillance data were not available at the census tract level for most of Louisiana,

we estimated asthma burden using the inpatient discharge data available through LDH. To

minimize the need for suppression, inpatient discharge data was aggregated for the three most

recent years available (2017-2019) and average annual crude rates were calculated for cases

where asthma (ICD-10 code J45) was the primary diagnosis, as well as where asthma was any

diagnosis. While the former indicated the level of asthma exacerbation in a community that

required medical intervention (an indicator for uncontrolled asthma), the latter served as an

estimate of asthma prevalence in the community. In order to calculate these, we divided the total

number of cases for each CT (2017-2019) by the total population of the CT during that period,

and then divided that value by 3 to yield the average annual crude rates.

Similar to the environmental vulnerability datasets, percentile scores were also calculated for

COVID-19 incidence and asthma data **(Table 1)**, which were then classified in five categories.

<u>Correlation analysis</u>

Spearman's Rank Correlation was utilized to analyze the correlation between various social and

environmental vulnerability factors, COVID-19 incidence, and the measures of asthma risk by

CT. This was performed by first ranking the values in each dataset using RANK.AVG function

in MS Excel 2016, followed by applying the PEARSON function to compare two datasets.

Significance was set at alpha less than 0.05 ($\alpha < 0.05$), with degrees of freedom (df) equal to two

less than the total number of data points represented in both datasets. Only CTs for which all data

were available (1035 out of 1145 total CTs in LA) were used for this analysis. Spearman's *rho*

($\rho$) was used to identify the direction and strength of correlation. If the p-value was less than

0.05 ($p < 0.05$), strength of correlation was defined as follows: absolute value of $\rho$ = 0.01-0.30 =

weakly correlated; 0.31-0.70 = moderately correlated; and 0.71-1.00 = strongly correlated. The

direction of correlation was determined by the sign (positive or negative) on $\rho$ value.

**Results**

<u>Relationship between Social and Environmental Vulnerability, COVID-19, and Asthma</u>

CTs were defined as 'high' for social vulnerability if the Overall SVI was greater than or equal

to 0.75, and 'high' for environmental vulnerability if the RHI, $PM_{2.5}$ level or Ozone level was at

or over the 75[th] percentile for all CTs in the state. When compared to all CTs, high SVI was

associated with higher average cumulative COVID-19 incidence rates early in the pandemic (3-

month and 6-month time points). $PM_{2.5}$ was associated with lower COVID-19 incidence rates at

the 3-month time point, but at 12 months, it was associated with higher COVID-19 incidence.

Ozone levels were associated with higher COVID-19 incidence all time points (except at 6

months), whereas RHI showed the opposite pattern, being associated significantly lower

COVID-19 incidence at time points except at 6 months **(Figure 1A)**. Additionally, the COVID-

19 incidence rates were higher in areas of high SVI + high Ozone compared to areas with only

high SVI. This additive effect over SVI was not observed from any other environmental factor.

With regard to average hospitalization rate for asthma and estimated asthma prevalence, higher

rates of both were associated with high SVI and high Ozone levels **(Figure 1B)**. High SVI

appeared to have an additive effect on high Ozone, with areas of high SVI + high Ozone having

higher asthma hospitalization rate and estimated prevalence than areas that just had high Ozone.

Taken together, these descriptive statistics indicated a possible relationship between social and

environmental vulnerability, COVID-19, and asthma.

**Figure 1. COVID-19 and Asthma in areas with high social and/or environmental**

**vulnerability. (A)** High SVI was associated with higher average cumulative COVID-19

incidence rates early in the pandemic, whereas $PM_{2.5}$ was associated with higher rates later in the

pandemic. Cumulative COVID-19 incidence was elevated in areas of high Ozone at all points

during the pandemic. **(B)** Asthma hospitalization as well as estimated asthma prevalence

followed a similar pattern with higher rates of both being associated with high SVI and high

Ozone. Error bars indicate 95% confidence intervals; asterisks (*) indicate areas that are

significantly higher than "All CTs" and pounds (#) indicate areas significantly lower than "All

CTs" ($p < 0.05$; T-test).

These possible relationships were then further explored in a Spearman rank correlation **(Table 2-**

**3)**. Cumulative COVID-19 incidence rate at the 3-month time point was moderately correlated

with the Ozone levels ($\rho = 0.32$; $p < 0.0001$), SVI for minority status and language barriers

($\rho$=0.44; p<0.0001), asthma hospitalization ($\rho$=0.32; p<0.0001), and estimated prevalence of

asthma ($\rho$=0.50; p<0.0001). Asthma hospitalization and estimated prevalence were moderately

correlated with SVI for socioeconomic status, SVI for minority and language barriers and

Overall SVI ($\rho$=0.30-0.37; p<0.0001). Since one of the strongest correlation coefficients was

observed between the estimated prevalence of asthma and cumulative COVID-19 incidence at

the 3-month time point, we conducted a second Spearman rank correlation to test whether similar

social vulnerability measures may underlie both health indicators **(Table 3)**. According to the

results, positive correlations of moderate strength were observed between both health indicators

and percent minority population ($\rho$=0.51-0.52; p<0.0001) and percent households without a

vehicle ($\rho$=0.37-0.44; p<0.0001).

**Table 2. Spearman's rho (correlation coefficient) for social and environmental**

**vulnerability, asthma, and COVID-19.** With regards to COVID-19, strongest correlation was

observed between the cumulative incidence during the first quarter of the pandemic in LA (Mar-

Jun 2021) and the estimated prevalence of asthma, SVI due to minority status and language

barriers, asthma hospitalization crude rate, and the percentile score for ozone levels. With

regards to asthma, SVI due to socioeconomic status, minority status and language barriers, and

overall SVI were also significantly correlated with both asthma hospitalization rate as well as the

estimated prevalence.

|  |  | Hosp. Asthma | EPV Asthma | CV19 (3m) | CV19 (6m) | CV19 (9m) | CV19 (12m) |
|---|---|---|---|---|---|---|---|
| **Environmental Factors** | **RHI** | -0.02 | 0.04 | -0.12 | -0.05 | -0.14 | -0.14 |
|  | **Ozone** | 0.21 | 0.29 | **0.32** | 0.03 | 0.09 | 0.11 |
|  | **PM2.5** | 0.08 | 0.02 | -0.07 | -0.02 | 0.12 | 0.08 |
| **Social Vulnerability** | **SVI_SES** | **0.36** | **0.34** | 0.22 | 0.15 | -0.16 | -0.24 |
|  | **SVI_HC** | 0.26 | 0.26 | 0.09 | 0.16 | 0.01 | -0.02 |
|  | **SVI_ML** | **0.30** | **0.37** | **0.44** | 0.29 | -0.01 | -0.06 |
|  | **SVI_HT** | 0.13 | 0.09 | 0.11 | 0.19 | 0.02 | -0.07 |
|  | **SVI_OV** | **0.37** | **0.36** | 0.27 | 0.26 | -0.06 | -0.15 |
| **Asthma** | **Hosp. Asthma** |  | **0.61** | **0.32** | 0.21 | 0.02 | 0.01 |
|  | **EPV Asthma** | **0.61** |  | **0.50** | 0.28 | 0.03 | 0.07 |
| **COVID-19 (cumulative)** | **CV19 (3m)** | **0.32** | **0.50** |  |  |  |  |
|  | **CV19 (6m)** | 0.21 | 0.28 |  |  |  |  |
|  | **CV19 (9m)** | 0.02 | 0.03 |  |  |  |  |
|  | **CV19 (12m)** | 0.01 | 0.07 |  |  |  |  |

*Abbreviations:* RHI = NATA Respiratory Hazard Index; Ozone = Ozone levels; PM2.5 = $PM_{2.5}$ levels; SVI_SOCEC = Social Vulnerability Index (due to socioeconomic factors); SVI_HC = Social Vulnerability Index (due to household composition); SVI_ML = Social Vulnerability Index (due to minority status and language barriers); SVI_HT = Social Vulnerability Index (due to housing and transportation factors); SVI_OV = overall Social Vulnerability Index; Hosp. Asthma = Asthma hospitalization crude rate; EPV Asthma = Estimated prevalence of asthma; CV19 = Cumulative COVID-19 incidence at 3, 6, 9 and 12 months

| *Legend based on rho value:* | *Strength and direction of correlation:* |
|---|---|
| >0.7 to 1.0 | Strongly positive correlation |
| >0.3 to 0.7 | Moderately positive correlation |
| >0 to 0.3 | Weakly positive correlation |
| *Rho*=0 or p ≥ 0.05 | No significant correlation |
| <0 to -0.3 | Weakly negative correlation |
| < -0.3 to -0.7 | Moderately negative correlation |
| < -0.7 to -1.0 | Strongly negative correlation |

**Table 3. Spearman's rho (correlation coefficient) for specific social vulnerability factors, asthma and COVID-19.** Statistically significant correlations were observed between several vulnerability factors and cumulative COVID-19 incidence up to June 2020 (CV19 [3m]) and/or estimated prevalence of asthma (EPV Asthma). However, the factors that showed moderately

212  strong correlation coefficients with both health indicators were percent minority populations and

213  percent households without a vehicle.

| | EPV Asthma | CV19 (3m) | *Abbreviations:* PP/H/U = Percent Population/Household/Housing Unit |
|---|---|---|---|
| **PP_AGE17** | 0.01 | -0.06 | AGE17 = Age 17 and under; |
| **PP_UNINSUR** | 0.23 | 0.18 | UNINSUR = No health insurance; |
| **PP_POV** | **0.31** | 0.23 | POV = Below poverty; |
| **PP_UNEMP** | 0.29 | 0.16 | UNEMP = Unemployed; |
| **PP_NOHSDP** | 0.28 | 0.17 | NOHSDP = No high school diploma; |
| **PP_AGE65** | -0.01 | 0.01 | AGE65 = Age 65 and over; |
| **PP_DISABL** | 0.29 | 0.05 | DISABL = Disabled; |
| **PP_MINRTY** | **0.51** | **0.52** | MINRTY = Minority; |
| **PP_LIMENG** | 0.10 | 0.18 | LIMENG = Limited English; |
| **PH_SNGPNT** | 0.27 | 0.20 | SNGPNT = Single parent households; |
| **PU_CROWD** | 0.06 | 0.01 | CROWD = Crowded housing units; |
| **PH_NOVEH** | **0.44** | **0.37** | NOVEH = Households without vehicle; |
| **PP_GROUPQ** | -0.06 | 0.07 | GROUPQ = Living in group quarter; |
| *Legend based on rho value:* | | | *Strength and direction of correlation:* |
| **>0.7 to 1.0** | | | Strongly positive correlation |
| **>0.3 to 0.7** | | | Moderately positive correlation |
| >0 to 0.3 | | | Weakly positive correlation |
| *Rho*=0 or p ≥ 0.05 | | | No significant correlation |
| <0 to -0.3 | | | Weakly negative correlation |
| **< -0.3 to -0.7** | | | Moderately negative correlation |
| **< -0.7 to -1.0** | | | Strongly negative correlation |

<u>Identifying EJ Communities of Concern with respect to COVID-19 and Asthma</u>

Surveying the spatial distribution of statewide social vulnerability data showed that while most

LA parishes (59/64) had one or more CTs with high overall SVI, i.e. SVI>=0.75 **(Figure 2A)**,

environmental vulnerability factors were more likely to be high (at or above the 75th percentile)

in certain specific parts of the state. Ozone levels, for instance, were higher in the north,

northwest and southeast **(Figure 2B)**; PM$_{2.5}$ levels were higher in the north, northwest and east

**(Figure 2C)**; and RHI were higher in the northwest, southwest and eastern parts of the state

(map not shown). In terms of IEQ issues, at least one mold concern was reported from 542 CTs

in LA **(Figure 2D)**, 301 of which (55%) also had high RHI, high PM$_{2.5}$ or high Ozone levels.

These CTs, where both indoor and outdoor environments may be of concern, were present in

pockets but distributed across the state. A similar inspection of the health measures (asthma and

COVID-19) revealed that they were most often noted as 'high' (at or above the 75[th] percentile

for the state) in northwestern and southeastern parts of the state **(Figure 2E-F)**.

**Figure 2. Spatial analysis of social and environmental vulnerability, asthma and COVID-**

**19**. Darker colors for each measure indicate higher percentile ranks. Most counties in LA had

census tracts with SVI **(A)** above the median. Ozone **(B)** and PM$_{2.5}$ **(C)** were higher in the

northwest and southeastern parts of the state. Indoor environmental quality concerns (mostly in

the form of mold complaints) were also reported from all over the state **(D)**. Estimated

prevalence of asthma **(E)** was often above the median in census tracts located in the

northwestern, western, southern and southeastern parishes. At the 12-month time point,

cumulative COVID-19 incidence rates **(F)** were higher in the northern, northwestern and

southeastern parts of the state.

Next, we attempted to identify EJ communities in Louisiana using the social and environmental

vulnerability metrics. We used Overall SVI from CDC/ATSDR as the overarching metric for

social vulnerability. For environmental vulnerability, we chose RHI, PM$_{2.5}$ and Ozone levels.

IEQ reports were not considered because of the sparse nature of the data. Thus, we intersected

Overall SVI, RHI, PM$_{2.5}$ and Ozone datasets **(Figure 3A)**, and defined a CT as an area of EJ

concern if along with high SVI, it also had either high RHI, high PM$_{2.5}$ or high Ozone levels.

Mapping these areas revealed that the 137 CTs that fit these criteria were mostly located in the

major urban centers with a few spread across the more rural areas of the state **(Figure 3B)**.

**Figure 3. Identifying census tracts with EJ concerns.** Census Tracts were defined as being of

EJ concern if the Overall SVI was at or above the 75$^{th}$ percentile, along with at least one of the

environmental variables (RHI, PM$_{2.5}$ or Ozone) at or above the 75$^{th}$ percentile for the state **(A).**

Map shows the number of the CTs of EJ concern by Parish **(B).** Most of them were located in the

major urban areas, with a few spread across the rural areas of the state.

Finally, we surveyed the health metrics of these EJ CTs and sought to identify areas of high

asthma and COVID-19 burden **(Figure 4A)**. Out of the 137 EJ CTs, 75 (55%) had high

estimated asthma prevalence (at or above the 75$^{th}$ percentile for the state), 28 (20%) had a high

cumulative COVID-19 rates at the 12-month time point, and 15 CTs (11%) had a high burden of

both asthma and COVID-19. The 75 EJ CTs with a high burden of asthma were mostly located

in the major urban centers of the state, with a few in the more rural areas **(Figure 4B)**. A similar

pattern was observed for the 15 EJ CTs with a high burden of both asthma and COVID-19, with

a third of them being located in the northwestern part of the state in Caddo Parish **(Figure 4C).**

**Figure 4. Identifying EJ communities with high COVID-19 and asthma burdens. (A)** Of the

137 census tracts that were deemed of EJ concern, 75 (55%) had estimated prevalence of asthma

at or above the 75th percentile for the state. The majority of these CTs were located in the major

urban areas, with others spread across the more rural areas of the state **(B)**. A similar pattern was

observed for CTs that were of concern for EJ, asthma as well as COVID-19. About a third of the

15 EJ CTs deemed of more asthma and COVID-19 concerns were located in the northwestern

part of the State in Caddo Parish **(C).**

**Discussion**

Over the past year, COVID-19 has taken a disproportionate toll on minority populations across

the country, shining a spotlight on the health, socioeconomic and environmental disparities faced

by them. According to the CDC, asthma is one of the underlying conditions that, when moderate

to severe, may be a risk factor for severe illness with COVID-19. It is also a respiratory health

condition that disproportionately affects racial and ethnic minorities [14]. Here we show that

these trends in health disparities, previously assessed nationally and/or at the county level [3, 9,

14, 15], hold true at the census tract level in Louisiana. Specifically, our data show a moderately

strong positive correlation between SVI due to minority status/language barrier and three health

data variables: asthma hospitalization; estimated asthma prevalence; and cumulative COVID-19

incidence at 3 months **(Table 2)**. Interestingly, SVI measures were either negatively or not

significantly correlated COVID-19 incidence at the 9- and 12-month time points, indicating that

social vulnerability factors may have played a greater role in COVID-19 spread early in the

pandemic, but may have been of diminishing importance as the pandemic wore on **(Figure 1,**

**Table 2)**.

Regardless, an assessment of specific social vulnerability factors, asthma and COVID-19 is

likely to reveal important characteristics of minority populations that have contributed to long-

standing health disparities along racial and socioeconomic lines. One such characteristic may be

lack of access to a vehicle. (**Table 3**). This may be of particular importance in Louisiana, which

lacks extensive public transport infrastructure in most places, and may be a factor that prevents

people from accessing areas away from their immediate vicinity to acquire healthcare services,

or other commodities conducive to health (such as healthy foods, if living in a food desert). In

urban areas, where public transport is available, people without vehicles may not be able to

socially distance in the confined space while using public transportation. Indeed, a connection

between higher subway ridership and COVID-19 infection risk was observed in a previous study

in New York [16], and has been suggested previously in an earlier sub-county study in Louisiana

measuring Area Deprivation Index and COVID-19 risk [17]. Thus, the possible impact of not

having access to a vehicle (among other vulnerability factors) on chronic health conditions, as

well as COVID-19 spread, warrants further investigation, especially in Louisiana's minority

communities that face other socioeconomic and environmental stressors as well.

Minorities, especially African-Americans, are also more likely to live in EJ communities and be

exposed to environmental pollutants that can influence their health [18]. Asthma is a respiratory

condition that has several known environmental triggers such as ozone and particulate matter.

Various studies have also shown that there may be a connection between air pollutants affecting

respiratory health (mainly, particulate matter) and COVID-19 incidence as well as mortality [9,

15, 19, 20]. In our study, however, we found higher rates of COVID-19 incidence, asthma

hospitalizations and estimated asthma prevalence in CTs with higher ozone levels (**Figure 1,**

**Table 2**). According to the US EPA, long-term exposure to ozone pollution can damage the

lungs and make it likely to acquire respiratory tract infections [21, 22]. Ozone is also known to

aggravate asthma and may be one of the factors that leads to developing asthma [23-26]. Of note,

a 2018 study showed that exposure to even low levels of ozone (below the limit set by the

National Ambient Air Quality Standards) had adverse effects on the respiratory health of

African-American children with poorly-controlled asthma [27]. Since ground-level ozone is

formed from precursors present in traffic exhaust and industrial emissions, it is possible that

ozone levels may be higher in the more urban, industrialized areas of Louisiana than the more

rural regions, as has been previously documented by CDC for the entire United States [28]. If

this assumption holds true, it would explain our results indicating that CTs with EJ concerns are

largely located in the more urban areas of the state **(Figure 3)**. Thus, any possible contribution of

ozone to asthma prevalence, asthma exacerbation as well as the COVID-19 burden in Louisiana

bears investigation in the areas of concern.

Since about 60% of the CTs with EJ concerns related to outdoor air also reported IEQ concerns

(primarily, mold in homes), the 116 tracts identified to be of concern with regards to asthma

**(Figure 4A)** may benefit from educational outreach that includes both clinical and environmental

management of asthma [29-32]. To accommodate individuals with asthma who may not have

access to vehicles for easy transportation, we propose either a local community-based or a home-

based education model that ideally incorporates an environmental evaluation of their home. Such

initiatives are not only likely to reduce respiratory health disparities in LA, but they may also

open the door to further research that establish the factors underlying both COVID-19 incidence

and asthma prevalence **(Table 2)**, the potential environmental impact on both health indicators,

as well as any increased burden of asthma due to COVID-19 in EJ communities. Considering the

fact that much of the adverse impact from COVID-19 on vulnerable populations may have been

through pre-existing disparities in chronic health outcomes [1, 2], such initiatives focusing on

improving health and environmental equity are likely to yield long-term gains.

Our study is the first to our knowledge that evaluates the intersection of social and environmental

vulnerability, COVID-19 and asthma statewide at the census tract level in LA. Thus, it represents

an important milestone in identifying and understanding specific communities that are at the

crossroads of environmental justice issues as well as health disparities. However, there are some

important limitations that should be considered. First, it is an ecological study, and therefore,

while providing useful observational data to help generate hypotheses, it cannot be used to test

any hypothesis related to population health. Second, though we chose the most current

environmental vulnerability metrics available from EPA EJSCREEN, we cannot be certain that

they model current environmental conditions. At the same, much of the impact of air pollution

on respiratory health is due to long-term exposure [23-27], not just short-term; therefore, it is still

instructive to consider these data even if it is a few years old as of 2021. Third, there are multiple

different models for estimating the burden of air pollution; all have their own pros and cons, and

all are relatively more unstable at smaller geographic levels (such as sub-county geographies).

The environmental vulnerability metrics and visualization scheme presented here were based on

those used by a well-established resource to study environmental justice concerns (EJSCREEN).

However, we cannot rule out that using different models of air pollution estimates or applying

the same models but at a different spatial resolution may yield different results. Finally, due to

the lack of outpatient asthma surveillance data at the sub-county level, we used inpatient

discharge data to estimate the burden of asthma hospitalization as well as estimated asthma

prevalence. These data are subject to various limitations that result from incomplete reporting,

changes in hospital administration (such as closures or merges), inequities in healthcare access,

choice of diagnosis codes, etc. Complete metadata related to asthma hospitalizations can be

found through the LDH website (https://ldh.la.gov/assets/oph/Center-EH/envepi/EPHT/Infotab-

Metadata/Asthma_5-1-2019.pdf).

**Acknowledgements**

We extend our sincere thanks to the LDH Section of Infectious Disease Epidemiology and

Bureau of Health Informatics for their tireless efforts and dedication at gathering and publishing

reliable health data for the State of Louisiana, while simultaneously tackling the COVID-19

pandemic. Authors in this study have been supported through the CDC Environmental Public

Health Tracking Cooperative Agreement (AB, KF, AW); the EPA State Environmental Justice

Cooperative Agreement (AB); and the State General Fund (CM, CS, SS, KA).

**References**

1.  Alcendor DJ. Racial Disparities-Associated COVID-19 mortality among minority populations in the US. J Clin Med. 2020; 9:2442.

2.  Moore JT, Pilkington W, Kumar D. Diseases with health disparities as drivers of COVID-19 outcome. J Cell Mol Med. 2020; 24: 11038-11045.

3.  Cyrus E, Clarke R, Hadley D, Bursac Z, Trepka MJ, Dévieux JG, et al. The impact of COVID-19 on African American communities in the United States. Health Equity. 2020; **4:**476-483.

4.  Abedi V, Olulana O, Avula V, Chaudhary D, Khan A, Shahjouei S, et al. Racial, economic, and health inequality and COVID-19 infection in the United States. J Racial Ethn Health Disparities. 2020; **1**:1–11.

5.  Fitzpatrick AM, Gillespie SE, Mauger DT, Phillips BR, Bleecker ER, Israel E, et al.

Racial disparities in asthma-related health care use in the National Heart, Lung, and

Blood Institute's Severe Asthma Research Program. J Allergy Clin Immunol. 2019;

143:2052-2061.

6.  Moorman JE, Akinbami LJ, Bailey CM, Zahran HS, King ME, Johnson CA, et al.

National Surveillance of Asthma: United States, 2001–2010. National Center for Health

Statistics. Vital Health Stat. 2012; **3**. Available from:

https://www.cdc.gov/nchs/data/series/sr_03/sr03_035.pdf.

7.  Nishimura KK, Galanter JM, Roth LA, Oh SS, Thakur N, Nguyen EA, et al. Early-life air

pollution and asthma risk in minority children: The GALA II and SAGE II studies. Am J

Respir Crit Care Med. 2013; **188**:309-318.

8.  Pacheco CM, Ciaccio CE, Nazir N, Daley CM, DiDonna A, Choi WS, et al. Homes of

low-income minority families with asthmatic children have increased condition issues.

Allergy Asthma Proc. 2014; **35**:467-74.

9.  Chakraborty J. Convergence of COVID-19 and chronic air pollution risks: Racial/ethnic

and socioeconomic inequities in the U.S. Environ Res. 2021; **193**:110586.

10.  Centers for Disease Control and Prevention/ Agency for Toxic Substances and Disease

Registry/ Geospatial Research, Analysis, and Services Program. CDC Social

Vulnerability Index 2018 Database, Louisiana [cited 2021 May 17]. Available from:

https://www.atsdr.cdc.gov/placeandhealth/svi/data_documentation_download.html

11.  Office of Air Quality Planning and Standards, Environmental Protection Agency.

Technical Support Document: EPA's 2014 National Air Toxics Assessment [cited 2021

May 17]. Available from: https://www.epa.gov/sites/production/files/2018-

09/documents/2014_nata_technical_support_document.pdf.

12. Reff A, Phillips S, Eyth A, Mintz D. Bayesian space-time downscaling fusion model

(downscaler)–derived estimates of air quality for 2011. U.S. Environmental Protection

Agency, Office of Air Quality Planning and Standards, Air Quality Assessment Division.

2015; EPA/454/S-15/001. Available from:

https://nepis.epa.gov/Exe/ZyPDF.cgi?Dockey=P100OORX.PDF.

13. Zhang Z, Reponen T, Hershey GK. Fungal exposure and asthma: IgE and non-IgE-

mediated mechanisms. Curr Allergy Asthma Rep. 2016; **16**:86.

14. Centers for Disease Control and Prevention. 2018 National Health Interview Survey

(NHIS) Data [cited 2021 May 17]. Available from:

https://www.cdc.gov/asthma/nhis/2018/data.htm

15. Terrel K and James W. Racial disparities in air pollution burden and COVID-19 deaths in

Louisiana, USA, in the context of long-term changes in fine particulate pollution.

Environ. Justice. 2020; doi: https://doi.org/10.1089/env.2020.0021.

16. Carrion D, Colicino E, Pedretti NF, Arfer KB, Rush J, DeFelice N, et al. Assessing

capacity to social distance and neighborhood-level health disparities during the COVID-

19 pandemic. 2020; medRxiv. doi: 10.1101/2020.06.02.20120790.

17. Madhav KC, Oral E, Straif-Bourgeois S, Rung AL, Peters ES. The effect of area

deprivation on COVID-19 risk in Louisiana. PLoS One. 2020; **15**:e0243028.

18. Bullard RD. Race and environmental justice in the United States. Yale J Int Law. 1993;

Available from: https://digitalcommons.law.yale.edu/yjil/vol18/iss1/12.

19. Wu X, Nethery RC, Sabath MB, Braun D, Dominici F. Air pollution and COVID-19 mortality in the United States: Strengths and limitations of an ecological regression analysis. Sci Adv. 2020; **6:**eabd4049.

20. Hendryx M, Luo J. COVID-19 prevalence and fatality rates in association with air pollution emission concentrations and emission sources. Environ. Pollut. 2020; **265**(Part A):115126.

21. Al-Hegelan M, Tighe RM, Castillo C, Hollingsworth JW. Ambient ozone and pulmonary innate immunity. Immunol Res. 2011; **49:** 173–191.

22. Kesic MJ, Meyer M, Bauer R, Jaspers I. Exposure to ozone modulates human airway protease/antiprotease balance contributing to increased Influenza A infection. PLoS One. 2012; **7**: e35108.

23. To T, Zhu J, Stieb D, Gray N, Fong I, Pinault L, et al. Early life exposure to air pollution and incidence of childhood asthma, allergic rhinitis and eczema. Eur Respir J. 2020; **55**:1900913.

24. Chatkin J, Correa L, Santos U. External environmental pollution as a risk factor for asthma. Clinic Rev Allerg Immunol. 2021; Available from: https://doi.org/10.1007/s12016-020-08830-5.

25. Tiotiu AI, Novakova P, Nedeva D, Chong-Neto HJ, Novakova S, Steiropoulos P, et al. Impact of air pollution on asthma outcomes. Int J Environ Res Public Health. 2020; **17**: 6212.

26. Anenberg SC, Henze DK, Tinney V, Kinney PL, Raich W, Fann N, et al. Estimates of the global burden of ambient $PM_{2.5}$, ozone, and NO2 on asthma incidence and emergency room visits. Environ Health Perspect. 2018; **126**: 107004.

27. Hernandez M, Dhingra R, Burbank A, Todorich K, Loughlin C, Frye M, et al. Low level

ozone has both respiratory and systemic effects in African-American adolescents with

asthma despite asthma controller therapy. J Allergy Clin Immunol. 2018; **142**:1974-1944.

28. Strosnider H, Kennedy C, Monti M, Yip F. Rural and urban differences in air quality,

2008–2012, and community drinking water quality, 2010–2015 — United States.

MMWR Surveill Summ. 2017; **66**:1–10.

29. Marshall ET, Guo J, Flood E, Sandel MT, Sadof MD, Zotter JM. Home visits for children

with asthma reduce Medicaid costs. Prev Chronic Dis. 2020; **17**:190288.

30. Campbell JD, Brooks M, Hosokawa P, Robinson J, Song L, Krieger J. Community

Health Worker home visits for Medicaid-enrolled children with asthma: Effects on

asthma outcomes and costs. Am J Public Health. 2015; **105**:2366-72.

31. National Center for Healthy Housing. Case Studies: The benefits of home visits for

children with asthma. Available from: https://nchh.org/resource-library/case-studies_the-

benefits-of-home-visits-for-children-with-asthma.pdf

32. Bryant-Stephens T, Reed-Wells S, Canales M, Perez L, Rogers M, Localio AR, et al.

Home visits are needed to address asthma health disparities in adults. J Allergy Clin

Immunol. 2016; **138**:1526-1530.

**A)** Overall SVI

**B)** Ozone Levels

**C)** PM2.5 Levels

**D)** Indoor Mold Concerns

**E)** Est. Asthma Prevalence

**F)** COVID-19 (12 months)

**A)** Intersection of Social & Environmental Vulnerability

High RHI
High PM2.5
High Ozone
High SVI

**B)** No. of CTs of EJ Concern by Parish

Total # CT
137

Caddo

19 5 3 2 1
3 3
2 1
3 1
1
1

East Baton Rouge

3
5
3

123
1 2 2

9
Calcasieu

1 1

Orleans
24
7 Jefferson

1
2
1
2

**A)**

EJ Areas          High Estimated Asthma Prevalence

49    60    154

13    56

High Cumulative COVID-19 incidence (12 months)

**B)**

No. of EJ CTs by Parish with high burden of asthma

**C)**

No. of EJ CTs by Parish with high burden of asthma and COVID-19