Phase 1 Safety and Pharmacokinetics Studies of BRII-196 and BRII-198,
SARS-CoV-2 Spike-Targeting Monoclonal Antibodies

Yao Zhang[1,3,¶], Xiaohua Hao[2,¶], Ji Ma[3], Mingming Wang[3], Yanyan Li[3], Yang Liu[3],

Dong Zhao[2], Wen Zhang[2], Chunming Li[3], Li Yan[3], Qing Zhu[1,3,*], Fujie Zhang[2,*]

[1] TSB Therapeutics, Beijing, China

[2] Beijing Ditan Hospital, Capital Medical University, Beijing, China

[3] Brii Biosciences Inc., Durham, NC, US

[*:] Correspondening authors:

Qing Zhu

Email address: qing.zhu@briibio.com

Fujie Zhang

Email address: treatment@chinaaids.cn

[¶] These authors contributed equally to this work.

Running title: Safety and PK of COVID-19 mAbs in human

Keywords. SARS-CoV-2; monoclonal antibody therapy; BRII-196; BRII-198

Clinical Trials Registration. NCT04479631 and NCT04479644

Funding: These studies were sponsored by TSB Therapeutics.

Abstract

Background. BRII-196 and BRII-198 are two anti-SARS-CoV-2 monoclonal

neutralizing antibodies with modified Fc region that extends half-life and are being

developed as cocktail therapy for the treatment of COVID-19. Safety, tolerability,

pharmacokinetics, and immunogenicity of BRII-196 and BRII-198 were investigated

in healthy adults.

Methods. Single ascending doses of BRII-196 and BRII-198 were evaluated in

parallel in the first-in-human, placebo-controlled phase 1 studies. A total of 32 healthy

adults were randomized and received a single intravenous infusion of 750, 1500, and

3000 mg of BRII-196 (n=12), BRII-198 (n=12), or placebo (n=8) and were followed

for 180 days.

Results. All infusions were well tolerated at infusion rates between 0.5 mL/min to 4

mL/min with no dose-limiting adverse events, deaths, serious adverse events, or any

systemic or local infusion reactions. Most treatment-emergent adverse events were

isolated asymptomatic laboratory abnormalities of Grade 1-2 in severity. Each mAb

displayed pharmacokinetics expected of Fc-engineered human IgG1 with mean

terminal half-lives of approximately 46 days and 76 days, respectively, with no

evidence of significant anti-drug antibody development.

40   **Conclusions**. BRII-196 and BRII-198 were well-tolerated. Clinical results support

41   further development as therapeutic or prophylactic options for SARS-CoV-2

42   infection.

Background

As of March 2021, globally, over 120 million people have been infected by SARS-CoV-2, resulting in more than 2.6 million death by Coronavirus Disease 2019 (COVID-19). The number of cases continues to rise[1].

The development of safe and effective therapeutics represents a key strategy to curb COVID-19 mortality and morbidity. Several approved medications (e.g., chloroquine, hydroxychloroquine, lopinavir/ritonavir) have been "repurposed" and evaluated, none of which have been shown associated with improved clinical outcomes[2, 3]. There have been limited treatment options, including remdesivir, dexamethasone, and convalescent plasma, that demonstrated clinical efficacy in subsets of patients with COVID-19[4-7].

Neutralizing monoclonal antibodies (mAbs) represent a promising therapeutic option in preventing and treating known and emerging infectious diseases, including viruses[8]. Recent clinical trials of mAbs reported positive results indicating virological and clinical benefits among outpatients with COVID-19 [9, 10]. In addition, several COVID-19 vaccines offering>90% protection have become available for emergency use [11, 12]. However, recent data showed that several mutations identified in the receptor binding domain (RBD) of spike protein in the emerging new variants conferred reduced activity to neutralization by the antibodies that were developed as a therapy for treatment or elicited by vaccines for prevention [13, 14]. These emergent variants present new challenges for mAb therapy and threaten the protective efficacy of current vaccines and mAbs. Therefore, additional mAb treatment options, especially a cocktail of antibodies targeting distinct epitopes, are needed as a critical strategy to reduce the generation and selection of resistant viruses during treatment

and ensures binding and neutralization of the antibodies to new variants.

BRII-196 and BRII-198, two recombinant human IgG1 mAbs that target the distinct

epitope regions in RBD in coronavirus spike glycoproteins non-competitively, are

under clinical development as a cocktail for the treatment of COVID-19 in outpatients.

BRII-196 and BRII-198 are derived directly from human B cells of patients who

recovered from COVID-19 [15] with effective neutralizing activity against not only

the original isolate of SARS-CoV-2 but also the new variants of concerns, including

B.1.351 (as known South African variant) and P1 from Brazil [16, 17]. BRII-196 and

BRII-198 are engineered with a triple-amino-acid (M252Y/S254T/T256E [YTE])

substitution in the fragment crystallizable (Fc) region to allow an extended half-life

[18, 19] and reduced Fcγ receptor binding for potential risk of Fc-medicated antibody

dependent enhancement (ADE) (unpublished data).

The back-to-back first-in-human phase 1 studies in healthy adult participants were

designed and conducted to evaluate the safety, tolerability, and pharmacokinetics (PK)

of BRII-196 and BRII-198 individually before initiating clinical studies of BRII-196

and BRII-198 in combination in patients.

Methods

Study design and participants

BRII-196-001 and BRII-198-001 are two first-in-human phase 1, randomized,

single-blind, placebo-controlled, single ascending dose escalation studies in which

BRII-196 and BRII-198 were evaluated respectively among healthy adults. These

studies were conducted at a single phase 1 unit in China from July 2020 to February

2021 (registered at ClinicalTrials.gov under registration number NCT04479631 and

NCT04479644). The study protocol, amendments, and informed-consent forms were

reviewed and approved by the Institutional Ethics Committee (IEC). The IEC

included Guo'an Wu, Wangyan Jia, Xiulan Li, Xiangmei Zhao, Zhiyun Yang,

Xingwang Li, Xianbo Wang, Hanqiu Zhan, Hui Zeng, Yajie Wang, Yun'ao Zhou, and

Gang Wan from Beijing Ditan Hospital, Capital Medical University; Kairong Wang

and Yangang Tian from Beijing Lvli Law Firm; and Jianguo Sun as a community

representative. The study was conducted in accordance with the International

Conference on Harmonisation Guidance for Good Clinical Practice guidelines and all

applicable local regulatory requirements and laws. Participants provided written

informed consent before any study-related procedures were performed.

Inclusion and exclusion criteria for the two studies were identical. Eligible

participants were healthy male or female adults aged 18 to 49 years, had a body

weight ≤100 kg and a body mass index of 18-24 kg/m^2, and were in good health

determined by no clinically significant findings from medical history, physical

examination including vital signs, electrocardiogram (ECG), and clinical laboratory

assessments. Women were not pregnant or lactating. Exclusion criteria included a

history of SARS-CoV-2 infection or exposure, a history of severe allergic reactions,

use of any medications that were started within 14 days before randomization, a

positive test for human immunodeficiency virus (HIV) or hepatitis B virus or hepatitis

C virus, a history of drug or alcohol abuse within 1 year before the screening, or any

history of a medical or psychiatric condition that would place them at risk or interfere

with study participation. Participants were also excluded if they were unwilling to use

necessary contraception during the study. A complete list of exclusion criteria is

included in Supplementary Material 1.

Study Procedures

Three dose-level cohorts (750, 1500, and 3000 mg) were included to evaluate each

mAb and were initiated in a dose-escalation design. In each cohort, participants were

randomized at a 3:1 ratio to receive either the single mAb (n=3 or 6) or placebo (n=1

or 2). mAbs or placebo was dispensed into normal saline and was administered via

intravenous infusion on Day 1. The infusion started at the initial rate of 0.5 mL/min,

gradually increasing to a maximum rate of 4 mL/min. The participants were masked

to the treatment assignment and were admitted to the phase 1 unit for monitoring for 7

days post-dosing. The first participant in each cohort who received mAb infusion was

monitored for 24 hours before administering study intervention to the subsequent

participants. A Safety Review Committee made Dose-escalation recommendations

upon review of predefined safety data.

Vital signs, physical examinations, safety laboratory tests including hematology and

chemistry, and ECG were obtained before and after study drug administration on days

2, 4, 6, 8, 15, 22, 31, 61, 91, 121, 151, and at the end of the study on day 181. All

adverse events (AEs), including infusion reactions and serious adverse events (SAEs)

occurring throughout the trial, were recorded and assessed by study physicians at the

site. AEs were coded with the Medical Dictionary for Regulatory Activities

(MedDRA) version 23.1, and severity was graded using the Common Terminology

Criteria for Adverse Events (CTCAE) version 5.0, November 27, 2017.

Pharmacokinetics and Immunogenicity Assessments

The pharmacokinetics parameters were evaluated using blood samples collected

before dosing, at 3 and 8 hours post dose, and days 2, 4, 6, 8, 15, 22, 31, 61, 91, 121,

151, and 181. BRII-196 or BRII-198 serum concentrations were measured using

validated enzyme linked immunosorbent assays (ELISA) with the lower limit of

quantitation (LLOQ) at 150 ng/mL. Serum samples for anti-drug antibody (ADA)

assays were collected at the following time points: pre-dose on day 1 and at days 15,

31, and 181. Two validated titer-based 3-stage ELISA assays were used to detect

presences of ADAs from subjects over the study durations.

Study Outcomes

Primary endpoints were incidence of adverse events (AEs) and change from pre-dose

baseline in clinical assessments, including vital signs, ECG readings, and laboratory

results. Secondary endpoints included the pharmacokinetics profiles and presence of

ADAs to BRII-196/BRII-198 in samples collected after dosing for up to 180 days.

Statistical Analysis

The sample size of each study was consistent with a phase 1 first-in-human study. The

safety analysis included all participants who were randomized and received any dose

of the study drug. Categorical and continuous data were summarized descriptively.

Participants were analyzed according to the study drug they received. Participants

receiving placebo in different cohorts of each study were pooled in the analysis.

The PK parameters were estimated by non-compartmental analyses using WinNonlin

module in the Phoenix Platform (version 8.3.1.5014, Certara Inc., Princeton, NJ

08540). Calculations were performed prior to rounding, and nominal sampling times

were used in the pharmacokinetic analysis. All pharmacokinetic parameters and

summary statistics are reported to 3 significant digits except for T_{max} (reported in

median values), which is reported to 1 decimal place.

Data Availability:

The datasets generated during and/or analysed during the current study are available

from the corresponding author on reasonable request.

Results

From July 12, 2020, to August 10, 2020, a total of 135 participants were screened,

with 16 and 17 eligible and enrolled for BRII-196-001 and BRII-198-001 studies,

respectively (**Figure 1**). The demographics of the study population are summarized in

Table 1, consisting of 28 (84.8%) men and 5 women with a median age of 33.0 years

and a median BMI of 21.9 kg/m^2.

A total of 12 participants received a single dose of BRII-196 IV infusion, 12

participants received a single dose of BRII-198 IV infusion, and 8 participants

received placebo (normal saline). One participant randomized in BRII-198-001 study

did not receive the infusion because of elevated blood pressure before dosing. All

participants received their full planned dose except for 3 participants assigned to 750

mg of BRII-196 and 1 participant assigned to placebo for whom the received volume

was approximately 10% less than the intended volume due to the remaining volume

left within the infusion set. All participants dosed have completed study visits. Total

infusion duration for BRII-196 or BRII-198 at investigated dose levels were 1.53-1.70

hours (750 mg), 2.27-2.43 hours (1500 mg), and 3.58-3.75 hours (3000 mg)

(Supplementary Table 1).

Safety

Safety

BRII-196 and BRII-198 administration was safe and well-tolerated. No deaths,

serious AEs, treatment-emergent adverse events (TEAEs) leading to early termination,

or infusion-related reactions occurred during the study. Adverse events reported in

BRII-196-001 and BRII-198-001 are summarized in **Table 2** and **Table 3**, respectively.

26 (81%) participants had at least one treatment-emergent adverse events, including

18 (75%) participants receiving BRII-196 or BRII-198 compared to 8 (100%)

participants receiving placebo. Overall incidences of adverse events were balanced

between groups receiving BRII-196 or BRII-198 and placebo. The most common

TEAEs were isolated asymptomatic laboratory abnormalities of CTCAE Grade 1 or 2

in severity that did not require medical intervention and typically normalized or

returned to baseline level within four weeks (Supplementary Table 2). Seven out of 24

participants had 8 Grade 1 or 2 adverse events that were considered by the

investigators as related to BRII-196 or BRII-198, including increased blood total

bilirubin, increased alanine aminotransferase, decreased neutrophil count, and

decreased white blood cell count (Supplementary Table 3). No safety issues were seen

with respect to vital signs or ECG measurements. No dose-dependent pattern of any

adverse events was observed.

Pharmacokinetics

Pharmacokinetic data were available for a total of 12 participants each for BRII-196

and BRII-198 over the 181-day study duration. Following a single intravenous

infusion of BRII-196 or BRII-198 at 750, 1500, and 3000 mg, mean serum PK

parameters, including C_{max} and AUC, increased in an approximately

dose-proportional manner (**Table 4**, **Figure 2**). The mean systemic serum clearances

were 75.4 and 57.0 mL/day for BRII-196 and BRII-198, respectively. The mean

terminal half-lives ($t_{1/2}$) of BRII-196 and BRII-198 were 44.6-48.6 days and 72.2-83.0

days. The relatively shorter terminal half-life of BRII-196 correlated with slightly

higher systemic clearance.

ADA Response

In BRII-196-001 study, a total of 4 out of 12 participants had positive ADA samples in

the screening assay, but all tested negative in the follow-up confirmatory assay. In

BRII-198-001 study, one participant who received BRII-198 had positive ADA

samples in the screening assay but later confirmed negative in the tier-2 confirmatory

assay.

Discussion

BRII-196 and BRII-198 are fully human mAbs that bind to noncompeting epitope

regions on the RBD of SARS-CoV-2 S glycoprotein that interacts with the

angiotensin-converting enzyme 2 (ACE2) to enter human cells [20]. Therefore, they

neutralize virus infection by preventing virus binding to ACE2, blocking virus entry

and subsequent virus infection as demonstrated in vitro and in vivo [15, 21]. The

combined use of 2 independently neutralizing antibodies is likely to reduce the risk of

treatment failure from the development of virus escape mutants or new variants that

may arise during an outbreak. Currently approved COVID -19 vaccines were

constructed against the original virus, of which attenuation of serum neutralization

activity has been observed[14, 22]. In a recent trial conducted in South Africa, the

ChAdOx1 nCoV-19 vaccine did not protect against mild-to-moderate COVID-19 due

to the B.1.351 variant [23]. Pseudo viruses harboring mutations found in B.1.351

South Africa origin, P.1 Brazil origin, B.1.526 New York origin, L452R, and the spike

protein from the California origin variant lineage B.1.427/B.1.429, had significantly

reduced susceptibility to bamlanivimab [24], a mAb authorized for emergency use. In

addition, its approved cocktail with etesevimab under EUA was also tested inactive

against the live virus of B.1.351 South Africa [17]. As some of these variants are

beginning to spread around the world, it has posed a real threat to vaccine and

antibody therapies. By contrast, BRII-196 and BRII-198 cocktail retained their

neutralizing activity against all variants tested, including B1.1.7 UK variant, B1.357

South Africa variant, and P1 Brazilian variants[16, 17], thereby could provide more

treatment options to patients.

In these first-in-human clinical studies, intravenous administration of BRII-196 and

BRII-198 were both safe and well-tolerated in healthy adult participants receiving a

single dose up to 3000 mg, with no infusion reactions or AEs leading to adjustment or

discontinuation of the infusion. Study participants were followed up to 180 days

post-dosing considering the expected prolonged half-lives of the mAbs. Except for

one SAE, one Grade 4 elevation of creatine phosphokinase, and one Grade 3 elevation

of triglycerides, all deemed as not related to study drugs, all AEs reported were Grade

1-2 in severity, with the majority being isolated asymptomatic transient laboratory

abnormalities. There were no safety concerns regarding vital signs or ECGs. The

safety observations in these phase 1 studies are consistent with the expected safety

profile of human IgG1 mAbs in healthy subjects with no endogenous target.

With YTE modification, both BRII-196 and BRII-198 demonstrated anticipated 2-3

folds half-life extension in healthy adult subjects. In comparison with BRII-198, the

relatively shorter half-life observed with BRII-196 correlates with slightly higher

systemic clearance. As both antibodies showed comparable increased binding

affinities to human and cynomolgus monkey FcRns at pH 6.0 in vitro (unpublished

data), different molecular properties such as charge variants, glycan profiles, and

overall thermal stability might contribute to the observed differences. With up to a

3-week optimal treatment window for neutralizing antibodies for COVID-19 patients,

both antibodies are expected to provide comparable target coverage at the same dose

level.

BRII-196 and BRII-198 have several advantages that may facilitate their use for the

treatment and prevention of SARS-COV-2 infections. First, mAbs targeting specific

viral epitopes, including BRII-196 and BRII-198, belong to a platform technology

associated with a favorable safety profile. Findings regarding safety and tolerability in

phase 1 trials of BRII-196 and BRII-198 are consistent with the expectation, which

support an expedited development towards approval for future clinical use. Second,

targeting distinct viral epitopes in the RBD of SARS-COV-2 spike protein

non-competitively, BRII-196 and BRII-198 can be used in combination to provide

additive antiviral effect, reduce the generation of resistant viruses to treatment, and

retain activity against emergent naturally occurring variants as mentioned above.

Third, BRII-196 and BRII-198 are engineered with YTE substitution in the Fc region

to allow an extended half-life, demonstrated in the pharmacokinetic findings from

both phase 1 studies, with which BRII-196 and BRII-198 can potentially maintain an

adequate drug coverage over an extended period of time and subsequently are better

positioned for prophylactic use. Lastly, while YTE improves the half-life of the

antibody, it also reduced binding activity of BRII-196 and BRII-198 against human

Fcγ receptors, thereby minimizing the potential risk of Fc-mediated antibody

dependent enhancement of diseases.

In summary, BRII-196 and BRII-198 were generally safe and well-tolerated in healthy

adult participants. Pharmacokinetic and immunogenicity assessments demonstrated

linear pharmacokinetics with prolonged half-lives and no ADA development up to

180 days after a single dose. These findings support further development of BRII-196

and BRII-198 for the treatment and prophylaxis of SARS-COV-2 infection in the

pandemic outbreak. BRII-196 and BRII-198 combination is currently under

development in ACTIV-2, a platform phase 2/3 clinical study, for treatment of

COVID-19 in outpatients [25].

Acknowledgment: The authors thank all the study participants in these clinical

studies, without whom the studies would not have been possible. We thank Dr Yibo

Zhou, Jianmin Jing, and Yonghong Mang, for their contributions in the studies. We

also thank D.Margolis for critically reviewing the manuscript and Dr. Chan Gao for

her assistance with formatting/editing and submission.

Financial support: These trials were sponsored by TSB Therapeutics, a subsidiary

company of Brii Biosciences. Y.Z., J.M., Y.L., Y.Y.L., M.W., C.L., L.Y., and Q.Z. are

employees of Brii Biosciences. Y.Z. and Q.Z. are employees of TSB Therapeutics.

F.J.Z. received support from TSB for participation as a principal investigator in the

study.

Author contributions: Brii Biosciences was involved in the study design, collection,

analysis and interpretation of data, the writing of this report, and the decision to

submit this manuscript for publication. F.J.Z., X.H.H, D.Z., and W.Z. reviewed

protocol, and were responsible for study implementation and data collection at the

study site. All authors at Brii Bioscienecs had access to all study data for review,

participated in interpretation of the data and in drafting and revising the article for

important intellectual content, and provided final approval of the submitted

manuscript. All authors assume responsibility for the integrity and completeness of

the reported data. Y.Z., J.M., Y.L., and Q.Z prepared the manuscript.

Potential conflicts of interest. Y.Z., J.M., Y.L., Y.L., M.W., C.L., L.Y., and Q.Z. are

employees of Brii Biosciences. Y.Z. and Q.Z. are employees of TSB Therapeutics. All

other authors report no potential conflicts. All authors have submitted the ICMJE

Form for Disclosure of Potential Conflicts of Interest.

References

1.   Johns_Hopkins_Coronavirus_Resource_Center.   COVID-19   Map.   Available   at:

https://coronavirus.jhu.edu/map.html. Accessed Mar 17 2021.

2. Cao B, Wang Y, Wen D, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with

Severe Covid-19. N Engl J Med **2020**; 382:1787-99.

3. SolidarityTrialConsortium WHO, Pan H, Peto R, et al. Repurposed Antiviral Drugs for

Covid-19 - Interim WHO Solidarity Trial Results. N Engl J Med **2021**; 384:497-511.

4. Libster R, Pérez Marc G, Wappner D, et al. Early High-Titer Plasma Therapy to Prevent

Severe Covid-19 in Older Adults. N Engl J Med **2021**; 384:610-8.

5. Beigel JH, Tomashek KM, Dodd LE, et al. Remdesivir for the Treatment of Covid-19 - Final

Report. N Engl J Med **2020**; 383:1813-26.

6. Group RC, Horby P, Lim WS, et al. Dexamethasone in Hospitalized Patients with Covid-19.

N Engl J Med **2021**; 384:693-704.

7. Simonovich VA, Burgos Pratx LD, Scibona P, et al. A Randomized Trial of Convalescent

Plasma in Covid-19 Severe Pneumonia. N Engl J Med **2021**; 384:619-29.

8. Jin Y, Lei C, Hu D, Dimitrov DS, Ying T. Human monoclonal antibodies as candidate

therapeutics against emerging viruses. Front Med **2017**; 11:462-70.

9. Chen P, Nirula A, Heller B, et al. SARS-CoV-2 Neutralizing Antibody LY-CoV555 in

Outpatients with Covid-19. N Engl J Med **2021**; 384:229-37.

10. Weinreich DM, Sivapalasingam S, Norton T, et al. REGN-COV2, a Neutralizing Antibody

Cocktail, in Outpatients with Covid-19. N Engl J Med **2021**; 384:238-51.

11. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA

Covid-19 Vaccine. N Engl J Med **2020**; 383:2603-15.

12. Baden LR, El Sahly HM, Essink B, et al. Efficacy and Safety of the mRNA-1273

SARS-CoV-2 Vaccine. N Engl J Med **2021**; 384:403-16.

13. Liu H, Wei P, Zhang Q, et al. 501Y.V2 and 501Y.V3 variants of SARS-CoV-2 lose binding

to Bamlanivimab in vitro. bioRxiv [Preprint] February 16, 2021 [cited 2021 Apr 8]. Available

from: https://doi.org/10.1101/2021.02.16.431305

14. Wu K, Werner AP, Koch M, et al. Serum Neutralizing Activity Elicited by mRNA-1273

Vaccine. N Engl J Med 2021.

15. Ju B, Zhang Q, Ge J, et al. Human neutralizing antibodies elicited by SARS-CoV-2

infection. Nature 2020; 584:115-9.

16. Wang R, Zhang Q, Ge J, et al. SARS-CoV-2 variants resist antibody neutralization and

broaden host ACE2 usage. bioRxiv [Preprint] March 15, 2021 [cited 2021 Apr 8]. Available

from: https://doi.org/10.1101/2021.03.09.434497.

17. Wang P, Nair MS, Liu L, et al. Antibody Resistance of SARS-CoV-2 Variants B.1.351 and

B.1.1.7. bioRxiv [Preprint] January 26, 2021 [cited 2021 Apr 8]. Available from:

https://doi.org/10.1101/2021.01.25.428137.

18. Robbie GJ, Criste R, Dall'acqua WF, et al. A novel investigational Fc-modified humanized

monoclonal antibody, motavizumab-YTE, has an extended half-life in healthy adults.

Antimicrob Agents Chemother 2013; 57:6147-53.

19. Saunders KO. Conceptual Approaches to Modulating Antibody Effector Functions and

Circulation Half-Life. Front Immunol 2019; 10:1296.

20. Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, Function, and

Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell 2020; 181:281-92.e6.

21. Ge J, Wang R, Ju B, et al. Antibody neutralization of SARS-CoV-2 through ACE2 receptor

mimicry. Nature communications 2021; 12:250.

22. Liu Y, Liu J, Xia H, et al. Neutralizing Activity of BNT162b2-Elicited Serum. New England

Journal of Medicine 2021; 0:null.

23. Madhi SA, Baillie V, Cutland CL, et al. Efficacy of the ChAdOx1 nCoV-19 Covid-19

Vaccine against the B.1.351 Variant. New England Journal of Medicine 2021; 0:null.

24. Moruf A. Fact Sheet For Health Care Providers Emergency Use Authorization (EUA) Of

Bamlanivimab. U.S. Food and Drug Administration, 2021:26.

25. Large clinical trial will test combination monoclonal antibody therapy for mild/moderate

COVID-19.   Available   at:

https://www.nih.gov/news-events/news-releases/large-clinical-trial-will-test-combination-mono

clonal-antibody-therapy-mild-moderate-covid-19. Accessed March 24 2021.

Figure 1 Trial Profiles

Figure 2. Observed Mean (±Standard Deviation) Serum Concentration - Time Profiles of BRII-196 and BRII-198 Following A Single Intravenous Infusion Administration to The Healthy Adult Subjects

Table 1   Participant Demographics and Baseline Characteristics

	BRII-196-001				BRII-198-001			
	750 mg (n=3)	1500 mg (n=6)	3000 mg (n=3)	Placebo (n=4)	750 mg (n=3)	1500 mg (n=7)	3000 mg (n=3)	Placebo (n=4)
Median age, years	22.0	32.5	34.0	33.5	30.0	33.0	30.0	37.5
Men	3 (100%)	4 (67%)	3 (100%)	3 (75%)	2 (67%)	7 (100%)	3 (100%)	3 (75%)
Women	0	2 (33%)	0	1 (25%)	1 (33%)	0	0	1 (25%)
Race								
Han, Chinese	3 (100%)	6 (100%)	2 (67%)	3 (75%)	3 (100%)	6 (86%)	3 (100%)	4 (100%)
Other, Chinese	0	0	1 (33%)	1 (25%)	0	1 (14%)	0	0
Median body-mass index, kg/m^2	22.80	21.15	21.60	22.90	23.70	21.00	22.30	23.10

Table 2   Summary of Adverse Events of BRII-196-001 Study

	BRII-196 750mg (n=3)	BRII-196 1500mg (n=6)	BRII-196 3000mg (n=3)	Placebo (n=4)
Participants with any TEAE	3 (100%)	5 (83%)	2 (67%)	4 (100%)
Participants with Grade 1 TEAE	3 (100%)	5 (83%)	2 (67%)	4 (100%)
Participants with Grade 2 TEAE	1 (33%)	3 (50%)	0	2 (50%)
Participants with Grade 3 or above TEAE	0	0	0	0
Participants with serious TEAE	0	0	0	0
Participants with infusion reaction TEAE	0	0	0	0
Participants with TEAEs related to study drug	1 (33%)	3 (50%)	0	0
Participants with any TEAE leading to discontinuation of study drug	0	0	0	0
Death	0	0	0	0
Number of participants with AEs by MedDRA SOC [a]				
Investigations	3 (100%)	4 (67%)	2 (67%)	4 (100%)
Infections and infestations	1 (33%)	0	0	2 (50%)

TEAE=treatment-emergent adverse event. SOC=system organ class.

Participants who experienced the same AE on more than one occasion (based on the specific category) are counted once in each relevant category. Percentages are based on the number of participants in the treatment group.

[a] Only SOCs experienced by ≥2 participant or with the maximum severity category ≥ Grade 2 are included.

Table 3 Summary of Adverse Events of BRII-198-001 Study

	BRII-198 750mg (n=3)	BRII-198 1500mg (n=6)	BRII-198 3000mg (n=3)	Placebo (n=4)
Participants with any TEAE	3 (100%)	5 (83%)	0	4 (100%)
Participants with Grade 1 TEAE	2 (67%)	5 (83%)	0	3 (75%)
Participants with Grade 2 TEAE	1 (33%)	0	0	1 (25%)
Participants with Grade 3 or above TEAE	0	2[a] (33%)	0	1[b] (25%)
Participants with serious TEAE	0	0	0	1[b] (25%)
Participants with infusion reaction TEAE	0	0	0	0
Participants with TEAEs related to study drug	2 (67%)	1 (17%)	0	0
Participants with any TEAE leading to discontinuation of study drug	0	0	0	0
Death	0	0	0	0
Number of participants with AEs by SOC [c]				
Investigations	3 (100%)	4 (67%)	0	3 (75%)
Gastrointestinal disorders	1 (33%)	1 (17%)	0	0
Injury, poisoning and procedural complications	0	0	0	1 (25%)

TEAE=treatment-emergent adverse event. SOC=system organ class.

Participants who experienced the same AE on more than one occasion (based on the specific category) are counted once in each relevant category. Percentages are based on the number of participants in the treatment group.

[a] 1 participant had an elevated blood creatine phosphokinase of Grade 4 after excessive exercise, and 1 participant had an increased blood triglycerides of Grade 3 on Day 181. Both events were not considered as related to the study drug by the investigators and normalized within 1-3 weeks.

[b] 1 participant receiving placebo had a serious TEAE (bilateral traumatic foot fracture leading to hospitalization, which was also reported as a Grade 3 TEAE). The participant recovered after surgery.

[c] Only SOCs experienced by ≥ 2 participant or with the maximum severity category $\geq$ Grade 2 are included.

Table 4   BRII-196 and BRII-198 Pharmacokinetic Parameters Following A Single Intravenous Infusion Administration to The Healthy Adult Participants

Antibody	Dose (mg)	Number of Subject	C_{max} (µg/mL)	T_{max} (hour)	$t_{1/2}$ (day)	CL (mL/day)	V_{ss} (L)	AUC_{last} (day*µg/mL)	AUC_{inf} (day*µg/mL)
BRII-196	750	3	268 (19.9)	4.6	48.6 (4.56)	84.7 (10.5)	5.34 (0.500)	8240 (764)	8940 (1040)
	1500	6	584 (75.1)	5.4	46.0 (8.09)	72.4 (8.40)	4.50 (0.208)	19300 (2430)	20900 (2380)
	3000	3	1300 (91.7)	6.6	44.6 (6.75)	72.2 (4.11)	4.35 (0.289)	38900 (2510)	41700 (2300)
BRII-198	750	3	244 (35.5)	4.7	75.9 (8.97)	59.4 (5.54)	6.17 (0.760)	10400 (911)	12700 (1240)
	1500	6	509 (107)	5.4	72.2 (8.45)	57.4 (9.50)	5.76 (0.981)	20900 (2950)	26800 (4700)
	3000	3	1017 (149)	6.8	83.0 (37.6)	53.9 (8.90)	6.07 (2.20)	42800 (9680)	56600 (8900)

SD=standard deviation, C_{max}=observed maximum serum concentration, T_{max}=time to reach observed maximum serum concentration, $t_{1/2}$=terminal half-life, CL=systemic clearance, V_{ss}=volume of distribution at steady state, AUC_{last}=area under the concentration–time curve from time zero to the last measurable concentration, AUC_{inf}=area under the concentration–time curve from time zero to infinity. All parameters are reported in three significant figures except T_{max} that is reported with median values.