Herd immunity induced by COVID-19 vaccination programs and suppression of

epidemics caused by the SARS-CoV-2 Delta variant in China

Hengcong Liu[1]*, Juanjuan Zhang [1,2]*, Jun Cai[1], Xiaowei Deng[1], Cheng Peng[1],

Xinghui Chen[1], Juan Yang[1,2,3], Qianhui Wu[1], Xinhua Chen[1], Zhiyuan Chen[1], Wen

Zheng[1], Cécile Viboud[4], Wenhong Zhang[1]†, Marco Ajelli[5,6]† and Hongjie Yu[1,2,3]†

1. Department of Infectious Diseases, Huashan Hospital, School of Public Health,

Fudan University, Shanghai, China

2. Key Laboratory of Public Health Safety, Fudan University, Ministry of

Education, Shanghai, China

3. Shanghai Institute of Infectious Disease and Biosecurity, Fudan University,

Shanghai, China

4. Division of International Epidemiology and Population Studies, Fogarty

International Center, National Institutes of Health, Bethesda, MD, USA

5. Department of Epidemiology and Biostatistics, Indiana University School of

Public Health, Bloomington, IN, USA

6. Laboratory for the Modeling of Biological and Socio-technical Systems,

Northeastern University, Boston, MA, USA

*These authors contributed equally to this work.

†These authors are joint senior authors contributed equally to this work.

Corresponding authors: Hongjie Yu, Shanghai Institute of Infectious Disease and

Biosecurity, School of Public Health, Fudan University, Shanghai 200032, China

E-mail: yhj@fudan.edu.cn

Word count (main text): 4,202

Word count (abstract): 228

Abstract

Background

To allow a return to a pre-COVID-19 lifestyle, virtually every country has initiated a vaccination program to mitigate severe disease burden and control transmission. However, it remains to be seen whether herd immunity will be within reach of these programs.

Methods

We developed a data-driven model of SARS-CoV-2 transmission for China, a population with low prior immunity from natural infection. The model is calibrated considering COVID-19 natural history and the estimated transmissibility of the Delta variant. Three vaccination programs are tested, including the one currently enacted in China and model-based estimates of the herd immunity level are provided.

Results

We found that it is unlike to reach herd immunity for the Delta variant given the relatively low efficacy of the vaccines used in China throughout 2021, the exclusion of underage individuals from the targeted population, and the lack of prior natural immunity. We estimate that, assuming a vaccine efficacy of 90% against the infection, vaccine-induced herd immunity would require a coverage of 93% or higher of the Chinese population. However, even when vaccine-induced herd immunity is not reached, we estimated that vaccination programs can reduce SARS-CoV-2 infections

by 53-58% in case of an epidemic starts to unfold in the fall of 2021.

Conclusions

Efforts should be taken to increase population's confidence and willingness to be

vaccinated and to guarantee highly efficacious vaccines for a wider age range.

Keywords: Covid-19, herd immunity, vaccination program, SIR model, Delta variant

Introduction

The first-wave of novel coronavirus disease 2019 (COVID-19) in China subsided

quickly after the implementation of strict containment measures and travel restrictions

starting in March 2020 [1]. As of August 18, 2021, the COVID-19 pandemic has caused

over 208 million reported cases and 4.5 million deaths globally [2]. The pandemic is far

from over, as SARS-CoV-2 has undergone some significant mutations and a number

of variants have become widespread due to increased transmissibility and/or immune

escape characteristics – e.g., variants Alpha, Beta, Gamma, and Delta. Throughout the

globe, a rapid surge of Delta variant cases suggests a clear competitive advantage

compared with Alpha, Beta, and Gamma; more than 90% of daily sequences from

GISAID are ascribable to the Delta variant since July 2021 [3]. Despite of no major

epidemics, China has been experiencing several minor local outbreaks caused by

imported cases of Delta variant, including the outbreaks in Guangzhou, Nanjing, and

Zhengzhou city [1,4]. To suppress transmission, a large share of the world needs to have

immunity to SARS-CoV-2, especially to the Delta variant.

Effective vaccines against COVID-19 represent the most viable option to suppress

SARS-CoV-2 transmission globally. The effectiveness of vaccination programs

depends on several key factors, including vaccine supply, willingness to receive the

vaccine, vaccine efficacy, and the age groups targeted by the vaccination effort.

However, current vaccination programs are all based on vaccines developed against

the historical SARS-CoV-2 lineage, and the efficacy seems be reduced against the

Delta variant. In China, home of about 1. billion people (~18% of the world

population), 1.89 billion doses have been administered as of August 12, 2021 [5]. This

figure corresponds to 67.2% of the whole population and 87.6% of the target

population (i.e., individuals aged 18 years and above). However, it remains to be seen

if and when the vaccine coverage may reach a level sufficient to achieve herd

immunity. Countries around the globe are facing the same question.

The classical herd immunity level is defined as $1-1/R_0$, where R_0 is the basic

reproduction number – the average number of infections generated by a typical

infectious individual in a fully susceptible population [6]. For a vaccine with efficacy

VE that gives life-long protection, the level of herd immunity required to stop

transmission is $(1-1/R_0)$/VE. However, this estimate is an oversimplification of a

complex phenomenon as it ignores the heterogeneities of actual human population

(e.g., social mixing patterns, age-specific susceptibility to infection) [7,8] as well as of

vaccination (e.g., lifelong immunity, sterilizing vaccine). To overcome this limitation,

here we integrate contact survey specific of the Chinese population [9] as well as

official demographic statistics to develop an age-structured stochastic model to

simulate SARS-CoV-2 transmission (Fig.S1 in *SI Appendix*). We then use this model

to evaluate whether herd immunity is achievable against the Delta variant or not via

mass vaccination and to explore the way forward to achieve suppression of

transmission.

Methods

SARS-CoV-2 transmission and vaccination model

We built a compartmental model of SRAS-CoV-2 transmission and vaccination, based

on an age-structured stochastic SIR scheme, accounting for heterogeneous contact

patterns by age [9] and heterogeneous susceptibility to infection by age as estimated

using contact tracing data in Hunan province of China. In the model, the population is

divided into three epidemiological categories: susceptible, infectious, and removed,

stratified by 17 age groups. Susceptible individuals can become infectious after

contact with an infectious individual according to the age-specific force of infection.

The rate at which contacts occur is determined by the mixing patterns of each age

group. The average generation time was set to 7 days. We consider a basic

reproductive number (R_0) of 6.0 according to estimates for the SARS-CoV-2 Delta

variant. Simulations are initiated with 40 infectious individuals, corresponding to the

number of cases first detected in a local outbreak in Beijing on June 11, 2020.

We consider a 2-dose vaccine, that only susceptible individuals are eligible for

vaccination (we recall that natural immunity is close to 0 in China as of July 2021),

and that the duration of vaccine-induced immunity lasts longer than the time horizon

considered in this study (i.e., 1 year). Details about the model and parameters are

reported in Sec. 1 of *SI Appendix*.

The model allows the explicit simulation of the vaccination strategy currently used in

China: random distribution of vaccines to adults aged 18+ years (strategy 1). Two of

the SARS-CoV-2 vaccines (i.e., BBIBP-Corv and CoronaVac) used in China have

been licensed for children aged 3 to 17 years [10]; however, as of August 12, 2021

children are not included among the target population for vaccination in most

provinces. To explore the contribution of vaccinating children aged 3-17 years to

achieving herd immunity, we test two alternative strategies: i) same as strategy 1, but

extended to individuals aged 3+ years starting from September 1, 2021 (strategy 2);

ii) random distribution of vaccines to individuals aged 3+ years since the start of

vaccination programs (strategy 3).

Baseline scenario

As the baseline scenario, we considered the following assumptions:

137   i)   **Epidemic seeding**: An epidemic is assumed to be triggered by forty SARS-

138   CoV-2 infections on September 1, 2021; vaccines have been rolling out in

139   China since November 30, 2020.

140   ii)   **Vaccination strategy:** We test three different vaccination strategies:

141   a)   Strategy 1—random distribution of vaccines to adults aged 18+ years

142   b)   Strategy 2—same as strategy 1, but the vaccination is extended to individuals

143   aged 3+ years starting from September 1, 2021

144   c)   Strategy 3—random distribution of vaccines to individuals aged 3+ years

145   since the start of the vaccination program.

146   Note that in all scenarios, we consider that a fraction of the population (about

147   2% - Sec. 4 of *SI Appendix*) is not eligible to receive the vaccine (e.g.,

148   pregnant women, individuals with allergies or other conditions preventing

149   them to safely receive the vaccine).

150   iii)   **Vaccine capacity**: We simulated the daily vaccine administration capacity

151   based on the vaccine capacity data throughout the entire vaccination campaign

152   in China [5]. We found that the daily vaccine administration capacity

153   exponentially increased during the initial phase of the campaign before

154   stabilizing at 294,234 doses per day on June 2, 2021 for the Shanghai

155   population (about 24 million individuals; details are reported in Sec. 5 of *SI*

156   *Appendix*).

157   iv)   **Vaccine efficacy (VE)**: The vaccine schedule requires two doses and VE is

158   estimated at 54.3% against the infection for the Delta variant. This estimate is

based on the efficacy measured against the historical lineages and the

reduction of neutralizing antibodies estimated for Delta variant in clinical

studies (see Tab. S1 of *SI Appendix* for details).We also explore higher VE

values as sensitivity analyses (Tab. S1 of *SI Appendix*). We also test a two-

dose schedule with a 14-day interval. In addition, COVID-19 vaccines may

not be equally effective across age groups in preventing infection. To

understand the impact of this assumption, we also tested a relative VE of 50%

and 75% for individuals aged 3-17 and 60+ years as compared to VE for

individuals aged 18-59 years.

v)   **Vaccine action**: We considered two ways in which VE could be below 100%:

an "all-or-nothing" vaccine (baseline analysis), in which the vaccine provides

full protection to a fraction VE of individuals who are vaccinated and no

protection to the remaining 1-VE vaccinated individual. The second option we

considered is a "leaky" vaccine in which all vaccinated individuals have a

certain level of protection to the infection corresponding to VE.

vi)   **Initial immunity**: As of July 2021, there is essentially no population

immunity from natural infection in China. For the sake of generalizability of

results to other countries with ongoing transmission, we have explored a

scenario where 30% of the population has initial natural immunity.

vii)   **Susceptibility to infection by age**: Children under 15 years of age were

estimated to have a lower susceptibility to SARS-CoV-2 infection as

compared to adults (i.e., individuals aged 15 to 64 years), while individuals

aged 65+ years had the highest susceptibility to infection.

viii)   **Immunity duration**: We let the transmission model run for one year,

assuming a life-long protection from natural infection or vaccination.

Comprehensive sensitivity analyses to evaluate the impact of the baseline assumptions

on our results are carried out as well (Tab. S1 in *SI Appendix*).

Alternative vaccination scenarios

We test three alternative scenarios to explore the potential for vaccination-induced

herd immunity, where: i) epidemic start is delayed from September 1, 2021 to October

1, 2021, and November 1, 2021; ii) the initial reproduction number varies between 1.1

and 6 to account for different intensity of NPIs (R_0^{NPIs}); iii) combinations of scenarios i

and ii. For scenario ii), we do not explicitly model single non-pharmaceutical

interventions (NPIs) such as case isolation, contact tracing, wearing masks, social

distancing, improved hygiene. Instead, the synergetic effect of these measures is

considered as a reduction of the reproduction number.

Data analysis

For each scenario, 100 stochastic simulations were performed. The output of these

simulations determined the distribution of the number of infections. We defined 95%

credible intervals as quantiles 0.025 and 0.975 of the estimated distributions.

We used the next-generation matrix approach to estimate the reproduction number, R_e.

Herd immunity is considered as achievable when R_e <1. Details are reported in Sec. 2-

3 in *SI Appendix*.

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data

interpretation, or writing of the report. The corresponding authors had full access to

all the data in the study and had final responsibility for the decision to submit for

publication.

Results

Baseline scenario

By forward simulating one year of epidemic and assuming no vaccine hesitancy,

continued vaccination efforts would lead to a final coverage of 97.6% of the target

population, which corresponds to 86.4% of the total population for strategy 1 (Fig.

1A). For strategy 2 and 3, the estimated coverage of the total population increases to

93.0% and 95.0%, respectively (Fig. 1B-C). Under any scenario, the mean incidence

of newly infected individuals never reaches 250 over 10,000 residents (Fig. 1D-F).

We estimated that the effective reproduction number at the time the infection is

seeded (R_e) is still well above the epidemic threshold, namely 4.50 (95%CI: 3.80-

4.98), 4.49 (95%CI: 3.90-4.89), and 3.67 (95%CI: 3.64-3.68) for strategy 1-3,

respectively. These estimates suggest that the vaccine coverage on September 1, 2021

is not enough to prevent onward transmission, regardless of the vaccination strategy.
R_e is estimated to cross the epidemic threshold (i.e., 1) on October 15, October 16,
and October 21, 2021 for strategy 1-3, respectively, due to the accumulation of
immune individuals both through the continue vaccination efforts and natural
infections (Fig. 1G-I). The estimated infection attack rates are 47.3% (95%CI: 45.8-
48.6%), 44.0% (95%CI: 42.3-45.9%), and 42.3% (95%CI: 39.5-44.0%) for strategy 1-
3, respectively (Fig.1J-L).

Although vaccine-induced immunity is not enough to prevent viral circulation, all the
scenarios considered are associated with substantial mitigation of COVID-19 burden.
We estimate the infection attack rate for the three vaccination strategies to decrease by
more than 50% with respect to a reference scenario with no interventions. Strategy 2-
3 achieve the highest reduction (55.3% and 57.0%) thanks to the inclusion of the age
group 3-17 years in the target population of the vaccination program (Fig. 2A-B).

These results were based on the assumption of an "all-or-nothing" vaccine (i.e., a
vaccinated individual will either develop full protection with probability given by the
vaccine efficacy or zero protection). To test the robustness of our findings to this
assumption, we also tested a "leaky" vaccine (i.e., the susceptibility to infection of
any vaccinated individual is reduced by a factor equal to the vaccine efficacy [11]) and
we obtained similar results with respect to the baseline analysis (Fig. S3 in *SI
Appendix*). Moreover, the obtained results are confirmed when the initial number of

seeds is varied in the range from 10 to 100 (Fig. S4 in *SI Appendix*), and when equal

susceptibility to infection by age is assumed (Fig. S5 in *SI Appendix*). Finally, we also

conducted a counterfactual analysis where we assume that a part of the population

was already immune before the start of the vaccination campaign (similar to the

situation in Western countries). Under this assumption, we found that a 30% initial

immunity proportion would not lead to R_e below the epidemic threshold for strategy 3

before September 1, the start date of the simulations (Fig. S6 in *SI Appendix*).

As regard the parameters regulating the vaccination process, we found that the

(overall) vaccine efficacy has the largest impact, followed by the vaccine efficacy of

individuals aged 3-17 and 60+ relative to individuals aged 18-59 years (Fig. S7 and

S8 in *SI Appendix*). On the other hand, the time between vaccination and maximum

protection and the time interval between the first and second dose have a more

moderate impact on the overall effectiveness of the analyzed vaccination strategies

(Fig. S9 and S10 in *SI Appendix*).

Scenario 1: Delaying the start of the epidemic

The findings presented thus far suggest that herd immunity against Delta variant

cannot be built through vaccination by September 1, 2021. Next, we test to what

extent the start of a new epidemic wave needs to be delayed (e.g., by keeping strict

restriction for international travels) to allow the immunity to build up in the

population, potentially reaching herd immunity levels. According to the daily vaccine

capacity used in the baseline scenario (based on the history of daily vaccination

capacity data up to May 23, 2021), we estimate that R_e remains above the epidemic

threshold for all three strategies even if the seeding of an epidemic is delayed to

November 1, 2021 - before that time, the vaccination coverage has reached the

maximum in the target population (i.e., 97.6%, 97.8%, and 97.8% for strategy 1-3)

(Fig. 3A and Fig. S11 in *SI Appendix*). In addition, in this case, strategies 2 and 3 lead

to a higher reduction of the infection attack rate with respect to the scenario with no

intervention (less than 4,200 per 10,000 individuals for an epidemic starting on

November 1, 2021 - Fig. 3B-C).

Scenario 2: Adopting NPIs in case of a new outbreak

The results presented so far suggest that herd immunity against Delta variant is not

achievable at any time point. Adopting NPIs as a response to an epidemic outbreak

can lower the transmission potential of the virus. It is thus worth investigating the

synergetic effect of vaccination programs combined with NPIs of different intensity. It

is important to note that we do not explicitly model every single measure to limit

transmission (e.g., case isolation, contact tracing, wearing masks, social distancing,

improved hygiene). These measures are implicit as concerted strategies that result in a

decreased reproduction number. We define the value of the reproduction number in a

fully susceptible population and under a certain level of NPIs as R_0^{NPIs}. We explored

R_0^{NPIs} in the range 1.1-6.0. Values between 1 and 2 are showed in the main text, while

larger values are shown in *SI Appendix* (Fig. S12).

The mean net reproduction number (which accounts both for immunity and

interventions) on September 1, 2021 for strategy 1 and 2 can be reduced to below 1

only when $R_0^{NPIs} \leqslant 1.3$, while for strategy 3 R_0^{NPIs} can be up to 1.6 (Fig. 3D). By forward

vaccinating and simulating 1 year of epidemic, substantial infections could be reduced

(close to 100% for $R_0^{NPIs} < 1.6$) thanks to the synergetic effect of vaccination and NPIs

(Fig. 3E-F and Fig. S13 in *SI Appendix*).

Scenario 3: Delaying the start of the epidemic and adopting NPIs

To further improve the potential for vaccination-induced herd immunity and reduce

COVID-19 burden, here we tested the combination of the two scenarios mentioned

above: delaying the start of the epidemic and adopting NPIs of different level of

intensity in response to a new outbreak.

Should an epidemic start in October-November 2021 and moderate NPIs (R_0^{NPIs} in the

range 1.5-2.0) are adopted, strategy 2 and 3 can succeed in blocking transmission

(Fig. 4B-C). In the case of strategy 1 (i.e., the vaccination policy excluding underage

individuals), strict NPIs ($R_0^{NPIs} < 1.3$) should be adopted to prevent a major epidemic

wave (Fig. 4A and Fig. S14 in *SI Appendix*).

The effectiveness of age-targeted vaccination strategies depend on the age-mixing

patterns of the population [12]. To test the robustness of our findings, we tested an

alternative contact matrix for China and we found consistent results (Fig. S15-16 in *SI*

Appendix).

Herd immunity threshold

Till now, herd immunity is unattainable for any vaccination strategy considering the

relatively low efficacy (54.3%) of the analyzed vaccine in preventing the infection

from the Delta. We thus explored the potential of herd immunity for the three

vaccination strategies given a higher efficacy (95%) (Fig. S17).

We estimated that R_e decreases below 1.0 only for strategy 2 and 3 (Fig. S17A). The

estimated herd immunity threshold under these two strategies is 94.0% and 88.1%

respectively, which suggests that level of immunity needed to lead the effective

reproduction number below the epidemic threshold is lower if vaccination is extended

to individuals aged 3-17 years early on.

We also estimated the infection attack rate under different vaccination coverages

under the assumption that vaccination stops at the time the epidemic is seeded. This

purely hypothetical scenario shows that when the adult population (18+ years of age)

is vaccinated (strategy 1), despite a fairly high estimated reproduction number (2.9),

the estimated infection attack rate is relatively low (13.2%) (Fig. S17B). In fact, given

the age-targeted vaccination program and the lack of natural immunity, the

susceptible population is mostly concentrated in the young population. The high

number of contacts in younger age groups, combined with the high vaccination

coverage in the rest of the population, lead to a fairly high reproduction number but,

at the same time, the infections are focused on a small segment of the population only

(young individuals) and thus the overall infection attack rate remains fairly low.

We also explored whether herd immunity is achievable or not and what is the herd

immunity threshold by estimating R_e under the assumption of having access to a

vaccine with a different efficacy (60%-100%) and exploring different scenarios on the

vaccine coverage.

Our results show that, for a vaccine with an efficacy lower than 85%, herd immunity

is unattainable, even in the extreme case where the vaccine coverage is 100% (Fig. 5).

Vaccine-induced herd immunity may only be achievable with higher VE and high

coverage. For example, for a vaccine with 90% efficacy against infection from the

Delta variant, more than 93% of the population would need to be vaccinated to reach

herd immunity (Fig. 5). In the presence of NPIs, the net reproduction number can be

reduced below the unit for lower vaccine efficacy and coverage values (Fig. S18 in *SI*

Appendix).

Discussion

Our study evaluates the feasibility of reaching herd immunity against the SARS-CoV-

2 Delta variant through vaccination, considering heterogeneity in population age

structure, age-specific contact patterns, vaccine efficacy, as well as biological

characteristics of SARS-CoV-2. Our findings show that herd immunity is unlikely to

be reached against the Delta variant given the relatively low efficacy of the current

vaccines (developed against the historical SARS-CoV-2 lineage), also in the presence

of prior natural immunity up to 30%. Even considering vaccines with higher efficacy,

our results show the key role of extending the vaccination program to school-aged

children in order to increase the potential of reaching herd immunity and reduce the

infection attack rate. If we consider a protection against the Delta variant of 90%

(which goes beyond current vaccines), herd immunity would require the vaccination

of 93% of the whole population. The adoption of NPIs could prevent the spread of a

major epidemic wave even when the herd immunity level is not reached, but such an

option obviously entails social and economic costs. Further, all strategies considered

in this study would mitigate the overwhelming majority of infections.

Our study explored if and when vaccination-induced herd immunity can be reached in

China. Under the hypotheses that the circulating strain has the same transmissibility

as Delta variant and that the vaccination campaign will not slow down due to vaccine

hesitancy, herd immunity seems to remain unreachable even in the extreme case

where the vaccine coverage is 100%. Nonetheless, it is important to remark that the

effectiveness of the vaccination program is impacted both by the natural immunity

accumulated in the population (which is close to 0 in China as of July 2021) and the

age structure of the population. In fact, in populations with a higher natural immunity

level and a lower proportion of children, herd immunity may be achievable.

Our findings point to the importance of adopting NPIs and/or self-precautionary

measures until herd immunity is reached or the burden of the epidemic becomes

manageable. These measures can either help delay the seeding of the infection (e.g.,

strict border control measures) or, should an epidemic start to unfold, mitigate its

burden (e.g., social distancing, contact tracing, testing, wearing masks, hygiene

practices, limiting contacts). However, questions remain about which NPIs need to be

implement, their intensity, and timing. Future studies are needed to address these

questions.

A key role to determine the success of a vaccination campaign is played by the

willingness-to-vaccine of the population. According to previous surveys on COVID-

19 vaccine hesitancy, vaccine acceptance in China was estimated to vary between

60.4% and 91.3% for general population aged 18 years and above [13-16], and may be

even lower for older adults [16] and healthcare workers [17]. Similar estimates were

obtained for several other countries including the UK (71.5%) [18] and the US (75.4%)

[18]. Given these levels of vaccine hesitancy, achieving high levels of coverage may

remain an elusive target. Efforts to increase population's confidence and willingness

to be vaccinated will thus be of paramount importance to allow a return to a pre-

COVID-19 lifestyle. Our study shows that the spread of the more transmissible Delta

variant has substantially increased the herd immunity threshold to a level that may not

be feasible in any population, so that mitigation strategies become even more relevant.

Previous studies have estimated the herd immunity threshold either through natural infection or vaccination under the assumption of an homogenously mixed population [19-21], but heterogeneity in contact structure, age structure of the population, susceptibility to infection by age, and order in which individuals are vaccinated are all key factors shaping the herd immunity level [8]. The developed model is based on social mixing patterns estimated for the Shanghai population [9] and on China-specific data on COVID-19 epidemiology (population immunity, etc.). Nevertheless, the introduced modeling framework is flexible and can be tailored to other countries. We tested a scenario somehow resembling the situation in the USA, where we considered naturally immunity and the adoption of BNT162b2/Pfizer vaccine, whose efficacy against the Delta variant was estimated at 79%. Also, in this scenario, we estimate that herd immunity may not be reached (Fig. S19 in *SI Appendix*). Moreover, vaccination hesitancy may jeopardize the vaccination effort in the US and other Western countries as well.

This study is prone to the limitations pertaining to modeling exercises. First, VE against infections from the Delta variant was inferred instead of directly measures from epidemiological observations. Moreover, VE for children have not been estimated for the vaccines in use in China; therefore, we have assumed the same VE as in adults based on immunogenicity studies [22]. Given such a lack of field evidence, we have conducted a sensitivity analysis where a lower vaccine efficacy is assumed

for children. The overall conclusions of the study do not change. Still, further data on

age-specific vaccine efficacy could help refine the obtained estimates.

Second, we assumed that immunity induced either from infection or vaccination lasts

more than the time horizon considered in the simulations (i.e., 1 year). There are both

evidence from laboratory studies and the field suggesting that the protection lasts

several months [23]. Despite these preliminary pieces of evidence, the duration of the

immunity remains a research area of paramount importance and intrinsically linked to

viral evolution. It is also possible that waning immunity will continue to provide

protection against severe disease but only partial against infection or transmission,

which affects the herd immunity threshold. Overall, the duration and quality of

immunity will determine the periodicity of COVID-19 outbreaks globally [24,25].

Third, in the baseline scenario, we referred to an inactivated SARS-CoV-2 vaccine

(BBIBP-CorV) taken to be 54.3% efficacious against the Delta variant infection.

However, several other vaccines (including CoronaVac, WBIP-CorV, Ad5-nCoV, and

ZF2001) are licensed and have been used in China. We varied vaccine efficacy up to

79% in sensitivity analyses. The main conclusion about the potential of herd

immunity and the need to extend the vaccination campaign to children as well as to

use more efficacious vaccines is unaltered.

In conclusion, based on the current evidence, reaching vaccine-induced herd

immunity in a population with little/no natural immunity is challenging. Key steps

will be the authorization of COVID-19 vaccines for children, and minimize vaccine

hesitancy. These, together with highly efficacious vaccines or booster vaccinations,

will be even more crucial given the possible emergence of new, even more

transmissible, SARS-CoV-2 variants. Importantly, even if herd immunity is unlikely

to be reached, vaccination will still dramatically reduce COVID-19 burden.

Contributors

H.Y. conceived and designed the study. H.Y. and M.A. supervised the study. M.A.

designed the model. H.L. developed the model. H.L., J.Z., J.C., J.Y., X.D. analyzed

the model outputs. H.L., C.P., X.D, and Z.C. prepared the tables and figures. H.L. and

J.Z. prepared the first draft of the manuscript. W.Z., Q.W. and X. C. participated in

data collection. X.C and Z.C. updated the relative literatures. H.Y., M.A. W.Z. and C.

V. revised the content critically. All authors contributed to review and revision and

approved the final manuscript as submitted and agree to be accountable for all aspects

of the work.

Declaration of interests

H.Y. has received research funding from Sanofi Pasteur, GlaxoSmithKline, Yichang

HEC Changjiang Pharmaceutical Company, and Shanghai Roche Pharmaceutical

Company. M.A. has received research funding from Seqirus. None of those research

funding is related to COVID-19. All other authors report no competing interests.

Disclaimer

This article does not necessarily represent the views of the NIH or the US government.

Acknowledgments

The study was supported by grants from the Key Program of the National Natural Science Foundation of China (82130093), European Union Grant 874850 MOOD (MOOD 000), and the National Institute for Health Research (NIHR) (grant no. 16/137/109) using UK aid from the UK Government to support global health research. We also acknowledge grant from Shanghai Key Laboratory of Infectious Diseases and Biosafety Emergency Response (20dz2260100). The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the UK Department of Health and Social Care.

Data sharing

The data and code that support the findings of this study will be made available in GitHub upon manuscript acceptance.

Figure 1. Time series of vaccine coverage, daily incidence of new SARS-CoV-2

Delta variant infections, effective reproductive number, R_e, and fraction of

immune population. **A** Age-specific vaccine coverage over time for strategy 1.

Vaccination program is assumed to be initiated on November 30, 2020 (i.e., the time

that the vaccine doses administrated was first officially reported in China). The dotted

lines correspond the start of epidemic. The inserted table shows the age-specific

coverage for the two key time points (the start of epidemic (i.e., September 1, 2021),

and the time that the coverage keeps constant (i.e., September 24), respectively). The

line corresponds to the mean value, while the shaded area represents 95% quantile

intervals (CI). **B** As A, but for strategy 2. **C** As A, but for strategy 3. **D** Daily

incidence of new SARS-CoV-2 infections per 10,000 individuals for strategy 1 (mean

and 95% CI). **E** As D, but for strategy 2. **F** As D, but for strategy 3. **G** Effective

reproduction number R_e over time (mean and 95% CI), as estimated using the Next-

Generation matrix method from the time series of susceptible individuals for strategy

1. The shaded area in gray indicates the epidemic threshold $R_e = 1$. The numbers

around the shaded area indicate when R_e cross this threshold (i.e., October 15) for

strategy 1. **H** As G, but for strategy 2. **I** As G, but for strategy 3. **J** Proportion of

immune population due to either natural infection or vaccination over time for

strategy 1. **K** As J, but for strategy 2. **L** As J, but for strategy 3.

Figure 2. Burden of COVID-19 in the baseline scenario. A Cumulative number of

infections per 10,000 individuals after 1 simulated year for *reference scenario* and

three vaccination strategies (mean and 95% CI). *Reference scenario* indicates no

vaccination and no NPIs with R_0=6.0 at the beginning of transmission. Infections

consist of unvaccinated and vaccinated individuals. The bar corresponds to the mean

value, while the vertical line represents 95% quantile intervals. **B** Reduction in

infections (mean and 95% CI) with respect to the *reference scenario* in different age

groups and the total population. The reduction is defined as the estimated number of

infections after 1 year since the introduction of the initial infected individuals under

reference scenario minus the one under the vaccination strategy, relative to the

estimated number under *reference scenario*. The 95% CI of the reduction may cross 0

as the burden between reference scenario and vaccination scenario is approximately

the same in some simulations. We thus trimmed the lower limit of 95% CI at 0

through the manuscript.

Figure 3. Impact of delaying the start of the epidemic and adopting NPIs. A

Estimated effective reproduction number (R_e, mean and 95% CI) as a function of

vaccine coverage at the time the infection is seeded (i.e., September 1, October 1 and

November 1). Colors refer to the scenario of delaying the start of the epidemic to

different date. The shaded area in gray indicates $R_e \leq 1$. **B** Cumulative number of

infections per 10,000 individuals after 1 simulated year for *reference scenario* and

three vaccination strategies (mean and 95% CI). *Reference scenario* indicates no

vaccination and no NPIs with R_0=6.0 at the beginning of transmission. **C** Reduction in

infections (mean and 95% CI) with respect to the *reference scenario*. **D** As A, but for

estimated net reproduction number (R_t, mean and 95% CI) adopting different intensity

of NPIs, R_0^{NPIs}. **E** As B, but for the scenario of adopting different intensity of NPIs. **F**

As C, but for the scenario of adopting different intensity of NPIs.

532

533 **Figure 4. Impact of delaying the start of the epidemic start and adopting NPIs on**

534 **estimated net reproduction number R_t.** **A** Estimated net reproduction number (R_t)

535 as a function of R_0^{NPIs} and epidemic start date for strategy 1. The bold line in black

536 indicates the herd immunity threshold $R_t = 1$. **B** As A, but for strategy 2. **C** As A, but

537 for strategy 3.

Figure 5. The impact of vaccine efficacy and vaccine coverage on estimated

effective reproduction number R_e. The bold line in black indicates the herd

immunity threshold R_e =1. R_e is estimated in the scenario that all individuals are

eligible to vaccination and vaccinated 2 doses before the epidemic starts.

Reference

1.   Chinese Center for Disease Control and Prevention. Update on COVID-19 as of 24:00 on
August 9, 2021. 2021. http://2019ncov.chinacdc.cn/2019-nCoV/ (accessed August 10 2021).
2.   World Health Organization. COVID-19 Weekly Epidemiological Update, 2021.
3.   Outbreak.info. B.1.617.2 Lineage Report. 2021. https://outbreak.info/situation-
reports?pango=B.1.617.2&loc=IND&loc=GBR&loc=USA&selected=Worldwide (accessed Aug 10
2021).
4.   Zhang M, Xiao J, Deng A, et al. Transmission Dynamics of an Outbreak of the COVID-19
Delta Variant B.1.617.2 – Guangdong Province, China, May – June 2021. *China CDC Weekly*
2021; **3**: 584-6.
5.   National Health Commision of the People's Republic of China. Update on the doses of
COVID-19 vaccine administration. 2021. http://www.nhc.gov.cn/xcs/yqjzqk/list_gzbd.shtml.
6.   Diekmann O, Heesterbeek H, Britton T. Mathematical Tools for Understanding Infectious
Disease Dynamics: Princeton University Press; 2013.
7.   May RM, Anderson RM. Spatial heterogeneity and the design of immunization programs.
Mathematical Biosciences 1984; **72**(1): 83-111.
8.   Britton T, Ball F, Trapman P. A mathematical model reveals the influence of population
heterogeneity on herd immunity to SARS-CoV-2. *Science* 2020; **369**(6505): 846-9.
9.   Zhang J, Klepac P, Read JM, et al. Patterns of human social contact and contact with animals
in Shanghai, China. *Sci Rep* 2019; **9**(1): 15141.
10.   The State Council Information Office PRC. BBIBP-Corv and CoronaVac are extended to
children aged 3-17 years. 2021. http://www.gov.cn/xinwen/gwylflkjz160/index.htm.
11.   Halloran ME, Longini IM, Struchiner CJ. Modes of Action and Time-Varying VES. Design
and Analysis of Vaccine Studies. New York, NY: Springer New York; 2010: 131-51.
12.   Yang J, Marziano V, Deng X, et al. Despite vaccination, China needs non-pharmaceutical
interventions to prevent widespread outbreaks of COVID-19 in 2021. *Nature Human Behaviour*
2021.
13.   Wang J, Jing R, Lai X, et al. Acceptance of COVID-19 Vaccination during the COVID-19
Pandemic in China. *Vaccines (Basel)* 2020; **8**(3).
14.   Chen M, Li Y, Chen J, et al. An online survey of the attitude and willingness of Chinese
adults to receive COVID-19 vaccination. *Hum Vaccin Immunother* 2021: 1-10.
15.   Wang C, Han B, Zhao T, et al. Vaccination willingness, vaccine hesitancy, and estimated
coverage at the first round of COVID-19 vaccination in China: A national cross-sectional study.
Vaccine 2021; **39**(21): 2833-42.
16.   Gan L, Chen Y, Hu P, et al. Willingness to Receive SARS-CoV-2 Vaccination and Associated
Factors among Chinese Adults: A Cross Sectional Survey. *Int J Environ Res Public Health* 2021;
18(4).
17.   Zhang HJ, Ding LL, Pan XJ, et al. Willingness to receive novel coronavirus vaccine and
factors influencing willingness among healthcare workers in Zhejiang province. *Chin J Vacc*
Immun 2021.
18.   Lazarus JV, Ratzan SC, Palayew A, et al. A global survey of potential acceptance of a
COVID-19 vaccine. *Nat Med* 2021; **27**(2): 225-8.
19.   Hodgson D, Flasche S, Jit M, Kucharski AJ, Group CC-W, Centre for Mathematical Modelling
of Infectious Disease C-WG. The potential for vaccination-induced herd immunity against the

SARS-CoV-2 B.1.1.7 variant. *Euro Surveill* 2021; **26**(20).

20. Kwok KO, Lai F, Wei WI, Wong SYS, Tang JWT. Herd immunity - estimating the level

required to halt the COVID-19 epidemics in affected countries. *J Infect* 2020; **80**(6): e32-e3.

21. Omer SB, Yildirim I, Forman HP. Herd Immunity and Implications for SARS-CoV-2 Control.

JAMA 2020; **324**(20): 2095-6.

22. Zhang Y, Zeng G, Pan H, et al. Safety, tolerability, and immunogenicity of an inactivated

SARS-CoV-2 vaccine in healthy adults aged 18-59 years: a randomised, double-blind, placebo-

controlled, phase 1/2 clinical trial. *Lancet Infect Dis* 2021; **21**(2): 181-92.

23. Wang Z, Muecksch F, Schaefer-Babajew D, et al. Naturally enhanced neutralizing breadth

against SARS-CoV-2 one year after infection. *Nature* 2021; **595**(7867): 426-31.

24. Lavine JS, Bjornstad ON, Antia R. Immunological characteristics govern the transition of

COVID-19 to endemicity. *Science* 2021; **371**(6530): 741-5.

25. Saad-Roy CM, Wagner CE, Baker RE, et al. Immune life history, vaccination, and the

dynamics of SARS-CoV-2 over the next 5 years. *Science* 2020; **370**(6518): 811-8.