regentrans: a framework and R package for using genomics to study regional pathogen transmission

Sophie Hoffman, BS*[1]; Zena Lapp, PhD*[1]; Joyce Wang, PhD[2]; Evan S Snitkin, PhD[2,3]

[1]Department of Computational Medicine and Bioinformatics, University of Michigan, 1150 W. Medical Center Dr. Ann Arbor, MI, 48109-5680
[2]Department of Microbiology and Immunology, University of Michigan, 1150 W. Medical Center Dr. Ann Arbor, MI, 48109-5680
[3]Department of Medicine, Division of Infectious Diseases, University of Michigan, 1150 W. Medical Center Dr. Ann Arbor, MI, 48109-5680

*These authors contributed equally to this work.

Corresponding author:
Evan Snitkin
esnitkin@umich.edu

Keywords

R package, regional transmission, genomics, whole genome sequencing, software, pathogen spread

Repositories

https://github.com/Snitkin-Lab-Umich/regentrans/

Abstract

Increasing evidence of regional pathogen transmission networks highlights the importance of investigating the dissemination of multidrug-resistant organisms (MDROs) across a region to identify where transmission is occurring and how pathogens move across regions. We developed a framework for investigating MDRO regional transmission dynamics using whole-genome sequencing data and created regentrans, an easy-to-use, open source R package that implements these methods (https://github.com/Snitkin-Lab-Umich/regentrans). Using a dataset of over 400 carbapenem-resistant *Klebsiella pneumoniae* isolates collected from patients in 21 long-term acute care hospitals over a one-year period, we demonstrate how to use our framework to gain insights into differences in inter- and intra-facility transmission across different facilities and over time. This framework and corresponding R package will allow investigators to better understand the origins and transmission patterns of MDROs, which is the first step in understanding how to stop transmission at the regional level.

Impact statement

Increasing evidence suggests that pathogen transmission occurs across healthcare facilities.
Genomic epidemiologic investigations into regional transmission shed light on potential drivers
of regional prevalence and can inform coordinated interventions across healthcare facilities to
reduce transmission. Here we present a framework for studying regional pathogen transmission
using whole-genome sequencing data, and a corresponding open-source R package,
regentrans, that implements these methods to streamline analyses and make them more
accessible to other researchers and public health practitioners. We also discuss how these
methods can be extended to study transmission in other settings.
53
54 **Data summary**
55
56 The authors confirm all supporting data, code and protocols have been provided within the
57 article or through supplementary data files.
58
● The regentrans R package can be downloaded from GitHub: https://github.com/Snitkin-
Lab-Umich/regentrans/
● The manuscript figures are generated from regentrans example data and can also be
found on GitHub: https://github.com/Snitkin-Lab-
Umich/regentrans/tree/master/vignettes/manuscript_figures
● The example data used in the package and manuscript is from BioProject accession no.
PRJNA415194. The specific SRA accession numbers can be found in supplementary file
S1. The metadata corresponding to these sequences can be found on the SRA Run
Selector (isolate column) and as example data in the regentrans package.
● The KPNIH1 sequence was used as the reference genome (SRA accession number
SRZ080789)
70
71 **Introduction**
72
Multidrug-resistant organisms (MDROs) are a global public health threat due to limited
treatment options paired with widespread global transmission [1]. Healthcare facilities in
particular, where critically ill patients reside in close proximity to one another, are hotspots of
MDRO transmission [2]. Furthermore, increasing evidence suggests that substantial
transmission occurs not only within facilities, but also between facilities in regional healthcare
networks, and that intra- and inter-facility transmission does not occur evenly across these
networks [3, 4]. This observation, paired with limited resources for state and regional public
health efforts, necessitates the identification of optimal intervention locations to reduce overall
regional prevalence. Investigating MDRO transmission from a regional perspective can shed
light on the origin and spread of MDROs, providing critical information for precision public health
interventions to allocate resources to maximally reduce transmission across a region [5, 6].
84
Understanding where and how recent transmission is occurring is an integral first step in
developing interventions to curb transmission at the regional level. A powerful tool for studying
regional pathogen transmission is whole-genome sequencing, which allows us to investigate
pathogen movements at very high resolution [4, 6]. Several studies have used whole-genome

sequencing, sometimes paired with additional epidemiological metadata, to gain insights into
locations [6, 7] and drivers [3, 4, 8] of elevated intra- or inter-facility transmission. These types of
analyses have the potential to transform our public health response to MDROs if they are
regularly performed at, or in collaboration with, regional public health centers. However, regional
transmission analyses require careful consideration of the biological and epidemiological
context, and the methods can be time consuming to implement.
95
Here, we aim to reduce the barrier for performing regional transmission analyses by providing a
framework for studying regional pathogen transmission using whole-genome sequencing data,
suggesting techniques for making analysis decisions, and presenting regentrans, an open-
source R package that implements these methods. We discuss methods to study transmission
within and between healthcare facilities using whole-genome sequencing data from single-
colony isolates from patients, and discuss how these methods can be applied to study
transmission within and between other locations such as zip codes. The methods presented
here focus on studying recent transmission in a clonal set of isolates and can be applied to
investigate overall transmission or transmission patterns over time, and to compare the
transmission dynamics of different strains circulating in a region. We believe that these tools will
help investigators more easily perform regional pathogen transmission analyses, and thus
potentially guide interventions to reduce transmission.
108
109 **Investigating regional transmission patterns**
110
111 Below we describe the questions, data, and methods for studying regional pathogen
112 transmission. More details about using regentrans to implement these methods can be found in
113 the package vignette, a step-by-step guide to using the package for an analysis.
114
115 **Questions**
116
117 Our framework for studying regional pathogen transmission aims to help investigators
118 interrogate the following questions (**Table 1**):
119
120   1. Is transmission occurring within and/or between facilities?
121   2. What facilities is transmission occurring within/between?
122   3. Have transmission dynamics changed over time?
123   4. Is transmission occurring along paths of higher patient/person flow?
124   5. Are there any observable geographic trends in prevalence/transmission?
125
126 **Data**
127
128 <u>Input data</u>
129
Whole-genome sequences of isolates combined with corresponding metadata can be used to
identify the genetic relatedness between isolates and subsequently investigate intra- and inter-
facility transmission (**Figure 1**). Note that users can input data from multiple isolates from a

given patient as the corresponding regentrans functions use this information in generating the
output data (see the "Handling multiple isolates from a given patient" section for more details).
Below we describe each component of the input data in more detail.
136
137 *A note on study design and sampling density*
138
The data can come from studies such as prospective observational studies, point-prevalence
surveys, and regional surveillance across different facilities in a region. While comprehensive
sampling of culture or surveillance isolates from all facilities in a region is ideal, inferences can
still be made using a subset of isolates and a subset of facilities. In general, users must
consider how the study design, missing samples, and missing facilities may influence the results
and use this to guide their interpretations. We find it useful to perform thought experiments
considering what the results might look like under counterfactual scenarios where more
comprehensive data was included.
147
148 *Genetic data*
149
150 The core functionality of regentrans assumes that users have whole-genome sequences of
151 each isolate to be included in the analysis. As the methods we present here are focused on
152 identifying recent transmission events, we suggest that they be used only on closely related
153 isolates, e.g. ones within the same sequence type (ST) or clonal complex. However,
154 comparisons between the transmission dynamics of different groups can be performed.
155
Depending on the method being used to study transmission, the genetic data required is either
a phylogeny of all isolates, a genetic distance matrix, or a variant alignment. We highly
recommend using a recombination-filtered variant alignment to ensure that variants included in
the analysis arose from vertical gene transfer rather than horizontal gene transfer, as an
assumption of the described methods is that isolates with a more recent common ancestor are
more closely related at the variant level. Users can also remove variant sites known to be under
positive selection. We suggest using Gubbins [9] to mask recombinant sites, IQ-TREE [10] to
generate a maximum-likelihood phylogeny from the recombination-masked alignment, and the
ape package [11] in R to calculate a pairwise genetic distance matrix. The genetic distance
matrix can be any type of genetic distance as long as smaller numbers indicate more similarity
between two isolates. Two examples of genetic distances are pairwise single nucleotide variant
(SNV) distances (can be calculated with ape::dist.dna()) and patristic distances, or the sum of
the branch lengths that connect two isolates on a phylogeny (can be calculated with
ape::cophenetic.phylo()). While patristic distances properly account for ancestral relationships
between strains, we often use pairwise SNV distances as they are easier to interpret and are
the current standard with respect to transmission inference. Concordance between results using
these two distances would also increase confidence in using pairwise SNV distances for
interpretability.
174
175 *Isolate metadata*
176

177 For each isolate, users need to know the facility at which the isolate was collected and the
178 patient from whom the isolate was collected. If only one isolate was collected from each patient,
179 patient information is not required. Furthermore, isolate collection dates are needed if
180 researchers are interested in studying transmission over time.
181
182 *Facility-level connectedness data*
183
184 Researchers can investigate the relationship between genetic distance and patient transfer
185 between facilities, which requires a patient transfer network that minimally includes all facilities
186 represented in the dataset. Additionally, it is possible to visualize and quantify potential
187 geographic trends in prevalence and transmission, for which information on the physical
188 distance, or driving time, between facilities is required.
189
190 <u>Datasets used in the package and for analyses</u>
191
192 *Genomic data*
193
The genomic data used for this manuscript, and included in the regentrans package, were
generated from whole-genome sequences of clinical isolates obtained from 21 long-term acute
care hospitals across the U.S. [4]. The data was processed as in [12]. Briefly, trimmed Illumina
short reads were aligned to the KPNIH1 reference genome (BioProject accession no.
PRJNA73191) using the Burrows-Wheeler short-read aligner (bwa v0.7.17) [13] and
recombinant sites were masked using Gubbins v2.3.4 [9]. We used the Gubbins variant output
fasta file to generate a pairwise single nucleotide variant (SNV) distance matrix using the
dist.dna() function (method = 'N', pairwise.deletion = TRUE, as.matrix = TRUE) and a patristic
distance matrix using the cophenetic.phylo() function, both in ape v5.5 [11]. IQ-TREE v1.6.12
[10] was used to generate a whole-genome phylogeny of all isolates. All non-recombinant
variants, including variants that may have been under positive selection, were included when
calculating pairwise SNV distances and generating the phylogeny. For all analyses, the data
was subset to include only ST258 isolates. Sequence types were determined using Kleborate
v0.4.0 [14].
208
209 *Isolate metadata*
210 We used information on the facility and the patient the isolate was taken from, as well as the
211 isolate collection date.
212
213 *Patient transfer data*
214
215 Aggregate patient transfer data from all hospitals in the state of California was used to calculate
216 paths of maximum patient flow between the 11 Los Angeles area facilities in our dataset. The
217 data and methods are described in [4].
218
219 *Geographic data*
220

221 The example geographic data was de-identified by adding random horizontal, vertical, and
222 rotational shifts using the R package tangles v0.8.1 [15].
223
224 **Methods**
225
226 <u>Q0: How do you choose pairwise genetic distance thresholds?</u>
227
Several of the methods discussed below rely on interpreting, comparing, or thresholding
pairwise genetic distances between isolates to make inferences. It is generally understood that
small pairwise distances between isolates implies recent transmission [16–18], but that this
method is not entirely accurate due to within-host evolution and variable mutation rates [19, 20].
To identify recent transmission pairs using pairwise distances, investigators must choose a
threshold to determine what pairs are considered closely related [16–18]. The threshold for
"closely related" depends on the pathogen mutation rate and genome size, and the setting.
Within a given time period, a higher mutation rate or a larger genome size will lead to higher
pairwise distances. Additionally, the mutation rate of pathogens in outbreak settings is often
higher than endemic settings [21]. Thus, for a given pathogen, closely related isolate pairs in an
outbreak setting will likely have a higher pairwise genetic distance than closely related pairs in
an endemic setting. For this reason, knowledge of the epidemiologic context of the isolates, and
the species or sequence type itself, is essential for correctly interpreting pairwise genetic
distance distributions. Here, we discuss two methods for identifying reasonable pairwise SNV
distance thresholds. Similar methods can be used to identify thresholds for other pairwise
genetic distance metrics, such as pairwise patristic distances.
244
One way to choose a pairwise SNV distance threshold is to use the genome length and
mutation rate. For instance, in the context of the dataset we use here, *K. pneumoniae* ST258
isolates from an endemic setting, we could calculate a pairwise SNV distance threshold of 11
SNVs based on the KPNIH1 reference genome length of 5,394,056 base pairs and a mutation
rate of 1.03e-6 mutations per base pair per year [22] (2 isolates * 5,394,056 bases * 1.03e-6
mutations per base per year per isolate). Mutation rates can be estimated for measurably
evolving populations with a temporal signal [21–23]. However, mutation rate estimates are less
reliable for lineages with high among-lineage rate variation or phylogenetic and temporal
clustering [24], and are unreliable for datasets with no temporal signal [24]. Therefore, it is often
difficult or impossible to reliably calculate the evolutionary rate of the pathogen for the particular
dataset being used, and more general estimates of mutation rate may not be translatable.
256
If a mutation rate cannot be reliably calculated, or to perform a sensitivity analysis,
one way to identify potential pairwise SNV distance thresholds is to use the assumption that
pairs of individuals from the same facility will be enriched for recent transmission relative to
pairs from different facilities. In particular, this method requires you to plot the fraction of isolate
pairs from the same facility for increasing pairwise SNV distances. A decrease in the fraction of
intra-facility isolate pairs at a given SNV distance suggests a potentially reasonable threshold.
Performing this analysis on our data indicated that using SNV distance thresholds of ≤ 10 and ≤
6 appear to be reasonable (**Figure 2**). We chose to use these two thresholds for our sensitivity

analysis as they are more directly supported by our data compared to the more general
mutation rate analysis using a mutation rate that was not calculated from the same dataset.
However, in this case the results would likely be similar as the mutation rate analysis suggests
using a threshold of 11 SNVs, which is similar to the 10 SNV threshold we use here.
269
270 When performing analyses where choosing a pairwise SNV distance threshold is necessary, we
271 suggest that investigators evaluate the robustness of their results by comparing their findings for
272 different pairwise SNV distance thresholds. Here, we compare the results using a threshold of 6
273 and 10, but a wider range of values can be used, particularly in instances with more uncertainty
274 about what threshold should be used.
275
276 Q1: Is transmission occurring within and/or between facilities?
277
One of the first questions an investigator might ask about isolates collected from a certain
region is if transmission is occurring within particular facilities and/or between facilities.
Inspecting pairwise genetic distances between all isolates can provide information about the
extent of recent transmission both within and between facilities, which will often manifest as an
enrichment in closely related isolate pairs between patients (i.e. isolate pairs with small pairwise
genetic distances; see note above on what to consider closely related). In our example dataset,
we observe an enrichment in both closely related intra-facility pairs and inter-facility pairs
(pairwise SNV distance of ≤ 10 or ≤ 6), suggesting that recent transmission is occurring both
within and between facilities (**Figure 3**).
287
One limitation of the pairwise genetic distance method for investigating transmission is that it
may miss transmission events occurring over longer time periods when there is, for instance,
sparse sampling and more evolutionary divergence between isolates. Another limitation is that
researchers must define a threshold for what is considered closely related, and different
thresholds may lead to different conclusions. Therefore, we recommend performing a sensitivity
analysis across several pairwise genetic distance thresholds and genetic distance metrics to
investigate the robustness of the results to different thresholds.
295
296 Q2: What facilities is transmission occurring within/between?
297
Once we have investigated the extent of transmission occurring within and between facilities on
a general scale, we can dig deeper into identifying certain facilities and facility pairs with closely
related isolates. Intra-facility transmission can be investigated using phylogenetic context, and
inter-facility transmission can be investigated by taking a population-level approach computing
genetic flow between facilities. These methods are both threshold-free. Intra- and inter-facility
transmission can also be investigated by identifying probable recent transmission events
between facilities, which requires a pairwise genetic distance threshold to be defined.
305
306 *Investigating intra-facility transmission using the phylogeny*
307

One way to investigate the extent of intra-facility transmission is to identify maximum subclades
that all originate from distinct patients from the same facility and quantify the size of these
clusters. Larger clusters indicate more intra-facility transmission, as those isolates are all more
closely related to one another than to isolates from other facilities. Furthermore, the range of
sampling dates of each cluster can provide insight into whether the cluster is from an intra-
facility outbreak (smaller date range) or from sustained intra-facility transmission (larger date
range). In our dataset, we see that some facilities (e.g. facility D) exhibit extensive intra-facility
transmission as evidenced by large cluster sizes, while others exhibit relatively little intra-facility
transmission (e.g. facility K) (**Figure 4**). All large clusters also have large date ranges,
suggesting that they are from sustained intra-facility transmission rather than a shorter-term
outbreak. However, it is also important to note that, under these circumstances of sustained
transmission, isolates within a cluster may be distantly related, particularly if there are few
exportation events from that facility to other facilities (e.g. if the facility is more geographically
isolated). This may be considered a strength of the method if users are interested in
investigating transmission across longer timescales.
323
There are a few key limitations to investigating intra-facility transmission based on phylogenetic
cluster size that must be considered when interpreting results. First, cluster sizes may be biased
when there is skewed sampling across facilities [8], where larger clusters may be identified for
over-represented facilities and smaller clusters for under-represented facilities. For instance, if
unsampled strains from under-represented facilities were to be included, they may increase the
cluster size of pre-existing clusters from that facility or decrease the size of clusters from
another facility by being interspersed among those samples. Therefore, we recommend careful
consideration of potential sampling bias prior to comparing cluster sizes across facilities.
Second, if there is frequent intra- and inter-facility transmission, recent transmission events
between facilities may not be identified as these clusters will be broken up by recent inter-facility
transmission events. We provide the option to reduce the cluster "pureness" to attempt to
accommodate this limitation. Finally, the clustering method does not directly allow investigators
to assess the extent of inter-facility transmission. The genetic flow and pairwise genetic distance
methods described below address these limitations of the cluster size method.
338
339 *Genetic flow between facilities*
340
Identifying the extent to which variants are shared among isolates at different facilities by
calculating genetic flow between facilities (Fsp) [8] provides a threshold-free population-level
approach to investigating the extent of inter-facility transmission. Fsp is a measure of
population-level similarity between facilities, where an Fsp value closer to 0 indicates more
genetically similar populations and a value closer to 1 indicates more genetically different
populations. This method is particularly useful in endemic scenarios when the relationship
between individual isolates may be relatively diffuse due to frequent patient transfer over time.
Using our dataset, we found that certain facility pairs (e.g. facilities G and P) have more genetic
flow between them compared to others, indicating that there is likely more transmission between
those facilities (**Figure 5**). Facilities that have high Fsp compared to other facilities (e.g. J) have
populations that are genetically distant from all other facilities, which indicates that they are

likely less involved in the inter-facility transmission network. Note that one limitation of using Fsp
is that small sample sizes may overestimate genetic differences, although this is less of a
concern when using many variant sites [27].
355
356   *Pairwise genetic distance threshold*
357
Using a pairwise genetic distance threshold, the number of closely related pairs within and
between facilities can be determined and used to identify facilities and with more or less putative
spread. For instance, we observed that not only are there differing extents of spread at different
facilities, but some facilities have more closely related intra-facility pairs than inter-facility pairs
(e.g. facility D), while other have more even amounts of closely related intra- and inter-facility
pairs (e.g. facility H) (**Figure 6A**). These findings are generally concordant with our findings from
the phylogenetic clustering and genetic flow analyses as well (e.g. more intra-facility
transmission in facility D, isolates from facility H being related to isolates from other facilities).
Furthermore, as there are limitations to choosing genetic distance cutoffs, we highly recommend
doing a sensitivity analysis by choosing several different thresholds and using different genetic
distance metrics to see how robust the results are to these changes.
369
One limitation of visualizing the number of closely related pairs within and between facilities is
that it does not provide insight into the potential transmission networks among patients in these
pairs. For instance, we do not know how often each patient isolate is included in these pairs, or
whether the pairs are closely related to each other. This question can be probed by visualizing
the connectedness of patients with closely related isolates. For instance, in our data we see that
there is an isolated network of closely related isolates from facility G, as well as a network of
closely related isolates from facility D that is connected to isolates from other facilities (**Figure
6B**).
378
379   The phylogenetic clustering, genetic flow, and pairwise genetic distance methods provide
380   complementary information related to intra- and inter-facility transmission. We suggest
381   evaluating the strengths and limitations of each method with respect to the dataset being used
382   and the questions of interest to determine which method(s) to use for a given application.
383   Concordance between methods provides additional confidence in the results.
384
385   Q3: Have transmission dynamics changed over time?
386
All of the methods described above can be applied to discrete time windows to gain insight into
whether transmission dynamics have remained stable or changed over time. For instance, in an
outbreak setting we observed an increase in intra-facility transmission followed by an increase
in inter-facility transmission [7]. In an endemic setting, these trends may remain more stable
over time. In our data, we observe an increase in the total number of pairs from 2014 to 2015,
but no change in the distribution of closely related intra- or inter-facility isolate pairs (**Figure 7**).
393
394   Q4: Does transmission correlate with patient/person flow?
395

In addition to only using genomic information to study transmission, inter-facility transmission
can be studied in the context of patient flow between facilities. While sometimes investigators
may have access to patient-level information regarding prior facility exposures, this information
is often not available. In this case, aggregate patient transfer data can be used to study the
relationship between patient flow and transmission. The simplest way to do this is to determine
whether there is a relationship between direct flow between facilities and either the number of
closely related pairs defined by pairwise SNV distances, the actual values of pairwise SNV
distances, or genetic flow (Fsp). To take into account potential indirect transfers that may
influence transmission, a more complex algorithm can be used to identify paths of maximum
patient flow between facilities, and then this can be compared to metrics of genomic relatedness
[4]. These analyses can provide insight into whether patient flow may be driving transmission
between facilities. For instance, when subsetting our data to the 11 Los Angeles area facilities
for which we have information on inter-facility patient transfer events, we observed a negative
correlation between patient flow and genetic flow (Fsp), indicating that facilities connected by
more patient flow often have more similar populations (**Figure 7A**). We also observed a
negative correlation between patient flow and minimum pairwise SNV distance between
facilities (**Figure 7B**), and a positive correlation between patient flow and the number of closely
related isolate pairs between facilities (**Figure 7C**). Taken together, this suggests that patient
flow may, in part, drive inter-facility transmission. However, it is important to note that these
correlations are driven by facilities with high and low patient flow. This indicates that genomic
data adds information about regional genomic transmission that is not captured by the static
patient transfer network, particularly for facility pairs with intermediate patient flow between
them.
419
420 Q5: Are there any observable geographic trends in prevalence/transmission?
421
Finally, it is often useful to visualize the geographic distribution of closely related isolates. This
can provide insight into whether inter-facility transmission is concentrated in a certain
geographic region, or is more diffuse. For instance, we can see in our data that facilities that are
geographically more proximate tend to have more transmission between them (**Figure 9**), as
indicated by a positive correlation between geographic distance and genetic flow (Fsp; **Figure
9B**) and a negative correlation between geographic distance and number of closely related
isolate pairs for a given facility pair (**Figure 9D**). Interestingly, we do not see a correlation
between geographic distance minimum pairwise SNV distance between facilities (**Figure 9C**),
possibly because this metric is more susceptible to noise.
431
432 **Package details**
433
434 **Usage**
435
436 The `regentrans` package provides three main functions for investigating genetic similarity
437 between intra- and inter-facility isolates in a dataset:
438
439   1. Identifying clusters of isolates originating from the same facility (`get_clusters()`),

440　　　2.　Identifying intra- and inter-facility pairwise similarity between isolates
441　　　　(`get_pair_types()`).
442　　　3.　Calculating population-level similarity between facilities (`get_genetic_flow()`).
443

The corresponding output of these functions, sometimes in conjunction with other functions, can
then be used to answer the questions described above. The required and optional inputs to
each function can be found in **Table 1**. Many functions require a phylogenetic tree read in by
ape::read.tree() and/or a genetic distance matrix that can be calculated using ape::dist.dna(),
which requires a DNAbin object input that can be read in using ape::read.fasta(), or
ape::cophenetic.phylo(), which requires a phylogenetic tree input.
450

451　Our introductory vignette provides examples of how to read in data, use each function, and plot
452　the corresponding output for interpretation.
453

454　**Underlying methods**
455

456　Please refer to the references noted in **Table 1** for a detailed description of the methods and
457　algorithms used in the underlying functions [3, 4, 6, 7, 25].
458

459　**Handling multiple isolates from a given patient**
460

Inclusion of multiple samples from an individual is desirable to capture accumulated genetic
diversity; however, this requires careful consideration as it is important to ensure that observed
similarities are not due to similarities within a given patient, but rather between different patients.
For each of the three methods for investigating intra- and inter-facility genetic similarity, we first
define unique patient-facility combinations to ensure that we capture events where a patient has
transferred from one facility to another. However, we handle multiple isolates from the same
patient and the same facility differently for each method. We describe these differences below.
Also note that if users would like to process the data in a different or more nuanced way, they
may do so prior to inputting the data into the function.
470

471　Phylogenetic clustering
472

473　Prior to identifying phylogenetic clusters of isolates, we subset to include only one isolate per
474　patient per facility to ensure that all intra-facility clusters contain only one isolate per patient.
475

476　Intra- and inter-facility pairwise genetic distances
477

478　For each pair of unique patient-facility combinations, we include only the most closely related
479　isolate pair.
480

481　Between-facility genetic similarity
482

As Fsp only accommodates biallelic sites, we first subset to include only biallelic sites. Then, for
each patient-facility combination, we merge all sequences into a "meta-sequence" where, for
each variant position the population-level minor allele is prioritized over the population-level
major allele. In other words, all of the minor alleles for a patient-facility combination are merged
into one sequence. This meta-sequence is used to calculate Fsp values between facilities. This
method allows us to capture all minor alleles for a given patient within a given facility without
biasing the population-level prevalence of a minor allele by including multiple isolates from the
same patient.
491
Package implementation
492
493
494 regentrans is implemented in R [28] and is available on GitHub (https://github.com/Snitkin-Lab-
495 Umich/regentrans). Our package depends on several other packages including tidyverse [29]
496 packages (dplyr and tidyr), ape [11], phytools [30], igraph [31], and future.apply [32]. The
497 ggplot2 [29], ggtree [33], ggraph [34], and pheatmap [35] packages are used in the vignette for
498 plotting.
499
Additional possible uses
500
501
While our expertise in studying regional transmission largely lies in investigating bacterial
pathogen transmission within and between healthcare facilities, the methods implemented in
regentrans can be used for many additional applications. Firstly, they are generalizable to the
analysis of not only other bacterial pathogens (e.g. MRSA), but also other organisms including
fungal (e.g. *Candida auris*) and viral (e.g. SARS-COV-2) pathogens, and even non-pathogenic
microbes. Additionally, rather than investigating transmission between facilities, users could
investigate transmission between different zip codes, different rooms or wards within a hospital,
or even transmission between patient and environmental sources. As one example, Popovich
and Thiede *et al.* [36] identified transmission signatures within a large urban jail by comparing
pairwise SNV distances of community-onset MRSA to MRSA acquired within the jail.
512
Cautionary notes on interpretation
513
514
It is important to emphasize that there are several limitations to the methods we describe here.
First, none of these methods include the use of epidemiological data to confirm or corroborate
putative transmission links. Therefore, while we can gain useful insight into the likely extent of
transmission within and between facilities, we cannot understand the nuances of actual
transmission events. If epidemiological data is available, we highly recommend incorporating
this information into the analysis to provide further insights into putative transmission pathways
(examples: [6, 36]). Additionally, as mentioned above, for methods where choosing a threshold
of genomic relatedness is required, care in choosing the threshold and investigating the
sensitivity of the threshold on your interpretation of the results is warranted as the results may
change drastically depending on what threshold is chosen [37]. Finally, the sampling schemes
or time frames used in the study can influence the output of the methods presented here [38,

39]. Therefore, as always, the strengths and limitations of the dataset being used must be
considered carefully when interpreting the results.
528
529   **Conclusion**
530
Investigating regional pathogen transmission can provide insight into transmission dynamics
and guide infection prevention and control. Here we provide a framework for studying regional
pathogen transmission within and between healthcare facilities using whole-genome
sequencing data, and implement these methods in the easy-to-use R package regentrans.
regentrans allows users to interrogate the transmission dynamics of pathogens using various
metrics of genomic relatedness, including genetic-threshold and threshold-free approaches.
Using several complementary methods to investigate intra- and inter-facility transmission allows
investigators to gain a better understanding of the robustness of their findings and provide
different insights into transmission dynamics in the region of interest. Therefore, we believe that
this tool will be a useful resource for researchers and public health practitioners interested in
investigating regional pathogen transmission.
542
543   **Authors and contributors**
544
545   All authors developed methodology and reviewed and edited the manuscript. SH, ZL, and ESS
546   authors conceptualized the project. SH and ZL curated the data, designed the software, and
547   visualized the results. ZL, JW, and ESS acquired funding. ZL and ESS wrote the original
548   manuscript draft. ESS supervised the project.
549
550   **Conflicts of interest**
551
552   The authors declare that there are no conflicts of interest.
553

554   **Funding information**
555
ESS received support from the National Institutes of Health under Grant No. 1R01AI148259-01.
ZL received support from the National Science Foundation Graduate Research Fellowship
Program under Grant No. DGE 1256260. JW received support from the Canadian Institutes of
Health Research fellowship Grant No. 201711MFE-396343-165736 and the Michigan Institute
for Clinical and Health Research (MICHR) Postdoctoral Translational Scholars Program (PTSP).
Any opinions, findings, and conclusions or recommendations expressed in this material are
those of the authors and do not necessarily reflect the views of the National Science
Foundation. The funding bodies had no role in the design of the study or collection, analysis,
and interpretation of data, or in writing the manuscript.

565
566   **Ethical approval**
567
568   N/A
569

Consent for publication

N/A

Acknowledgements

We would like to thank Emily Benedict for testing out the alpha version of our package.

References

1. **Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, *et al.*** Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *The Lancet Infectious Diseases*. Epub ahead of print 21 December 2017. DOI: 10.1016/S1473-3099(17)30753-3.
2. **Blanco N, O'Hara LM, Harris AD**. Transmission pathways of multidrug-resistant organisms in the hospital setting: a scoping review. *Infection Control & Hospital Epidemiology* 2019;40:447–456.
3. **Wang J, Foxman B, Pirani A, Lapp Z, Mody L, *et al.*** Application of combined genomic and transfer analyses to identify factors mediating regional spread of antibiotic resistant bacterial lineages. *Clin Infect Dis*. DOI: 10.1093/cid/ciaa364.
4. **Han JH, Lapp Z, Bushman F, Lautenbach E, Goldstein EJC, *et al.*** Whole-Genome Sequencing To Identify Drivers of Carbapenem-Resistant Klebsiella pneumoniae Transmission within and between Regional Long-Term Acute-Care Hospitals. *Antimicrobial Agents and Chemotherapy*;63. Epub ahead of print 1 November 2019. DOI: 10.1128/AAC.01622-19.
5. **Paul P, Slayton RB, Kallen AJ, Walters MS, Jernigan JA**. Modeling Regional Transmission and Containment of a Healthcare-associated Multidrug-resistant Organism. *Clin Infect Dis*. DOI: 10.1093/cid/ciz248.
6. **Snitkin ES, Won S, Pirani A, Lapp Z, Weinstein RA, *et al.*** Integrated genomic and interfacility patient-transfer data reveal the transmission pathways of multidrug-resistant Klebsiella pneumoniae in a regional outbreak. *Science Translational Medicine* 2017;9:eaan0093.
7. **Lapp Z, Crawford R, Miles-Jay A, Pirani A, Trick WE, *et al.*** Regional Spread of blaNDM-1-Containing Klebsiella pneumoniae ST147 in Post-Acute Care Facilities. *Clinical Infectious Diseases*. Epub ahead of print 17 May 2021. DOI: 10.1093/cid/ciab457.
8. **Donker T, Reuter S, Scriberras J, Reynolds R, Brown NM, *et al.*** Population genetic structuring of methicillin-resistant Staphylococcus aureus clone EMRSA-15 within UK reflects patient referral patterns. *Microb Genom*;3. Epub ahead of print 4 July 2017. DOI: 10.1099/mgen.0.000113.
9. **Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, *et al.*** Rapid phylogenetic analysis of large samples of recombinant bacterial whole genome sequences using Gubbins. *Nucleic Acids Res* 2015;43:e15–e15.
10. **Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ**. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol* 2015;32:268–274.
11. **Paradis E, Schliep K**. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 2019;35:526–528.
12. **Lapp Z, Han JH, Choudhary D, Castaneda S, Pirani A, *et al.*** Fitness barriers to spread of colistin-resistant Klebsiella pneumoniae overcome by establishing niche in patient

population with elevated colistin use. *medRxiv* 2021;2021.06.11.21258758.
13. **Li H, Durbin R**. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics 2009;25:1754–1760.
14. **Lam MMC, Wick RR, Watts SC, Cerdeira LT, Wyres KL, *et al.*** A genomic surveillance
framework and genotyping tool for Klebsiella pneumoniae and its related species complex.
Nat Commun 2021;12:4188.
15. **Malone B**. *tangles: Anonymization of Spatial Point Patterns and Raster Objects.*
https://CRAN.R-project.org/package=tangles (2019, accessed 24 July 2021).
16. **Sherry NL, Lee RS, Gorrie CL, Kwong JC, Stuart RL, *et al.*** Pilot study of a combined
genomic and epidemiologic surveillance program for hospital-acquired multidrug-resistant
pathogens across multiple hospital networks in Australia. *Infection Control & Hospital*
Epidemiology 2021;42:573–581.
17. **Gouliouris T, Coll F, Ludden C, Blane B, Raven KE, *et al.*** Quantifying acquisition and
transmission of Enterococcus faecium using genomic surveillance. *Nature Microbiology*
2020;1–9.
18. **Coll F, Raven KE, Knight GM, Blane B, Harrison EM, *et al.*** Definition of a genetic
relatedness cutoff to exclude recent transmission of meticillin-resistant Staphylococcus
aureus: a genomic epidemiology analysis. *The Lancet Microbe* 2020;1:e328–e335.
19. **Stimson J, Gardy J, Mathema B, Crudu V, Cohen T, *et al.*** Beyond the SNP Threshold:
Identifying Outbreak Clusters Using Inferred Transmissions. *Mol Biol Evol* 2019;36:587–
603.
20. **Hawken SE, Yelin RD, Lolans K, Weinstein RA, Lin MY, *et al.*** Threshold-free genomic
cluster detection to track transmission pathways in healthcare settings. *medRxiv*
2021;2020.09.26.20200097.
21. **Duchêne S, Holt KE, Weill F-X, Le Hello S, Hawkey J, *et al.*** Genome-scale rates of
evolutionary change in bacteria. *Microb Genom*;2. Epub ahead of print 30 November 2016.
DOI: 10.1099/mgen.0.000094.
22. **Bowers JR, Kitchel B, Driebe EM, MacCannell DR, Roe C, *et al.*** Genomic Analysis of the
Emergence and Rapid Global Dissemination of the Clonal Group 258 Klebsiella
pneumoniae Pandemic. *PLoS ONE* 2015;10:e0133727.
23. **Drummond AJ, Pybus OG, Rambaut A, Forsberg R, Rodrigo AG**. Measurably evolving
populations. *Trends in Ecology & Evolution* 2003;18:481–488.
24. **Biek R, Pybus OG, Lloyd-Smith JO, Didelot X**. Measurably evolving pathogens in the
genomic era. *Trends in Ecology & Evolution* 2015;30:306–313.
25. **Duchene S, Featherstone L, Haritopoulou-Sinanidou M, Rambaut A, Lemey P, *et al.***
Temporal signal and the phylodynamic threshold of SARS-CoV-2. *Virus Evolution*;6. Epub
ahead of print 1 July 2020. DOI: 10.1093/ve/veaa061.
26. **Duchêne S, Geoghegan JL, Holmes EC, Ho SYW**. Estimating evolutionary rates using
time-structured data: a general comparison of phylogenetic methods. *Bioinformatics*
2016;32:3375–3379.
27. **Willing E-M, Dreyer C, Oosterhout C van**. Estimates of Genetic Differentiation Measured
by FST Do Not Necessarily Require Large Sample Sizes When Using Many SNP Markers.
PLOS ONE 2012;7:e42649.
28. **R Core Team**. R: A Language and Environment for Statistical Computing. https://www.R-
project.org/ (2020).
29. **Wickham H, Averick M, Bryan J, Chang W, McGowan L, *et al.*** Welcome to the
Tidyverse. *Journal of Open Source Software* 2019;4:1686.
30. **Revell LJ**. phytools: an R package for phylogenetic comparative biology (and other things).
Methods in Ecology and Evolution 2012;3:217–223.
31. **Csardi G, Nepusz T**. The Igraph Software Package for Complex Network Research.
InterJournal 2005;Complex Systems:1695.

32. **Bengtsson H, Team RC**. future.apply: Apply Function to Elements in Parallel using Futures. https://CRAN.R-project.org/package=future.apply (2020, accessed 26 August 2020).

33. **Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y**. ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and Evolution* 2017;8:28–36.

34. **Pedersen TL, RStudio**. *ggraph: An Implementation of Grammar of Graphics for Graphs and Networks*. https://CRAN.R-project.org/package=ggraph (2021, accessed 16 November 2021).

35. **Kolde R**. *pheatmap: Pretty Heatmaps*. https://CRAN.R-project.org/package=pheatmap (2019, accessed 15 April 2020).

36. **Popovich KJ, Thiede SN, Zawitz C, Aroutcheva A, Payne D, *et al.*** Genomic Epidemiology of MRSA During Incarceration at a Large Inner-City Jail. *Clin Infect Dis* 2021;ciaa1937.

37. **Hall MD, Holden MT, Srisomang P, Mahavanakul W, Wuthiekanun V, *et al.*** Improved characterisation of MRSA transmission using within-host bacterial sequence diversity. *eLife* 2019;8:e46402.

38. **Mooney SJ, Garber MD**. Sampling and Sampling Frames in Big Data Epidemiology. *Curr Epidemiol Rep* 2019;6:14–22.

39. **Murray M**. Sampling Bias in the Molecular Epidemiology of Tuberculosis. *Emerg Infect Dis* 2002;8:363–369.

Figures and tables

Table 1: Questions regentrans can help investigate, and corresponding regentrans functions (bold indicates main function for method)

	Question	Method	regentrans function	Reference
Q0	How do you choose pairwise genetic distance thresholds?	Visualize the fraction of intra-facility pairs for various pairwise genetic distances	get_pair_types(), **get_frac_intra()**	[4]
Q1	Is transmission occurring within and/or between locations?	Pairwise genetic distances within and between locations	**get_pair_types()**	[7]
Q2	What locations is transmission occurring within/between?	Phylogenetic clustering of isolates from the same location	**get_clusters()**	[6]

		Population-level similarity between locations	**get_genetic_flow()**	[8]
		Number of closely related pairs within and between locations	-	-
Q3	Have transmission dynamics changed over time?	Methods above but split over time	-	[7]
Q4	Is transmission occurring along paths of higher patient/person flow?	Compare patient/person flow between locations to inter-location pairwise genetic distances or genetic flow (Fsp)	**get_patient_flow()**, get_pair_types(), get_genetic_flow(), **summarize_pairs()**	[3]
Q5	Are there any observable geographic trends in prevalence/transmission?	Visualize geographic distribution of prevalence and closely related pairs or genetic flow (Fsp)	get_pair_types(), get_genetic_flow()	[4]

697
698
699

700
701 **Figure 1: regentrans input data.** Required and optional isolate genetic data and metadata for
702 using regentrans. Figure created on BioRender.com.
703

704
Figure 2: Data-specific method for choosing pairwise genetic distance thresholds. (A)
Fraction of intra-facility pairs for various pairwise single nucleotide variant (SNV) distances. **(B)**
Fraction of intra-facility pairs for various pairwise patristic distances grouped into 100 bins.
These plots can help identify drops in intra-facility pair fractions that may indicate a reasonable
pairwise distance threshold, assuming intra-facility transmission is more common than inter-
facility transmission. Note that it may be difficult to clearly identify a large drop at any given
threshold; therefore, we recommend performing sensitivity analyses with several thresholds.
Furthermore, users should consider the tradeoff between sensitivity and specificity when
deciding what thresholds to choose. This data can be generated using the get_frac_intra()
function.
715
716
717

718
Figure 3: Pairwise genetic distances between facilities suggest recent intra- and inter-
facility transmission. (A) Pairwise single nucleotide variant distances. Grey line is at a
pairwise SNV distance of 10, and the inset shows all pairs with a pairwise SNV distance less
than or equal to this threshold, which we consider indicative of recent transmission. See Q0 on
genetic distance thresholds in the main text for more details on how we chose these thresholds..
(B) Pairwise patristic distances. Grey line is at a pairwise patristic distance of 2.56e-6 mutations
per site per year, and the inset shows all pairs with a pairwise patristic distance less than or

equal to this threshold. These plots indicate that transmission is likely occurring both within and
between facilities, due to small pairwise genetic distances for both intra- and inter-facility pairs.
Data generated using the get_pair_types() function.
729

730
Figure 4: Clusters of isolates from the same facility suggest intra-facility transmission.
(A) Mapping isolate location on the phylogeny provides a visual for the extent of clustering by
facility. Here we can see clustering of isolates from the same facility in several subclades of the
phylogeny. **(B)** Quantification of the size of phylogenetic clusters from a single facility using
get_clusters(). The size of the points in panel B can provide insight into whether large clusters
are from an intra-facility outbreak (smaller point size) or sustained intra-facility transmission
(larger point size). Points in panel B are jittered for visualization purposes.

738
Figure 5: Some facility pairs have similar populations, indicating potential transmission between them. Genetic flow (Fsp) was calculated using the get_genetic_flow() function in regentrans. Rows and columns are facilities. Lower Fsp indicates more similar populations and thus more putative transmission. At least two isolates are required from a facility to perform the Fsp calculation as within-facility variation must be calculated, so facilities with only one isolate have been removed. The heatmap is clustered based on Fsp values.

745
746 **Figure 6: Some facilities have many closely related isolates, indicating potential intra-**
747 **and inter-facility transmission. (A)** Number of closely related isolate pairs for different genetic
748 distance metrics and thresholds. Only facilities with at least one pair of closely related isolates
749 are shown. **(B)** Network of patient connectedness. Patient nodes are connected by edges if they
750 share an isolate with a pairwise SNV distance ≤ 6.
751

752
Figure 7: The pairwise SNV distance distribution does not change over time. (A) Count of
pairwise single nucleotide variant (SNV) distances faceted by year. (B) Fraction of intra- vs.
inter-facility pairwise SNV distances faceted by year. Trends are similar across both years.

Figure 8: Facilities with more patient flow tend to have more similar *K. pneumoniae*
populations. (A) Patient flow and genetic flow (Fsp) are negatively correlated. (**(B)** Patient flow
and the minimum pairwise single nucleotide variant (SNV) distance (i.e. the SNV distance
between the most closely related isolates) are negatively correlated. **(C)** Patient flow and
number of closely related isolate pairs are positively correlated. Patient flow is the path of
maximum patient flow. For this analysis we considered indirect transfers as long-term acute
care hospitals are often not connected by direct transfers, but rather are connected by transfers
to an intermediate facility such as an acute care hospital. Mean patient flow was calculated as
the mean of the two directional patient flow metrics between two facilities. Lines were plotted
using ggplot::geom_smooth() with the 'lm' method.
767
768

770
Figure 9: Geographically close facilities are often connected by closely related isolate pairs. (A) Facilities are located as they are geographically in space but latitude and longitude are de-identified by horizontal, vertical, and rotational shifts. The smaller single nucleotide variant (SNV) threshold was chosen for visualization purposes. Size and color of points indicates sample size, size and color of edges indicates number of closely related pairs. **(B)** Physical distance between facilities is positively correlated with genetic flow (Fsp). **(C)** Physical distance between facilities is not correlated with minimum pairwise SNV distance. **(D)** Physical distance between facilities is negatively correlated with number of closely related isolate pairs (≤10 SNVs). The larger SNV threshold was chosen to have a wider distribution of number of closely related isolate pairs. Physical distance was calculated as the shortest distance between

781   the points of latitude and longitude for the facility pair. Lines in panels B, C, and D were plotted

782   using ggplot::geom_smooth() with the 'lm' method.