

The association between statin and COVID-19 adverse outcomes: National COVID-19 cohort in South Korea

Ronald Chow, MS, FRSPH,^{1,†} Jihui Lee, PhD,^{2,†} Hyerim Noh, BSc,^{3,†} Jongseong Lee, MPP,^{4,5} Hyun Joon Shin, MD, MS, MPH, ScD,^{1,6,7} Young-Geun Choi, PhD³

[†]These authors contributed equally to this work as co-first authors

¹Hanyang Impact Science Research Center, Seoul, South Korea;

²Department of Healthcare Policy & Research, Weill Cornell Medicine, New York, NY, USA;

³Department of Statistics, Sookmyung Women's University, Seoul, South Korea;

⁴School of Social Work, Columbia University, New York, NY, USA;

⁵Ministry of Health and Welfare, South Korea;

⁶Division of Cardiology, Department of Medicine, Lemuel Shattuck Hospital, Massachusetts Department of Public Health, Jamaica Plains, MA, USA;

⁷Division of General Internal Medicine, Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA.

Address correspondence to:

Young-Geun Choi, Assistant Professor

Department of Statistics, Sookmyung Women's University, 99 Cheongpa-Ro-47-Gil, Yongsan-Gu, Seoul 04310, South Korea

Tel: +82-2-710-9178; E-mail: ygchoi@sm.ac.kr

Hyun Joon Shin, Chief of Cardiology

Division of Cardiology, Department of Medicine, Lemuel Shattuck Hospital, Massachusetts Department of Public Health, 170 Morton St, Jamaica Plain, MA 02130, USA

Tel: +1-617-522-8110; E-mail: hyun.shin@mass.gov

Word count (main text): 2,630

Word count (summary): 304

Research in context

Evidence before this study: There is limited and conflicting data reporting on statin use among COVID-19 patients, and its association with COVID-19 outcomes

Added value of this study: We report no difference in COVID-19 outcomes between patients who used and did not use statins prior to COVID-19 diagnosis, except in hypertensive patients in which statins was shown to have a protective effect.

Implications of all the available evidence: As statins are not detrimental and relatively cheap, we encourage further investigation into statin for the prevention and treatment of COVID-19.

Summary

Background: There currently exists limited and conflicting clinical data on the use of statins amongst COVID-19 patients. Given the both paucity and lack of consensus among data on statin's efficacy and safety amongst COVID-19 patients, the current guideline is to continue statin in COVID-19 patients, who have previously been treated with statins. The aim of this paper was to compare hospitalized patients with COVID-19 who did and did not receive statins, in terms of COVID-19 outcomes.

Methods: We conducted population-based retrospective study using South Korea's nationwide healthcare database as of May 15 2020. We identified 4,349 patients hospitalized with COVID-19 and aged 40 years or older. The cohort entry was defined as the date of hospitalization. Statin users were individuals with inpatient and outpatient prescription records of statins in the 240 days before cohort entry, and non-users were those without such records during this period. Our primary outcome was a composite endpoint of all-cause death, intensive care unit (ICU) admission, mechanical ventilation use and cardiovascular outcomes (myocardial infarction (MI), transient cerebral ischemic attacks (TIA) or stroke). We conducted inverse probability of treatment weighting (IPTW)-adjusted logistic regression analysis to estimate odds ratio (OR) and corresponding 95% confidence intervals (CI), to compare outcomes between statin users and non-users.

Findings: 1,115 patients were statin users (mean age = 65.9 years; 60% female), and 3,234 were non-users (mean age = 58.3 years; 64% female). Statin use was not associated with increased risk of the primary outcome (IPTW OR 0.82; 95% CI: 0.60-1.11). Subgroup analysis showed a protective role of statins, for individuals with hypertension (IPTW OR 0.40; 95% CI: 0.23-0.69, p for interaction: 0.0087).

Interpretation: Given that statins are not detrimental and that it may be beneficial amongst hypertensive patients and relatively cheap, we would encourage further investigation into statin for the prevention and treatment of COVID-19.

Funding: YGC's work was partially supported by 2020R1G1A1A01006229 awarded by the National Research Foundation of Korea.

Introduction

On March 11, 2020, the World Health Organization (WHO) declared the COVID-19 outbreak a global pandemic^{1,2}. Since then, the daily number of global COVID-19 cases has increased from a few thousand in the beginning of March, to 50,000 at the end of March, to 100,000 in May, and to 200,000 in July³.

The pathophysiology of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which causes the COVID-19 disease, involves an overproduction of an early response proinflammatory cytokines, specifically tumour necrosis factor (TNF), IL-6 and IL-1 β)⁴. If unabated, this cytokine storm subsequently places COVID-19 patients at increased risk of vascular hyperpermeability, multiorgan failure, and death⁵.

As a result, statins have been suggested for use as a therapy for COVID-19. It is reported that statin can stabilize MYD88 at normal levels and reduce an ensuing cytokine storm⁶. Statins may also up-regulate ACE2, which is typically downregulated by SARS-CoV-2 and facilitates the infiltration of SARS-CoV-2⁷. Statins can cause side effects of myalgia, increased creatine phosphokinase, and rhabdomyolysis and corresponding acute kidney injury, albeit quite rare; all these are reversible upon discontinuation of statins⁸.

There currently exists limited and conflicting clinical data on the use of statins amongst COVID-19 patients. While Zhang *et al*⁹ and Rodriguez-Nava *et al*¹⁰ report that statin reduced the risk of mortality and/or disease severity, and Wang *et al*¹¹ reported an increased mortality among statin users, other studies¹²⁻¹⁸ report no significant difference between statin users and non-users. Meanwhile, a recent systematic review and meta-analysis of 110,078 patients reported a reduced risk of mortality among those administered statins after their COVID-19

hospitalization, but no difference in patients who were administered statins before hospitalization¹⁹. Although there is a lack of consensus among data on statin's efficacy and safety amongst COVID-19 patients, the current recommendation is for COVID-19 patients to continue any antecedent statin use²⁰.

Given the limited and conflicting data, there is a need for further investigation into the relationship between statin use and COVID-19 outcomes. The aim of this paper was to compare hospitalized patients with COVID-19 who did and did not receive statins, in terms of COVID-19 outcomes.

Methods

Data Source

We conducted a population-based cohort study using the Health Insurance Review and Assessment Service (HIRA) claims data released by the South Korean government. The HIRA is a sole nationwide governmental agency providing health insurance claims review, covering 98% of the Korean population (about 50 million), and employs a fee-for-service reimbursement system²¹. In March 2020, the HIRA released its claim database for all COVID-19 patients in South Korea. This population-level claims dataset includes information on sociodemographic characteristics, healthcare utilization history, diagnosis results (International Classification of Diseases, 10th Revision²²; ICD-10) and prescription from both inpatient and outpatient settings. COVID-19 diagnosis was determined through the South Korean Disease Control and Prevention Agency (KDCA) database. This database is the first nationwide dataset of COVID-19 patients. Details of the HIRA database are provided in Supplementary Material 1.

This study was approved by the Human Investigation Review Board of Public Institutional

Bioethics Committee designated by the South Korean Ministry of Health and Welfare, which waived the requirement of informed consent due to retrospective study design and anonymity of the HIRA database (IRB # P01-2020-1262-001).

Study Design and Participants

Between January 1, 2020 and May 15, 2020, 7,590 individuals tested positive for COVID-19. A positive COVID-19 result was defined as a positive result from South Korean Ministry of Food and Drug Safety–approved diagnostic tests that used a reverse-transcription polymerase chain reaction method²³. Individuals were excluded if they were younger than 40 years old, since younger patients are less likely to experience COVID-19 outcomes regardless of statin use. As well, we excluded individuals who were not hospitalized (Figure 1), for precise outcome assessment. As a result, 4,349 individuals were included in the sample cohort. Cohort entry was defined as the date of admission for COVID-19 hospitalization. Most of these patients were hospitalized until they fully recovered. A full recovery was defined as cessation of fever without medication use, and 2 consecutive negative test results within a 24 hour period²⁴.

Statin Exposure

We defined exposure by using inpatient and outpatient prescription records of statin from the HIRA database (Supplementary Material 2). Our definition follows an intention-to-treat approach. Patients prescribed statin within 240 days prior to cohort entry were regarded as statin users, and otherwise defined as non-users.

Outcomes

We set the primary outcome as a composite endpoint of all-cause death, intensive care unit

(ICU) admission, mechanical ventilation use and cardiovascular outcomes (myocardial infarction (MI), transient cerebral ischemic attacks (TIA) or stroke). Secondary outcomes were the individual components of the composite endpoint. These outcomes were defined using in-hospital ICD-10 diagnostic codes and the national procedure coding system (Supplementary Material 2). We measured study outcomes between the date of cohort entry and the earliest of the date of hospital discharge or end of study period (May 15, 2020), whichever occurred earlier.

Potential Confounders

We included sociodemographic and clinical factors that are considered to be associated with both statin use and risk of the outcomes. For sociodemographic factors, we included age in years, square of age, sex, and health insurance type at cohort entry. We included clinical variables of 20 comorbidities and use of 12 co-medications. The ascertainment period to define comorbidities was from 3 years prior (-1080d) through the start of the exposure ascertainment (-240d). The ascertainment period to define co-medications was from 2 years prior (-720d) through -240d. We defined comorbidity variables from in-hospital ICD-10 diagnostic codes and co-medications from in/out-hospital anatomical therapeutic chemical codes (Supplementary Material 2). When defining malignancy, we additionally used the expanded benefit codes in addition to diagnosis codes to reduce false-positives.

Statistical Analyses

We summarized the baseline characteristics for statin users and non-users using mean and standard deviation for continuous variables, and frequency and percentage for categorical variables. We investigated imbalance of distribution of covariates between statin users and non-users using absolute standardized difference (aSD). An $aSD \leq 0.1$ was preferred for balance.

We employed the inverse probability of treatment weighting (IPTW)-adjustment approach in order to reduce the confounding bias of statin users. Propensity score (PS) was estimated by a logistic regression with baseline sociodemographic and clinical characteristics as predictors (Table 1). We assigned the IPTW weight to each individual in the cohort, $1/PS$ to the statin users, and $1/(1-PS)$ to the non-users and constructed a balanced pseudo-population. Weighted univariable logistic regression models were fitted to estimate the effect of statin use on primary and secondary outcomes. For comparison, we also considered unweighted logistic regression: both univariable and multivariable, adjusted for age, sex, insurance type, history of diabetes and history of hypertension. For each logistic regression model, we reported the estimated odds ratio (OR) and its 95% confidence interval (CI).

Subgroup Analyses

We stratified the cohort by (1) age (<65 and ≥ 65 years) and (2) sex for the risk of the primary outcome. We also considered stratification by the history of (3) hyperlipidemia, (4) hypertension, and (5) diabetes mellitus. For each subgroup analysis, we recalculated PS by multivariable logistic regression. We obtained p for interaction by each subgroup.

Sensitivity Analyses

We conducted sensitivity analyses by extending the scope of the study population in two settings. The first setting was to redefine the study population by relaxing the hospitalization condition. That is, we included all confirmed positive individuals of at least 40 years of age regardless of their hospitalization record. The second setting was to extend the time window for exposure. To capture the patients who infrequently received statin prescription, we extended

the time window for ascertaining exposure from 240 to 360 days, prior to cohort entry.

To investigate the sensitivity of results against the choices of statistical methods, we considered the following alternative approaches. First, we repeated the main IPTW analysis with discarding individuals with extreme PS values, $PS < 0.025$ or $PS > 0.975$ (IPTW with trimming). Second, in our unweighted multivariable logistic regression models, we included the estimated PSs as an additional predictor to other covariates (outcome adjustment model). Third, we applied the stabilized mortality ratio weighting, 1 for the user group and $PS/(1-PS)$ for the non-user group, instead of IPTW (SMR weighting). Lastly, we considered propensity score matching (PS matching).

A two-sided p-value less than 0.05 was considered statistically significant. All analyses were conducted using R 3.5.2 (R Core Team, Vienna, Austria).

Results

A total of 4,349 adults of at least 40 years of age were hospitalized with COVID-19 in South Korea between January 1, 2020 and May 15, 2020 and included this analysis. Of these, 1,115 (27%) were statin users. Statin users were older than non-users on average (65.9 ± 11.2 vs 58.3 ± 12.3). Among statin users, 39.6% of statin users were males, whereas 36.1% of non-users were male. A larger proportion of statin users had comorbidities of chronic lung disease (50.6% vs 24.6%), diabetes mellitus (40.3% vs 10.7%), hyperlipidemia (65.4% vs 18.1%), and hypertension (53.5% vs 17.7%), relative to non-users. Statin users also used more antidiabetic (35.6% vs 6.4%) and antiplatelet (37.7% vs 7.9%) medications (Table 1).

Among 530 primary composite events of all-cause death, ICU admission, mechanical

ventilation use and cardiovascular adverse outcomes, 362 occurred in non-users (11.2%; 362/3234) and 168 (15.1%; 168/1115) in statin users. No difference was noted between statin users and non-users in the IPTW weighted analysis (IPTW adjusted OR 0.82; 95% CI: 0.60-1.11). Similarly, there was no difference between statin users and non-users for the individual components of all-cause death, mechanical ventilation, ICU admission, and cardiovascular adverse outcome (Table 2).

In the subgroup analyses, there existed a significant effect modification by hypertension status ($p = 0.0087$). Statin users who had hypertension experienced lower odds for the composite (IPTW OR 0.40, 95% CI: 0.23 – 0.69). There was no effect modification by age, sex, hyperlipidemia and diabetes mellitus (Figure 2).

The results from the sensitivity analyses remained generally consistent with the main analysis with no harmful association between statin use and the outcomes. When redefining the study population, changing the exposure ascertainment window and applying other statistical methods, the primary outcome, all-cause death, ICU admission and cardiovascular outcomes showed generally similar results. Statins showed a protective association with the primary outcome, when using SMR weighting (OR 0.64, 95% CI: 0.45 – 0.93) and PS matching (OR 0.74, 95% CI: 0.56 – 0.98). For all-cause deaths, statins showed protective association when using SMR weighting (OR 0.56, 95% CI: 0.35 – 0.89) and PS matching (OR 0.67, 95% CI: 0.46 – 0.98). As well, statins showed a protective association for all-cause death, when including all confirmed patients with COVID-19 (OR 0.81, 95% CI: 0.68 – 0.96). For ICU admission, statin showed protective association when using IPTW weighting with trimming (IPTW OR 0.67, 95% CI: 0.47 – 0.97), or redefining exposure window to 360 days (IPTW OR 0.67, 95% CI: 0.47 – 0.97) (Supplementary Material 3-7).

Discussion

This study reported on one of the largest investigations of COVID-19 patients with hospitalization in South Korea. It utilized a nationwide and completely enumerated dataset, and applied a propensity score-based weighting to control for potential confounding variables. In the main analysis, there was no significant difference between baseline (pre-hospitalization) statin use vs non-use, with respect to the outcomes of mortality, ICU admission, mechanical ventilation use and cardiovascular outcomes.

Even though our main analysis results showed null association between statin and all cause death, our sensitivity analysis of all cause death showed protective association with statin when using SMR weighting, PS matching or when including all confirmed patients with COVID-19. This finding is similar to report by Lee *et al*²⁵, which showed a protective association of statins for all cause death in the similar cohort while using PS matching with a several different methodological choices from our study (first, they included patients of all age, rather than restricting their analyses to patients who were 40 years old and older; second, statin exposure was defined by Lee *et al* as a prescription up to 90 days prior to COVID-19 diagnosis rather than 240 days prior to COVID-19 diagnosis; third, they used Cox survival analysis while we used logistic regression). It therefore is possible that we missed detecting real protective association of statin in the main analysis, especially since some of sensitivity analysis showed protective association of statin for primary outcome, all cause death and ICU admission based on varying statistical methods, changing the exposure window change to 360 days and inclusion of non-hospitalized COVID-19 patients.

Also, in the prespecified subgroup analysis, statin was found to be associated with lower odds of the primary endpoint in hypertensive patients. As hypertensive COVID-19 patients are known to have poorer prognosis than normotensive patients²⁶, this result is encouraging; statins may have a greater protective effect, for this sicker cohort of patients with hypertension. Further investigation is needed since statins may be cost-effective, with the potential benefit outweighing any potential toxicities²⁷. Mechanistically, statins is postulated to upregulate ACE2, countering the effect of SARS-CoV-2 where ACE2 is downregulated and thereby facilitates the infiltration of SARS-CoV-2⁷. Although statin can cause muscle and liver side effects, discontinuation of statins treatment is known reverse these side effects, alleviating concerns of significant and irreversible adverse effects.

This study has several strengths. First, we used a nationwide cohort of all hospitalized patients with COVID-19 in South Korea between January 1, 2020 and May 15, 2020. Since South Korea maintained a strict patient management system during this period, the use of this population-based cohort may mitigate any potential sampling bias issue. In addition, relative to the study by Lee *et al*²⁵, our study design aimed to further mitigate healthy user bias, by including chronic co-medications as confounders, which may have adjusted for past adherence²⁸. Furthermore, our main findings of no harmful association between statin and COVID-19 outcomes were supported by our sensitivity analyses which showed even protective association when we redefined study population, extended the ascertainment period of the statin use, and considered various statistical approaches. Given these neutral to protective association between statin and COVID-19 outcomes, our findings may support the current guideline which is to continue statin in COVID-19 patients, who have previously been treated with statins²⁰.

There are also several limitations of this study. First, there may have been outcome misclassification in cardiovascular outcome (MI, stroke, and TIA), because we defined our cardiovascular outcome variables by diagnostic codes in administrative claim data. HIRA reported that 82% of primary diagnosis codes in claims coincide with electronic medical records,²⁹ so more than 18% of outcome misclassification can occur while trying to capture all the diagnostic codes other than primary diagnosis codes. However, we expect a greater validity for MI, stroke, and TIA since our study population focused on hospitalized patients under careful management. Other outcomes such as all-cause death seem to be well classified because the HIRA COVID-19 database was linked to the national death records and outcomes defined from procedure codes (ICU admission and mechanical ventilation use) are also expected to be valid since the codes are mandatory in the reimbursement review process. Second, statin exposure was defined based on inpatient and outpatient prescription. We are not aware of detailed information on the exposure, for example, adherence to the prescription. Last, although we performed IPTW and thorough sensitivity analyses, due to the inherent limitation of claims data, there might be still a residual confounding by potential unmeasured confounders typically used in clinical studies (e.g., body mass index, baseline blood pressures, laboratory test values).

In conclusion, our analysis of a nationwide sample of South Korean, hospitalized, COVID-19 patients found that statin users overall did not experience poorer COVID-19 outcomes, relative to non-users. However, in hypertensive patients, a protective association between statin use and COVID-19 outcomes were shown, which calls for further investigation. Given that statins are relatively cheap and not detrimental in this setting, we would encourage further investigation into statin for the prevention and treatment of COVID-19.

Author Contributions: YGC and HJS conceived this project. YGC and HN had full access to data for analysis. RC, Jihui L and Jongseong L drafted the manuscript. All authors contributed significantly, edited the manuscript and are accountable for its reporting.

Acknowledgements: The authors thank healthcare professionals dedicated to treating COVID-19 patients in South Korea, the Ministry of Health and Welfare, the Health Insurance Review and Assessment Service (HIRA), and Ye-Jin Sohn (HIRA) for sharing invaluable national health insurance claims data.

Declaration of interest: All authors completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. The authors declare no competing interests.

Data sharing: No additional data available.

References

1. Chow R, Elsayeed S, Lock M. How robust are the results of one of the first positive trials exploring hydroxychloroquine for treatment of COVID-19? medRxiv 2020. DOI: 10.1101/2020.05.06.20093237.
2. Chow R, Simone CB, 2nd, Lock M. Hydroxychloroquine for the treatment of COVID-19: the importance of scrutiny of positive trials. *Ann Palliat Med* 2020. DOI: 10.21037/apm-20-1538.
3. World Health Organization. Timeline: WHO's COVID-19 response. World Health Organization. (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/interactive-timeline#!>).
4. Jose RJ, Manuel A. COVID-19 cytokine storm: the interplay between inflammation and coagulation. *Lancet Respir Med* 2020;8(6):e46-e47. DOI: 10.1016/s2213-2600(20)30216-2.
5. Meduri GU, Kohler G, Headley S, Tolley E, Stentz F, Postlethwaite A. Inflammatory cytokines in the BAL of patients with ARDS. Persistent elevation over time predicts poor outcome. *Chest* 1995;108(5):1303-14. DOI: 10.1378/chest.108.5.1303.
6. Dashti-Khavidaki S, Khalili H. Considerations for Statin Therapy in Patients with COVID-19. *Pharmacotherapy* 2020;40(5):484-486. DOI: 10.1002/phar.2397.
7. Fedson DS, Opal SM, Rordam OM. Hiding in Plain Sight: an Approach to Treating Patients with Severe COVID-19 Infection. *mBio* 2020;11(2). DOI: 10.1128/mBio.00398-20.
8. Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020;382(18):1708-1720. DOI: 10.1056/NEJMoa2002032.
9. Zhang XJ, Qin JJ, Cheng X, et al. In-Hospital Use of Statins Is Associated with a Reduced Risk of Mortality among Individuals with COVID-19. *Cell Metab*

- 2020;32(2):176-187.e4. DOI: 10.1016/j.cmet.2020.06.015.
10. Rodriguez-Nava G, Trelles-Garcia DP, Yanez-Bello MA, Chung CW, Trelles-Garcia VP, Friedman HJ. Atorvastatin associated with decreased hazard for death in COVID-19 patients admitted to an ICU: a retrospective cohort study. *Crit Care* 2020;24(1):429. DOI: 10.1186/s13054-020-03154-4.
 11. Wang B, Van Oekelen O, Mouhieddine TH, et al. A tertiary center experience of multiple myeloma patients with COVID-19: lessons learned and the path forward. *Journal of Hematology & Oncology* 2020;94:13.
 12. Argenziano MG, Bruce SL, Slater CL, et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020;369:m1996. DOI: 10.1136/bmj.m1996.
 13. Ayed M, Borahmah A, Yazdani A, Sultan A, Mossad A, Rawdhan H. Assessment of clinical characteristics and mortality-associated factors in COVID-19 Critical cases in Kuwait. *medRxiv* 2020. DOI: 10.1101/2020.06.17.20134007.
 14. De Spiegeleer A, Bronselaer A, Teo JT, et al. The Effects of ARBs, ACEis, and Statins on Clinical Outcomes of COVID-19 Infection Among Nursing Home Residents. *J Am Med Dir Assoc* 2020;21(7):909-14.
 15. Dreher M, Kersten A, Bickenbach J, et al. The Characteristics of 50 Hospitalized COVID-19 Patients With and Without ARDS. *Deutsches Arzteblatt International* 2020;117(16):271-8.
 16. Grasselli G, Greco M, Zanella A, et al. Risk Factors Associated With Mortality Among Patients With COVID-19 in Intensive Care Units in Lombardy, Italy. *JAMA Intern Med* 2020;180(10):1345-1355. DOI: 10.1001/jamainternmed.2020.3539.
 17. Tan WYT, Young BE, Lye DC, Chew DEK, Dalan R. Statin use is associated with lower disease severity in COVID-19 infection. *Sci Rep* 2020;10(1):17458. DOI:

- 10.1038/s41598-020-74492-0.
18. Yan H, Valdes AM, Vijay A, et al. Role of Drugs Used for Chronic Disease Management on Susceptibility and Severity of COVID-19: A Large Case-Control Study. *Clin Pharmacol Ther* 2020;108(6):1185-1194. DOI: 10.1002/cpt.2047.
 19. Chow R, Im J, Chiu N, et al. The protective association between statins use and adverse outcomes among COVID-19 patients: a systematic review and meta-analysis. *PLOS One* 2021;16:e0253576. DOI: <https://doi.org/10.1101/2021.02.08.21251070>.
 20. Kim AY, Gandhi RT. COVID-19: Management in hospitalized adults. UpToDate. (<https://www.uptodate.com/contents/covid-19-management-in-hospitalized-adults/print?search=statin> covid&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1 1/37).
 21. Kwon S. Thirty years of national health insurance in South Korea: lessons for achieving universal health care coverage. *Health Policy and Planning* 2009;24(1):63-71. DOI: 10.1093/heapol/czn037.
 22. World Health Organization. International Statistical Classification of Diseases and Related Health Problems 10th Revision. (<https://icd.who.int/browse10/2019/en>).
 23. World Health Organization. Laboratory testing for coronavirus disease 2019 (COVID-19) in suspected human cases: interim guidance, 2 March 2020. Geneva: World Health Organization, 2020 2020. (WHO/COVID-19/laboratory/2020.4) (<https://apps.who.int/iris/handle/10665/331329>).
 24. Ministry of Health and Welfare, Republic of Korea. About COVID-19: patient treatment & management.
 25. Lee HY, Ahn J, Park J, et al. Beneficial Effect of Statins in COVID-19-Related Outcomes-Brief Report: A National Population-Based Cohort Study. *Arterioscler*

- Thromb Vasc Biol 2021;41(3):e175-e182. DOI: 10.1161/atvbaha.120.315551.
26. Zheng Z, Peng F, Xu B, et al. Risk factors of critical & mortal COVID-19 cases: A systematic literature review and meta-analysis. *J Infect* 2020;81(2):e16-e25. DOI: 10.1016/j.jinf.2020.04.021.
 27. Chow R, Prsic EH, Shin HJ. Cost-Effectiveness Analysis of Statins for Treatment of Hospitalized COVID-19 Patients. *medRxiv* 2021:2021.04.29.21256335. DOI: 10.1101/2021.04.29.21256335.
 28. Shrank WH, Patrick AR, Brookhart MA. Healthy user and related biases in observational studies of preventive interventions: a primer for physicians. *J Gen Intern Med* 2011;26(5):546-50. DOI: 10.1007/s11606-010-1609-1.
 29. Park EC, Jang SI, Jeon SY, Lee SA, Lee JE, Choi DW. Assessment of level of agreement in disease codes between Health Insurance Claims Data and Medical Records. *Health Insurance Review & Assessment Service* 2018 (<http://www.alio.go.kr/informationResearchView.do?seq=2343982>).

Table 1. Baseline sociodemographic and clinical characteristics of COVID-19 patients with hospitalization ≥ 40 years of age between January 1, 2020 and May 15, 2020 in South Korea.

	Before IPTW			After IPTW ^s		
	Statin User* 1115	Non-User 3234	aSD	Statin User 4099	Non-User 4505	aSD
Age (years; mean+sd)	65.9±11.2	58.3±12.3	0.64	61.1±13.2	60.9±12.9	0.01
40-49	79 (7.1)	829 (25.6)		892 (21.8)	893 (19.8)	
50-59	252 (22.6)	1155 (35.7)		1080 (26.3)	1432 (31.8)	
60-69	379 (34.0)	684 (21.2)		1097 (26.8)	1066 (23.7)	
70-79	259 (23.2)	312 (9.6)		549 (13.4)	644 (14.3)	
80-89	131 (11.7)	202 (6.2)		415 (10.1)	380 (8.4)	
90+	15 (1.3)	52 (1.6)		67 (1.6)	90 (2.0)	
Sex						
Male	442 (39.6)	1167 (36.1)	0.07	1703 (41.5)	1673 (37.1)	0.09
Female	673 (60.4)	2067 (63.9)	0.07	2397 (58.5)	2832 (62.9)	0.09
Health insurance type						
Medical insurance	960 (86.1)	2871 (88.8)	0.08	3490 (85.1)	3952 (87.7)	0.08
Medical aid	155 (13.9)	363 (11.2)	0.08	610 (14.9)	553 (12.3)	0.08
Comorbidities						
Arrhythmia	46 (4.1)	62 (1.9)	0.13	112 (2.7)	116 (2.6)	0.01
Asthma	144 (12.9)	279 (8.6)	0.14	400 (9.8)	458 (10.2)	0.01
Atrial fibrillation	36 (3.2)	35 (1.1)	0.15	63 (1.5)	72 (1.6)	0.00
Autoimmune disease	117 (10.5)	272 (8.4)	0.07	303 (7.4)	396 (8.8)	0.05
Chronic lung disease	564 (50.6)	794 (24.6)	0.56	1425 (34.8)	1487 (33.0)	0.04
Coronary artery disease	187 (16.8)	118 (3.6)	0.44	320 (7.8)	349 (7.7)	0.00
Dementia	98 (8.8)	189 (5.8)	0.11	371 (9.1)	330 (7.3)	0.06
Diabetes mellitus	449 (40.3)	345 (10.7)	0.72	910 (22.2)	933 (20.7)	0.04
Heart failure	87 (7.8)	90 (2.8)	0.23	200 (4.9)	213 (4.7)	0.01
Hyperlipidemia	729 (65.4)	585 (18.1)	1.09	1335 (32.6)	1480 (32.9)	0.01
Hypertension	596 (53.5)	571 (17.7)	0.81	1197 (29.2)	1332 (29.6)	0.01
Kidney disease	28 (2.5)	28 (0.9)	0.13	65 (1.6)	71 (1.6)	0.00
Liver disease	44 (3.9)	146 (4.5)	0.03	295 (7.2)	214 (4.8)	0.10
Malignancy	61 (5.5)	146 (4.5)	0.04	223 (5.4)	248 (5.5)	0.00
Other cerebrovascular diseases	112 (10.0)	125 (3.9)	0.24	252 (6.1)	279 (6.2)	0.00
Peripheral vascular disease	144 (12.9)	207 (6.4)	0.22	398 (9.7)	368 (8.2)	0.05
Pneumonia including tuberculosis	95 (8.5)	204 (6.3)	0.08	411 (10.0)	339 (7.5)	0.09
Psychiatric disorders	357 (32.0)	690 (21.3)	0.24	1197 (29.2)	1132 (25.1)	0.09
Stroke or TIA	156 (14.0)	129 (4.0)	0.36	324 (7.9)	350 (7.8)	0.00
Thromboembolism	120 (10.8)	93 (2.9)	0.32	246 (6.0)	248 (5.5)	0.02
Co-medications						
Acetaminophen	610 (54.7)	1475 (45.6)	0.18	1970 (48.1)	2210 (49.1)	0.02
Antibiotics systemic	840 (75.3)	2167 (67.0)	0.18	2778 (67.8)	3114 (69.1)	0.03
Anticoagulants	93 (8.3)	76 (2.4)	0.27	173 (4.2)	207 (4.6)	0.02
Antidementia	146 (13.1)	263 (8.1)	0.16	490 (12.0)	457 (10.2)	0.06
Antidepressants	199 (17.8)	361 (11.2)	0.19	667 (16.3)	620 (13.8)	0.07
Antidiabetics	397 (35.6)	207 (6.4)	0.77	647 (15.8)	727 (16.1)	0.01
Antiplatelets	420 (37.7)	254 (7.9)	0.76	722 (17.6)	842 (18.7)	0.03
Antipsychotics	269 (24.1)	682 (21.1)	0.07	1100 (26.8)	1037 (23.0)	0.09
Antivirals	79 (7.1)	223 (6.9)	0.01	284 (6.9)	298 (6.6)	0.01
Anxiolytics	390 (35.0)	777 (24.0)	0.24	1329 (32.4)	1213 (26.9)	0.12
Immunosuppressant	630 (56.5)	1524 (47.1)	0.19	2061 (50.3)	2255 (50.1)	0.00
NSAIDs	917 (82.2)	2437 (75.4)	0.17	3091 (75.4)	3468 (77.0)	0.04

aSD = absolute standardized difference; IPTW = inverse probability of treatment weighting; TIA = transient cerebral ischemic attack; NSAIDs = nonsteroidal anti-inflammatory drugs.

Mean and standard deviation were reported for continuous variables. Frequency and percentage were reported for categorical variables.

*Patients prescribed within 240 days prior to cohort entry were regarded as statin users, and otherwise defined as non-users.

§Weighted cohort using the inverse probability of treatment weighting (IPTW).

Table 2. Risk of adverse clinical outcomes associated with statin use among COVID-19 \geq 40 years of age with hospitalization

	Number of Patients	Number of Events	Cumulative Incidence (%)	Odds Ratio (95% Confidence Interval)		
				Unadjusted*	Adjusted [§]	IPTW Weighted [#]
All-cause death, ICU admission, mechanical ventilation, MI, stroke, TIA						
Non-user	3234	362	11.2	1.00 (reference)	1.00 (reference)	1.00 (reference)
User	1115	168	15.1	1.41 (1.16-1.71)	0.87 (0.70-1.09)	0.82 (0.60-1.11)
All-cause death						
Non-user	3234	131	4.1	1.00 (reference)	1.00 (reference)	1.00 (reference)
User	1115	86	7.7	1.98 (1.50-2.62)	0.97 (0.70-1.34)	0.90 (0.58-1.39)
Mechanical ventilation						
Non-user	3234	66	2.0	1.00 (reference)	1.00 (reference)	1.00 (reference)
User	1115	57	5.1	2.59 (1.80-3.71)	1.41 (0.94-2.09)	1.24 (0.74-2.09)
ICU admission						
Non-user	3234	233	7.2	1.00 (reference)	1.00 (reference)	1.00 (reference)
User	1115	97	8.7	1.23 (0.96-1.57)	0.97 (0.73-1.28)	0.72 (0.49-1.05)
Cardiovascular adverse outcome (MI, stroke and TIA)						
Non-user	3234	31	1.0	1.00 (reference)	1.00 (reference)	1.00 (reference)
User	1115	15	1.3	1.41 (0.76-2.62)	1.17 (0.60-2.29)	0.79 (0.32-1.97)


MI = myocardial infarction; TIA = transient cerebral ischemic attack.

* Unweighted univariable logistic regression model

§ Unweighted multivariable logistic regression model adjusted for age, sex, insurance type, history of diabetes and history of hypertension


Inverse probability of treatment weighting (IPTW)-weighted univariable logistic model

Figure 1. Population-based cohort study design using the HIRA and KDCA database of South Korea



HIRA = Health Insurance Review and Assessment Service; KDCA = Korean Disease Control and Prevention Agency

Figure 2. Forest plot summarizing the risk of primary outcome associated with statin use when stratified for age, sex, history of hyperlipidaemia, hypertension, and diabetes mellitus.



Supplementary Material Legends

Supplementary Material 1. Data schema of the Health Insurance Review and Assessment Service database.

Supplementary Material 2. Diagnosis codes based on the Korean Standard Classification of Diseases, 7th Revision or International Classification of Disease, 10th Revision codes, National Procedure codes, and drug codes based on World Health Organization-Anatomical Therapeutic Chemical classification codes.

Supplementary Materials 3-7. The results of sensitivity analysis, by outcomes.

Supplementary Material 1. Data schema of the Health Insurance Review and Assessment Service (HIRA) database

1. Data Description

The database used in this study is the Health Insurance Review and Assessment Service (HIRA) claim data released by the South Korean government as the world's first de-identified COVID-19 nationwide data. Since South Korea have implemented a single National Health Insurance system, this administrative data includes the entire health insurance claims and clinical information of South Korean population. Information on non-essential medical treatment that are not covered by NHI was not collected based on fee-for-service payment system.

2. Data Scope

The list of patients of COVID-19 (using submitted claims data) is connected to their history of medical service use for the past 5 years (using finalized claims data, from January 2015 to February 2020), and the entire data set is de-identified. All benefit claims relevant to COVID-19 as of March 15, 2020.

3. Data Extraction Criteria

The databased include all health insurance types such as National Health Insurance, Medical Aid, Korea veterans service. COVID19 related claims consists of i) claims with "3/02" in classification code of MT043 (Target of medical cost support due to national disaster) based on the serial number of claim statement, ii) claims that contained "D6584" COVID-19 (Real-time polymerase chain reaction), iii) claims that contained COVID-19 related disease code (B342, B972, Z208, Z290, U18, U181, Z038, Z115, U071, U072), and iv) claims that contained other COVID-19 related fee code (COVID-19 related admission fee, management fee, IUD, isolation, public relief hospital, residential treatment center, screening center, negative pressure room, etc.).

4. Data Schema

The data based consists of COVID-19 related claims data, Healthcare service use history of the individuals who submitted COVID-19 related claims for the past 3 years. COVID-19 confirmation information and some related results such as was extracted from KCDA database. The included information are;

- Sociodemographic characteristics (i.e., sex, age, medical institution type)
- Diagnosis information (i.e., ICD-10 diagnosis codes, main diagnosis, sub diagnosis)
- Healthcare utilization information (i.e., number of visit, date of visit, duration of hospital stay)
- Medical expenditure information (i.e., out of pocket medical expenditure, reimbursed expenditure from NHI)
- Prescription information (i.e., national drug chemical code, number of supply, date of supply, dosage)

Note: COVID-19, coronavirus disease 2019; ICD-10, International Classification of Disease 10th Revision. Detailed information on data schema is available upon request.

Supplementary Material 2. Diagnosis codes based on the Korean Standard Classification of Diseases, 7th Revision or International Classification of Disease, 10th Revision codes, National Procedure codes, and drug codes based on World Health Organization-Anatomical Therapeutic Chemical classification codes.

Diagnoses	Codes
Inclusion criteria	
COVID-19 (KCD-7) Confirmed (Database-specific code)	B342, B972, Z208, Z290, U18, U181, Z038, Z115, U071, U072 'Y'
Study Endpoints	
All-cause death (Database-specific code)	'Y'
Intensive care unit admission (NPC)	AH110, AH190-192, AH194, AH195, AH210, AH29-299, AH390-AH396, AH398, AH399, AH501, AJ001, AJ003-011, AJ020, AJ021, AJ031, AJ043-046, AJ100, AJ102, AJ110, AJ112, AJ120, AJ122, AJ130, AJ132, AJ140, AJ142, AJ143, AJ150, AJ152 AJ160, AJ180, AJ190, AJ200, AJ202, AJ210, AJ212, AJ220, AJ222, AJ230, AJ240, AJ242, AJ250, AJ252, AJ260, AJ280, AJ290, AJ300, AJ302, AJ310, AJ312, AJ320, AJ322, AJ330, AJ332, AJ340, AJ342, AJ350, AJ352, AJ360, AJ380, AJ390, AJ500, AJ510, AJ520, AJ530, AJ540, AJ550, AJ560, AJ580, AJ590
Mechanical ventilation use (NPC)	M0850, M0857, M0858, M0860, M5830, M5850~5858, M5860, MM360, MM400
Cardiovascular adverse outcome	
Myocardial infarction	I21
Ischemic stroke	I63, I64, G463-468
TIA	G45
Comorbidities (ICD-10)	
Kidney disease	
Acute kidney failure	N17
Chronic kidney disease (CKD)	N18, N19
Asthma	J45, J46
Chronic lung disease	
Bronchiectasis	J47
Chronic obstructive pulmonary disease	J40, J41, J43, J44, E11
Interstitial pulmonary disease	J84
Liver disease	
Chronic liver disease	K70-K77
Viral hepatitis	B15-B19
Dementia	F00, F01, F02, F03, G30, G3100, G3182
Heart failure	
Heart failure	I110, I50
Valvular heart disease	I34, I35, I36, I37
Malignancy	C00-C97 (and expanded benefit coverage codes 'V027', 'V193', 'V194')
Arrhythmia	I44, I45, I47 (or Anti-arrhythmias drug uses ATC code 'C01B')
Other cerebrovascular diseases	G46, I65, I66, I67, I68, I69
Psychiatric disorders	F04-F99
Pneumonia including tuberculosis	
Tuberculosis	A15, A16, A17, A18, A19
Pneumonia	J12, J13, J14, J15, J16, J17, J18
Autoimmune disease	
Rheumatoid arthritis	M05, M06
Systemic connective tissue disorder	M30-M36
Noninfectious enteritis and colitis, ulcerative colitis, crohn's disease	K50-K52, R652, R653
Idiopathic thrombocytopenic purpura,	D693, D59

autoimmune hemolytic anemia	
Psoriatic arthritis, Juvenile Rheumatoid arthritis, Ankylosic spondylitis	M07, M08, M45
Other interstitial pulmonary disease	J84
Transplanted status, rejection	Z97, T86
Glomerular disease	N00-N08
Lupus	L93, M32
Intestinal malabsorption, celiac disease	K90
Multiple sclerosis, myasthenia gravis	G35, G700
Vasculitis	M05
Autoimmune thyroiditis, polyglandula failure, hepatitis	E063, E31, K754
Psoriasis	L40, L41, M07, M09
Sarcoidosis	D86, G532, M633 (or Immunosuppressant drug uses ATC codes 'L04A', 'H02', 'P01BA')
Diabetes mellitus	E10-E14
Hyperlipidemia	E78
Hypertension	I10-I15 (or Anti-hypertensive drug uses ATC codes 'C09A', 'C09B', 'C09C', 'C09D', 'C07', 'C08', 'C03', 'C01D', 'C02A', 'C02B', 'C02C')
Stroke or TIA	
Stroke	G45
TIA	I60-I64, G463, G464, G465, G466, G467, G468
Coronary artery disease	
Coronary artery disease	I20-I25
Atherosclerosis	I70
Peripheral vascular disease	I70-I79
Atrial fibrillation	I48
Thromboembolism	I26, I63, I74, I801, I802, I803, I809, I82
Study Drugs (ATC)	
Statin	
Lipid lowering agents	C10AA, C10AA01-08, C10BA, C10BA01-09, C10BX, C10BX01- 17, A10BH51, A10BH52
Simvastatin	C10AA01, C10BA02, C10BA04, C10BX01, C10BX04, A10BH51
Lovastatin	C10AA02, C10BA01
Pravastatin	C10AA03, C10BA03, C10BX02
Fluvastatin	C10AA04
Atorvastatin	C10AA05, C10BA05, C10BA08, C10BX03, C10BX06, C10BX08, C10BX11, C10BX12, C10BX15
Cerivastatin	C10AA06
Rosuvastatin	C10AA07, C10BA06, C10BA07, C10BA09, C10BX05, C10BX07, C10BX09, C10BX10, C10BX13, C10BX14, C10BX16, C10BX17, A10BH52
Pitavastatin	C10AA08
Co-medications (ATC)	
Antibiotics-systemic	J01
Antivirals	J05
Antidementia	N06D
Antidiabetics	A10
Anxiolytics	N05B
Antipsychotics	N05A
Antidepressants	N06A
NSAIDs	M01A
Acetaminophen	N02BE01, N02BE05, N02BE51, N02BE71
Immunosuppressant	L04A (L04AA, L04AB, L04AC, L04AD, L04AX), H02, P01BA
Anticoagulants	B01AA, B01AB, B01AE, B01AF, B01AX
Antiplatelets	A01AD05, C07FX02, C07FX03, C07FX04, C10BX01, C10BX02, C10BX04, C10BX05, C10BX06, C10BX08, C10BX12, M01BA03, N02AJ02, N02AJ07, N02AJ18, N02BA01, N02BA51, N02BA71, B01AC

ATC = Anatomical Therapeutic Chemical classification code; COVID-19 = coronavirus disease 2019; KCD-7 = Korean Standard Classification of Diseases 7th Revision; ICD-10 = International Classification of Disease 10th Revision; NPC = national procedure codes; TIA = transient cerebral ischemic attack; NSAIDS, nonsteroidal anti-inflammatory drugs.



Supplementary Material 3. Risk of primary outcome associated with statin use among COVID-19 patients with ≥ 40 years of age.

	Number of Patients	Cumulative Incidence (%)			Odds Ratio	(95% Confidence Interval)
		Non- Users	Users			
Confirmed-patients with COVID-19						
Statin user vs non-user	4613	10.7	14.7		0.77	(0.56-1.05)
Redefining the exposure searching window						
360 days before and including the date of cohort entry						
Statin user vs non-user	4613	10.7	14.7		0.71	(0.52-0.97)
Hospitalized-patients with COVID-19						
Statin user vs non-user	4349	11.2	15.1		0.82	(0.60-1.11)
Redefining the exposure searching window						
360 days before and including the date of cohort entry						
Statin user vs non-user	4349	11.2	15.0		0.77	(0.57-1.03)
Statistical Method based on propensity Score						
IPT weighting with trimming	4349	11.2	15.1		0.77	(0.57-1.03)
Outcome adjustment	4349	11.2	15.1		0.82	(0.64-1.06)
SMR weighting	4349	11.2	15.1		0.64	(0.45-0.93)
PS matching	4349	11.2	15.1		0.74	(0.56-0.98)





Supplementary Material 4. Risk of all-cause death associated with statin use among COVID-19 patients with ≥ 40 years of age.

	Number of Patients	Cumulative Incidence (%) Non- Users	Users		Odds Ratio	(95% Confidence Interval)	
Confirmed-patients with COVID-19							
Statin user vs non-user	4613	3.9	7.7		0.81	(0.64-0.96)	
Redefining the exposure searching window							
360 days before and including the date of cohort entry							
Statin user vs non-user	4613	3.8	7.8		0.75	(0.46-1.21)	
Hospitalized-patients with COVID-19							
Statin user vs non-user	4349	4.1	7.7		0.90	(0.58-1.39)	
Redefining the exposure searching window							
360 days before and including the date of cohort entry							
Statin user vs non-user	4349	4.0	7.8		0.87	(0.57-1.33)	
Statistical Method based on Propensity Score							
IPT weighting with trimming	4349	4.1	7.7		0.87	(0.57-1.33)	
Outcome adjustment	4349	4.1	7.7	0.84	(0.59-1.21)		
SMR weighting	4349	4.1	7.7	0.56	(0.35-0.89)		
PS matching	4349	4.1	7.7	0.67	(0.46-0.98)		

Supplementary Material 5. Risk of adverse mechanical ventilation associated with statin use among COVID-19 patients with ≥ 40 years of age.


	Number of Patients	Cumulative Incidence (%) Non- Users	Cumulative Incidence (%) Users		Odds Ratio	(95% Confidence Interval)
Confirmed-patients with COVID-19						
Statin user vs non-user	4613	1.9	4.8		1.23	(0.73-2.06)
Redefining the exposure searching window						
360 days before and including the date of cohort entry						
Statin user vs non-user	4613	1.9	4.7		1.18	(0.72-1.95)
Hospitalized-patients with COVID-19						
Statin user vs non-user	4349	2.0	5.1		1.24	(0.74-2.09)
Redefining the exposure searching window						
360 days before and including the date of cohort entry						
Statin user vs non-user	4349	2.0	5.0		1.17	(0.71-1.94)
Statistical Method based on Propensity Score						
IPT weighting with trimming	4349	2.0	5.1		1.18	(0.71-1.95)
Outcome adjustment	4349	2.0	5.1		1.22	(0.77-1.95)
SMR weighting	4349	2.0	5.1		1.09	(0.63-1.90)
PS Matching	4349	2.0	5.1		1.03	(0.63-1.70)

Supplementary Material 6. Risk of ICU admission associated with statin use among COVID-19 patients with ≥ 40 years of age

	Number of Patients	Cumulative Incidence (%)			Odds Ratio	(95% Confidence Interval)
		Non- Users	Users			
Confirmed-patients with COVID-19						
Statin user vs non-user	4613	6.8	8.4		0.70	(0.48-1.03)
Redefining the exposure searching window						
360 days before and including the date of cohort entry						
Statin user vs non-user	4613	6.9	8.3		0.67	(0.47-0.96)
Hospitalized-patients with COVID-19						
Statin user vs non-user	4349	7.2	8.7		0.72	(0.49-1.05)
Redefining the exposure searching window						
360 days before and including the date of cohort entry						
Statin user vs non-user	4349	7.2	8.6		0.67	(0.47-0.97)
Statistical Method based on Propensity Score						
IPT weighting with trimming	4349	7.2	8.7		0.67	(0.47-0.97)
Outcome adjustment	4349	7.2	8.7		0.79	(0.58-1.09)
SMR weighting	4349	7.2	8.7		0.67	(0.42-1.06)
PS matching	4349	7.2	8.7		0.77	(0.54-1.10)

0.2 1.0 5.0

Supplementary Material 7. Risk of cardiovascular outcomes (myocardial infarction, ischemic stroke and transient ischemic attack,) associated with statin use among COVID-19 patients with ≥ 40 years of age

	Number of Patients	Cumulative Incidence (%)			Odds Ratio	(95% Confidence Interval)
		Non- Users	Users			
Confirmed-patients with COVID-19						
Statin user vs non-user	4349	0.9	1.3		0.80	(0.33-1.96)
Redefining the exposure searching window						
360 days before and including the date of cohort entry						
Statin user vs non-user	4613	0.9	1.2		0.84	(0.36-1.97)
Hospitalized-patients with COVID-19						
Statin user vs non-user	4349	1.0	1.3		0.79	(0.32-1.97)
Redefining the exposure searching window						
360 days before and including the date of cohort entry						
Statin user vs non-user	4349	1.0	1.3		0.84	(0.36-1.97)
Statistical Method based on Propensity Score						
IPT weighting with trimming	4349	1.0	1.3		0.84	(0.36-1.98)
Outcome adjustment	4349	1.0	1.3		1.13	(0.51-2.50)
SMR weighting	4349	1.0	1.3		0.61	(0.16-2.26)
PS matching	4349	1.0	1.3		0.93	(0.36-2.39)