

The clinical feature of triple-negative breast cancer in Beijing, China

Hui Zhao¹, Yueqing Feng¹, Junzheng Yang^{1*}

¹Xinxiang Central Hospital, the Fourth Clinical College of Xinxiang Medical University,
Xinxiang, 453000, China

* Correspondence to Junzheng Yang (yangjunzheng606403@163.com), Xinxiang Central
Hospital, the Fourth Clinical College of Xinxiang Medical University,
Xinxiang, 453000, China

Abstract

Objective: To collect and analyze the clinical feature of triple-negative breast cancer (TNBC) in Beijing, to provide the information for the local oncologist to make sound treatment plans.

Methods: The clinical data of 280 breast cancer patients with TNBC admitted to the oncology hospital of China Academy of Medical Sciences were collected and divided into (recurrence and metastasis) group and non-(recurrence and metastasis) group. Breast cancer patients with TNBC were classified according to age distribution, family history of breast cancer, pathological type, histological grade, clinical stage, tumor thrombus, tumor size and lymph node metastasis. and 15 BRCA1 gene SNP loci were also measured by a high throughput Mass ARRAY time-of-flight mass spectrometry biochip system and compared the expression of 15 BRCA1 gene SNP loci between (recurrence and metastasis) group and non-(recurrence and metastasis) group. χ^2 test was used to analyze the difference between two groups, and $P < 0.05$ considered statistically significant.

Results: A total of 280 breast cancer patients with TNBC were enrolled in this study, median age 45 years old. There were 117 cases breast cancer patients with TNBC in (recurrence and metastasis) group, accounting for 41.79% in total breast cancer patients with TNBC and 163 cases breast cancer patients with TNBC in non-(recurrence and metastasis) group, accounting for 58.21% in total breast cancer

history of breast cancer, pathological type and histological grade between non-(recurrence and metastasis) group and (recurrence and metastasis) group ($P>0.05$); but there were significant differences in clinical stage, vascular tumor thrombus, tumor size and lymph node metastasis between two groups ($P<0.05$); and then we compared the expression of 15 BRCA1 gene SNP loci in (recurrence and metastasis) group and non-(recurrence and metastasis) group, found that BRCA1 gene rs 12516 CC loci (38.8% VS 44.4%), BRCA1 gene rs 12516 TT loci (15.6% VS 10.4%), BRCA1 gene rs 16940 CC loci (15.1% VS 10.4%), BRCA1 gene rs 16940 TT loci (39.0% VS 44.8%), BRCA1 gene rs 16941 AA loci (38.1% VS 44%), BRCA1 gene rs 16941 GG loci (15.0% VS 10.3%), BRCA1 gene rs16942 AA loci (39.0% VS 44.8%), BRCA1 gene rs16942 GG loci (15.1% VS 10.4%), BRCA1 gene rs799906 CC loci (15.9% VS 10.4%), BRCA1 gene rs799906 TT loci (38.7% VS 44.8%), BRCA1 gene rs799917 CC loci (38.7% VS 44.4%), BRCA1 gene rs799917 TT loci (15.7% VS 10.4%), BRCA1 gene rs1060915 CC loci (15.5% VS 10.4%), BRCA1 gene rs1060915 TT loci (39.1% VS 44.8%), BRCA1 gene rs1799966 AA loci (37.7% VS 44.4%), BRCA1 gene rs1799966 GG loci (15.1% VS 10.4%), BRCA1 Gene rs2070833 AA loci (3.1% VS 7.0%), BRCA1 Gene rs2070833 CC loci (56.3% VS 51.3%), BRCA1 gene rs3737559 GG loci (78.5% VS 84.5%), BRCA1 gene rs3737559 GA loci (19.0% VS 14.6%), BRCA1 gene rs8176199 AA loci (60.5% VS 64.6%), BRCA1 gene rs8176318 GG loci (38.4% VS 43.4%), BRCA1 gene rs8176318 TT loci (15.1% VS 10.6%), BRCA1 gene rs8176323 CC loci (38.6% VS 43.9%), BRCA1 gene rs8176323 GG loci (15.2% VS 10.5%), BRCA1 gene rs11655505 AA loci (14.9% VS 10.4%), BRCA1 gene rs11655505 GG loci (39.1% VS 44.8%) had a difference at the accident rate between recurrence and metastasis group and non-(recurrence and metastasis) group, but the frequencies of genotypes in the (recurrence and metastasis) group and non-(recurrence and metastasis) group were similar, there was no statistical significant correlation between the SNP genotype of the BRCA1 gene and the recurrence and metastasis risk of TNBC ($P>0.05$).

Conclusions: There were higher recurrence and metastasis (41.79%) in total 280 cases breast cancer patients with TNBC in Beijing area; breast cancer patients with

TNBC in Beijing area had a unique clinical feature no matter at clinical stage, vascular tumor thrombus, tumor size and lymph node metastasis or the expression of 15 BRCA1 gene SNP loci, those data may provide some information for clinical staff for TNBC treatment.

Keywords: TNBC; recurrence and metastasis; BRCA1; SNP

1 Introduction

Breast cancer is a kind of cancers with the higher incident rate in women in the world. The latest statistics showed that nearly 1.5 million women suffered from breast cancer every year, accounting for 25%-30% of all malignant tumors in women. Incidence rate of breast cancer has been increasing year by year with the improvement of living standard and lifestyle and the deterioration of living environment in China^[1]. Breast cancer is a highly heterogeneous disease, and there are great differences in tissue morphology, molecular phenotypes, clinical manifestations, therapeutic effects and prognosis^[2].

Triple negative breast cancer (TNBC) is a kind of subtype of breast cancer with negative estrogen receptor (ER), negative progesterone receptor (PR) and negative human epidermal growth factor receptor 2 (HER2). accounting for 15~20% in all breast cancer types^[3]. It has the characteristics of poor tumor differentiation, strong invasiveness, easy local recurrence and distant metastasis^[4]. The effective treatment methods are mainly chemotherapy and radiotherapy, and it is often insensitive to radiotherapy and chemotherapy. The present study found that TNBC is a group of tumors with different biological behavior. There are obvious differences in molecular biological characteristics, clinical manifestations and prognosis, and can be divided into more different subtypes^[5].

BRCA1 gene is located in human 17q21 with a length of about 100kb, including 24 exons and 22 introns, there are 22 exons with transcription and coding functions, and finally encode into phosphorylated protein containing 1863 amino acids, which plays a role in DNA damage repair, homologous recombination, cell cycle regulation, transcriptional activation and inhibition and centrosome replication^[6]. BRCA1 gene is the most common susceptibility gene for breast cancer. The proportion of breast

cancer with BRCA1 mutation in the lifetime is as high as 60%~80%, and the onset age is more than 20 years earlier than that of the general population^[7]. A large number of studies have shown that the proportion of TNBC in BRCA1 mutation carriers is higher, and often shows large size, poor differentiation, strong invasion, and easy to have local recurrence and distant metastasis; more than 30% of the TNBC are associated with mutations in the BRCA1 gene^[8].

Here, we investigated and analyzed the clinical features of TNBC patients in Beijing, compared the difference between (recurrence and metastasis) group and non-(recurrence and metastasis) group in age distribution, family history of breast cancer, pathological type, histological grade, clinical stage, vascular tumor thrombus, tumor size and lymph node metastasis and the expression of 15 BRCA1 gene SNP loci, which may provide some clues for clinic staff for TNBC treatment in Beijing.

2 Investigation subject

The investigation subjects in this study were TNBC breast cancer patients who were admitted to the Cancer Hospital of China Academy of Medical Sciences in the past 5 years. There were a total of 280 breast cancer patients, They were Han population of China, had complete demographic data, clinical pathological data and follow-up data. The patients in the study were divided into (recurrence and metastasis) group and non-(recurrence and metastasis) group according to whether there was recurrence or metastasis. Usually according to the criterion, patients who had no local recurrence or distant metastasis for more than 3 years from diagnosis to enrollment were considered there was no recurrence and metastasis.

All the patients in the study were diagnosed as TNBC patients, when expression of ER, PR and HER2 were negative considered as TNBC patients. The expression of ER and PR was determined by immunohistochemistry (IHC), The expression of HER-2 was determined by IHC or fluorescence in situ hybridization (FISH). When the positive ratio <10% in breast cancer cells considered ER negative and PR negative, the criteria for HER2 negative were immunohistochemical staining (-)/(+) or fluorescence in situ hybridization without amplification.

All enrolled patients were followed up for a long time by consulting the original

medical records, outpatient reexamination and telephone follow-up, and the tumor recurrence, metastasis and survival of enrolled patients were recorded in detail.

3 Investigation methods

TNBC breast cancer patients were classified according to age distribution, family history of breast cancer, pathological type, histological grade, clinical stage, vascular tumor thrombus, tumor size and lymph node metastasis and the expression of 15 BRCA1 gene SNP loci, and compared the difference between (recurrence and metastasis) group and non-(recurrence and metastasis) group.

For BRCA1 gene SNP determination, firstly BRCA1 gene was retrieved from the SNP public database provided and the human genome genetic variation integration map published by the thousand people genome project (<http://hapmap.ncbi.nlm.nih.gov/>), all SNP loci related to the known BRCA1 gene information in the Chinese Han population were selected, combined with the statistical characteristics of minimum allele frequency (MAF) and Hardy Weinberg equilibrium p value (HWPval), the above candidate SNP sites were screened to obtain the best labeled SNP. Selection requirements was $MAF > 0.1$ and $HWPval > 0.1$. Then the labeled SNP sites were replaced and supplemented according to the location of SNP sites (missense mutation sites in exon region, UTR region, promoter region and intron region were usually selected, and synonymous mutation was not selected) and the positive SNP sites related to BRCA1 gene reported in the literature at home and abroad. Finally, sequence primers were designed for the selected loci by software. Considered the interaction between different primers, 15 SNP loci were selected finally. The specific information of SNP loci of BRCA1 gene is shown in Table 1.

Table 1. The information of 15 SNP loci of BRCA 1

SNP loci	Functional location	Allele gene
rs16940	exon-synon	C/T
rs16941	exon	A/G
rs16942	exon	A/G
rs12516	3' UTR	C/T

rs799906	promoter	C/T
rs1799966	exon	A/G
rs799917	exon-missense	C/T
rs3737559	intron	A/G
rs1060915	exon-synon	C/T
rs2070833	intron	C/A
rs4793204	promoter	A/G
rs8176199	intron	C/A
rs8176318	3' UTR	G/T
rs8176323	3' UTR	C/G
rs11655505	promoter	A/G

Table 2. Primer sequences of 15 SNP loci of BRCA 1

SNP loci	Primer sequence	Amplification sequence
rs16940	Forward: ACGTTGGATGGACCAGTATGACTGAATCTC	AGCACTAGTCACCTTCT
	Reverse: ACGTTGGATGATTTAGCACAGGGCTACCTC	
rs16941	Forward: ACGTTGGATGTTAGTACTGGAACCTACTTC	TAACATTAGAGAAAATGTTT TTAAAG
	Reverse: ACGTTGGATGACAGTGAGCACAATTAGCCG	
rs16942	Forward: ACGTTGGATGCTGCTGTTTTTAGCAAAAGCG	ACTCCTGCTAAGCTCTCCT
	Reverse: ACGTTGGATGGTAACCCTGAGCCAAATGTG	
rs12516	Forward: ACGTTGGATGTATAGGTCAGGAACCACACG	TGTGTGCTGATTCTCCT
	Reverse: ACGTTGGATGAATCCCCAGATCTTGTACC	
rs799906	Forward: ACGTTGGATGTGATTGCCTGGCCAATGTCC	ATTGGGCCCAACAGG
	Reverse: ACGTTGGATGTTCCGAAGCCTCTAAGAGTC	
rs1799966	Forward: ACGTTGGATGTTGAAAAGTTGCAGAATCT	AGTATGAGCAGCAGCTGGAC
	Reverse: ACGTTGGATGTCCCTGCTCACACTTTCTTC	
rs799917	Forward: ACGTTGGATGCACCTGGCCAACCCCGTG	CCCTGCAGCACTTCGT
	Reverse: ACGTTGGATGTATCAGCGGCGACGGGAG	
rs3737559	Forward: ACGTTGGATGAGTAAGGACCTAGCTCCAAC	ACCAAGGCTCCATAATTAC
	Reverse: ACGTTGGATGTTGTGAGCAGGGACAAGAAC	
rs1060915	Forward: ACGTTGGATGATAATGGGAGTGCCCATACC	CAATCCTGGCACTTGTAGA
	Reverse: ACGTTGGATGAGAAGGGAGACTCAATCCTG	

rs2070833	Forward: ACGTTGGATGAGGTTTGCTGACATCATCTC Reverse: ACGTTGGATGCCATCAAGGTGCTTACAGTC	ACAATGAACCACAAACAATT
rs4793204	Forward: ACGTTGGATGAGGCAAGAGGATTGCTTGAG Reverse: ACGTTGGATGTCTCATTCTGCACTCAGGC	CGATCATAGCTTGCTGC
rs8176199	Forward: ACGTTGGATGCTCGTGATCGTATGTGTATG Reverse: ACGTTGGATGCCACGATGACAGAGATATCC	CGTATGTGTATGTGCGTGATT GT
rs8176318	Forward: ACGTTGGATGAGCAGGGCTTAGCTGCTTGT Reverse: ACGTTGGATGACTTAGCCACTGTGAACACG	TGTGAACACGACTTGG
rs8176323	Forward: ACGTTGGATGCTGAGCAGTTTTTACTCTGTG Reverse: ACGTTGGATGGGAGAATAGGTCAATTCATAC	GGATAGATATGGTATCCCAG
rs11655505	Forward: ACGTTGGATGAATGCAAAGACCGTCCGCTG Reverse: ACGTTGGATGGGAGAAACGGGACTAGTTAC	GGGACTAGTTACTGTCTTT

For PCR amplification reaction system as follows:

Reagent	Concentration	Volume (uL)
Water, HPLC grade	NA	6.5
PCR Buffer with 15mM MgCl ₂	10x	10
MgCl ₂	25mM	0.5
dNTP Mix	25mM	0.5
Primer Mix	0.5uM	0.5
HotStar Taq	5U/μL	0.5
DNA template	10ng/uL	1
Total		20

And PCR amplification procedure as follows:

Temperature (°C)	Time (second)	Cycle
94	900	1
94	20	
56	30	
72	60	45
72	180	1
4	∞	1

SAP reaction system as follows:

Reagent	Concentration	Volume (uL)
Water	NA	6
SAP Buffer	10x	2
SAP	1.7U/uL	2
Total		10

SAP procedure as follows:

Temperature (°C)	Time (minute)	Cycle
37	40	1
85	5	1
4	∞	1

SAP extension reaction system as follows:

Reagent	Concentration	Volume (uL)
Water	NA	3.5
iPLEX buffer plus	10x	5
iPLEX terminator	NA	0.5
Primer Mix	0.6-1.3uM	0.4
iPlex enzyme	NA	0.6
Total		10

SAP extension procedure as follows:

Temperature (°C)	Time	Cycle
94	30sec	1
94	5sec	1
52	5sec	40
80	5sec	5
72	3min	1
4	∞	1

4 Statistics

All data were analyzed by SPSS17.0 software. χ^2 test were used for analyzing the

difference between groups, $P < 0.05$ considered as statistical significance.

5 Results

A total of 280 TNBC breast cancer patients were enrolled in this study, median age were 45 years old (24~78 years old), there were 85 cases TNBC patients aged under 40 years old, and 195 cases TNBC patients aged above 40 years old, accounting for 30.36% and 69.64% in total TNBC breast cancer patients respectively. There were 117 cases TNBC patients occurred recurrence and metastasis, accounting for 41.79% in total 280 cases breast cancer patients with TNBC in Beijing area, There was no significant difference in age distribution, family history of breast cancer, pathological type and histological grade between relapse and metastasis group and disease-free survival group ($P > 0.05$). There were significant differences in clinical stage, vascular tumor thrombus, tumor size and lymph node metastasis ($P < 0.05$). In the (recurrence and metastasis) group, 58 cases (49.6%) had late clinical stage, 62 cases (53.0%) had vascular tumor thrombus, 74 cases (63.2%) had large tumor and 84 cases (71.8%) had lymph node metastasis; the data were shown at table 3.

Table 3. the demographic characteristics of TNBC Patients in the study

	non-(Recurrence and metastasis) group	(Recurrence and metastasis) group	P value
	Cases (%)	Cases (%)	
Age			0.696
≤40	48 (29.4)	37 (31.6)	
>40	115 (70.6)	80 (68.4)	
Family history			0.223
No	150 (92.0)	105 (89.7)	
Yes	10 (6.1)	12 (10.3)	
Unknown	3 (1.9)	0 (0.0)	
Clinical stage			0.000
I+II stage	141 (86.5)	54 (46.2)	
III+IV stage	19 (11.7)	58 (49.6)	
Unknown	3 (1.8)	5 (4.2)	
Pathological type			0.233

Invasive nonspecific carcinoma	131(80.4)	102(87.2)	
Other type	29 (17.8)	15(12.8)	
Unknown	3 (1.8)	0 (0.0)	
Histological grade			0.111
Level1+level 2	88 (54.0)	56 (47.9)	
Level 3	53 (32.5)	51 (43.6)	
Unknown	22 (13.5)	10 (8.5)	
Vascular tumor thrombus			0.017
No	98 (60.1)	52 (44.4)	
Yes	65 (39.9)	62(53.0)	
Unknown	0 (0.0)	3 (2.6)	
Tumor size			0.004
≤2cm	82 (50.3)	39 (33.3)	
>2cm	75 (46.0)	74 (63.2)	
Unknown	6 (3.7)	4 (3.4)	
Lymph node metastasis			0.000
No	112 (68.7)	28 (23.9)	
Yes	48 (29.5)	84 (71.8)	
Unknown	3 (1.8)	5 (4.3)	

And then we detected and analyzed the expression of 15 BRCA1 gene SNP loci in (recurrence and metastasis) group and non-(recurrence and metastasis) group, found that BRCA1 gene rs 12516 CC loci (38.8% VS 44.4%), BRCA1 gene rs 12516 TT loci (15.6% VS 10.4%), BRCA1 gene rs 16940 CC loci (15.1% VS 10.4%), BRCA1 gene rs 16940 TT loci (39.0% VS 44.8%), BRCA1 gene rs 16941 AA loci (38.1% VS 44%), BRCA1 gene rs 16941 GG loci (15.0% VS 10.3%), BRCA1 gene rs16942 AA loci (39.0% VS 44.8%), BRCA1 gene rs16942 GG loci (15.1% VS 10.4%), BRCA1 gene rs799906 CC loci (15.9% VS 10.4%), BRCA1 gene rs799906 TT loci (38.7% VS 44.8%), BRCA1 gene rs799917 CC loci (38.7% VS 44.4%), BRCA1 gene rs799917 TT loci (15.7% VS 10.4%), BRCA1 gene rs1060915 CC loci

(15.5% VS 10.4%), BRCA1 gene rs1060915 TT loci (39.1% VS 44.8%), BRCA1 gene rs1799966 AA loci (37.7% VS 44.4%), BRCA1 gene rs1799966 GG loci (15.1% VS 10.4%), BRCA1 Gene rs2070833 AA loci (3.1% VS 7.0%), BRCA1 Gene rs2070833 CC loci (56.3% VS 51.3%), BRCA1 gene rs3737559 GG loci (78.5% VS 84.5%), BRCA1 gene rs3737559 GA loci (19.0% VS 14.6%), BRCA1 gene rs8176199 AA loci (60.5% VS 64.6%), BRCA1 gene rs8176318 GG loci (38.4% VS 43.4%), BRCA1 gene rs8176318 TT loci (15.1% VS 10.6%), BRCA1 gene rs8176323 CC loci (38.6% VS 43.9%), BRCA1 gene rs 8176323 GG loci (15.2% VS 10.5%), BRCA1 gene rs11655505 AA loci (14.9% VS 10.4%), BRCA1 gene rs11655505 GG loci (39.1% VS 44.8%) had a difference at the accident rate between (recurrence and metastasis) group and non-(recurrence and metastasis) group, but the frequencies of genotypes in the (recurrence and metastasis) group and non-(recurrence and metastasis) group were similar, there was no statistical significant correlation between the SNP genotype of the BRCA1 gene and the recurrence and metastasis risk of TNBC ($P > 0.05$), the results were shown at table 4.

Table 4. the expression of 15 BRCA1 gene SNP loci in TNBC patients

SNP	Group A Cases (%)	Group B Cases(%)	P value
BRCA1 gene rs 12516			0.396
CC	62 (38.8)	51(44.4)	
TT	25 (15.6)	12(10.4)	
TC	73 (45.6)	52(45.2)	
BRCA1 gene rs 16940			0.422
CC	24 (15.1)	12(10.4)	
TT	62 (39.0)	52(44.8)	
TC	73 (45.9)	52(44.8)	
BRCA1 gene rs 16941			0.427
AA	61 (38.1)	51(44.0)	
GG	24 (15.0)	12 (10.3)	
AG	75 (46.9)	53 (45.7)	
BRCA1 gene rs16942			0.422

AA	62 (39.0)	52 (44.8)	
GG	24 (15.1)	12 (10.4)	
GA	73 (45.9)	52 (44.8)	
BRCA1gene rs799906			0.334
CC	26 (15.9)	12 (10.4)	
TT	63 (38.7)	52 (44.8)	
TC	74 (45.4)	52 (44.8)	
BRCA1gene rs799917			0.373
CC	61(38.4)	51(44.4)	
TT	25 (15.7)	12(10.4)	
TC	73 (45.9)	52(45.2)	
BRCA1gene rs1060915			0.389
CC	25 (15.5)	12 (10.4)	
TT	63 (39.1)	52(44.8)	
TC	73 (45.4)	52 (44.8)	
BRCA1gene rs1799966			0.391
AA	60 (37.7)	51 (44.4)	
GG	24 (15.1)	12 (10.4)	
GA	75 (47.2)	52 (45.2)	
BRCA1gene rs2070833			0.301
AA	5 (3.1)	8 (7.0)	
CC	90 (56.3)	59 (51.3)	
CA	65 (40.6)	48 (41.7)	
BRCA1gene rs3737559			0.348
AA	4 (2.5)	1 (0.9)	
GG	128(78.5)	98(84.5)	
GA	31(19.0)	17(14.6)	
BRCA1gene rs4793204			0.943
GG	115 (75.2)	83 (74.8)	
GA	38 (24.8)	28 (25.2)	
BRCA1gene rs8176199			0.772
AA	98 (60.5)	75 (64.6)	
CC	10 (6.2)	6 (5.2)	
CA	54 (33.3)	35 (30.2)	
BRCA1gene rs8176318			0.494
GG	61(38.4)	49 (43.4)	
TT	24(15.1)	12(10.6)	
GT	74(46.5)	52(46.0)	
BRCA1gene rs8176323			0.463
CC	61 (38.6)	50 (43.9)	
GG	24 (15.2)	12 (10.5)	

	GC	73 (46.2)	52 (45.6)	
BRCA1 gene rs11655505				0.444
AA		24 (14.9)	12 (10.4)	
GG		63 (39.1)	52 (44.8)	
AG		74 (46.0)	52 (44.8)	

Group A: non-(recurrence and metastasis) group; Group B: (recurrence and metastasis) group.

6 Discussion

TNBC has the poor prognosis and worse survival, endocrine therapy and targeted therapy is ineffective for TNBC treatment; the effective treatment methods are mainly chemotherapy and radiotherapy, but they are often insensitive to radiotherapy and chemotherapy^[9,10], therefore, it is of important clinical significance to understand the clinical characteristics and pathogenesis of TNBC patients for local oncologist.

In this study, we collected and analyzed the clinical data of a total 280 case of TNBC patients which admitted to the oncology hospital of China Academy of Medical Sciences. There were 85 cases TNBC patients aged under 40 years old, and 195 cases TNBC patients aged above 40 years old, accounting for 30.36% and 69.64% in total TNBC breast cancer patients respectively, patients there were 117 cases breast cancer patients occurred recurrence and metastasis, accounting for 41.79% in total 280 cases breast cancer patients with TNBC in Beijing area; clinical stage, vascular tumor thrombus, tumor size and lymph node metastasis in (recurrence and metastasis) group were significantly different from their in non-(recurrence and metastasis) group, those data demonstrated that the age of TNBC were 40 years old, and there were the higher recurrence and metastasis in TNBC patients in Beijing area.

And then we detected and analyzed expression of 15 BRCA1 gene SNP loci in (recurrence and metastasis) group and non-(recurrence and metastasis) group, found that there were many difference in several SNP loci, but the frequencies of genotypes in the (recurrence and metastasis) group and non-(recurrence and metastasis) group were similar, there was no statistical significant correlation between the SNP genotype of the BRCA1 gene and the recurrence and metastasis risk of TNBC ($P>0.05$). Those data demonstrated that the BRCA1 gene SNP loci in TNBC patients expression of BRCA1 gene in different TNBC patients, and the mechanism of TNBC was genetic

variation, recurrence and metastasis of negative breast cancer is relatively complicated.

7 Conflict of interests

The authors declare that they have no competing interests in this article.

8 Acknowledgements

We would like to thank Mr. Ma Fei from the Department of oncology, cancer hospital, Chinese Academy of Medical Sciences for his guidance and help in this study.

9 Funding

None.

10 References

- [1] Jie He, Ping Zhao, Wanqing Chen. Annual report of China tumor registration [M]. Beijing: Military Medical Science Press, 2012.
- [2] Hugh J, Hanson J, Cheang MC, et al. Breast cancer subtypes and response to docetaxel in node-positive breast cancer: use of an immunohistochemical definition in the BCIRG 001 trial[J]. *J Clin Oncol*, 2009, 27(8): 1168-1176.
- [3] Yao-Lung K, Dar-Ren C, Tsai-Wang C. Clinicopathological features of triple-negative breast cancer in Taiwanese women[J]. *Int J Clin Oncol*, 2011, 16(5) : 500-505.
- [4] Heitz F, Harter P, Lueck HJ, et al. Triple-negative and HER-2-overexpressing breast cancers exhibit an elevated risk and an earlier occurrence of cerebral metastases[J]. *Eur J Cancer*, 2009, 45(16) : 2792-2798.
- [5] Junling Ren, Chen Bin, Yabing Wang. Research progress of three negative breast cancer [J]. *Journal of oncology*, 2011, 16(10): 817-820 (In Chinese).
- [6] Heng Ji, Baoguo Wang. Advances in breast cancer resistance protein (BCRP/ABCG2) [J]. *oncology pharmacy*, 2013, 3(5): 326-330 (In Chinese).
- [7] Ping Zhang, Binghe Xu, Dongxin Lin. Association of small molecule RNA gene polymorphism and ESR1 gene polymorphism with susceptibility and prognosis of three negative breast cancer. [D]. [Ph.D. thesis]. Beijing: Peking Union Medical College, 2011: 1-74 (In Chinese).
- [8] Lingxiao Zhang, Baohe Xu. Relationship between polygene single nucleotide polymorphisms

and prognosis and efficacy of three negative breast cancer[D]. [doctoral dissertation]. Beijing: Peking Union Medical College, 2012: 1-67 (In Chinese).

[9] Gelmon, K. et al. Targeting Triple-Negative Breast Cancer: Optimising Therapeutic Outcomes. *Ann. Oncol.*, 2012, 23: 2223–2234.

[10] Bianchini, G., Balko, J. M., Mayer, I. A., Sanders, M. E. & Gianni, L. Triple-Negative Breast Cancer: Challenges and Opportunities of a Heterogeneous Disease. *Nat. Rev. Clin. Oncol.*, 2016, 13: 674–690.