Epidemiological characteristics of the B.1.526 SARS-CoV-2 variant

Wan Yang,[1][*] Sharon K. Greene,[2] Eric R. Peterson,[2] Wenhui Li,[3] Robert Mathes,[2] Laura Graf,[2] Ramona Lall,[2] Scott Hughes,[4] Jade Wang,[4] Anne Fine[2][#]

[1]Department of Epidemiology, Mailman School of Public Health, Columbia University;
[2]Bureau of Communicable Disease, New York City Department of Health and Mental Hygiene;
[3]Bureau of Vital Statistics, New York City Department of Health and Mental Hygiene;
[4]Public Health Laboratory, New York City Department of Health and Mental Hygiene
* Corresponding author: wy2202@cumc.columbia.edu
#current affiliation: Council of State and Territorial Epidemiologists

Abstract:

To characterize the epidemiological properties of the B.1.526 SARS-CoV-2 variant of interest, here we utilized nine epidemiological and population datasets and model-inference methods to reconstruct SARS-CoV-2 transmission dynamics in New York City, where B.1.526 emerged. We estimated that B.1.526 had a moderate increase (15-25%) in transmissibility and could escape immunity in 0-10% of previously infected individuals. In addition, B.1.526 substantially increased the infection-fatality risk (IFR) among adults 65 or older by >60% during Nov 2020 – Apr 2021, compared to baseline risk estimated for preexisting variants. Overall, findings suggest that new variants like B.1.526 likely spread in the population weeks prior to detection and that partial immune escape (e.g., resistance to therapeutic antibodies) could offset prior medical advances and increase IFR. Early preparedness for and close monitoring of SARS-CoV-2 variants, their epidemiological characteristics, and disease severity are thus crucial to COVID-19 response as it remains a global public health threat.

Main Text:

INTRODUCTION

The SARS-CoV-2 virus spread quickly worldwide in early 2020, causing the COVID-19 pandemic. As the virus spread, it also diversified, and multiple novel SARS-CoV-2 variants emerged in different populations, producing both local and global waves of infection. Several variants have been characterized as variants of concern (VOC) or of interest (VOI), based on evidence regarding their ability to increase transmissibility, evade immunity conferred by either prior infection or vaccination, or cause more severe disease. Accurately estimating the epidemiological characteristics and impact of these variants is thus important for informing public health response, such as monitoring effectiveness of vaccines and therapeutic antibodies. More broadly, such findings can also provide insights into the long-term trajectory of SARS-CoV-2 beyond the pandemic phase.

The B.1.526 variant (WHO designation: Iota)(*1*), a SARS-CoV-2 VOI, was identified during Nov 2020 and quickly became a predominant variant in the New York City (NYC) area (*2-4*). It has also been detected in all 52 states/territories in the US and at least 27 other countries (GISAID data (*5*), as of 6/9/2021). An initial laboratory study (*2*) suggested that this variant is to some extent resistant to two therapeutic monoclonal antibodies in clinical use and neutralization by convalescent plasma and vaccinee sera. However, another study (*4*) examined all sequenced B.1.526 cases in NYC identified as of April 5, 2021 (n = 3,679) and showed preliminary evidence that this variant did not increase risk for infection after vaccination or reinfection. Both studies may be limited due to the small number of specimens available for analysis as well as delay in observation and reporting. Given these discrepancies, here we utilize detailed population epidemiological data collected since the beginning of the COVID-19 pandemic in NYC (March 1, 2020 – April 30, 2021) and multiple model-inference methods to estimate the transmissibility, immune escape potential, and disease severity of B.1.526. Of note, we include the combination of B.1.526-S:E484K and B.1.526-S:S477N as B.1.526.

As shown in Fig 1 (overall study design), we first apply a network model-inference system to reconstruct underlying SARS-CoV-2 transmission dynamics in NYC, accounting for under-detection of infection. This analysis allows estimation of key population variables and parameters (e.g., the infection rate including those not detected as cases and transmission rate) at the neighborhood-level as well as citywide. As such, we are able to examine in detail the epidemiological dynamics in Washington Heights – Inwood (WHts), the neighborhood where B.1.526 was initially identified (*2*). Using these estimates and additional variant prevalence data, we further apply a city-level multi-variant, age-structured model to estimate the changes in transmissibility and immune escape potential for B.1.526. Lastly, we utilize findings from the first two model systems to estimate variant-specific infection fatality risk (IFR, i.e. the fraction of all persons with SARS-CoV-2 infection who died from the disease), for B.1.526 and B.1.1.7, separately. Overall, our study documents the emergence and impact of B.1.526. We close with a discussion on lessons learned from this SARS-CoV-2 VOI and implications for future COVID-19 pandemic response.

RESULTS

Epidemic dynamics of the second pandemic wave in NYC

NYC experienced a very large first pandemic wave during spring 2020. Similar to our previous work (*6*), the model-inference system here estimates that 16.6% of the population (95% CrI: 13.6 – 21.5%; or 1.1 – 1.8 million people) had been infected by the end of May 2020 (i.e., end of the first wave; Fig 2D). The city was able to gradually reopen part of its economy during summer 2020 after a 3-month long stay-at-home mandate for all non-essential workers. However, infection resurged beginning in the fall of 2020 and the city experienced a second pandemic

wave around Nov 2020 – April 2021 (Fig 2). Following the second wave, an estimated total of
41.7% (95% CrI: 35.4 – 49.3%; or 3.0 – 4.1 million people) had been infected by the end of Apr
2021, including all those infected during the first wave. Note these estimates accounted for
under-detection of infections (Fig S1), for which the overall infection-detection rate increased to
37.1% (95% CrI: 33.3 – 43.0%) during the 2nd wave from 15.1% (95% CrI: 11.7 – 18.5%) during
the 1st wave. This large number of infections occurred despite the non-pharmaceutical
interventions implemented throughout the pandemic and rollout of mass-vaccination starting
mid-Dec 2020. In addition, unlike the first wave that predominantly affected older age groups,
the second pandemic wave affected all age groups (Figs S2-3).
86
Transmission rate increased earliest in the neighborhood where B.1.526 was initially
identified
The emergence and rapid increase of B.1.526 coincided with the second pandemic wave in NYC.
While first reported in Feb 2021 (*2*), testing initially identified the B.1.526 variant in patient
samples dated back to early Nov 2020 from the city's WHts neighborhood (*2*). As such, we first
examined potential changes in the transmission rate there. Indeed, prior to the identification of
B.1.526, estimated neighborhood relative transmission rate (b_i in Eqn 1; see Methods) in WHts
gradually increased, remained at high levels during Nov 2020 – Feb 2021, and decreased to the
baseline level afterwards when B.1.526 became a predominant variant citywide (~40% of all
cases sequenced by end of Feb 2021). In comparison, the estimates were relatively stable for
other neighborhoods (Fig 3A), suggesting the changes in WHts were likely due to the early
spread of B.1.526. Averaging over this period, we estimate that the relative transmission rate in
WHts increased by 8.4% (95% CI: -5.8 – 22.5%). Concurrently, the citywide transmission rate
increased by 13.3% (95% CI: -21.1 – 47.8%; Fig 3B). These two preliminary estimates in
combination suggest that the transmission rate of B.1.526 likely is 22.8% (95% CI: -12.4 –
58.0%) higher than preexisting non-VOC/VOI variants, without accounting for potential change
due to immune evasion.
104
B.1.526 likely causes a moderate increase in transmissibility (15-25%) and slight immune
evasion (0-10%)
We further examine model estimations under a wide range of transmissibility and immune
escape settings for B.1.526. Under all three possible scenarios of initial prevalence (i.e., 0.5 –
2.5%, 1.5 – 3.5%, and 0.5 – 3.5%), model simulations consistently show that B.1.526 likely
increases transmissibility by 15-30% and can escape immunity in 0-10% of previously infected
persons (Fig 4 A-C). Overall, a higher initial prevalence (1.5 – 3.5% at the beginning of Nov
2020) combining with a 15-25% increase in transmissibility and 0-10% immune escape (Fig 4 A-
C, middle column; and Fig 4D) generated the most accurate estimates of cases, hospitalizations
and deaths as well as variant percentages during the second wave. Model simulations show that,
with this moderate increase in transmissibility and small immune escape, B.1.526 was able to
outcompete preexisting variants and gradually increase its percentage from Nov 2020 to March

2021; however, afterwards its percentage decreased with the surge of B.1.1.7, a more infectious
variant (Fig 4D, bottom right panel).
119

B.1.526 likely increases disease severity substantially
During the second wave, estimated IFR increased gradually in later months, particularly among
older age groups (Fig 5). During this period (Nov 2020 – Apr 2021), 16-35% of hospital beds
and 18-32% of intensive care unit (ICU) beds in NYC were available, suggesting lack of access
to healthcare was not a reason behind the IFR increases. In addition, the number of COVID-19-
related deaths declined following mass-vaccination in early 2021. Modeling accounting for
infections and deaths due to B.1.526, B.1.1.7, and non-VOC/VOI variants suggests that B.1.526
increased IFR in older adults: by 46% (95% CI: 7.4 – 84%) among 45-64 year-olds [absolute
IFR: 0.42% (95% CI: 0.31 – 0.54%) vs. 0.29% (95% CI: 0.15 – 0.44%) baseline risk]; 82% (95%
CI: 20 – 140%) among 65-74 year-olds [absolute IFR: 1.9% (95% CI: 1.2 – 2.5%) vs. 1.0%
(95% CI: 0.57 – 2.5%) baseline risk], and 62% (95% CI: 45 – 80%) among 75+ [absolute IFR:
6.7% (95% CI: 5.9 – 7.4%) vs. 4.1% (95% CI: 2.2 – 6.3%) baseline risk], during Nov 2020 –
Apr 2021; overall, B.1.526 increased the IFR by 60% (95% CI: 38 – 82%), compared to
estimated baseline risk (Table 1). The analysis restricting to Nov 2020 – Jan 2021 suggests
similar IFR increases (Table S3). These estimated IFR increases were lower than for B.1.1.7 but
comparable. Of note, the IFRs for B.1.1.7 estimated here were higher than but in line with those
reported in the UK [e.g., overall increase: 100% (75-130%) vs. 61% (42–82%) in the UK(*7*)].

DISCUSSION

The B.1.526 variant is one of the SARS-CoV-2 variants designated as a VOI by both the WHO
(*1*) and the US CDC (*8*). However, due to a lack of extensive genomic sequencing and contact
tracing data particularly during the early phase of its emergence, its key epidemiological
properties have not been well characterized. Utilizing multiple epidemiological datasets and
comprehensive modeling, here we have estimated the changes in transmissibility, immune
escape potential, and disease severity for B.1.526. Results suggest that, compared to preexisting
non-VOC/VOI variants, B.1.526 causes a moderate increase in transmissibility and minimal
immune evasion; however, it might substantially increase IFR in older adults. As such, continued
monitoring of the circulation of this variant is warranted.

Our study offers several lessons for future outbreak response. First, prior to the emergence of
B.1.526, the estimated transmission rate in WHts, where it likely emerged, was consistently
higher than other neighborhoods in NYC throughout the pandemic. Population characteristics
(e.g., household structure) that may contribute to this higher transmission rate need further
investigation; however, the higher transmission rate may have facilitated the spread of new
mutants between hosts and its emergence population-wide. It is thus important to closely monitor
populations with sustained higher transmission rates for new variants, particularly in areas

lacking robust and timely sequencing of samples from newly identified cases. In addition, the

estimated transmission rate in WHts further increased in conjunction with the emergence of

B.1.526; such changes thus may serve as an early indicator for in-depth epidemiological

investigation (e.g., to assess changes in circulating variants and transmissibility). A similar

approach has been applied in the UK, where subregions with higher estimated growth rates were

prospectively investigated, leading to identification of B.1.1.7 as a VOC (*9-12*).

Second, we did not find a higher B.1.526-related IFR among younger age groups (those under 45

years); this finding is consistent with the findings of Thompson et al. (*4*) based on analysis of all

sequenced cases, the majority of whom (67%) were under 45 years. However, for older ages, we

found substantially higher B.1.526-related IFRs (e.g. >60% higher for those above 65 years).

This latter finding appears to be consistent with the report by Annavajhala et al. (*2*) showing

resistance of B.1.526 to therapeutic antibodies. Over the course of the pandemic, SARS-CoV-2

IFR has decreased substantially (about a 3-fold difference between the two pandemic waves),

likely due to improved medical treatments (e.g., therapeutic antibodies), better patient

management, and earlier diagnosis. As older adults are more likely to suffer from severe

COVID-19 and thus receive therapeutic antibodies (*13, 14*), the resistance of B.1.526 may render

these treatments ineffective despite their prior success against other variants, leading to increases

in IFR among older adults. These findings highlight the importance of monitoring the efficacy of

therapeutics against different variants and timely update of treatments. In addition, a better

understanding of factors contributing to the higher IFRs in certain variants is warranted to inform

countermeasures (*15, 16*).

Lastly, our analyses suggest both B.1.526 and B.1.1.7 likely had been spreading in the

population for weeks or months prior to detection by the surveillance system (*2, 3, 17*).

Expanding genomic sequencing programs for SARS-CoV-2 and improving linkage to

epidemiologic data can improve detection of new VOIs/VOCs. Such efforts are underway (e.g.,

in the US) but more efforts and resources are urgently needed globally. In addition, to support

more timely detection and control, targeted screening of key subpopulations (e.g., those

prospectively identified from modeling as having high transmission rates) and viral traits (e.g.,

mutations linked to increased transmissibility and/or immune evasion as done in Annavajhala et

al.(*2*)) is needed, as well as timely sharing of key information globally. The documentation of

new VOIs/VOCs anywhere in the world then should prompt preparedness measures to detect and

rapidly respond to the introduction of those variants into local areas. More fundamentally, to

limit emergence of new VOIs/VOCs and end the COVID-19 pandemic, all populations

worldwide should have timely access to vaccination, and multiple layers of mitigation efforts are

needed until a sufficient portion of the population is protected by vaccination.

Our study also has several limitations. First, most of our analyses are based on population-level

data without variant-specific information, given limited variant testing during most of the study

period. We circumvented this data deficiency by analyzing estimates of a key subpopulation
(e.g., the WHts neighborhood where B.1.526 was initially detected) and leveraging prior
knowledge (e.g., estimated IFR prior to B.1.526 emergence). Second, our study did not
distinguish the two subclades within the B.1.526 lineage – one containing the E484K mutation
and the other containing the S477N mutation. Both the E484K and S477N mutations have been
shown to mediate immune escape (*16, 18-20*); in addition, the percentages of these two
subclades were similar during our study period, suggesting they likely have similar
epidemiological characteristics. Third, while it is likely that the emergence of B.1.526 led to the
increase in transmission rate in WHts at the time, we cannot rule out the possibility that other
factors contributed to this increase and in turn the emergence of B.1.526. Lastly, there is a likely
larger uncertainty in B.1.526-related and B.1.1.7-related IFR estimates for younger ages (those
under 45), due to the smaller number of deaths and larger uncertainty in baseline IFR estimates.
Future investigation addressing these issues is warranted should a large sample of variant-
specific data become available.
210
211   In summary, our study has reconstructed the early epidemic trajectory and subsequent rise of
212   B.1.526 in NYC and estimated its key epidemiological properties. Findings highlight the
213   importance of monitoring the viral diversity of SARS-CoV-2, epidemiological characteristics of
214   new variants, and disease severity, as COVID-19 remains a global public health threat.
215

MATERIALS AND METHODS

Study design and data

This study included three interconnected modeling analyses, synthesizing nine epidemiological
and population datasets (Fig 1). The first analysis applied a network model-inference system to
reconstruct underlying SARS-CoV-2 transmission dynamics in NYC, accounting for under-
detection of infection; it also enabled estimation of key population variables and parameters
(e.g., the infection rate including those not detected as cases and transmission rate). The second
analysis applied a city-level multi-variant, age-structured model to simulate and estimate the
changes in transmissibility and immune escape potential for B.1.526 based on the network
model-inference estimates and additional data (e.g., variant prevalence data). The last analysis
utilized estimates from the first two model systems to estimate variant-specific infection fatality
risk (IFR, i.e. the fraction of all persons with SARS-CoV-2 infection who died from the disease),
for B.1.526 and B.1.1.7, separately.
229

For the network model-inference system, we utilized multiple sources of epidemiological data,
including confirmed and probable COVID-19 cases, emergency department (ED) visits, and
deaths, as well as vaccination data. As done previously (*6*), we aggregated all COVID-19
confirmed and probable cases (*21, 22*) and deaths (*22*) reported to the NYC Department of
Health and Mental Hygiene (DOHMH) by age group (<1, 1-4, 5-14, 15-24, 25-44, 45-64, 65-74,

and 75+ year-olds), neighborhood of residence (42 United Hospital Fund neighborhoods in
NYC(*23*)) and week of occurrence (i.e., week of diagnosis for cases or week of death for
decedents). COVID-19-related ED visit data were obtained from the NYC syndromic
surveillance system, comprised of all 53 hospital EDs in the city (*24*). This system identified
individuals presenting at the EDs with COVID-like-illness (CLI; defined as having a fever and
cough or sore throat or respiratory illness, or pneumonia, or a COVID-19 discharge diagnosis
code, excluding those with a discharge diagnosis code of influenza only); in addition, CLI
patients were matched to electronic laboratory reports of SARS-CoV-2 tests with diagnosis
date within ±7 days of ED visit. We estimated the number of COVID-19-related ED visits as the
number classified as CLI multiplied by the percentage of those who tested positive for SARS-
CoV-2 RNA, stratified by the same age and neighborhood groups in weekly intervals. To
account for the impact of vaccination, we also included COVID-19 vaccination data (partially
and fully vaccinated, separately), aggregated to the same age/neighborhood strata by week.
248
249 In addition, as in our previous study (*6*), we used mobility data from SafeGraph (*25*) to model
250 changes in SARS-CoV-2 transmission rate due to non-pharmaceutical interventions. These data
251 were aggregated to the neighborhood level by week without age stratification.
252
For the multi-variant model analysis, we additionally utilized four city-level, weekly datasets: 1)
COVID-19 confirmed and probable cases, 2) hospitalizations (*26*), 3) deaths, and 4) the
percentage of different variants in NYC based on genomic sequencing of samples submitted to
the NYC DOHMH Public Health Laboratory and Pandemic Response Laboratory (*4, 27*). The
additional hospitalization and variant percentage data were published by the NYC DOHMH (*26,
27*) and accessed on June 22, 2021. We used the variant data from the week starting Jan 31, 2021
to the week starting April 25, 2021 in this analysis, because earlier weeks had very low sample
sizes (<200 samples sequenced per week).
261
262 This study was classified as public health surveillance and exempt from ethical review and
263 informed consent by the Institutional Review Boards of both Columbia University and NYC
264 DOHMH.
265
Network model-inference system
The network model-inference system used here is similar to the approach described in Yang et
al.(*6*); however, here we further accounted for waning immunity and vaccination and
additionally used COVID-19-related ED visit data for model optimization. Briefly, the model-
inference system uses an epidemic model (Eqn 1) to simulate the transmission of SARS-CoV-2
by age group and neighborhood, under implemented public health interventions and mass-
vaccination when vaccines became available starting Dec 14, 2020:
273

$$\begin{cases} \dfrac{dS_i}{dt} = \dfrac{R_i}{L} - (S_i \sum_{j=1}^{j=42} \dfrac{b_s b_j \beta_{city} m_{ij} I_j}{N_j}) - v_{i,1} - v_{i,2} \\[2mm] \dfrac{dE_i}{dt} = (S_i \sum_{j=1}^{j=42} \dfrac{b_s b_j \beta_{city} m_{ij} I_j}{N_j}) - \dfrac{E_i}{Z} \\[2mm] \dfrac{dI_i}{dt} = \dfrac{E_i}{Z} - \dfrac{I_i}{D} \\[2mm] \dfrac{dR_i}{dt} = \dfrac{I_i}{D} - \dfrac{R_i}{L} + v_{i,1} + v_{i,2} \end{cases} \qquad [1]$$

where S_i, E_i, I_i, R_i, and N_i are the number of susceptible, exposed (but not yet infectious),

infectious, and removed (either recovered, immune, or deceased) individuals and the total

population, respectively, from a given age group in neighborhood-i. β_{city} is the average citywide

transmission rate; b_s is the estimated seasonal trend (*6*). The term b_i represents the neighborhood-

level transmission rate relative to the city average. The term m_{ij} represents the changes in contact

rate in each neighborhood (for $i=j$) or spatial transmission from neighborhood-j to i (for $i{\neq}j$) and

was computed based on the mobility data (*6*). *Z, D, and L* are the latency period, infectious

period, and immunity period, respectively. The term $v_{i,1}$ represents the number of individuals in

neighborhood-i successfully immunized after the first dose of the vaccine and is computed using

vaccination data and vaccine efficacy (VE) for 1st dose; $v_{i,2}$ is the additional number of

individuals successfully immunized after the second vaccine dose (excluding those successfully

immunized after the first dose). Because 97% of vaccine doses administered in NYC during our

study period (through April 30, 2021) were the Pfizer-BioNTech or Moderna vaccines, we

assumed a VE of 85% fourteen days after the first dose and 95% seven days after the second

dose based on clinical trials and real-word data (*28-30*).

Using the model-simulated number of infections occurring each day, we further computed the

number of cases, ED visits, and deaths each week to match with the observations (*6*). Similar to

the procedure for cases and deaths described in Yang et al.(*6*), to compute the number of ED

visits, we multiplied the model-simulated number of new infections per day by the ED-

consultation rate (i.e. the fraction of model-simulated persons with new SARS-CoV-2 infections

presenting at the EDs), and further distribute these estimates in time per a distribution of time-

from-infection-to-ED-consultation (Table S1); we then aggregated the daily lagged, simulated

estimates to weekly totals for model inference.

Each week, the system uses the ensemble adjustment Kalman filter (EAKF)(*31*) to compute the

posterior estimates of model state variables and parameters based on the model (prior) estimates

and observed case, ED visit, and mortality data per Bayes' rule (*6*). In particular, using this

model-inference, we estimated the citywide transmission rate (β_{city}), neighborhood relative

transmission rate (b_i), and IFR by age group for each week, from the week starting March 1,

2020 (i.e. the beginning of the COVID-19 pandemic in NYC) to the week starting April 25,
2021.
307
308 **Multi-variant, age-structured model**
309 Due to model complexity, the model-inference system described above does not account for the
310 circulation of different variants. To model variants, we used a city-level multi-variant, age-
311 structured model (*32*), per Eqn 2:
312

$$
\begin{cases}
\dfrac{dS_i^A}{dt} = \dfrac{R_i^A}{L_i^A} - \sum_j b_s m c_{ij} \sum_a \dfrac{\beta_j^{Aa} S_j^A I_j^a}{N^a} - \varepsilon_i - v_{i,1}^A - v_{i,2}^A \\[2mm]
\dfrac{dE_i^A}{dt} = b_s m \sum_a \dfrac{\beta_i^{Aa} S_i^A I_i^a}{N^a} - \dfrac{E_i^A}{Z_i^A} + \varepsilon_i \\[2mm]
\dfrac{dI_i^A}{dt} = \dfrac{E_i^A}{Z_i^A} - \dfrac{I_i^A}{D_i^A} \\[2mm]
\dfrac{dR_i^A}{dt} = \dfrac{I_i^A}{D_i^A} - \dfrac{R_i^A}{L_i^A} + v_{i,1}^A + v_{i,2}^A
\end{cases}
\qquad [2]
$$

313
Model variables and parameters in Eqn 2 are similar to those in Eqn 1 with the same symbols.
For instance, β_i is the transmission rate for variant-*i*. However, instead of modeling the spatial
structure, Eqn 2 focuses on the interactions among different variants (indicated by the subscript,
i) and age structure (indicated by the superscript, a or A). For age structure, infection in age
group *A* by variant-*i* comes from all age groups, per the summation $b_s m \sum_a \frac{\beta_i^{Aa} S_i^A I_i^a}{N^a}$ (see the 2nd
line in Eqn 2). For variant interactions via cross-immunity, we use a status-based construct
similar to Yang et al. (*33*) and Gog and Grenfell (*34*). Specifically, c_{ij} measures the strength of
cross-immunity to variant-*i* conferred by infection of variant-*j* (e.g., close to 0 if it is weak and
c_{ii}=1 for infection by the same variant). To compute the depletion of susceptibility to variant-*i*
due to infection of variant-*j* ($i{\neq}j$), we multiply that infection by c_{ij}; that is, non-specific immunity
is scaled by the strength of cross-immunity c_{ij}. As such, the double summation
$\sum_j b_s m c_{ij} \sum_a \frac{\beta_j^{Aa} S_j^A I_j^a}{N^a}$ in Line 1 of Eqn 2 represents the depletion of susceptibility due to variant-
specific infection (when *i*=*j*) and non-specific infections (for all *i*≠*j*). The vaccination model
component $v_{i,1}^A$ and $v_{i,2}^A$ are also variant-specific and can additionally account for the reduction in
VE against the new variants if needed; however, here we used the same VE estimates for all
variants included (i.e., B.1.526, B.1.1.7, B.1.427 and B.1.429) based on observations (*4, 30, 35,*
36). Additionally, the term ε_i represents travel-related importation of infections of variant-*i* (see
Table S2).
332
We restricted this simulation to Nov 2020 – Apr 2021 (i.e., from the initial identification of
B.1.526 to before the further detection and increase of other variants such as Gamma and Delta).
In addition to preexisting variants of SARS-CoV-2 prior to Nov 2020 (i.e. "non-VOC/VOI
variants" for simplicity) and B.1.526, the model included B.1.1.7 and B.1.427/B.1.429
(combined for simplicity), based on available genomic surveillance data showing consistent
detection of these variants during the simulation period and very low levels for others if detected.

Model parameters for B.1.1.7 and B.1.427/B.1.429 were listed in Table S2. For simplicity, we
did not account for other VOCs/VOIs variants because their percentages were very low during
either analysis period (*27*).
342
Initial analysis based on the model-inference estimates suggested B.1.526 was around 20% more
infectious than non-VOC/VOI variants, without accounting for changes in immunity due to
potential immune escape (see details in Results). Therefore, in this analysis, we tested
combinations of change in transmissibility ranging from 10 – 30% increases and immune escape
ranging from 0 – 30%, both with a 5% increment and ±5% intervals (35 combinations in total).
For instance, for the combination centering at 10% transmissibility increase and 0% immune
escape, the model is initialized using values in the range of 5-15% (i.e., $10 \pm 5\%$) transmissibility
increase and 0-5% (i.e., $0 \pm 5\%$ and setting negatives to 0) immune escape. In addition, due to
uncertainty on the initial prevalence, we tested three different levels of initial seeding for the
week starting Nov 1, 2020, i.e., low (0.5 – 2.5%), high (1.5 – 3.5%), and wider range (0.5 –
3.5%). For reference, Washington Heights – Inwood (WHts), which is the neighborhood where
the first patients identified with B.1.526 resided and sought care, constituted 3.2% of the NYC
population in 2018. We initialized the model using the model-inference estimates (e.g.,
population susceptibility and transmission rates by age group; Table S2) and ran the model for
each parameter combination 10 times, each with 1000 realizations to account for model
stochasticity. Results are summarized from the 10,000 model realizations.
359
To identify the most plausible combination of transmissibility and immune escape properties for
B.1.526, we compared the model-estimated weekly number of cases, hospitalization, and deaths
as well as the percentage of the variants to available data. Evaluation was made based on 1)
accuracy, i.e., if the observation falls within the model-estimated interquartile range, it is deemed
accurate; 2) relative root-mean-square-error (RMSE) between the observed and the model-
estimated; and 3) Pearson correlation between the two time-series. Because results show that
model accuracy and relative RMSE had a wider spread among the combinations tested (i.e.,
more distinctive), we first subset those having accuracy within the highest 25th percentile and
relative RMSE within the lowest 25th percentile (2-4 out of 35 combinations remained for each
setting of initial prevalence); we then selected the one with the highest correlation in the subset
as the best-performing and most plausible combination.
371

Estimating the changes in IFR due to B.1.526
The network model-inference system enables estimation of the IFR by age group over time.
These estimates are made combining all variants and do not distinguish by variant. However, we
reasoned that the combined IFR is a weighted average of individual, variant-specific estimates
given the relative prevalence of each variant. Accordingly, we built two linear regression models
to estimate the variant-specific IFR. Model 1 restricted the analysis to Nov 2020 – Jan 2021

(when the relative prevalence of B.1.1.7 in NYC was likely <10%; n = 14 weeks) and only
included two categories of variants:

$$IFR_{combined} \sim IFR_{B.1.526} i_{B.1.526} + IFR_{baseline} i_{others} \qquad [3]$$

381   Model 2 extended the analysis to Nov 2020 – Apr 2021 (n = 26 weeks) and included both
382   B.1.526 and B.1.1.7, in addition to other variants:

$$IFR_{combined} \sim IFR_{B.1.526} i_{B.1.526} + IFR_{B.1.1.7} i_{B.1.1.7} + IFR_{baseline} i_{others} \qquad [4]$$

In both models, $IFR_{combined}$ is the overall IFR for each week, estimated using the model-inference
system; $i_{B.1.526}$, $i_{B.1.1.7,}$ i_{others} are the percentage of infection by the corresponding variant for each
week, estimated using the multi-variant age-structured model with the most plausible parameter
combination as data are not available. $IFR_{baseline}$ is the baseline IFR for the preexisting variants,
set to the average of model-inference estimates over the period of Oct – Nov 2020 (i.e., prior to
the increase of the new variants). The variant-specific IFRs, $IFR_{B.1.526}$ and $IFR_{B.1.1.7}$, are then
estimated using the regression models (n = 14 weekly data points for Model 1; and n = 26
weekly data points for Model 2). For either model, the change in IFR due to a given variant is
then computed as:

$$\Delta IFR = \frac{IFR_{variant} - IFR_{baseline}}{IFR_{baseline}} \times 100\% \qquad [5]$$

394   Both model analyses were performed for each age group or all ages combined, separately; we
395   also combined all those aged under 25 as the IFRs were similarly low for the four sub-age groups
396   (i.e. <1, 1-4, 5-14, and 15-24 year-olds).

Supplementary Materials

Table S1. Prior ranges for the network model-inference system. The prior ranges are similar to
Table S1 of Yang et al.(*6*) but include additional parameters in Eqn 1. The spatial, temporal, and
age resolution of each parameter or variable, estimated in the model-inference system, is
specified in the column "Resolution". Note posterior parameter estimates can extend outside the
specified prior ranges.

Table S2. Initial conditions used to simulate co-circulation of different variants in the multi-
variant, age-structured model. To partially account for changing infection-detection rate, ED-
consultation rate (EDR) and IFR, for these three parameters, we used the model-inference
estimates averaged over the entire simulation period (i.e. Nov 2020 – April 2021). For the initial
transmission rate (for the preexisting non-VOC/VOI variants), we used the model-inference
estimates averaged over the week of 10/25/2020 – the week of 11/7/2020 (i.e. the 3 weeks
around the start of simulation). For the rest of model state variables and parameters, we used
model-inference estimates made at the week of 10/25/2020. For B.1.1.7, we used the following

ranges based on estimates from Yang and Shaman (*32*): 40.3 – 52.3% higher transmissibility (related to estimates for the preexisting non-VOC/VOI variants listed below) and 0 – 10% immune escape; for comparison, contact tracing data from the UK showed that B.1.1.7 was 30-50% more infectious (*37*). For B.1.427/ B.1.429, we used the following ranges based on estimates from Deng et al.(*38*): 16 – 24% higher transmissibility and 0-10% immune escape (vs. 21.4 – 27.8% increase in transmission rate in Deng et al.(*38*) without accounting for changes in immunity due to potential immune escape).

Table S3. Estimated IFR for different variants and changes compared to the baseline risk estimated for preexisting variants during Oct – Dec 2020, using Eqn 3.

Fig S1. Estimated infection-detection rate by age group. Red lines show the estimated median infection-detection rate with surrounding areas indicating the 50% (darker color) and 95% (lighter color) CrI. For comparison, the grey bars show the number of cases reported for each week from the week of Oct 4, 2020 to Apr 25, 2021. Labels of x-axis show the week starts (mm/dd/yy).

Fig S2. Model-fit by age group. Boxes show model estimates (thick horizontal lines and box edges show the median, 25[th], and 75[th] percentiles; vertical lines extending from each box show 95% Crl) and red dots show corresponding.

Fig S3. Estimated cumulative infection rates by age group. Thick horizontal lines and box edges show the median, 25[th], and 75[th] percentiles; vertical lines extending from each box show 95% Crl.

References and Notes

1. World Health Organization, Tracking SARS-CoV-2 variants, 2021, https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/
2. M. K. Annavajhala *et al.*, A Novel and Expanding SARS-CoV-2 Variant, B.1.526, Identified in New York. *medRxiv*, 2021.2002.2023.21252259 (2021).
3. A. P. West *et al.*, Detection and characterization of the SARS-CoV-2 lineage B.1.526 in New York. *bioRxiv*, 2021.2002.2014.431043 (2021).
4. C. N. Thompson *et al.*, Rapid Emergence and Epidemiologic Characteristics of the SARS-CoV-2 B.1.526 Variant - New York City, New York, January 1-April 5, 2021. *MMWR. Morbidity and mortality weekly report* **70**, 712-716 (2021).
5. Global Initiative on Sharing All Influenza Data (GISAID), GISAID, 2021, https://www.gisaid.org
6. W. Yang *et al.*, Estimating the infection-fatality risk of SARS-CoV-2 in New York City during the spring 2020 pandemic wave: a model-based analysis. *The Lancet. Infectious diseases* **21**, 203-212 (2021).

453   7.   N. G. Davies *et al.*, Increased mortality in community-tested cases of SARS-CoV-2
454   lineage B.1.1.7. *Nature* **593**, 270-274 (2021).
455   8.   Centers for Disease Control and Prevention, SARS-CoV-2 Variant Classifications and
456   Definitions, 2021, https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html
457   9.   New and Emerging Respiratory Virus Threats Advisory Group, "Minutes of the
458   NERVTAG COVID-19 FortiethMeeting: 11 December 2020," (2020).
459   10.   New and Emerging Respiratory Virus Threats Advisory Group, "Minutes of the
460   extraordinary meeting of NERVTAG COVID-19 and SPI-M on SARS-CoV-2 variants:
461   21 December 2020," (2020).
462   11.   New and Emerging Respiratory Virus Threats Advisory Group, "NERVTAG/SPI-M
463   Extraordinary meetingon SARS-CoV-2 variant of concern 202012/01 (variant B.1.1.7),"
464   (2020).
465   12.   E. Volz *et al.*, Transmission of SARS-CoV-2 Lineage B.1.1.7 in England: Insights from
466   linking epidemiological and genetic data. *medRxiv*, 2020.2012.2030.20249034 (2021).
467   13.   National Institutes of Health, COVID-19 Treatment Guildlines: Anti-SARS-CoV-2
468   Monoclonal Antibodies, 2021,
469   https://www.covid19treatmentguidelines.nih.gov/therapies/anti-sars-cov-2-antibody-
470   products/anti-sars-cov-2-monoclonal-antibodies/
471   14.   Centers for Disease Control and Prevention, COVID-19: Treatment considerations for
472   healthcare providers, 2021, https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-
473   info.html
474   15.   T. N. Starr *et al.*, Prospective mapping of viral mutations that escape antibodies used to
475   treat COVID-19. *Science* **371**, 850-+ (2021).
476   16.   A. J. Greaney *et al.*, Complete Mapping of Mutations to the SARS-CoV-2 Spike
477   Receptor-Binding Domain that Escape Antibody Recognition. *Cell Host & Microbe* **29**,
478   44-57.e49 (2021).
479   17.   N. L. Washington *et al.*, Emergence and rapid transmission of SARS-CoV-2 B.1.1.7 in
480   the United States. *Cell* **184**, 2587-+ (2021).
481   18.   A. J. Greaney *et al.*, Comprehensive mapping of mutations in the SARS-CoV-2 receptor-
482   binding domain that affect recognition by polyclonal human plasma antibodies. *Cell Host
483   & Microbe* **29**, 463-476.e466 (2021).
484   19.   Z. Liu *et al.*, Identification of SARS-CoV-2 spike mutations that attenuate monoclonal
485   and serum antibody neutralization. *Cell host & microbe* **29**, 477-488.e474 (2021).
486   20.   W. T. Harvey *et al.*, SARS-CoV-2 variants, spike mutations and immune escape. *Nature
487   Reviews Microbiology* **19**, 409-424 (2021).
488   21.   Centers for Disease Control and Prevention, National Notifiable Diseases Surveillance
489   System (NNDSS) - Coronavirus Disease 2019 (COVID-19), 2021,
490   https://ndc.services.cdc.gov/conditions/coronavirus-disease-2019-covid-19/
491   22.   New York City Department of Health and Mental Hygiene, Defining confirmed and
492   probable cases and deaths, 2020, https://www1.nyc.gov/site/doh/covid/covid-19-
493   data.page
494   23.   NewYork City Department of Health and Mental Hygiene, NYC UHF 42
495   Neighborhoods, http://a816-dohbesp.nyc.gov/IndicatorPublic/EPHTPDF/uhf42.pdf
496   24.   R. Lall *et al.*, Advancing the Use of Emergency Department Syndromic Surveillance
497   Data, New York City, 2012-2016. *Public Health Rep* **132**, 23s-30s (2017).

498   25.   SafeGraph, Weekly Patterns: Foot Traffic Data To Understand The COVID-19
499   Pandemic, 2020, https://www.safegraph.com/weekly-foot-traffic-patterns
500   26.   New York City Department of Health and Mental Hygiene, NYC Coronavirus Disease
501   2019 (COVID-19) Data, https://github.com/nychealth/coronavirus-data
502   27.   New york City Department of Health and Mental Hygiene, Variants, 2021,
503   https://github.com/nychealth/coronavirus-data/tree/master/variants
504   28.   F. P. Polack *et al.*, Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *New*
505   *Engl J Med*, (2020).
506   29.   L. R. Baden *et al.*, Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N*
507   *Engl J Med* **384**, 403-416 (2021).
508   30.   E. J. Haas *et al.*, Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-
509   CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a
510   nationwide vaccination campaign in Israel: an observational study using national
511   surveillance data. *The Lancet* **397**, 1819-1829 (2021).
512   31.   J. L. Anderson, An ensemble adjustment Kalman filter for data assimilation. *Mon.*
513   *Weather Rev.* **129**, 2884-2903 (2001).
514   32.   W. Yang, J. Shaman, Development of a model-inference system for estimating
515   epidemiological characteristics of SARS-CoV-2 variants of concern. *Nature*
516   *Communications* **12**, 5573 (2021).
517   33.   W. Yang, E. H. Y. Lau, B. J. Cowling, Dynamic interactions of influenza viruses in Hong
518   Kong during 1998-2018. *PLoS Comput Biol* **16**, e1007989 (2020).
519   34.   J. R. Gog, B. T. Grenfell, Dynamics and selection of many-strain pathogens. *Proc Natl*
520   *Acad Sci U S A* **99**, 17209-17214 (2002).
521   35.   L. J. Abu-Raddad, H. Chemaitelly, A. A. Butt, C.-V. National Study Group for,
522   Effectiveness of the BNT162b2 Covid-19 Vaccine against the B.1.1.7 and B.1.351
523   Variants. *N Engl J Med* **385**, 187-189 (2021).
524   36.   W. F. Garcia-Beltran *et al.*, Multiple SARS-CoV-2 variants escape neutralization by
525   vaccine-induced humoral immunity. *Cell* **184**, 2372-2383 e2379 (2021).
526   37.   Public Health England, "Investigation of novel SARS-CoV-2 variant, Variant of Concern
527   202012/01, Technical briefing 3," (2020).
528   38.   X. Deng *et al.*, Transmission, infectivity, and antibody neutralization of an emerging
529   SARS-CoV-2 variant in California carrying a L452R spike protein mutation. *medRxiv*,
530   2021.2003.2007.21252647 (2021).
531   39.   W. Yang, wan-yang/covid_voc_study: covid_voc_study_yang_shaman_2021_NatComm
532   (v1.0). Zenodo., 2021, https://doi.org/10.5281/zenodo.5715611
533   40.   New York City Department of Health and Mental Hygiene.
534   41.   J. Mossong *et al.*, Social contacts and mixing patterns relevant to the spread of infectious
535   diseases. *PLoS Med* **5**, e74 (2008).
536   42.   Q. Li *et al.*, Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–
537   Infected Pneumonia. *New Engl J Med*, (2020).
538   43.   J. T. Wu, K. Leung, G. M. Leung, Nowcasting and forecasting the potential domestic and
539   international spread of the 2019-nCoV outbreak originating in Wuhan, China: a
540   modelling study. *Lancet* **395**, 689-697 (2020).
541   44.   R. Li *et al.*, Substantial undocumented infection facilitates the rapid dissemination of
542   novel coronavirus (SARS-CoV-2). *Science* **368**, 489-493 (2020).

543   45.   J. Zhang *et al.*, Evolving epidemiology and transmission dynamics of coronavirus disease
544   2019 outside Hubei province, China: a descriptive and modelling study. *The Lancet.*
545   *Infectious diseases* **20**, 793-802 (2020).
546   46.   W. Yang, J. Shaman, A simple modification for improving inference of non-linear
547   dynamical systems. *arXiv*, 1403.6804 (2014).
548   47.   R. Verity *et al.*, Estimates of the severity of coronavirus disease 2019: a model-based
549   analysis. *The Lancet. Infectious diseases* **20**, 669-677 (2020).

Acknowledgments: We thank Columbia University Mailman School of Public Health for high performance computing, Safe Graph (safegraph.com) for providing the mobility data, and Sasikiran Kandula at Columbia University for compiling the mobility data used in this study. We thank the NYC DOHMH Incident Command System Surveillance and Epidemiology Section for processing, cleaning, and managing COVID-19 surveillance data, the NYC DOHMH Public Health Laboratory and Pandemic Response Laboratory for generating and analyzing sequence data, and Iris Cheng, Mohammed Almashhadani, Charles Ko, and Jaimie Shaff from the NYC DOHMH for providing the vaccination data and helpful suggestions on the manuscript.

Funding: This study was supported by the National Institute of Allergy and Infectious Diseases (AI145883) and the NYC DOHMH.

Author contributions: WY designed the study, conducted the analysis, and wrote the first draft; SKG contributed to study coordination and specification of COVID-19 case data, and provided input on parameter estimation; ERP led aggregation and provision of COVID-19 case data; LG contributed to management of COVID-19 case data; WL provided the COVID-19-associated mortality data; RM and RL oversaw the collection of and provided the COVID-19 ED data; SH and JW provided input on SARS-CoV-2 variants and interpretation of the NYC variant percentage data; AF oversaw data collection and management processes at DOHMH. All authors contributed to the final draft.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: The COVID-19 case and mortality data were used with permission under a Data Use and Nondisclosure Agreement between the New York City Department of Health and Mental Hygiene and Columbia University. The New York City Department of Health and Mental Hygiene also has a comprehensive, publicly available data website here: https://github.com/nychealth/coronavirus-data. Additional data sources are detailed in the manuscript. Model code for the multi-variant, age-structured model and a simpler

model-inference system using the ensemble adjustment Kalman filter (EAKF; see ref (*32*)) is available on Zenodo (*39*).

Figures

Fig 1. Study design. This study included three modeling analyses: 1) spatial network model-inference to construct the transmission dynamics and estimate key population variables and parameters by United Hospital Fund (UHF) neighborhood of residence and age group; 2) city-level multi-variant, age-structured modeling to simulate and estimate the changes in transmissibility and immune escape potential for B.1.526; and 3) linear regression models to estimate variant-specific infection fatality risk (IFR), for B.1.526 and B.1.1.7, separately. Nine datasets (listed in the black open boxes) were used as model inputs or to evaluate the accuracy of model estimates (indicated for each dataset below). Models used are shown in the blue filled boxes and model outputs are listed in the blue open boxes (key estimates reported in detail in the Results are bolded). Connections among the analyses are indicated by the arrows and associated annotations.

Fig 2. Model fit and key estimates. Upper panel shows model-fit to weekly number of cases (A), ED visits (B), and deaths (C), for all ages combined. Lower panel shows key model-inference estimates of weekly number of infections including those not detected as cases (D), cumulative number of infections in NYC overall (E), and cumulative infection rate by neighborhood (F). Boxes show model estimates (thick horizontal lines and box edges show the median, 25^{th}, and 75^{th} percentiles; vertical lines extending from each box show 95% CrI) and red dots show corresponding. For the weekly estimates, week starts (mm/dd/yy) are shown in the x-axis labels. Star (*) in the map indicates the location of the Washington Heights – Inwood neighborhood.

Fig 3. Changes in transmission rate. (A) Changes in neighborhood-level relative transmission rate. (B) Changes in citywide transmission rate. Vertical dashed lines indicate the earliest date B.1.526 was identified as reported in Annavajhala et al. Labels of x-axis show the week starts (mm/dd/yy).

Fig 4. Comparison of different combinations of changes in transmissibility and immune escape property for B.1.526. Left panel shows the overall accuracy (A), relative RMSE (B), and correlation (C) of model estimates under different transmissibility and immune escape settings. White crosses (x) indicate the best-performing parameter combination. Right panel shows model estimates using the overall best-performing parameter combination (i.e., 1.5-3.5% initial prevalence, 15-25% higher transmissibility, and 0-10% immune escape). Lines and surrounding areas show model-simulated median estimates and interquartile range; dots show corresponding observations; colors indicate different variants as specified in the legend. Note that these model simulations used same infection-detection rate, hospitalization-rate and IFR (i.e., average during Nov 2020 – Apr 2021); that is, they did not account for changes in case ascertainment or disease

severity by week during this period, due to, e.g., increases in disease severity by the new variants. As such, there were larger deviations from the observations during later months of the simulation with more infections by the new variants.

Fig 5. Estimated infection fatality risk. Red lines show the estimated median IFR with surrounding areas indicating the 50% (darker color) and 95% (lighter color) CrI. For comparison, the grey bars show the number of deaths reported for each week from the week of Oct 4, 2020 to Apr 25, 2021. X-axis labels show the week starts (mm/dd/yy).

Table 1. Estimated IFR for different variants and changes compared to the baseline risk estimated for preexisting variants during Oct – Dec 2020, using Eqn 4.

Table and Figures

Table 1. Estimated IFR for different variants and changes compared to the baseline risk estimated for preexisting variants during Oct – Dec 2020, using Eqn 4.

Age	IFR, baseline (%)	IFR, B.1.526 (%)	IFR, B.1.1.7 (%)	Changes, B.1.526 (%)	Changes, B.1.1.7 (%)	Model goodness-of-fit (R^2)
<25	0.004 (0.0021, 0.0059)	0.004 (0.0039, 0.0041)	0.004 (0.0039, 0.0042)	-0.61 (-3.5, 2.3)	0.97 (-2.8, 4.8)	1
25-44	0.04 (0.021, 0.059)	0.037 (0.034, 0.04)	0.043 (0.039, 0.047)	-6 (-13, 1.4)	8.4 (-1.9, 19)	1
45-64	0.29 (0.15, 0.44)	0.42 (0.31, 0.54)	0.51 (0.35, 0.67)	46 (7.4, 84)	76 (22, 130)	0.97
65-74	1 (0.57, 2.5)	1.9 (1.2, 2.5)	3.3 (2.4, 4.2)	82 (20, 140)	210 (130, 300)	0.96
75+	4.1 (2.2, 6.3)	6.7 (5.9, 7.4)	8 (7, 9)	62 (45, 80)	95 (71, 120)	0.99
all	0.35 (0.2, 0.58)	0.56 (0.48, 0.63)	0.72 (0.61, 0.82)	60 (38, 82)	100 (75, 130)	0.99

Fig 1. Study design. This study included three modeling analyses: 1) spatial network model-inference to construct the transmission dynamics and estimate key population variables and parameters by United Hospital Fund (UHF) neighborhood of residence and age group; 2) city-level multi-variant, age-structured modeling to simulate and estimate the changes in transmissibility and immune escape potential for B.1.526; and 3) linear regression models to estimate variant-specific infection fatality risk (IFR), for B.1.526 and B.1.1.7, separately. Nine datasets (listed in the black open boxes) were used as model inputs or to evaluate the accuracy of model estimates (indicated for each dataset below). Models used are shown in the blue filled boxes and model outputs are listed in the blue open boxes (key estimates reported in detail in the Results are bolded). Connections among the analyses are indicated by the arrows and associated annotations.

Fig 2. Model fit and key estimates. Upper panel shows model-fit to weekly number of cases (A), ED visits (B), and deaths (C), for all ages combined. Lower panel shows key model-inference estimates of weekly number of infections including those not detected as cases (D), cumulative number of infections in NYC overall (E), and cumulative infection rate by neighborhood (F). Boxes show model estimates (thick horizontal lines and box edges show the median, 25[th], and 75[th] percentiles; vertical lines extending from each box show 95% CrI) and red dots show corresponding. For the weekly estimates, week starts (mm/dd/yy) are shown in the x-axis labels. Star (*) in the map indicates the location of the Washington Heights – Inwood neighborhood.

Fig 3. Changes in transmission rate. (A) Changes in neighborhood-level relative transmission rate. (B) Changes in citywide transmission rate. Vertical dashed lines indicate the earliest date B.1.526 was identified as reported in Annavajhala et al. Labels of x-axis show the week starts (mm/dd/yy).

Fig 4. Comparison of different combinations of changes in transmissibility and immune escape property for B.1.526. Left panel shows the overall accuracy (A), relative RMSE (B), and correlation (C) of model estimates under different transmissibility and immune escape settings. White crosses (x) indicate the best-performing parameter combination. Right panel shows model estimates using the overall best-performing parameter combination (i.e., 1.5-3.5% initial prevalence, 15-25% higher transmissibility, and 0-10% immune escape). Lines and surrounding areas show model-simulated median estimates and interquartile range; dots show corresponding observations; colors indicate different variants as specified in the legend. Note that these model simulations used same infection-detection rate, hospitalization-rate and IFR (i.e., average during Nov 2020 – Apr 2021); that is, they did not account for changes in case ascertainment or disease severity by week during this period, due to, e.g., increases in disease severity by the new variants. As such, there were larger deviations from the observations during later months of the simulation with more infections by the new variants.

Fig 5. Estimated infection fatality risk. Red lines show the estimated median IFR with surrounding areas indicating the 50% (darker color) and 95% (lighter color) CrI. For comparison, the grey bars show the number of deaths reported for each week from the week of Oct 4, 2020 to Apr 25, 2021. X-axis labels show the week starts (mm/dd/yy).

Supplemental Tables and Figures

Table S1. Prior ranges for the network model-inference system. The prior ranges are similar to Table S1 of Yang et al.(*6*) but include additional parameters in Eqn 1. The spatial, temporal, and age resolution of each parameter or variable, estimated in the model-inference system, is specified in the column "Resolution". Note posterior parameter estimates can extend outside the specified prior ranges.

Parameter/ variable	Symbol	Resolution	Prior range	Source/rationale
Initial exposed	$E(t=0)$	neighborhood- and age-group specific, estimated for the beginning of the Week of March 1, 2020	300 – 8000 total citywide, scaled by population size for each age group and neighborhood	Large uncertainties, used very wide range
Initial infectious	$I(t=0)$	neighborhood- and age-group specific, estimated for the beginning of the Week of March 1, 2020	150 – 4000 total citywide, scaled by population size for each age group and neighborhood	Assumed to be half the initial exposed
Initial susceptible	$S(t=0)$	neighborhood- and age-group specific, estimated for the beginning of the Week of March 1, 2020	$N - E - I$	Assumed all were susceptible except for those initially exposed/infectious
Population size in each age group and neighborhood	N	neighborhood- and age-group specific	N/A	NYC intercensal population estimates for 2018 (*40*)
Citywide transmission rate	β_{city}	Citywide, age-group specific, estimated for each week	[0.5, 1] per day overall; scaled by contact rate for each age group based on contact data from the POLYMOD study(*41*) (averaged across 8 countries)	Based on R_0 estimates of around 1.5-4 for SARS-CoV-2 (*42-44*)
Scaling of neighborhood	b_i	neighborhood- and age-group specific, estimated for each week	[0.8, 1.2] for age groups under 65 years;	Around 1; larger variation for elderly groups based on data

transmission rate			[0.5, 1.5] for age groups 65 or older	
Latency period	Z	Citywide, age-group specific, estimated for each week	[2, 5] days	Incubation period: 5.2 days (95% CI: 4.1, 7) (*42*); latency period is likely shorter than the incubation period
Infectious period	D	Citywide, age-group specific, estimated for each week	[2, 5] days	Time from symptom onset to hospitalization: 3.8 days (95% CI: 0, 12.0) in China,(*45*) plus 1-2 days viral shedding before symptom onset. We did not distinguish symptomatic/asymptomatic infections.
Immunity period	L	Citywide, age-group specific, estimated for each week	[2.5, 3.5] years	Based on estimated immunity period for endemic human coronaviruses (see Appendix of Yang et al.(*6*))
Multiplicative factor for mobility; see Yang et al.(*6*) for detail	m_1	Citywide, age-group specific, estimated for each week	[1, 2] for <1 year; [0.5, 1.5] for three age groups 1-24 years; [0.1, 1.5] for age group 25-44; [1, 2.5] for age groups 45 or older	Initial model testing showed transmission rates for younger age groups were more sensitive to changes in mobility whereas the two oldest age groups were not sensitive to mobility. For age groups with contact rates lower than the average (based on the POLYMOD study (*41*)), we raised the diagonal elements in the mobility

				matrix to the power of the relative contact rate (<1) to account for insensitivity of transmission rate in these age groups to mobility.
Multiplicative factor for neighborhood connectivity; see Yang et al.(*6*) for detail	m_2	Citywide, age-group specific, estimated for each week	[0.5, 2]	Likely around 1 but with large uncertainties
Mean of time from viral shedding to diagnosis; see Yang et al.(*6*) for detail	T_m	Citywide, age-group specific, estimated for each week	[3, 8] days	From a few days to a week from symptom onset to diagnosis,(*45*) plus 1-2 days of viral shedding (being infectious) before symptom onset
Standard deviation (SD) of time from viral shedding to diagnosis; see Yang et al.(*6*) for detail	T_{sd}	Citywide, age-group specific, estimated for each week	[1, 3] days	To allow variation in time to diagnosis
Infection-detection rate; see Yang et al.(*6*) for detail	r	Citywide, age-group specific, estimated for each week	Starting from [0.001, 0.05] at time 0 and allowed to increase over time using space re-probing(*46*)	Large uncertainties
Infection fatality risk (IFR); see Yang et al.(*6*) for detail		Citywide, age-group specific, estimated for each week	$[5, 15] \times 10^{-5}$ for ages under 25; $[5, 15] \times 10^{-4}$ for ages 25-44; $[5, 15] \times 10^{-3}$ for ages 45-64; [0.01, 0.1] for ages	Based on previous estimates(*47*) but extend to have wider ranges

		65-74; [0.02, 0.2] for ages 75+;	
Time from diagnosis to death; see Yang et al.(*6*) for detail	Citywide	Gamma distribution with mean of 9.36 days and SD of 9.76 days	Based on *n*=15,686 COVID-19 confirmed deaths in NYC as of May 17, 2020.
ED consultation rate (EDR)	Citywide, age-group specific, estimated for each week	[0.001, 0.02] for ages under 25; [0.003, 0.03] for ages 25-44; [0.006, 0.06] for ages 45-64; [0.01, 0.15] for ages 65-74; [0.02, 0.25] for ages 75+;	Based on the ratio of total ED visits and estimated infections during March – Dec 2020
Time-from-infectiousness-to-ED or hospitalization	Citywide, for all ages	Gamma distribution with mean of [5, 7] days and SD of [2, 4] days	

Table S2. Initial conditions used to simulate co-circulation of different variants in the multi-variant, age-structured model. To partially account for changing infection-detection rate, ED-consultation rate (EDR) and IFR, for these three parameters, we used the model-inference estimates averaged over the entire simulation period (i.e. Nov 2020 – April 2021). For the initial transmission rate (for the preexisting non-VOC/VOI variants), we used the model-inference estimates averaged over the week of 10/25/2020 – the week of 11/7/2020 (i.e. the 3 weeks around the start of simulation). For the rest of model state variables and parameters, we used model-inference estimates made at the week of 10/25/2020. For B.1.1.7, we used the following ranges based on estimates from Yang and Shaman (*32*): 40.3 – 52.3% higher transmissibility (related to estimates for the preexisting non-VOC/VOI variants listed below) and 0 – 10% immune escape; for comparison, contact tracing data from the UK showed that B.1.1.7 was 30-50% more infectious.(*37*) For B.1.427/ B.1.429, we used the following ranges based on estimates from Deng et al.(*38*): 16 – 24% higher transmissibility and 0-10% immune escape (vs. 21.4 – 27.8% increase in transmission rate in Deng et al.(*38*) without accounting for changes in immunity due to potential immune escape).

variant	parameter	lower bound	upper bound
B.1.526	Low initial prevalence (%)	0.5	2.5
B.1.526	High initial prevalence (%)	1.5	3.5
B.1.526	Wide initial prevalence (%)	0.5	3.5
B.1.1.7	Increase in transmission rate	0.403	0.5227
B.1.1.7	Immune escape	0	0.1
B.1.427/9	Increase in transmission rate	0.16	0.24
B.1.427/9	Immune escape	0	0.1
non-VOC/VOI	Travel-related importation ε_i	Nominally set to 1 per week for the entire city (N = 8.4 million people)	
B.1.526	Travel-related importation ε_i	Set to 0 as it emerged locally	
B.1.1.7	Travel-related importation ε_i	For the entire city (N = 8.4 million), set to 1 per 2 days for 11/1 – 11/15/20 to reflect lower initial seeding, 1.5 per day for 11/16 -12/31/20 to reflect higher seeding during the holidays, and 2 per day for 1/1 – 4/30/21 to reflect higher seeding due to increases in these variants in the US. Same settings were used for B.1.1.7 and B.1.427/9, because once local transmission is established, travel-related importation plays a nominal role.	
B.1.427/9	Travel-related importation ε_i		
non-VOC/VOI	β_{11} (per day, same below)	0.14	0.21
non-VOC/VOI	β_{22}	0.15	0.2
non-VOC/VOI	β_{33}	0.16	0.22

non-VOC/VOI	β_{44}		0.16	0.23
non-VOC/VOI	β_{55}		0.24	0.36
non-VOC/VOI	β_{66}		0.17	0.25
non-VOC/VOI	β_{77}		0.17	0.22
non-VOC/VOI	β_{88}		0.19	0.25
non-VOC/VOI	β_{12}		0.071	0.1
non-VOC/VOI	β_{13}		0.018	0.027
non-VOC/VOI	β_{14}		0.0074	0.011
non-VOC/VOI	β_{15}		0.023	0.034
non-VOC/VOI	β_{16}		0.011	0.015
non-VOC/VOI	β_{17}		0.0075	0.011
non-VOC/VOI	β_{18}		0.0052	0.0075
non-VOC/VOI	β_{21}		0.074	0.1
non-VOC/VOI	β_{23}		0.019	0.026
non-VOC/VOI	β_{24}		0.0078	0.011
non-VOC/VOI	β_{25}		0.024	0.033
non-VOC/VOI	β_{26}		0.011	0.015
non-VOC/VOI	β_{27}		0.0079	0.011
non-VOC/VOI	β_{28}		0.0054	0.0074
non-VOC/VOI	β_{31}		0.02	0.028
non-VOC/VOI	β_{32}		0.02	0.028
non-VOC/VOI	β_{34}		0.0098	0.014
non-VOC/VOI	β_{35}		0.013	0.018
non-VOC/VOI	β_{36}		0.0072	0.01
non-VOC/VOI	β_{37}		0.0051	0.0071
non-VOC/VOI	β_{38}		0.0066	0.0091
non-VOC/VOI	β_{41}		0.0085	0.013
non-VOC/VOI	β_{42}		0.0085	0.013
non-VOC/VOI	β_{43}		0.013	0.02
non-VOC/VOI	β_{45}		0.014	0.021
non-VOC/VOI	β_{46}		0.01	0.015
non-VOC/VOI	β_{47}		0.0037	0.0054
non-VOC/VOI	β_{48}		0.0069	0.01
non-VOC/VOI	β_{51}		0.11	0.17
non-VOC/VOI	β_{52}		0.11	0.17
non-VOC/VOI	β_{53}		0.092	0.14
non-VOC/VOI	β_{54}		0.073	0.11
non-VOC/VOI	β_{56}		0.07	0.11
non-VOC/VOI	β_{57}		0.042	0.064
non-VOC/VOI	β_{58}		0.041	0.062

non-VOC/VOI	β_{61}	0.053	0.076
non-VOC/VOI	β_{62}	0.053	0.076
non-VOC/VOI	β_{63}	0.043	0.063
non-VOC/VOI	β_{64}	0.051	0.074
non-VOC/VOI	β_{65}	0.053	0.076
non-VOC/VOI	β_{67}	0.058	0.084
non-VOC/VOI	β_{68}	0.052	0.076
non-VOC/VOI	β_{71}	0.032	0.041
non-VOC/VOI	β_{72}	0.032	0.041
non-VOC/VOI	β_{73}	0.022	0.028
non-VOC/VOI	β_{74}	0.0096	0.012
non-VOC/VOI	β_{75}	0.023	0.029
non-VOC/VOI	β_{76}	0.032	0.042
non-VOC/VOI	β_{78}	0.066	0.085
non-VOC/VOI	β_{81}	0.028	0.036
non-VOC/VOI	β_{82}	0.028	0.036
non-VOC/VOI	β_{83}	0.03	0.038
non-VOC/VOI	β_{84}	0.022	0.028
non-VOC/VOI	β_{85}	0.027	0.034
non-VOC/VOI	β_{86}	0.047	0.06
non-VOC/VOI	β_{87}	0.073	0.093
all	Z_1 (days, same below)	2.9	4
all	Z_2	3.3	4.3
all	Z_3	3.4	4.4
all	Z_4	3.4	4.4
all	Z_5	3.5	4.4
all	Z_6	3.5	4.5
all	Z_7	3.3	4.2
all	Z_8	3.3	4.2
all	D_1	2.1	2.9
all	D_2	2.7	3.6
all	D_3	3.2	4.1
all	D_4	3.4	4.4
all	D_5	3.1	4.1
all	D_6	3.2	4.2
all	D_7	2.8	3.7
all	D_8	2.5	3.3
all	IFR_1	3.10E-05	5.00E-05
all	IFR_2	3.00E-05	4.90E-05
all	IFR_3	3.10E-05	4.90E-05

all	IFR$_4$	3.00E-05	4.90E-05
all	IFR$_5$	3.00E-04	0.00048
all	IFR$_6$	0.0029	0.004
all	IFR$_7$	0.013	0.016
all	IFR$_8$	0.046	0.055
all	EDR$_1$	0.0087	0.013
all	EDR$_2$	0.0059	0.0084
all	EDR$_3$	0.0023	0.0033
all	EDR$_4$	0.0046	0.006
all	EDR$_5$	0.01	0.012
all	EDR$_6$	0.019	0.023
all	EDR$_7$	0.031	0.039
all	EDR$_8$	0.057	0.07
all	*Infection detection rate, r_1*	0.14	0.19
all	*Infection detection rate, r_2*	0.21	0.26
all	*Infection detection rate, r_3*	0.23	0.29
all	*Infection detection rate, r_4*	0.28	0.34
all	*Infection detection rate, r_5*	0.39	0.47
all	*Infection detection rate, r_6*	0.36	0.42
all	*Infection detection rate, r_7*	0.33	0.41
all	*Infection detection rate, r_8*	0.34	0.41

Table S3. Estimated IFR for different variants and changes compared to the baseline risk estimated for preexisting variants during Oct – Dec 2020, using Eqn 3.

Age	IFR, baseline (%)	IFR, B.1.526 (%)	Changes, B.1.526 (%)	Model fit, R^2
<25	0.004 (0.0021, 0.0059)	0.004 (0.0038, 0.0041)	-0.03 (-3.8, 3.8)	1
25-44	0.04 (0.021, 0.059)	0.035 (0.031, 0.038)	-12 (-21, -3.1)	0.97
45-64	0.29 (0.15, 0.44)	0.42 (0.24, 0.59)	43 (-17, 100)	0.67
65-74	1 (0.57, 2.5)	1.5 (0.53, 2.5)	46 (-50, 140)	0.46
75+	4.1 (2.2, 6.3)	6.1 (5.1, 7)	47 (25, 69)	0.94
all	0.35 (0.2, 0.58)	0.5 (0.4, 0.61)	43 (13, 73)	0.89

Fig S1. Estimated infection-detection rate by age group. Red lines show the estimated median infection-detection rate with surrounding areas indicating the 50% (darker color) and 95% (lighter color) CrI. For comparison, the grey bars show the number of cases reported for each week from the week of Oct 4, 2020 to Apr 25, 2021. Labels of x-axis show the week starts (mm/dd/yy).

Fig S2. Model-fit by age group. Boxes show model estimates (thick horizontal lines and box edges show the median, 25th, and 75th percentiles; vertical lines extending from each box show 95% CrI) and red dots show corresponding.

Fig S3. Estimated cumulative infection rates by age group. Thick horizontal lines and box edges show the median, 25th, and 75th percentiles; vertical lines extending from each box show 95% Crl.