Distribution of rare *LOXL1* missense alleles, haplotypes, and diplotypes suggests association with reduced risk of glaucoma-related exfoliation syndrome.

Inas F. Aboobakar[1], Rob P. Igo, Jr.[2,3], Tyler G. Kinzy[4,5], Jessica N. Cooke Bailey[4,5], Chiea Chuen Khor[6,7], Tin Aung[7], Robert Ritch[8], Arthur J. Sit[9], Richard K. Lee[10], Margaret Pericak-Vance[11], Jae H. Kang[12], Jonathan L. Haines[2,3], Louis R. Pasquale[8], Janey L. Wiggs[1]

Affiliations

1. Department of Ophthalmology, Harvard Medical School, Massachusetts Eye and Ear, Boston, MA

2. Department of Population and Quantitative Health Sciences, Case Western Reserve University, Cleveland, OH

3. Cleveland Institute for Computational Biology, Case Western Reserve University, Cleveland, OH

4. Department of Pharmacology & Toxicology, Brody School of Medicine, East Carolina University, Greenville, NC, USA

5. Center for Health Disparities, East Carolina University, Greenville, NC, USA

6. Genome Institute of Singapore, Singapore

7. Singapore Eye Research Institute, Singapore

8. Department of Ophthalmology, Icahn School of Medicine at Mount Sinai, New York, New York.

9. Department of Ophthalmology, Mayo Clinic, Rochester, MN

10. Department of Ophthalmology, Bascom Palmer Eye Institute, University of Miami, Miami FL

11. Hussman Institute of Human Genetics, University of Miami, Miami, FL

12. Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Boston, MA.

ABSTRACT/SUMMARY

Purpose: Common *LOXL1* protein-altering variants are significant genetic risk factors for exfoliation syndrome (XFS) and the related secondary glaucoma (XFG). A rare *LOXL1* missense allele was associated with protective effects in a Japanese cohort, suggesting that other rare alleles may also exhibit protective effects. The goal of this study is to assess the contributions of rare *LOXL1* variants to XFS/XFG risk in cases and controls from the United States (US).

Methods: *LOXL1* rare variants (minor allele frequency (MAF) < 1%) were identified from Human exome BeadArray (Illumina) data for 1118 XFS/XFG cases and 3661 controls (Mass Eye and Ear (MEE) cohort) and from exome sequence data for 284 cases and 37,499 controls in *All of Us* (AoU) The distribution of rare variants, haplotypes, and diplotypes was examined using Fisher exact test.

Results: Four rare *LOXL1* missense alleles, more common in controls, were identified in MEE, P= 7.6E-4), and 456 variants identified in AoU were found only in controls (P=0.045). The rare protective alleles were preferentially located (P= 5.8E-45) in a LOXL1 intrinsic disordered region (IDR) potentially involved in LOXL1 aggregation. Haplotypes that included the rare or minor variants were more common in controls compared to cases in both MEE (Odds Ratio (OR)= 0.21 (95% Confidence interval (CI): 0.19-0.24), P=1.7E-173) and AoU (OR=0.28 (95% CI: 0.23-0.34), P=4.4E-41), and heterozygous diplotypes were also significantly associated with reduced risk overall in both MEE (OR= 0.45 (95% CI: 0.52-0.39), P= 1.7E-89) and AoU (OR=0.26 (95% CI: 0.18-0.33, P=7.6E-28). Diplotypes comprised of only homozygous genotypes were associated with increased disease risk in both MEE and AoU (OR= 4.16 (95% CI: 3.60-4.76), P= 4.2E-89 and 5.26 (95% CI: 4.12-6.73), P=1.87E-42, respectively) and excess homozygosity and decreased heterozygosity was correlated with disease risk for common *LOXL1* variants across multi-ethnic populations (P=1.0E-6).

Conclusions: Using exome array and exome sequence data from XFS/XFG cases and controls from the United States, we identified rare protective *LOXL1* missense variants and show that the distribution of the corresponding haplotypes and diplotypes was associated with lower risk of XFS/XFG. Diplotype results also demonstrated that *LOXL1* allelic heterozygosity was inversely associated, while homozygosity was associated with increased disease risk. These results suggest that *LOXL1* MAF variation among populations, with corresponding variation in genotype heterozygosity and homozygosity, determines the XFS/XFG association.

INTRODUCTION

Exfoliation syndrome (XFS) is a systemic disorder characterized by progressive accumulation of abnormal fibrillar protein aggregates that can obstruct drainage of fluid from the eye, raising intraocular pressure (IOP) and causing exfoliation-related secondary glaucoma (XFG). XFG is the most common secondary glaucoma worldwide and is a major cause of irreversible blindness (Nazarali, 2018). XFS is also associated with premature cataract formation and complications during cataract surgery. Systemic conditions have also been associated with XFS, including cardiovascular disease (Chung, 2018), sleep apnea (Shumway, 2021), and obstructive pulmonary disease (Taylor, 2019).

LOXL1 (lysyl oxidase-like 1) is a major genetic risk factor for XFS/XFG in all populations examined (Aung, 2017), with *LOXL1* risk variants occurring in up to 98% of patients (Fan, 2011). Aggregated LOXL1 protein is one component of the exfoliation fibrillar material (Zenkel, 2014; Sharma, 2009) that has been identified in extracellular spaces throughout the body, but in particular in the ocular anterior segment (Schlötzer-Schrehardt, 2009). The LOXL1 protein includes several intrinsic disordered regions (IDRs) that could be sites of initiation of protein aggregates, and recent studies suggest that deletion of the most disordered regions reduces protein aggregation (Bernstein, 2019).

Two of the major *LOXL1* risk variants, (G153D and R141L) are common missense alleles located within the IDR with a high likelihood of aggregation (Bernstein, 2019). (**Figure 1**). Of interest, in most populations studied, the common allele 'G' at 153 and the common 'R' allele at 141 are associated with increased risk; however, in East Asian populations the 'L' allele at 141 is associated with increased risk (Hayashi 2008), and in South Africans, the 'D' allele at 153 is associated with increased risk (Williams 2010). The R141 'L' and G153 'D' are the 'common' alleles in East Asian and South African populations, respectively. Several studies have investigated the role of *LOXL1* genetic variants including effects on

protein function (Sharma 2016), gene expression (Pasutto, 2017), and dysregulation related to a long noncoding RNA also located within the LOXL1 genomic region (Hauser 2015). However, none of these studies have observed a consistent effect for the associated risk allele in all populations, which limits the determination of the underlying disease-causing mechanism(s).

A rare *LOXL1* protein-altering variant has been associated with significantly reduced risk of XFS in Japanese cases and controls (Aung, 2017). This finding, as well as the observation that the less frequent allele at G153D and R141L is protective in all populations studied so far (i.e., the heterozygous genotype is consistently associated with reduced risk), suggests that other rare or low-frequency alleles may also be protective. To investigate this hypothesis, we examined the effects on risk of rare *LOXL1* coding variants in two cohorts from the United States: a cohort of XFS cases and controls with European genetic ancestry from Mass Eye and Ear (MEE), as well as a cohort of XFS cases and controls with European genetic ancestry from the *All of Us* (AoU) Study.

METHODS

Massachusetts Eye and Ear cohort: This study adhered to the tenets of the Declaration of Helsinki and has been reviewed and approved by the Institutional Review Boards of the Massachusetts Eye and Ear, Harvard School of Public Health, and the Brigham and Women's Hospital. Informed consent was obtained from the participants after explanation of the nature and possible consequences of the study. XFS/XFG cases were recruited from MEE (Harvard), Mayo Clinic, Bascom Palmer Eye Institute (University of Miami), the Nurses' Health Study (NHS), and Health Professionals Follow-up Study (HPFS). The controls were previously genotyped samples from the NHS and HPFS, and were over the age of 40, self-reported Caucasians with European ancestry, and located throughout the United States (Kang 2016). Controls have no evidence of XFS by clinical exam or medical record or have repeatedly denied a self-

report of glaucoma on biennial questionnaires administered over two decades. We also accessed

genotype data from the MEE and NEIGHBOR controls (Wiggs, 2012; Margeta 2020).

All XFS cases were of self-reported European ancestry and were over the age of 40 and have

documentation of characteristic ocular exfoliation material at the pupil margin or surface of the ocular

lens, either through clinical exam or medical records. Clinical examination included measurement of

visual acuity and intraocular pressure, slit lamp biomicroscopy, and fundoscopy. Cases also had visual

field assessment primarily using the Humphrey automated visual fields. Individuals were excluded if

other types of glaucoma (pigment dispersion, steroid-induced, uveitis) were evident on exam. All cases

provided blood samples for DNA extraction.

Genotyping and quality control

Cases were genotyped at the Center for Inherited Disease Research (CIDR; http://www.cidr.jhmi.edu/)

using the Illumina OmniExpress+Exome platform that includes 700,000 common SNPs (MAF >0.3) and

250,000 rare or low frequency functional exonic SNPs collected from 12,000 exomes

(http://genome.sph.umich.edu/wiki/Exome_Chip). The XFS/XFG controls were previously genotyped

using the Omni Express+Exome platform. The NEIGHBOR control data set was also genotyped using the

Illumina HumanExome BeadArray (Illumina, Inc., San Diego, CA) at CIDR.

Quality control for human exome array genotype data (Igo 2016) for all cases and controls was carried

out as follows. The Illumina Genome Studio (Illumina) and PLINK (Purcell 2007) were used for all quality

controls (QC) steps except where noted. Basic QC for samples included screens for call rate (≥98.5%) and

high (≥95%) concordance with a previous Illumina 660K panel run on the same sample, where available

(about 80% of samples). We verified recorded sex in the clinical records with genotyped sex by two

criteria: mean fluorescence intensity on the X and Y chromosomes, plus genotype heterozygosity on the X chromosome and call rate on the Y, allowing male and female samples to have heterozygous X-linked and successful Y-linked genotypes, respectively. We tested samples for pairwise relationships and unexpected duplication using KING (Manichaikul, 2010).

We verified European ancestry from the first two principal components derived from genotypes at 9000 ancestry-informative markers by means of the SNP weights program (Chen, 2013), including representative HapMap CEU, YRI, CHB, and JPT samples as reference populations. Moreover, we conducted a principal components analysis over 52,040 independent (pairwise $r^2 < 0.1$), common (MAF ≥0.005) SNPs using the smartpca program in EIGENSOFT (Price, 2006) to detect finer population structure.

Initial QC screens for markers included call rate (≥98%) and consistency with Hardy-Weinberg proportions ($P > 10^{-6}$ by Fisher's exact test). We screened markers for differences in allele frequency between whole-genome-amplified DNA samples and all other samples by the Fisher's exact test and removed from analysis all markers with $P < 0.0001$. All pseudoautosomal, Y-linked, and mitochondrial SNPs were subject to review in Illumina Genome Studio (Illumina, Inc.), and if necessary, were manually re-clustered. We also confirmed genotype clustering for rare (MAF <0.02) variants from fluorescence intensity data by means of zCall, run with a stringent z score threshold of $z = 21$ for calling heterozygous genotypes. Every rare SNP with two or more additional heterozygous calls by zCall than by GenCall was reviewed in Genome Studio, and if necessary, cluster locations were adjusted manually. Data for four rare coding variants (G120S, S159A, A160P, S427F) included on the genotyping platform were extracted for analyses.

All of Us: Informed consent was obtained for all participants enrolled in AllofUs. The AllofUs IRB follows the regulations and guidance of the National Institutes of Health Office for Human Research Protections for all studies, ensuring that the rights and welfare of research participants are overseen and protected uniformly (AllofUs Research Program, 2019; AllofUs Research Program, 2024). Cases were older than 55 years and had an ICD9/10 code for XFS or XFG (366.11, 365.52, H40.14, H40.143, H40.149, H40.141, H40.142, H40.1490, H40.1430, H40.1431, H40.1432, H40.1413, H40.1411, H40.1433, H40.1421, H40.1434, H40.1422, H40.1491, H40.1423, H40.1412, H40.1494, H40.1492, H40.1420, H40.1493, H40.1410, H40.1424, H40.1414). Controls were older than 75 years and had no ICD-9/-10 code for any form of glaucoma, glaucoma suspect, or ocular hypertension. After data quality control (relatedness, sex mismatch) , 371 XFS/XFG cases and 37,522 controls were identified. Since the majority of the cases were of European genetic ancestry, other genetic ancestries were removed, leaving 284 cases and 37,499 controls.

1190 rare variants (Minor allele frequency (MAF) <1%) identified in exome sequence data from the region corresponding to *LOXL1* coding sequence (Chr15, 73926784-73951837) were filtered to remove variants with allele counts of <2 in cases and controls and missingness >2%, leaving 47 variants which were mainly missense alleles and all only present in controls and with 456 controls carrying at least one rare allele.

Other genetic ancestry-defined groups: For replication purposes, data from genetic ancestry-defined groups where the variant of interest was identified in Aung et al. (Aung, 2017), was included.
UK Biobank: While diagnostic codes for XFS or XFG are not included in the UK Biobank clinical showcase, and therefore cannot be used for case or phenotype selection, the individuals with overall glaucoma codes (H40) would include these individuals. Using publicly available data (Genebass), we investigated overall burden results for the glaucoma groups as well as single variant associations.

<u>Multi-ethnic ancestry groups from literature review</u>: To investigate deviations from Hardy-Weinberg equilibrium for the common risk alleles (G153D and R141L) we used study-specific data for 40 independent studies published as a meta-analysis by Li et al., (2021), that included European Caucasians, Asians, and Africans ancestry groups.

Analyses

Distribution of the four rare *LOXL1* variants in MEE cases compared to controls and the 456 AoU variants in cases and controls was assessed using Fisher's exact test. Haplotypes were constructed using IMPUTE2 (Howie 2009), and haplotype distribution between cases and controls was also assessed using Fisher's exact test for both cohorts. Diplotypes were identified using the haplotype data for each study participant. Chi-square and Fisher's exact test were used to assess Hardy Weinberg disequilibrium and variation percent homozygosity and heterozygosity with disease risk for common *LOXL1* risk alleles. To assess Hardy Weinberg equilibrium for the common risk alleles (G153D and R141L), genotype results from individual studies included in a large meta-analysis (Li, 2021) were tallied for each genetic ancestry group (Europeans, Asians and Africans). Using the frequency for the common and minor allele for each variant, the expected Hardy Weinberg distribution was calculated using p2 +2pq + q2 =1. The observed genotype distribution was compared to the expected using the chi-square test.

RESULTS

For the MEE cohort, after sample and genotyping quality control, genotype data was available for 1,118 cases and 3,661 XFS/XFG controls. Additionally, genotype data was available for 2,606 primary open angle glaucoma (POAG) subjects without XFS/XFG from the NEIGHBOR study (Wiggs 2013). Cases and controls have similar proportions of females (**Supplementary Table 1**). The cases are on average older than both control cohorts; however, the overall difference was not significant (P= 0.26). Similar

age and sex distribution was observed for the AoU cases (N=284) and controls (N=37499)

(Supplementary Table 1).

Distribution of rare missense alleles in XFS cases and controls suggest protective effects. In the MEE

cohort we identified four *LOXL1* missense alleles with minor allele frequencies (MAFs) less than 1%:

G120S, S159A, A160P and S427F (**Table 1**). Three of these have Combined Annotation Dependent

Depletion (CADD) (Kircher 2014) scores >20, indicating potential deleterious effects on protein

structure/function. Two missense alleles (S159A and A160P) are located within the *LOXL1* intrinsic

disordered region (IDR) with high propensity to protein aggregation (**Figure 1**). All four missense

variants were observed more often in controls compared to cases (**Table 1**) for both the XFS/XFG

(combined P= 7.6E-4) and NEIGHBOR control datasets (combined P= 2.2E-3). Using data from an

international XFS meta-analysis (Aung 2017), we assessed the allelic distributions for these variants in

other populations. While these rare variants are not universally present, we were able to observe

consistent protective effects in an Icelandic case-control study for A160P and S427F and in a South

African cohort for S159A (**Table 2**). G120S was not associated (P> 0.05) with XFS in any other population

(1 of 458 cases from Italy and 1 of 2827 cases from Japan and not in any controls from either population

(267 and 3013, respectively).

In AoU controls, 47 variants were found in 456 individuals (**Supplementary Table 2**). These are

mainly missense alleles and were found only in controls and not in any of the 284 cases (P= 0.045).

Three of the four *LOXL1* missense alleles identified in the MEE cohort were among the 47 variants (456

controls) identified in AllofUs controls: A160P (213 controls), S159A (19 controls) and G120S (36

controls) (**Supplementary Table 2**).

While XFS/XFG cases and controls could not be identified in the UK biobank, we did assess

contributions of *LOXL1* rare coding variants in the UK Biobank glaucoma groups ('Glaucoma H40',

'Glaucoma Custom', 'Glaucoma', and 'Glaucoma surgery'), (**Supplementary Table 3**) using the publicly available website Genebass (Karczewski, 2022). We examined each glaucoma group for aggregated variant effects (SKAT-O and Gene Burden) in 3 variant groups: loss of function alleles, missense alleles and synonymous alleles. For each phenotype group, nominal evidence of association was only observed for the missense alleles and in each case the effect was protective (**Supplementary Table 3**). Five alleles showed nominal single variant effects (**Supplementary Table 4**), including A160P with protective effects (P=5.29e-3 β= -3.29e-1).

LOXL1 rare haplotypes are more common in controls than in cases. To further explore the protective effects of the rare missense alleles, in the MEE cohort, we investigated the distribution of haplotypes defined by the 4 missense alleles as well as the two common *LOXL1* variants, G153D and R141L. All haplotypes that include a rare variant allele or the minor allele for G153D and R141L are found more often in controls compared to cases (OR = 0.33, P=1.7E-173) (**Table 3**). In addition, only the haplotype that includes the common allele for all 6 missense variants was associated with increased disease risk (OR = 4.76, P= 1.7E-173).

We also investigated haplotypes in the AoU data where only the haplotype homozygous for common 'G' alleles at G153D and R141L, without the addition of any rare alleles, was associated with disease risk (OR 3.22, P=6.2E-43) (Haplotype 1, **Table 5**).

LOXL1 heterozygous diplotypes are more common in controls than in cases. As rare alleles are more likely to be present as heterozygous genotypes, in the MEE cohort, we examined the distribution of the 6-variant diplotypes (derived from the haplotypes in Table 3) among XFS/XFG cases and controls. Similar to the haplotype distribution, the homozygous diplotype comprised of the common alleles (Diplotype A, **Table 4**) is associated with increased disease risk (OR= 6.78, P=4.7E-157). All diplotypes

that include heterozygous genotypes are protective with the exception of diplotype F, which was associated with an OR of 1.64 (**Table 4**) but was also very rare, and the distribution between cases and controls was not statistically significant (P= 0.55). Overall, heterozygous diplotypes are significantly protective (OR= 0.45, P=4.2E-89) (**Table 4**). We also observed similar protective effects for heterozygous diplotypes in the AllofUs data (**Table 6**) and an overall significant protective effect for heterozygous diplotypes (OR= 0.26, P=7.62E-28) (**Table 7**).

Two diplotypes comprised of only homozygous genotypes had higher frequency in controls: the GA|GA diplotype in AllofUs (**Table 6**) and the TG|TG diplotype in Allof Us and corresponding diplotype B in the MEE cohort (**Table 4 and Table 6**). Both of these diplotypes include the homozygous minor allele for one of the two common variants.

Distribution of heterozygous variants with protective effects in LOXL1 disordered domains. We used the disordered domain predictor PONDR (Peng, 2005) to define the LOXL1 disordered regions (**Figure 1**). The variants with protective effects were preferentially located in the LOXL1 disordered region extending from amino acid residue 138 to 166 (**Table 8**) (P= 5.81E-45).

Excess homozygosity and decreased heterozygosity for common variants in XFS/XFG cases.
Considering the consistent protective effects for heterozygous variants, we hypothesized that in case/control cohorts, an excess of homozygosity for the common G153D and R141L variants could be observed in cases (with a decrease in heterozygosity) and an excess of heterozygosity could be evident in controls. To investigate this hypothesis, we used genotype data from 40 independent previously published studies for 3 genetic ancestry groups (European, Asian, and African) summarized in Li et al (Li, 2021), and calculated the expected and observed distribution of genotype groups using Hardy Weinberg equilibrium and population-specific allele frequencies. We found that in all ancestral groups there was

an excess of homozygosity and a decrease in heterozygosity for G153D genotypes in cases (Supplemental Table 5A). A similar trend was observed for the G153D in AllofUs (Europeans) and in the MEE cohort (Supplemental Table 5B). The combined data for G153D in cases (N=5748) showed a significant excess of homozygosity for both the major and minor alleles and a significant decrease in heterozygosity (Table 9). For R141L, similar deviations were observed for the Korean and Japanese cases N=1039 (Table 10) but not for any other genetic ancestry or the AoU or MEE cohorts (Supplementary table 6). In controls, only the data from the MEE cohort showed an excess of heterozygote genotypes for G153D (Supplementary Table 5B).

DISCUSSION

Common *LOXL1* coding variants are major XFS/XFG genetic risk factors, but despite the strong association with disease risk, the underlying pathogenic mechanism(s) are not known. The discovery of a protective rare *LOXL1* variant in a Japanese case-control study provided the first insight that *LOXL1* protein-altering variants could reduce disease risk. In this study, we have expanded this initial observation to include other rare *LOXL1* missense alleles that are all more commonly found in controls compared to cases. In particular, A160P shows protective effects in the MEE cohort, an Icelandic dataset, AoU, and the UK Biobank. We also show that haplotypes and diplotypes that include these rare coding variants are more common in controls.

In case/control studies, deviation from Hardy Weinberg distribution in cases for the common variants G153D and R141L suggests that overall, heterozygosity is protective and homozygosity is associated with increasing disease risk. We did not observe a consistent significant deviation from expected patterns in controls, which could be due to several factors, including undiagnosed cases in

controls. Interestingly, the only control group with excess heterozygosity was the MEE cohort, where controls had eye exams and did not report having a physician-diagnosis of glaucoma.

The risk effect for homozygous variants is supported by the observation that only the diplotype that includes common homozygous variants (diplotype A in the MEE cohort and diplotype GG|GG in AoU) is associated with increased disease risk. However, the homozygous diplotypes that include homozygosity for the minor alleles G153D [D] and R141L [L], (diplotype B in MEE and diplotypes GA|GA and TG|TG in AllofUs), while rare, are found more often in controls, potentially suggesting that homozygosity alone is not sufficient for disease development. It is not clear what factors could lead to the higher frequency of these diplotypes in controls; however, it is intriguing that the homozygous 'minor' allele is involved in both diplotypes demonstrating this effect. It is possible that other, as yet undetected rare variants, are present on these diplotypes that are underlying the protective effects, especially considering that these diplotypes are not common in any population. Interestingly, both homozygous 'D' and homozygous 'L' are more common in combined cases than expected for G153D and in the Korean/Japanese datasets for R141L (**Tables 9, 10**).

Rare variants with protective effects are primarily located in a LOXL1 disordered region that includes the 'common' variants G153D and R141L, and includes a region that has been shown to reduce aggregation when deleted from the protein (Bernstein 2019). This observation, coupled with the protective effects of heterozygosity, suggests that protein variation within this region could reduce protein aggregation. IDRs can contribute to pathogenic aggregation for several neurodegenerative diseases including Alzheimer's, ALS, Parkinson's, Huntington's, and prion-related diseases, among others (Eftekharzadeh 2016, Flores, 2025). Understanding the nature of aggregation induced by IDRs is an area of intense investigation (Flores, 2025). Protein variants may increase or decrease aggregation (Candelise

2021), with specific examples including a missense allele protective for Alzheimer's disease shown to reduce amyloid beta aggregation (Das, 2017) and TDP-43 variants that influence the rate of protein aggregation related to ALS (Jiang, 2016). The strong homozygous and heterozygous effects observed in this study suggest that a matched amino acid sequence within the LOXL1 IDR supports aggregation (increases risk) while a mismatch (heterozygous) reduces aggregation and reduces risk.

In some populations, the risk allele at the common *LOXL1* variants (G153D and R141L) are reversed making it difficult to evoke a specific biological mechanism that explains disease risk. For example, a molecular interaction with the allele associated with risk or allele-specific gene expression influencing risk would have to be population dependent. While possible, it seems unlikely that a gene variant could underlie a very different function in one population compared to another. The results from this study, which showed a strong correlation between homozygosity and disease risk combined with overall protective effects observed for heterozygous diplotypes, supports a novel hypothesis that the observed risk allele reversal in some populations is simply the result of variation in minor allele frequency due to population effects and leading to variation in relative homozygosity and heterozygosity of risk variants. For example, R141 allele 'L' is the minor allele in most populations and is also associated with protective effects in most populations. However, in East Asians, where 'L' is the risk allele (Hayashi 2008), the 'L' allele is the more frequent allele and therefore is more likely to be homozygous and associated with disease risk. A similar argument can be made for the reversed risk observed for the G153D variant in South Africans (Williams 2010). This hypothesis also explains the protective effects of rare *LOXL1* protein-coding variants, as these are likely to be heterozygous due to their low minor allele frequencies. Further work will be required to confirm this hypothesis including testing heterozygous versus homozygous variants for protective functional effects.

There are several limitations of our study. First, the MEE analyses are based on exome array data, which does not comprehensively sample all *LOXL1* variants. The comprehensive sequence data fromAoU does help mitigate this concern; however, the number of cases in AoU is relatively small. Further work using larger numbers of cases with exome data will be necessary to further define the effects of rare coding variants on disease risk. Second, for the datasets available for this study, the number of controls is larger than the number of cases, which may provide more power to detect rare variants in controls. However, the decrease in heterozygosity for the common variants suggests that the higher frequency of heterozygous variants in controls is not entirely due to the larger sample size. Additionally, the significant enrichment of the AoU variants in the IDR argues against random distribution of rare variants in controls. Third, the differences in average ages for cases and controls may influence these results; however, the overall difference in age was not significant, and the distribution of diplotypes among case and control age groups was not significantly different (Supplementary Figure 1). Finally, because only one haplotype including a rare variant did not also include the protective allele for the common variants, we could not examine the effects of rare variants independent of the common risk alleles.

In summary, we identified rare protective *LOXL1* protein coding variants and showed that the distribution of the haplotypes and diplotypes that include these variants suggest an association with reduced risk of XFS/XFG. We also showed that only a homozygous diplotype comprised of all common alleles was significantly associated with disease risk and that excess homozygosity of the common risk variants G153D [G] and R141L [R] was observed in cases across ancestral groups. These results support a novel hypothesis that the reversal of *LOXL1* risk alleles observed in some populations is dependent on variation of minor allele frequencies among populations and that *LOXL1* allelic homozygosity is associated with increased disease risk, potentially due to effects on protein aggregation involving

intrinsic disordered regions. Further work identifying additional protective variants and testing effects

on protein aggregation will be necessary to confirm this hypothesis.

ACKNOWLEDGEMENTS

Supported by NIH/NEI R01 EY020928, EY015473, and an unrestricted challenge grant from Research to Prevent Blindness (NYC to Icahn School of Medicine at Mount Sinai). Dr. Pasquale is supported by The Barry Family Center for Ophthalmic Artificial intelligence and The Glaucoma Foundation in New York City.

REFERENCES

All of Us Research Program, I. et al. The "All of Us" Research Program. N Engl J Med 507 381, 668-676 (2019).

All of Us Research Program Genomics, I. Genomic data in the All of Us Research Program. Nature 627, 340-346 (2024).

Aung T, et al., Genetic association study of exfoliation syndrome identifies a protective rare variant at LOXL1 and five new susceptibility loci Nat Genet. 2017 Jul;49(7):993-1004.

Bernstein AM, Ritch R, Wolosin JM. LOXL1 folding in exfoliation glaucoma. Adv Protein Chem Struct Biol. 2019;118:273-288.

Candelise N, Scaricamazza S, Salvatori I, Ferri F, Valle C, Manganelli V, Garofalo T, Sorice M, Misasi R. Protein Aggregation Landscape in Neurodegenerative Diseases: Clinical Relevance and Future Applications Int J Mol Sci. 2021 Jun 2;22(11):6016.

Chen CY, Pollack S, Hunter DJ, Hirschhorn JN, Kraft P, Price AL. Improved ancestry inference using weights from external reference panels. *Bioinformatics*. 2013; 29: 1399–140.

Chung H, Arora S, Damji KF, Weis E. Association of pseudoexfoliation syndrome with cardiovascular and cerebrovascular disease: a systematic review and meta-analysis. Can J Ophthalmol. 2018;53(4):365-372.

Das P, Chacko AR, Belfort G. Alzheimer's Protective Cross-Interaction between Wild-Type and A2T Variants Alters Aβ 42 Dimer Structure ACS Chem Neurosci. 2017 Mar 15;8(3):606-618.

Eftekharzadeh B, Hyman BT, Wegmann S. Structural studies on the mechanism of protein aggregation in age related neurodegenerative diseases. Mech Ageing Dev. 2016 Jun; 156():1-13.

Fan BJ, Pasquale LR, Rhee D, Li T, Haines JL, Wiggs JL LOXL1 promoter haplotypes are associated with exfoliation syndrome in a U.S. Caucasian population. Invest Ophthalmol Vis Sci. 2011 12;52(5):2372-8.

Flores E, Acharya N, Castañeda CA, Sukenik S. Single-point mutations in disordered proteins: Linking sequence, ensemble, and function. Curr Opin Struct Biol . 2025 Apr:91:102987.

Hauser MA, Aboobakar IF, Liu Y, Miura S, Whigham BT, Challa P, Wheeler J, Williams A, Santiago-Turla C, Qin X, Rautenbach RM, Ziskind A, Ramsay M, Uebe S, Song L, Safi A, Vithana EN, Mizoguchi T, Nakano S, Kubota T, Hayashi K, Manabe S, Kazama S, Mori Y, Miyata K, Yoshimura N, Reis A, Crawford GE, Pasutto F, Carmichael TR, Williams SE, Ozaki M, Aung T, Khor CC, Stamer WD, Ashley-Koch AE, Allingham RR. Genetic variants and cellular stressors associated with exfoliation syndrome modulate promoter activity of a lncRNA within the LOXL1 locus. Hum Mol Genet. 2015 Nov 15;24(22):6552-63.

Hayashi H, Gotoh N, Ueda Y, Nakanishi H, Yoshimura N. Lysyl oxidase-like 1 polymorphisms and exfoliation syndrome in the Japanese population. Am J Ophthalmol. 2008 Mar;145(3):582-585.
Howie BN, Donnelly P and Marchini J (2009) A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genetics* 5(6): e1000529.

Igo RP Jr, Cooke Bailey JN, Romm J, Haines JL, Wiggs JL. Quality Control for the Illumina HumanExome BeadChip. Curr Protoc Hum Genet. 2016 Jul 1;90:2.14.1-2.14.16.

Jiang L-L, Zhao J, Yin X-F, He W-T, Yang H, Che M-X, Hu H-Y. Two mutations G335D and Q343R within the amyloidogenic core region of TDP-43 influence its aggregation and inclusion formation. Sci Rep 2016, 6, 23928.

Kang JH, Wu J, Cho E, Ogata S, Jacques P, Taylor A, Chiu CJ, Wiggs JL, Seddon JM, Hankinson SE, Schaumberg DA, Pasquale LR. Contribution of the Nurses' Health Study to the Epidemiology of Cataract, Age-Related Macular Degeneration, and Glaucoma. Am J Public Health. 2016 Sep;106(9):1684-9.

Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. Nat Genet. 2014 Mar;46(3):310-5.

Karczewski KJ, Solomonson M, Chao KR, Goodrich JK, Tiao G, Lu W, Riley-Gillis BM, Tsai EA, Kim HI, Zheng X, Rahimov F, Esmaeeli S, Grundstad AJ, Reppell M, Waring J, Jacob H, Sexton D, Bronson PG, Chen X, Hu X, Goldstein JI, King D, Vittal C, Poterba T, Palmer DS, Churchhouse C, Howrigan DP, Zhou W, Watts NA, Nguyen K, Nguyen H, Mason C, Farnham C, Tolonen C, Gauthier LD, Gupta N, MacArthur DG, Rehm HL, Seed C, Philippakis AA, Daly MJ, Davis JW, Runz H, Miller MR, Neale BM. Systematic single-variant and gene-based association testing of thousands of phenotypes in 394,841 UK Biobank exomes. Cell Genom. 2022 Aug 15;2(9):100168.

Li X, He J, Sun J. LOXL1 gene polymorphisms are associated with exfoliation syndrome/exfoliation glaucoma risk: An updated meta-analysis. PLoS One 2021 Apr 28;16(4):e0250772.

Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen WM. Robust relationship inference in genome-wide association studies. *Bioinformatics*. 2010; 26: 2867–2873.

Margeta MA, Letcher SM, Igo RP Jr, Cooke Bailey JN, Pasquale LR, Haines JL, Butovsky O, Wiggs JL; NEIGHBORHOOD consortium. Association of *APOE* With Primary Open-Angle Glaucoma Suggests a Protective Effect for *APOE* epsilon4. Invest Ophthalmol Vis Sci. 2020 Jul 1;61(8):3.

Nazarali S, Damji F, Damji KF. What have we learned about exfoliation syndrome since its discovery by John Lindberg 100 years ago? Br J Ophthalmol. 2018 Oct;102(10):1342-1350.

Pasutto F, Zenkel M, Hoja U, Berner D, Uebe S, Ferrazzi F, Schödel J, Liravi P, Ozaki M, Paoli D, Frezzotti P, Mizoguchi T, Nakano S, Kubota T, Manabe S, Salvi E, Manunta P, Cusi D, Gieger C, Wichmann HE, Aung T, Khor CC, Kruse FE, Reis A, Schlötzer-Schrehardt U. Pseudoexfoliation syndrome-associated genetic variants affect transcription factor binding and alternative splicing of LOXL1. Nat Commun. 2017 May 23;8:15466.

Peng K, Vucetic S, Radivojac P, Brown CJ, Dunker AK, Obradovic Z. Optimizing long intrinsic disorder predictors with protein evolutionary information. *J Bioinform Comput Biol*. 2005; 3(1): 35–60.

Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet*. 2006; 38: 904–909.

Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007; 81: 559–575.

Schlötzer-Schrehardt U. Molecular pathology of pseudoexfoliation syndrome/glaucoma--new insights from LOXL1 gene associations. Exp Eye Res. 2009 Apr; 88(4):776-85.

Sharma S, Chataway T, Burdon KP, Jonavicius L, Klebe S, Hewitt AW, Mills RA, Craig JE. Identification of LOXL1 protein and Apolipoprotein E as components of surgically isolated pseudoexfoliation material by direct mass spectrometry. Exp Eye Res. 2009 Oct;89(4):479-85.

Sharma S, Martin S, Sykes MJ, Dave A, Hewitt AW, Burdon KP, Ronci M, Voelcker NH, Craig JE. Biological effect of LOXL1 coding variants associated with pseudoexfoliation syndrome. Exp Eye Res. 2016 May;146:212-223.

Shumway C, Curtin K, Taylor S, Sundar KM, Wirostko BM, Ritch R. Association between Obstructive Sleep Apnea and Exfoliation Syndrome: The Utah Project on Exfoliation Syndrome. Ophthalmol Glaucoma. 2021 May-Jun;4(3):260-267.

Taylor SC, Bernhisel AA, Curtin K, Allingham RR, Ritch R, Wirostko BM. Association between Chronic Obstructive Pulmonary Disease and Exfoliation Syndrome: The Utah Project on Exfoliation Syndrome. Ophthalmol Glaucoma. 2019 Jan-Feb;2(1):3-10.

Wiggs JL, Hauser MA, Abdrabou W, Allingham RR, Budenz DL, Delbono E, Friedman DS, Kang JH, Gaasterland D, Gaasterland T, Lee RK, Lichter PR, Loomis S, Liu Y, McCarty C, Medeiros FA, Moroi SE, Olson LM, Realini A, Richards JE, Rozsa FW, Schuman JS, Singh K, Stein JD, Vollrath D, Weinreb RN, Wollstein G, Yaspan BL, Yoneyama S, Zack D, Zhang K, Pericak-Vance M, Pasquale LR, Haines JL. The NEIGHBOR consortium primary open-angle glaucoma genome-wide association study: rationale, study design, and clinical variables. J Glaucoma. 2013 Sep;22(7):517-25.

Wiggs JL, Yaspan BL, Hauser MA, Kang JH, Allingham RR, Olson LM, Abdrabou W, Fan BJ, Wang DY, Brodeur W, Budenz DL, Caprioli J, Crenshaw A, Crooks K, Delbono E, Doheny KF, Friedman DS, Gaasterland D, Gaasterland T, Laurie C, Lee RK, Lichter PR, Loomis S, Liu Y, Medeiros FA, McCarty C, Mirel D, Moroi SE, Musch DC, Realini A, Rozsa FW, Schuman JS, Scott K, Singh K, Stein JD, Trager EH, Vanveldhuisen P, Vollrath D, Wollstein G, Yoneyama S, Zhang K, Weinreb RN, Ernst J, Kellis M, Masuda T, Zack D, Richards JE, Pericak-Vance M, Pasquale LR, Haines JL. Common variants at 9p21 and 8q22 are

associated with increased susceptibility to optic nerve degeneration in glaucoma. PLoS Genet. 2012;8(4):e1002654.

Williams SE, Whigham BT, Liu Y, Carmichael TR, Qin X, Schmidt S, Ramsay M, Hauser MA, Allingham RR. Major LOXL1 risk allele is reversed in exfoliation glaucoma in a black South African population. Mol Vis. 2010 Apr 21;16:705-12.

Zenkel M, Schlötzer-Schrehardt U. The composition of exfoliation material and the cells involved in its production. J Glaucoma. 2014 Oct-Nov;23(8 Suppl 1):S12-4.

Tables and Figures

Table 1. Distribution of rare *LOXL1* coding variants in XFS/XFG cases and controls in the MEE cohort

SNP [Minor allele]	CADD Score	Case (N=1118) MAC	Ctrl (N=3661) MAC	OR (95% CI) P	NBR Ctrl (N=2606) MAC	AllofUs Ctrl N= (37522) MAC	Combined Ctrls (N=43789) MAC	OR (95% CI) P
rs201884374 G120S [A]	25.8	0	2	NA	3	36	41	NA 0.63
rs78803776 S159A [G]	20.6	0	2	NA	2	19	43	NA 0.60
rs138183635 A160P [C]	26.6	1	19	0.16 (0.02, 1.19) 0.06	8	213	240	0.19 (0.03, 1.38) 0.03
rs189632283 S427F [T]	9.2	1	21	0.15 (0.02, 1.28) 0.04	8	5	34	0.17 (0.02, 1.25) 0.99
Total		2	44	0.15 (0.04, 0.62) 7.6E-4	21	273	338	0.17 (0.04, 0.70) 2.6E-3

Rare variant distribution in 1118 XFS/XFG cases and 3661 controls from the MEE cohort using Fisher's exact test. Variants were also identified in the NEIGHBOR controls (N= 2606) and AllofUS controls (N= 37522). No variants were identified 284 AllofUs cases. Abbreviations: SNP, single nucleotide polymorphism; CADD, combined annotation dependent depletion; Ctrl, control; OR, odds ratio; NBR, NEIGHBOR; MAC, minor allele count; MAF, minor allele frequency.

Table 2. Rare *LOXL1* coding variant association in replication cohorts

	SNP [Minor allele]			
	rs201884374 G120S [A]	rs78803776 S159A [G]	rs138183635 A160P [C]	rs189632283 S427F [T]
Population	OR (P value)	OR (P value)	OR (P value)	OR (P value)
South Africa 90 cases 250 controls	NP	0.21 (0.0004)	NP	NP
Iceland 195 cases 14,474 controls	NP	NP	0.17 (0.0014)	0.16 (0.026)

rs78803776[G] was only present in the South African sample, while rs138183635[C] and rs189632283[T] were only present in the Icelandic sample. Abbreviations: OR, Odds ratio; NP, not present.

Table 3. *LOXL1* haplotypes in the MEE cohort with common and rare coding variants and XFS/XFG association.

Hap #	G120S	R141L	G153D	S159A	A160P	S427F	Cases (N=2236)	Controls (N=7322)	P	OR (95% CI)
							N	N		
1	A	T	G	T	G	C	0	2	NA	NA
2	G	G	A	T	C	C	1	19	0.06	0.16 (0.02, 1.19)
3	G	G	A	T	G	C	34	1219	7.90E-106	0.08 (0.06, 0.11)
4	G	G	G	G	G	C	0	2	NA	NA
5	G	G	G	T	G	C	1869	3785	1.7E-173	4.76 (4.20, 5.36)
6	G	T	G	T	G	C	331	2274	3.74E-56	0.39 (0.34, 0.44)
7	G	T	G	T	G	T	1	21	0.04	0.15 (0.02, 1.28)
All minor							367	3537	1.7E-173	0.21 (0.19, 0.24)

The minor (rare) allele at each variant is highlighted in yellow. Only haplotype #5 (highlighted in orange) that includes all the major (common) alleles is associated with disease risk. Abbreviations: NA, not applicable; OR, odds ratio.

Table 4. *LOXL1* diplotypes and XFS/XFG association in the MEE cohort

Diplotype	G120S	R141L	G153D	S159A	A160P	S427F	Cases	Freq.	Controls	Freq.	P	OR (95% CI)
A (H5/H5)	G	G	G	T	G	C	786	0.703	947	0.259	4.7E-157	6.78 (5.87,7.
	G	G	G	T	G	C						
B (H6/H6)	G	T	G	T	G	C	18	0.016	448	0.122	6.4E-34	0.13 (0.08,0.
	G	T	G	T	G	C						
Sum Homozygous							**804**	**0.719**	**1395**	**0.38**	**4.2E-89**	**4.16 (4.76,3.**
C (H5/H6)	G	**G**	G	T	G	C	290	0.259	1220	0.333	3.0E-06	0.7 (0.80,0.
	G	**T**	G	T	G	C						
D (H5/H3)	G	G	**G**	T	G	C	15	0.013	659	0.18	1.8E-68	0.06 (0.10,0.
	G	G	**A**	T	G	C						
E (H7/H6)	G	T	G	T	G	**T**	1	8.9E-4	7	1.9E-3	0.69	0.47 (3.78,0.
	G	T	G	T	G	**C**						
F (H2/H3)	G	G	A	T	**C**	C	1	8.9E-4	2	5.5E-4	0.55	1.64 (0.15,1
	G	G	A	T	**G**	C						
Sum one Heterozygous							**307**	**0.27**	**1888**	**0.516**	**5.7E-47**	**0.36 (0.41,0.**
G (H3/H6)	G	**G**	**A**	T	G	C	7	6.3E-3	343	0.093	4.7E-32	0.06 (0.13,0.
	G	**T**	**G**	T	G	C						
H (H5/H2)	G	G	**G**	T	**G**	C	0	0	6	0.002	0.34	NA
	G	G	**A**	T	**C**	C						

							Cases		Controls		P	OR
I (H7/H5)	G	**T**	G	T	G	**T**	0	0	13	0.004	0.048	NA
	G	**G**	G	T	G	**C**						
J (H6/H4)	G	**T**	G	**T**	G	C	0	0	1	2.7E-4	NA	NA
	G	**G**	G	**G**	G	C						
K (H3/H4)	G	G	**A**	**T**	G	C	0	0	1	2.7E-4	NA	NA
	G	G	**G**	**G**	G	C						
L (H1/H5)	**A**	T	G	T	G	C	0	0	1	2.7E-4	NA	NA
	G	**G**	G	T	G	C						
Sum 2 heterozygous							7	0.0063	365	0.0997	1.2E-34	0.06 (0.13,0.
M (H2/H6)	G	**G**	**A**	T	**C**	C	0	0	11	0.003	0.078	NA
	G	**T**	**G**	T	**G**	C						
N (H3/H7)	G	**G**	**A**	T	G	**C**	0	0	2	5.5E-4	NA	NA
	G	**T**	**G**	T	G	**T**						
Sum 3 heterozygous							0	0	13	0.0036	0.048	**NA**
Sum 2 + 3 heterozygous							7	6.3E-3	378	0.10	1.0E-39	0.05 (0.10,0.
Sum all heterozygous							314	0.28	2266	0.62	4.2E-89	0.45 (0.52,0.

Heterozygous genotypes are shown in bold type. Only the homozygous diplotype A (H5/H5) is significantly associated with risk. Overall 72% of cases have a homozygous diplotype compared to 38% of controls. The diplotypes that include two heterozygous genotypes are found in 7 cases and 342 controls while the diplotypes with three heterozygous genotypes are only found in controls.

Table 5. *LOXL1* haplotypes in the AllofUs cohort with common and rare coding variants and XFS/XFG association.

Haplotype Number	Haplotype	Frequency Cases (n=568)	Frequency Controls (n=74998)	OR	95% CI	P
1	GG	447 (78.7%)	38456 (51.3%)	3.22	2.61-3.92	6.2E-43
2	GG(+)	0 (0%)	139 (0.2%)	--	--	0.41
3	GG(++)	0 (0%)	2 (0.003%)	--	--	0.07
4	GA	19 (3.3%)	12039 (16.1%)	0.14	0.09-0.22	5.1E-22
5	GA(+)	0 (0%)	255 (0.3%)	--	--	0.21
6	TG	102 (18.0%)	24006 (32.0%)	0.37	0.29-0.45	7.6E-14
7	TG(+)	0 (0%)	101 (0.1%)	--	--	0.54
8	TA	0 (0%)	0 (0%)			
9	TA(+)	0 (0%)	0 (0%)			
All minor*		121 (21.3%)	36542 (48.7%)	0.28	0.23-0.34	4.41E-41

*Minor alleles include the rare alleles (MAF <1%) and the alternate alleles for the two common variants ([D] at G153D and [L] at R141L).

Table 6. *LOXL1* diplotypes and XFS/XFG association in the AllofUs cohort

Haplotype Numbers	Diplotype	Frequency Cases (n=284)	Frequency Controls (n=37499)	OR	95% CI	P
1/1	GG|GG	186 (65.5%)	9873 (26.3%)	5.26	4.12-6.73	1.8E-42
1/2	GG|GG(+)	0 (0%)	62 (0.2%)	--	--	0.55
1/3	GG|GG(++)	0 (0%)	2 (0.005%)	--	--	0.13
1/4	GG|GA	8 (2.8%)	6184 (16.5%)	0.07	0.03-0.14	3.6E-13
1/5	GG|GA(+)	0 (0%)	105 (0.3%)	--	--	0.33
2/5	GG(+)|GA	0 (0%)	32 (0.09%)	--	--	0.89
1/6	GG|TG	67 (23.6%)	12307 (32.8%)	0.29	0.22-0.38	9.3E-4
1/7	GG|TG(+)	0 (0%)	50 (0.1%)	--	--	0.65
4/4	GA|GA	3 (1.1%)	983 (2.6%)	0.16	0.05-0.51	2.0E-3
4/5	GA|GA(+)	0 (0%)	49 (0.1%)	--	--	0.66
5/5	GA(+)|GA(+)	0 (0%)	1 (0.003%)	--	--	0.08
4/6	GA|TG	5 (1.8%)	3790 (10.1%)	0.07	0.03-0.17	9.6E-8
4/7	GA|TG(+)	0 (0%)	18 (0.05%)	--	--	0.80
6/2	TG|GG(+)	0 (0%)	45 (0.1%)	--	--	0.70
6/5	TG|GA(+)	0 (0%)	99 (0.3%)	--	--	0.35
6/6	TG|TG	15 (5.3%)	3866 (10.3%)	0.21	0.12-0.35	6.9E-3
6/7	TG|TG(+)	0 (0%)	33 (0.1%)	--	--	0.87

Diplotypes were comprised on individual haplotypes defined in Table 5. Only the diplotype homozygous for the 'G' allele for both G153D and R141L is significantly associated with disease risk.

Table 7. Summary of homozygous and heterozygous diplotypes among cases and controls in AllofUs

Number of haplotypes per individual with homozygous or heterozygous diplotypes	Frequency Cases (n=284)	Frequency Controls (n=37499)	OR	95% CI	P
All homozygous	204 (72%)	14,722 (39%)	3.95	3.04- 5.11	3.0E-28
1 heterozygous	75 (26%)	18,553 (49%)	0.37	0.28-0.48	2.4E-16
2 heterozygous	5 (0.18%)	4125 (11%)	0.15	0..06- 0.35	4.6E-9
1+2 heterozygous	80 (28%)	22,678 (60%)	0.26	0.18-0.33	7.6E-28

Summary of the diplotypes shown in Table 7 according to number of heterozygous genotypes.

Predicted Disorder Segment (Amino acid residues)	Average strength
1-8	0.7046
56-109	0.9336
138-166	0.7926
212-226	0.8237
251-276	0.8527
286-380	0.9036

Figure 1. LOXL1 intrinsic disordered domains. Six disordered domains are predicted by the program PONDR (Predictor of Natural Disordered Regions) (Peng, 2005). Disordered probably greater than 50% is considered to be significant. The regions resulting in reduced aggregation when deleted were residues 28-95 and residues 125-185, Bernstein et al. (Adv Protein Chem Struct Biol. 2019;118:273-288).

Table 8. Enrichment of rare variants in controls in predicted disordered segments

Predicted Disorder Segment (Amino acid residue)	Number of rare alleles N=456	P-value
1-8	3	0.36
56-109	40	0.82
138-166	241	5.8E-45
212-226	0	--
251-276	0	--
286-380	56	0.11

Rare alleles include missense, inframe insertion, inframe deletion and start loss. The full list of alleles included in this analysis are in Supplementary Table 2.

Table 9. rs3825942 (G153D) genotype distribution in cases (N=5748) from the combined datasets

All genotypes	GG	GA	AA	
Observed	5383	311	49	
Expected	5297	397	7.5	P=1.0e-6
All Homozygous	GG+AA			
Observed	5514			
Expected	5414	P=1.9E-4		
Homozygous minor allele	AA+GG*			
Observed	56			
Expected	10	P=6.3E-9		
Heterozygous	GA			
Observed	311			
Expected	392	1.8E-3		

The genotype counts for all ancestry groups (excluding the African group see below) was combined with the counts for the MEE cohort and the AllofUs cohort. The expected genotype distribution was calculated using the Hardy Weinberg equation ($p^2 + 2pq + q^2 = 1$) and using the combined allele frequency for each allele. The observed genotype distribution was determined from the combined genotype data. The difference between expected and observed was calculated using Chi square. The individual genotype distribution for each datasets is included in Supplementary Table 5.

Except for the minor allele homozygosity analysis, the African ancestry group was not included in this combined calculation due to the allele reversal. Individual data for the African group is included in Supplementary Table 5. *The African data was included in the minor allele homozygosity analysis with genotype 'GG' included as this is the minor allele genotype in this ancestry group.

Table 10. rs1048661 (R141L) genotype distribution in Japanese and Korean cases (N=1039)

All genotypes	GG	GT	TT	
Observed	65	171	803	
Expected	22	257	759	P=1.0E-6
All Homozygous	GG+TT			
Observed	868			
Expected	781	P=2.9E-6		
Homozygous minor allele	GG			
Observed	65			
Expected	22	2.7E-6		
Heterozygous	GT			

Observed	171			
Expected	257	3.7E-6		

The observed combined homozygous and heterozygous genotype distribution for the Korean and Japanese ancestral groups is compared to the expected. The expected genotype distribution was calculated using the Hardy Weinberg equation ($p^2 + 2pq + q^2 = 1$) and the combined allele frequency for each allele. The difference between expected and observed was calculated using Chi square. Only the Korean and Japanese ancestry groups were included as they are the only groups that show deviation from Hardy Weinberg equilibrium for this variant (see Supplementary Table 6).

Supplementary Tables

Supplementary Table 1. XFS/XFG case and control demographics

	MEE cases	MEE controls	NEIGHBOR controls	AllofUs cases	AllofUs controls	P value
N	1118	3661	2606	284	37522	
% Female	69	65	64	60	51	0.003
Age (average)	72.9 $\pm$ 9.6	54.0 $\pm$ 7.8	65.5 $\pm$ 13.5	80.5 $\pm$ 8.1	80.2 $\pm$ 4.6	0.28

Supplementary Table 2. LOXL1 alleles identified in the _All of Us_ cohort

Genome	alleles	Variant Type	AA Change	Number in Controls
chr15:73949555	['A', 'G']	Missense	Thr567Ala	16
chr15:73949465	['G', 'A']	Missense	Val537Met	11
chr15:73947894	['A', 'G']	Missense	Ile532Val	4
chr15:73947882	['C', 'A']	Missense	Pro528Thr	2
chr15:73947152	['A', 'G']	Missense	Ser479Gly	2
chr15:73947147	['A', 'G']	Missense	Lys477Arg	3
chr15:73947143	['C', 'T']	Missense	His476Tyr	2
chr15:73947137	['G', 'A']	Missense	Glu474Lys	5
chr15:73947117	['C', 'T']	Missense	Ala467Val	3

chr15:73946553	['C', 'G']	Missense	Gln450Glu	2
chr15:73946518	['G', 'A']	Missense	Arg438Gln	2
chr15:73946491	['A', 'G']	Missense	Gln429Arg	2
chr15:73946481	['G', 'A']	Missense	Val426Met	5
chr15:73946479	['G', 'A']	Missense	Arg425His	2
chr15:73946467	['G', 'A']	Missense	Arg421His	4
chr15:73942941	['CGGA', 'C']	In frame deletion Glu399del	4
chr15:73942923	['A', 'T']	Missense	Tyr391Phe	2
chr15:73927868	['C', 'A']	Missense	Pro362His	2
chr15:73927836	['G', 'C']	Missense	Gln351His	3
chr15:73927835	['A', 'C']	Missense	Gln351Pro	2
chr15:73927823	['A', 'C']	Missense	Asn347Thr	2
chr15:73927781	['C', 'G']	Missense	Pro333Arg	12
chr15:73927775	['A', 'C']	Missense	Tyr331Ser	3
chr15:73927760	['C', 'A']	Missense	Ala326Glu	3
chr15:73927748	['G', 'T']	Missense	Glu322Val	7
chr15:73927748	['GGCCGCCGCGCGCGCTGGAGCCGCCCTACCT', 'G']	In frame insertion Pro318_Pro324dup	10
chr15:73927747	['G', 'C']	Missense	Glu322Arg	3
chr15:73927693	['G', 'A']	Missense	Asp304Asn	5
chr15:73927686	['T', 'G']	Missense	His301Gln	2
chr15:73927639	['T', 'G']	Missense	Phe286Val	2
chr15:73927261	['G', 'C']	Missense	Ala160Pro	213
chr15:73927258	['T', 'G']	Missense	Ser159Ala	19
chr15:73927722	['C', 'G']	Missense	Ser147Cys	6

3					
chr15:73927193	['C', 'T']		Missense	Thr137Met	3
chr15:73927141	['G', 'A']		Missense	Gly120Ser	36
chr15:73927134	['C', 'G']		Missense	His117Gln	2
chr15:73927060	['C', 'T']		Missense	Arg93Trp	2
chr15:73927042	['C', 'G']		Missense	Arg87Gly	11
chr15:73927019	['G', 'A']		Missense	Gly79Asp	5
chr15:73927004	['G', 'C']		Missense	Arg74Pro	2
chr15:73927000	['T', 'G']		Missense	Ser73Ala	2
chr15:73926967	['G', 'T']		Missense	Val62Leu	3
chr15:73926967	['G', 'A']		Missense	Val62Met	3
chr15:73926950	['A', 'C']		Missense	Asn56Thr	12
chr15:73926875	['G', 'A']		Missense	Gly31Glu	5
chr15:73926817	['G', 'A']		Missense	Ala12Thr	2
chr15:73926785	['T', 'C']		Start loss	Met1Thr	3

Supplementary Table 3. *LOXL1* exome variant association in UK Biobank

Dataset	SKAT-O			Burden		
	LoF N=30	Missense N=361	Synonymous N=189	LoF N=30	Missense N=361	Synonymous N=189
Glaucoma H40 (11588 cases, 383253 controls)	P= 0.577 β= -2.3e-2	P= 3.39e-3 β= -8.37e-3	P= 0.374 β= 5.82e-3	P= 0.587 β= -2.3e-2	P= 1.82e-3 β= -8.7e-3	P= 0.371 β= 5.82e-3
Glaucoma custom 8912 cases, 385929, controls	P= 0.783 β= -1.69e-2	P= 8.55e-3 β= -9.03e-3	P= 0.632 β= 2.06e-3	P= 0.565 β= -1.69e-2	P= 4.33e-3 β= -9.03e-3	P= 0.781 β= 2.06e-3

Glaucoma 4063 cases, 390720 controls	P= 0.504 β= 8.353-3	P= 0.013 β= -1.24e-2	P= 0.859 β= -1.62e-3	P= 0.841 β= 8.35e-3	P= 7.93e-3 β= -1.24e-2	P= 0.879 β= 2.01e-3
Glaucoma Surgery 560 cases, 394223 controls	P= 1.73e-3 β= 0.178	P= 0.127 β= -2.23e-2	P= 0.654 β= 2.01e-3	P= 0.0473 β= 1.78e-1	P= 7.25e-2 β= -2.23e-2	P= 0.941 β= -1.62e-3

Gene burden results (SKAT-O and Burden) in the UK Biobank were assessed using GeneBass (Karczewski, 2022) for all 4 glaucoma phenotype groups and three variant classification groups (Loss of function, LoF, Missense and Synonymous). N= the number of variants in each group. For all glaucoma groups and for both gene burden methods the missense variants are consistently protective.

Supplementary Table 4. Individual variant results for *LOXL1* variants in the UK Biobank

Variant	IDR	Glaucoma H40	Case AC N= 11588	Control AC N= 383253
Asn56Thr	Y (56-109)	P= 5.04e-2 β= -7.53e-1	2	234
Ala160Pro	Y (138-166)	P=5.29e-3 β= -3.29e-1	48	2430
Pro293Ser	Y (286-380)	P=5.1e-2 β= 4.3	1	2
Cys401Tyr	N	P=4.74e-2 β= 4.13	1	2
Leu544Ser	N	P=4.58e-2 β= 4.17	1	2

Variants with allele count (AC) of at least 2 in cases+controls, and p values for association of <0.05 are listed. Asn56Thr, Ala160Pro, Pro293Ser are all located within LOXL1 intrinsic disordered regions (IDR).

Supplementary Table 5. Observed and expected genotype distribution for G153D (rs3825942)

A. Meta-analysis published cases

Caucasian	Cases N=2138				Controls N=5039			
	GG	GA	AA		GG	GA	AA	
Observed	2006	114	18		3539	1338	162	
Expected	1990	144	2.6	P=5.3e-4	3513	1388	137	P=0.211
Asian	Cases N=2208				Controls N=2939			
	GG	GA	AA		GG	GA	AA	
Observed	2022	156	25		2035	793	111	

Expected	1996	197	4.9	P=9.9e-5	2010	840	88	P=0.125
African	Cases N=93				Controls N=97			
	GG	GA	AA		GG	GA	AA	
Observed	7	11	75		39	42	16	
Expected	1.7	21.6	69.75	P=0.032	37	45	14	P=0.867

B. MEE cohort

	Cases N=1118				Controls N=3661			
	GG	GA	AA		GG	GA	AA	
Observed	1087	28	3		2637	1022	2	
Expected	1084	33	0.25	P=0.25	2707	881	71	P=1.0e-6

C. *All of Us*

	Cases N=284				Controls N=37521			
	GG	GA	AA		GG	GA	AA	
Observed	268	13	3		26,253	10,233	1034	
Expected	265	18	.3	P=0.21	26,222	10,288	1009	P=0.78

Observed and expected genotype distribution for the G153D variant in A) European, Asian, and African ancestry groups from Li et al. (Li, 2021), B) the MEE cohort and C) the AllofUs cohort. Expected genotype distribution was calculated using the Hardy Weinberg equation (p^2 +2pq+q^2 =1) and using observed allele frequencies for each group. The observed genotype distribution was determined from genotype data for each group. The difference between expected and observed was calculated using Chi square. The genotype distribution for the combined datasets is shown in Table 8 in the main text.

Supplementary Table 6. Observed and expected genotype distribution for R141L (rs1048661)

A. Meta-analysis published cases

Caucasian	Cases N=2596				Controls N=5039			
	GG	GT	TT		GG	GT	TT	
Observed	1727	805	64		2530	2437	612	
Expected	1745	766	84	P=0.15	2504	2467	607	P=0.33
Asian*	Cases N=1120				Controls N=1474			
	GG	GT	TT		GG	GT	TT	
Observed	738	324	58		635	647	192	
Expected	723	352	43	P=0.17	622	668	179	P=0.63
Japanese and Korean	Cases N=1039				Controls N=1323			
	GG	GT	TT		GG	GT	TT	
Observed	65	171	803		311	684	328	
Expected	22	257	759	P=1.0e-6	322	661	338	P-0.69
African	Cases N=93				Controls N=97			
	GG	GT	TT		GG	GT	TT	

Observed	92	1	0		70	24	3	
Expected	78	2	0	P=0.53	69	25	2	P=0.89

B. MEE cohort

	Cases N=1118				Controls N=3661			
	GG	GT	TT		GG	GT	TT	
Observed	802	297	19		1615	1591	455	
Expected	807	285	25	P=0.58	1585	1648	428	P=0.34

C. *All of Us*

	Cases N=284				Controls N=37521			
	GG	GT	TT		GG	GT	TT	
Observed	197	72	15		17306	16314	3901	
Expected	190	84	9.2	P=0.29	17350	16329	3842	P=0.77

Observed and expected genotype distribution for the R141L variant in A) European, Asian, and African ancestry groups from Li et al. (Li, 2021) with the Korean and Japanese separated from the other Asian groups as the R141L is the main risk allele in this group while G153D is main risk allele in the other ancestry groups. B) the MEE cohort and C) the AllofUs cohort. Expected genotype distribution was calculated using the Hardy Weinberg equation ($p^2 +2pq+q^2 =1$) and using observed allele frequencies for each group. The observed genotype distribution was determined from genotype data for each group. The difference between expected and observed was calculated using Chi square. The combined genotype distribution (all homozygous versus heterozygous) for the Korean and Japanese datasets is shown in Table 9 in the main text.

Supplementary Figure 1. *LOXL1* **diplotype distribution among cases and controls according to age**.
Total diplotypes with 0 heterozygous genotypes (ie homozygous) (blue), 1 heterozygous genotype
(orange), 2 heterozygous genotypes (purple) and 3 heterozygous genotypes (green) were summed for
cases and controls according to age groups: 40-59, 60-69, 70-89 and compared with the distribution
among cases and controls overall. Similar distribution of diplotypes was observed for all age groups
(P>0.05) with the exception of the rare diplotypes with 3 heterozygous variants which were found only
in controls primarily in the 60-69 age range.