Altered interpersonal distance regulation in autism spectrum disorder

Kinga Farkas[1,2*], Orsolya Pesthy[3,4], Anna Guttengéber[4], Anna Szonja Weigl[5], András Veres[6],

Anna Szekely[7], Eszter Komoróczy[1], Bálint Szuromi[1], Karolina Janacsek[7,2,8], János M.

Réthelyi[1], Dezső Németh[2,4,5]

[1] *Department of Psychiatry and Psychotherapy, Semmelweis University, Budapest, Hungary*

[2] *Brain, Memory and Language Research Group, Institute of Cognitive Neuroscience and Psychology, Research*

Centre for Natural Sciences, Budapest, Hungary

[3] *Doctoral School of Psychology, ELTE Eötvös Loránd University, Budapest, Hungary*

[4] *Institute of Psychology, ELTE Eötvös Loránd University, Budapest, Hungary*

[5] *Lyon Neuroscience Research Center (CRNL), Université de Lyon, Centre Hospitalier Le Vinatier, Lyon,*

France

[6] *Obimon Systems Ltd., Budapest, Hungary*

[7] *MTA-ELTE Lendület Adaptation Research Group, Institute of Psychology, ELTE Eötvös Loránd University,*

Budapest, Hungary

[8] *Centre for Thinking and Learning, Institute for Lifecourse Development, School of Human Sciences, Faculty of*

Education, Health and Human Sciences, University of Greenwich, London, United Kingdom

* Corresponding author

E-mail: farkas.kinga@med.semmelweis-univ.hu (KF)

Keywords: autism, social distance, interpersonal space, personal space, heart rate variability

Author contributions

KF, OP, AG, DN contributed to the design and implementation of the research and developed the theoretical formalism⁄background. KF, AW, OP performed the computations. KF designed the figures. KF, OP, AG, BS, DN wrote the manuscript. OP and AG carried out the experiment. BS and EK contributed to sample preparation. DN, KJ, and RJ supervised the project. All authors provided critical feedback and helped shaping the research, analysis, and manuscript.

Abstract

Interpersonal distance regulation is an essential element of social communication. Its impairment in autism spectrum disorder (ASD) is widely acknowledged among practitioners, but only a handful of studies reported empirical research in real-life settings focusing mainly on children. Interpersonal distance in adults with ASD and related autonomic functions received less attention. Here, we measured interpersonal distance along with heart rate variability (HRV) in adults with ASD, and tested the modulatory effects of eye-contact and attribution. Twenty-two adults diagnosed with ASD and 21 matched neurotypical controls participated in our study from 2019 October to 2020 February. Our experimental design combined the modified version of the stop distance paradigm with HRV measurement controlling for eye contact between the experimenter and the participant to measure interpersonal distance. Our results showed a greater preferred distance in ASD. These results were altered with eye contact. Moreover, we found lower baseline HRV and reduced HRV reactivity in ASD; however, these autonomic measurements could not predict preferred interpersonal distance. Our study highlights the importance of interpersonal space regulation in ASD and the need for sophisticated experimental designs to grasp the complexity and underlying factors of distance regulation in typical and atypical populations.

Introduction

Autism spectrum disorder (ASD) is a neurodevelopmental condition characterized by persistent difficulties in social communication and social interaction across multiple contexts, such as abnormal social approach or failure to initiate or respond to social interactions; and restricted, repetitive patterns of behaviour, interests, or activities [1]. At the neural level, cortical [2,3], subcortical [4,5], and autonomic [6,7] neural alterations can be observed, including developmental, structural and functional differences [8–10] in parallel to the pervasive cognitive [11,12], behavioural and physiological disturbances in ASD. However, one of the key components of social behavior, namely interpersonal distance regulation, has received relatively less attention in ASD research (see exceptions: [13–16]) even though its impairment in ASD is widely acknowledged among practitioners. Our study aims to measure the interpersonal distance regulation and a related physiological parameter (heart rate variability, HRV) during this task and test the modulatory effect of two relevant factors in social communication: eye contact and attribution of self or the other, in autism spectrum disorder.

Finding the appropriate social distance can be seen as the first step of physical social interactions. It is widely believed among practitioners that people with ASD keep a greater or abnormal distance [17], violations of personal space occur also more often in ASD in childhood [18]. However, it is challenging to measure this phenomenon experimentally with high ecological validity, and the results are inconsistent. In autistic participants, preference for both closer [15,19–21] or farther distance [16,22] can be found in the literature. Among these studies, only one measured interpersonal distance in adult ASD; they found no difference between study groups [18]. Studies applying electrophysiological or imaging methods usually present video recordings of an approaching individual [14] or use virtual reality displays [23,24] to measure interpersonal distance regulation in ASD. We argue that

76   virtual displays might be useful in training or therapeutic settings, but they cannot take into

77   account several sensory modalities of the scene while measuring the behavioural and

78   physiological reactions in interpersonal interactions. In the present study, we measured the

79   interpersonal distance among adult participants with ASD in an experimental setting with

80   personal presence as close as possible to real-life situations.

81   Despite its relevance, empirical studies on interpersonal distance regulation of

participants with ASD were conducted only in the past few years. The nomenclature of the

concept is still not unified. *Personal space* [16,19,22], *social-* or *physical distance* [18], and

interpersonal space and *-distance* [13,14,16] are all commonly used. When measuring the

physical distance between two people in one dimension, we use the term *interpersonal*

distance. The Stop Distance Paradigm [25] is the most commonly used method for measuring

interpersonal distance regulation. This method is considered an ecological measure of

permeability and flexibility of interpersonal space regulation [13]. It can be combined with

various factors. Previous studies suggested that the presence of eye contact [19], the role

played during the experiment (active approaching or passive indicating) [22], and social

interactions as experimental interventions [14,16] might influence interpersonal distance.

Most of the studies examined children or adolescents with autism [16,19,22] and found larger

preferred interpersonal distance in ASD, except one study [19]. The only study using the Stop

Distance Paradigm among adults found no difference between groups in terms of

interpersonal distance preferences [18]. Since different results were gained in different

conditions, the question arises as to what extent these factors contribute to the outcome

behaviour.

98   Social communication and interactions are tremendously complex processes that can

be altered in autism at cognitive, behavioural, and physiological levels. At the highest level,

theory of mind difficulties can be observed in ASD [26,27]. To capture this phenomenon

during the interpersonal distance measurement in a simplified way, we added a new

condition. First, participants had to make a decision based on their own personal preference,

next, they were asked to estimate the comfortable distance for the experimenter. This latter

situation requires third order mentalization, the mobilisation of theory of mind ability. We

called this the attribution dimension. The processing of facial expressions, particularly that of

the eye region, is highly relevant in the regulation of social behaviour, including interpersonal

distance. Facial emotion processing and emotion recognition is altered in autism [8,28–31].

Constraining eye contact led to an exaggerated increase of amygdala activation, while

decreased eye contact was associated with diminished amygdala response to faces in ASD

[4,32–34]. In addition, unconsciously avoided eye contact in ASD results in further

difficulties of reading socially important signals [35,36]. These results suggest that altered

amygdala functioning, including the regulation of eye contact, might have a substantial role

in the disturbances of several aspects of social behaviour, such as personal proximity or

interpersonal space regulation [25,37,38]. Therefore, in addition to the attribution dimension,

eye contact and no eye contact conditions were introduced to investigate the effect of these

two factors, which are relevant for social communication.

Physiological response to sensory, social, and emotional stimuli is suggested to be

altered in ASD in general, however, the methodology used is highly variable and the results

are inconsistent [39]. Since the classic electrophysiological experiment of Hutt et al. showed

hyperarousal in children with ASD [40], the majority of studies that measured autonomic

regulation (pupillometry, skin conductance, or cardiac measures) found atypical resting-state

functions indicating either hyper- or hypoarousal in ASD according to a recent review [41].

Recent studies aimed to overcome these limitations by measuring heart rate variability

(HRV): heart rate is affected by both sympathetic and parasympathetic modulatory effects;

thus, its variability might be a good marker of autonomic regulation, as higher HRV reflects

parasympathetic activity [42]. Furthermore, a study found an association between HRV and

cognitive flexibility in healthy individuals [43]. A recent meta-analysis showed that heart rate

variability is reduced in ASD: baseline HRV and HRV reactivity during social stress were

significantly lower in participants with ASD, but HRV reactivity performing cognitive tasks

did not differ [44]. The reduced variability in the heart rate indicates an altered

parasympathetic-sympathetic balance in ASD, suggesting the predominance of sympathetic

activity and less flexible switching between autonomic states in ASD compared to

neurotypicals. For these reasons, we measured interpersonal distance along with heart rate

variability to examine their putative alterations and their relationship in ASD.

In this study, our main goal was to establish a comprehensive design to measure

interpersonal distance and autonomic functions in ASD. Our first hypothesis was that in adult

ASD we observe greater interpersonal distance. Second, we hypothesized that interpersonal

distance is modulated by eye contact and attribution. Finally, we aimed to determine the role

of autonomic functions in interpersonal distance regulation in ASD, expecting decreased

baseline HRV and reduced HRV reactivity during the interpersonal distance task in ASD. It

was also hypothesized that autonomic regulation, as characterized by HRV, could predict the

preferred interpersonal distance in both study groups.

Materials and methods

Participants

In total, 45 adults participated in our research. Two neurotypical participants were excluded due to errors during data collection. The final sample consisted of forty-three participants, 22 were diagnosed with autism spectrum disorder without intellectual disability or language impairment, and 21 were neurotypical controls. The two groups did not differ in age, gender, and education (Table 1). All participants with ASD were diagnosed by trained clinicians, the diagnoses were confirmed with Autism Diagnostic Interview-Revised (ADI-R) and Autism Diagnostic Observation Schedule, IV-module (ADOS-IV.) [45,46]. Twelve participants had one or more comorbid disorders (attention deficit hyperactivity disorder (5), obsessive-compulsive disorder (3), generalized anxiety disorder (2), bipolar disorder (1), depression (1), and schizophrenia (1)). Participants with ASD were recruited from the outpatient unit of the Department of Psychiatry and Psychotherapy, Semmelweis University. Neurotypical control participants were recruited by advertisement.

Participants (and legal guardians if applicable) provided written informed consent and did not receive financial compensation for their attendance. The study was conducted in accordance with the Declaration of Helsinki, and it was approved by the Regional and Institutional Committee of Science and Research Ethics, Semmelweis University, Budapest, Hungary (SERKEB No.: 145/2019) from October 2019 to February 2020. The experiment took place at the Laboratory of Brain, Memory and Language Lab, Eötvös Loránd University, Budapest.

163 **Table 1. Demographics and clinical characteristics.**
164

		ASD (*N*=22)		NTP (*N*=21)		Statistic	*p*		*effect size*
		N (M/F)		*N* (M/F)		χ^2	*p*		
Gender		18/4		14/7		1.296	0.255		
		Mean	*SD*	*Mean*	*SD*	*Mann-Whitney U*	*p*	*r*	*95% CI lower, upper*
Age	(years)	27.59	7.25	25.86	6.44	191.00	0.336	0.173	-0.173, 0.481
Education	(years)	15.64	3.69	16.00	3.33	248.50	0.677	-0.076	-0.401, 0.267
AQ		31.09	6.63	15.05	5.63	18.50	<0.001	0.920	0.845, 0.959
MZQ		51.68	9.53	38.29	9.30	71.50	<0.001	0.690	0.462, 0.833
AAS	anxious	22.64	5.77	16.81	6.06	114.50	0.005	0.504	0.203, 0.718
	avoidant	41.23	8.80	31.71	8.12	93.50	<0.001	0.595	0.324, 0.776
ASRS	Part A	13.23	4.02	9.52	3.84	110.50	0.003	0.522	0.225, 0.730
	Part B	26.64	9.19	16.62	5.83	88.00	<0.001	0.619	0.358, 0.790
STAI-T		55.46	11.63	44.29	9.09	105.50	0.002	0.543	0.254, 0.743
ADOS-IV.		9.96	3.35	-	-	-	-	-	-
ADI-R		34.82	7.74	-	-	-	-	-	-
		N (Y/N/n.a.)		*N* (Y/N/n.a.)		χ^2	*p*		
Caffeine	regular	18/2/2		15/5/1		1.558	0.212		
	within 12h	13/7/2		13/6/2		0.051	0.821		
Smoking		2/18/2		4/16/1		0.784	0.376		
Exercise		18/4/0		17/3/1		0.076	0.782		

165 ASD: Autism Spectrum Disorder, NTP: Neurotypical Participant, N: sample size, SD: standard deviation, r:
166 rank biserial correlation, 95% CI: 95% confidence interval, M: male, F: female, Y: yes, N: no, n.a.: not
167 available, AQ: Autism-Spectrum Quotient, MZQ: Mentalization Questionnaire, AAS: Adult Attachment Scale,
168 ASRS: Adult ADHD Self-Report Scale, STAI-T: State-Trait Anxiety Inventory - Trait, ADOS-IV: Autism
169 Diagnostic Observation Schedule - IV. module, ADI-R: Autism Diagnostic Interview-Revised
170

171 **Experimental paradigm - interpersonal distance task**

In our study, we measured social distance regulation. Participants underwent an

interpersonal distance measurement, a modified version of the stop-distance paradigm [25].

In all conditions, the participant and the experimenter started from the opposite endpoints of

the tape measure (five meters) stuck on the ground. They were asked to consciously focus on

comfortable social distance overall eight times with the following order. First, (1) participants

were approaching actively and were asked to stop where it was still comfortable for them.

Next, (2) participants were approaching actively and were asked to stop where they thought it

was still comfortable for the experimenter. Then (3) participants stood passively and were

asked to stop the experimenter where it was still comfortable for them, finally, (4)

participants stood passively and were asked to stop the experimenter where they thought it

was still comfortable for the experimenter. Participants repeated this procedure twice, with

and without eye contact: either the experimenter was looking at the participant (eye contact

condition), or at the papers she was holding (no eye contact condition). The order of these

two conditions was randomised across participants (Fig 1). During the statistical analysis

active and passive conditions were pooled (averaged) together.

Fig 1. Experimental setting. The modified version of the stop distance Paradigm. First, (1)
participants were approaching actively and were asked to stop where it was still comfortable
for them. Next (2) participants were approaching actively and were asked to stop where they
thought it was still comfortable for the experimenter. Then (3) participants stood passively
and were asked to stop the experimenter where it was still comfortable for them; finally (4)
participants stood passively and were asked to stop the experimenter where they thought it
was still comfortable for the experimenter. Participants repeated this procedure twice with
and without eye contact; the order of the latter two conditions was randomised across
participants.

196 Heart rate monitor

A wearable device Polar H10 was placed on participants' chests which recorded heart

rate (HR) during the whole experiment. We measured cardiac interbeat intervals (RR

intervals) using Polar H10 heart rate monitor chest strap (Polar Electro Oy, Kempele,

Finland) [47], which has been shown to be a valid device to measure RR interval signals [48].

The HR monitor was connected to a Samsung Galaxy Tablet via Bluetooth. We used the Elite

HRV application to export the recorded RR intervals as .txt files. We measured heart rate

variability (HRV) at two different conditions for a duration of 60 seconds: 1) at baseline and

2) during the intentional interpersonal distance task (1 minute after starting the distance task)

using the Root Mean Square of Successive RR interval Differences (RMSSD) method [49].

Additionally, we calculated RMSSD at the preceding ten-second time window of trigger

points set by researchers. These triggers corresponded to the time when participants arrived at

their final location of each condition.

Distance measuring: Obimon Prox

In order to synchronize the distance data with the HRV data, both the experimenter

and the participant wore a distance measuring device. The Obimon Prox [50] measures the

distance and the relative orientation between two wearable devices in real-time. The devices

use Ultra Wide Band (UWB) technology and the Symmetrical Double Sided Two-Way

Ranging (SDS-TWR) method [51] to both determine the distance between each other by

emitting very short and low power radio transmissions and measure the so-called time-of-

flight (ToF) with very high precision between transmission and reception. The resolution of

the measurement is in the range of a few centimetres, while the absolute precision is

approximately 10 centimetres. The relative orientation is defined as the difference between

the angles between the two devices taking the Earth's magnetic field as a reference. For

increased precision, the device uses sensor fusion involving magnetometer, accelerometer,

and gyroscope sensors. The results of the measurements are collected over Bluetooth LE

wireless technology to a laptop computer and evaluated in real-time.

Procedure

Participants wore a Polar H10 and Obimon Prox device during the whole experiment.

They placed the wearable device on themselves before the experiment started, then they

waited five minutes while calibrating and registering 60 seconds of resting heart rate and

HRV.

Next, participants completed the interpersonal distance task. Then, after a short break,

they completed a computerized neurocognitive test battery - measuring working memory,

executive functions, attention, inhibition, implicit learning, faux pas. These results are not

reported in this paper. Finally, they completed computerized versions of self-report

questionnaires (AQ: Autism-Spectrum Quotient, MZQ: Mentalization Questionnaire, AAS:

Adult Attachment Scale, ASRS: Adult ADHD Self-Report Scale, STAI-T: State-Trait

Anxiety Inventory – Trait; see Table 1 and Supporting information, S1 File).

To avoid the sensory over-reactivity effect, experimenters did not wear any jewellery,

perfume, and had been asked not to eat spicy food before the experiment, they wore simple,

casual, non-coloured clothes (jeans and black T-shirt). The room was curtained and

artificially evenly lit.

Data preprocessing and analysis

Preparation of HRV data was carried out using Python 3.7 with NumPy 1.20.1 [52],

pandas 1.2.3 [53], and SciPy 1.6.1 [54] data processing packages. Since the samples were

measured at a different rate for the Polar H10 (one sample per second) and the Obimon prox

(one sample per milliseconds) devices, we resampled the Obimon data by taking the median

for each second. Missing data were dropped from the analysis. To synchronize HRV with the

proximity data we needed to obtain the timestamps for each file containing the RR intervals.

The first timestamp was obtained from the name of the file which indicated the start time of

the recording. Since the exported files only contained the RR intervals without a timestamp

for each sample, the interval values themselves were used to create the time elapsed since the

first sample. As RR intervals annotate the time between two successive heartbeats, it was

possible to append the value of the RR interval to the time of the previous sample. After

obtaining the timestamps, data points were replaced with the median if they indicated RR of

1200 milliseconds (ms) or above, or if their absolute Z score was higher than 2. Triggers

added to the distance data (see Distance measuring) were adjusted manually if needed. HRV

was estimated as the root mean square of successive RR interval differences (RMSSD) since

this measurement is relatively resistant to by-products caused by breathing [55], and can be

obtained for a shorter (10 seconds) period of time [56]. Calculations were done by the

following formula (1):

$$RMSSD = \sqrt{\frac{1}{N-1}\sum_{i=0}^{n}(RR_{i+1} - RR_i)^2}$$ (1)

Baseline HR and HRV were measured and calculated for 60 seconds (s) at baseline,

reactive HR and HRV were measured during interpersonal distance task one minute after

starting the explicit paradigm (from +60 s to +120 s).

Furthermore, we calculated RMSSD around time points where interpersonal distance

data were reported. To calculate RMSSD for each explicit condition, eight local minimums of

the distance data were determined from data recorded by Obimon Prox. These eight time

points indicate the shortest distances between the participant and the experimenter,

corresponding to the time point when the reported distance was reached. RMSSD was

calculated for an interval starting 10 seconds prior to reaching the reported distance.

Statistical analysis

Statistical analysis was accomplished using R Version 3.6.3 [57], RStudio Version

1.2.1335 [58], and JASP Version 0.14.0.0 [59]. First, to measure if the two study groups do

differ regarding age, gender, education, caffeine intake, smoking, exercise and scores on

questionnaires, we conducted nonparametric Mann-Whitney U (Wilcoxon rank-sum) tests

and a Chi-square test. To measure the effect of different conditions and study groups on the

distance and HRV data, mixed-design ANOVA tests were applied, while in the case of

significant interaction effects post hoc tests with Bonferroni correction were used. As there

was already a baseline difference in HRV between the two groups, HRV values were

standardised in the interpersonal conditions for the further comparisons. Associations

between distance, HRV, and scores of psychometric questionnaires and diagnostic tests were

analysed with Spearman's rank-order correlations. Analyses were performed and

visualizations were created with R-packages *dplyr* [60], *ggplot2*[61], *psych* [62], *gridExtra*

[63], *ggpubr* [64], *readxl* [65], *corrplot* [66], *Hmisc* [67], *varian* [68].

Results

Is preferred interpersonal distance different in ASD?

Participants with ASD set significantly greater distances than NTP participants on a

group level. For the descriptive statistics of all conditions see supplementary material (Table

S1). To test if the different conditions had different modulatory effects in the two study

groups, we used two-way mixed-design ANOVA on the interpersonal distance as dependent

variable, where the between-subject factor was the Group (ASD/NTP), within-subject factor

Condition (Eye contact, Attribution). The Group main effect $F(1,41)=8.999$, $p=.005$,

$\eta^2_p=0.180$ was significant, participants with ASD preferred larger distances in general (mean

difference=35.980, 95% CI [11.757, 60.203])   (Fig 2). According to the post-hoc power

analysis, the group difference was detected with 98% power. Levene's test showed that the

variances were equal.

Fig 2. Interpersonal distance in cm. Dots represent the mean of distance data of eight
conditions for each individual. The top and the bottom of the box show the upper (Q3) and
lower (Q1) quartiles, the line dividing the box represents the median, and notches show a
95% confidence interval around the median. Asterix indicates significant group difference.
Orange: neurotypical participants, blue: participants with ASD.
301

Does eye contact or attribution affect interpersonal distance?

The main effect of the conditions of eye contact, or attribution resulted in the

following. Eye-contact ($M_{eye}(SD)$ = 88.756 (47.616), $M_{no\_eye}(SD)$ = 83.442 (40.116),

$F(1,41)_{eye\_cont}$ =3.005, $p=.091$, η^2_p =0.068, 95% CI [-0.859, 11.267]) showed a trend, and

attribution ($M_{self}(SD)$ = 84.913 (53.165), $M_{other}(SD)$ = 87.285 (37.378), $F(1,41)_{attrib}=0.248$,

p=.621, η^2_p =0.006, 95% CI [-12.691, 7.671]) did not have a significant main effect,

indicating that participants attributed similar comfortable personal distance to the

experimenter as to themselves (Fig 4 a-b).

310   In the further analysis, we tested if the conditions influence each other's effect. We

found a trend in Eye-contact × Attribution interaction (F(1,41)=3.011, p=.090, η^2_p =0.068)

but no significant Group × Condition interaction. Post-hoc test revealed that there was the

biggest difference between NTP and ASD participants when they had to maintain eye contact

and they had to set the comfortable distance for themselves (t=3.426, p=.030, 95% CI [2.248,

90.457]) after adjusting p-values for multiple comparisons by Bonferroni-correction in post

hoc tests.

Are heart rate and heart rate variability altered in ASD?

318   In general, participants with ASD had a slightly higher heart rate ($M_{baseline}$=90.65,

SD=12.95; $M_{interpersonal}$=96.66, SD=12.49) than NTP participants ($M_{baseline}$=87.06, SD=15.74;

$M_{interpersonal}$=91.68, SD=15.08) (Figure 3), however, these differences were not significant.

Measuring the effect of the conditions we used mixed-design ANCOVA, with the between-

subject variable of Group (ASD/NTP), and within-subject variable Condition

(baseline/interpersonal). As caffeine intake could influence the heart rate, we included actual

caffeine use as a covariate, but it did not change the results. The Group main effect was not

statistically significant ($F(1,35)$=0.875, p=.356, η^2_p =0.024), the main effect of Conditions

was significant (F(1,35)=38.068, p<.001, η^2_p =0.521), but the Condition × Group interaction

was not (F(1,35)=0.647, p=.427, η^2_p =0.018). It means, that in both groups we measured the

highest HR during the intentional interpersonal distance task, and it was significantly higher

than baseline (t_{ASD}=5.866, p<.001, 95% CI [2.554, 9.454], t_{NTP}=3.846, p=.003, 95% CI

[1.260, 7.977]) according to the post-hoc tests, where *p* values were adjusted by using

Bonferroni-correction. Levene's test showed that the variances were equal (Figure 3a).

332   Heart rate variability (HRV) was higher in the NTP group ($M_{baseline}$=48.26, *SD*=26.68;

$M_{interpersonal}$=31.60, *SD*=16.11) than in ASD ($M_{baseline}$=32.90, *SD*=15.97; $M_{interpersonal}$=26.57,

SD=8.79). Again, measuring the different effects of the conditions we used mixed-design

ANOVA, with the same within- and between-subject variables described above. The group

main effect ($F(1,35)$=3.470, *p*=.071, η^2_p =0.090) showed a trend. The condition main effect

($F(1,35)$=22.744, *p*<.001, η^2_p =0.394) and Group × Condition interaction ($F(1,35)$=4.598,

p=.039, η^2_p =0.116) were significant. Post-hoc test showed significant difference between

the baseline and interpersonal condition (*t*=4.769, *p*<.001, 95% CI [6.600, 16.383]), but this

difference originates from the significant difference within the NTP group (t_{NTP}=4.956,

p<.001, 95% CI [7.258, 26.059]), whereas HRV did not differ significantly in ASD group

between the two conditions (t_{ASD}=1.831, *p*=.276, 95% CI [-3.333, 15.982]) (Figure 3b). The

more pronounced difference between baseline HRV and HRV during the task (reflective

HRV) suggests a greater reactive capacity of autonomic regulation in NTPs.

Fig 3. Heart rate and heart rate variability. Panel A: Baseline and reactive (interpersonal
conditions) heart rate in beat per minute (bpm). Panel B: Baseline and reactive (interpersonal
conditions) heart rate variability (RMSSD). Error bars: standard error of the mean. Asterix
indicates significant group difference. Orange line: neurotypical participants, blue line:
participants with ASD.

Does eye contact or attribution affect interpersonal heart rate variability?

351   HRV during the interpersonal distance task (measured before the time point reported

distance was reached) was numerically higher in the NTP group, but the difference was not

significant between groups. For descriptive statistics see Supplementary material (Table S2).

As the baseline HRV was higher in NTP group, we used standardized HRV here. When we

tested the effect of the conditions and their interactions, neither the main effect of eye contact

or attribution, and their interaction with the group, nor the group effect ($F(1,32)=0.0002$,

$p=.988$) were significant using repeated-measures ANOVA. (Fig 4 c-d). Autonomic

functioning might be influenced by smoking, exercise, regular caffeine consumption, or the

actual caffeine intake before the experiment. There was no difference between groups (see

Table 1), however, including these variables as covariates did not change the results.

The interpersonal distance (Panels a-b) and heart rate variability data measured by the

RMSSD method (Panels c-d) are presented in Fig 4 to introduce their characteristics in

different conditions in the two study groups.

Fig 4. Interpersonal distance and HRV differences in different conditions. Panel a-b
Interpersonal distance in cm. Panel a: With or without eye contact, Panel b: Attribution to self
or the other. Panel c-d Heart rate variability in explicit conditions at a reported distance.
Panel c: With or without eye contact, Panel d: Attribution to self or the other. Error bars:
standard error of the mean. Asterix indicates significant group difference. Orange line:
neurotypical participants, blue line: participants with ASD.

Is there any correlation between HRV, distance, and psychometric data?

Exploratory analysis

Correlation analysis was highly exploratory, due to the small sample size, but it might

be suitable for further hypothesis generation. We found a relatively high consistency of

distance and HRV data (Fig 5), suggesting both the measured distance and HRV data indicate

reliably a similar construct. Intentional interpersonal distance and baseline HRV data show a

weak negative correlation in NTP participants, suggesting that higher heart rate variability

fosters prosocial activity and neurotypical participants with greater autonomic regulation

could approach the experimenter more (Fig 5, upper triangle). In the case of participants with

ASD, however, we found less consistent association between distance and HRV results (Fig

5, lower triangle). Correlation between the mean interpersonal distance and HRV during the

interpersonal distance task, was not significant, however, it tend to point in different

directions in the two groups (Fig 6).

Fig 5. Correlations between interpersonal distance and heart rate variability at the
baseline and during the intentional interpersonal distance conditions. Dist = distance,
HRV = heart rate variability, preHRV = 10s RMSSD, Eye = eye contact, No eye = no eye
contact, Active = active moving, Passive = standing, Self = attribution to self, Other =
attribution to the other conditions. Upper triangle: neurotypical participants, lower triangle:
participants with ASD. Warm colors refer to positive, cold colors refer to negative Spearman
rank correlation *rho* values, grey asterisk marks the significant *p* values after (fdr) correcting
for multiple comparisons.

Fig 6. Correlation between mean distance (in cms) and HRV (60 s) during interpersonal
distance task in the two groups. Orange line: neurotypical participants, blue line:
participants with ASD

Results of psychometric questionnaires showed weak or no association with distance

and HRV results (Fig 7); however, the association with psychometric questionnaires in ASD

showed a different pattern than in NTP. High trait anxiety level, poor mentalization, and

attachment were weakly associated with greater interpersonal distance ASD (Fig 7, lower

triangle, first column), but neither of these correlations remained significant after correction

for multiple comparisons, only HRV at baseline and during the interpersonal condition, AQ

and mentalization scores were correlated in ASD group.

Fig 7. Correlations between interpersonal distance, heart rate variability at the baseline
and during the interpersonal distance conditions, and psychometric data. HRV = heart
rate variability, AQ = Autism-spectrum Quotient, AAS = Adult Attachment Scale, MZQ =
Mentalization Questionnaire, STAI-T = State-Trait Anxiety Inventory, Trait, ADI = Autism
Diagnostic Interview-Revised, ADOS = Autism Diagnostic Observation Schedule. Upper
triangle: neurotypical participants, lower triangle: participants with ASD. Warm colors refer
to positive, cold colors refer to negative Spearman rank correlation *rho* values, grey asterisk
marks the significant *p* values after (fdr) correcting for multiple comparisons.

Discussion

Our study aimed to investigate interpersonal distance regulation and the underlying

autonomic response regulation in autism spectrum disorder. To this end, we introduced a

methodology combining interpersonal distance measurement and physiological parameter

registration in an interpersonal experimental setting in groups of adult participants with ASD

and their matched neurotypical controls. We found increased interpersonal distance in ASD.

The difference was the most pronounced when the experimenter maintained eye contact and

requested to determine the comfortable distance for themselves during the interpersonal

distance task. Moreover, we found decreased baseline heart rate variability and decreased

HRV reactivity in ASD.

Interpersonal distance was measured using a modified version of the stop distance

paradigm to assess how far participants prefer to stand from another person and whether there

is a difference between ASD and NTP group in this respect. Participants were directly

instructed to define a still comfortable distance from the experimenter. Usually, in a stop

distance paradigm, the participant and the experimenter are facing each other at the endpoints

of a 300 to 600 cm long line along which the participants set their preferred interpersonal

distance. In our experiment, we have chosen 500 cm as the initial distance, which includes all

four distance zones of interpersonal space (intimate, personal, social and public) according to

Hall's proxemic rules [69]. During this task, participants set distances on average within the

personal space (far zone ~75 - 120 cm, close zone ~45 - 75 cm, the zones between the

intimate distance and personal distance). However, as expected, participants with ASD set

significantly greater distances than NTPs: around the far end of the personal space, or even

farther. The social space (the zone between personal and social distance 120 - 370 cm) is

reserved for strangers or new acquaintances [69]. We can speculate that this display can

change the non-verbal message conveyed by a set distance in real-life interpersonal

situations.

435   Prior to our experiment, five studies examined interpersonal space regulation in ASD,

four of which studied children or adolescent populations. These findings uniformly suggest

that interpersonal space regulation is altered in ASD in childhood, autistic children preferred

significantly larger interpersonal distance than neurotypical control participants [16,22]. A

study examining adolescents with ASD also concluded that their space regulation was

altered. Interestingly, this conclusion was derived from opposing results: adolescents with

ASD preferred shorter interpersonal distance than neurotypical controls [19]. Although, in

this study from Japan, neurotypical participants preferred longer personal distances (cca. 130

- 150 cm depending on condition) than in other cohorts, and the distances preferred by ASD

participants were comparable to our and Kennedy et al.'s results, the only adult study

observed no differences between participants with ASD and neurotypical controls [18] (cca.

70 – 100 cm). This raises the possibility of cultural differences in social rules and customs,

including personal space arrangement. Our results suggested that adult individuals (a

homogenous white Caucasian, Central European sample) with ASD prefer greater

interpersonal distance than neurotypicals.

450   Several factors can influence the interpersonal distance between the experimenter and

the participants [13,19,22]. Eye contact has been shown to affect the preferred interpersonal

distance of ASD and neurotypical adolescents: in eye contact conditions when participants

held passive roles, they preferred larger interpersonal distance, regardless of which study

group they belonged to, and this effect did not emerge when holding active roles [19]. In our

study we found a weak modulatory effect of eye contact, participants with ASD tend to set

slightly greater distances in conditions with eye contact. We can speculate that eye contact

might cause distress and evoke avoidance in ASD, instead of an appetitive or comforting

positive reinforcement, in line with the "eye avoidance" hypothesis [33,34]. Future studies

directly examining this phenomenon and testing whether and how eye contact plays a role,

seem warranted.

Reciprocal social interactions are impaired in ASD, leading to a weaker adaptation to

another person's perspective. Therefore, we also introduced a condition that requires higher

order mentalization. We tested what the participant thought was a comfortable distance for

the experimenter. Considering the theory of mind difficulties in ASD, we hypothesized that

participants with ASD will show less difference along the attribution dimension than NTPs.

We found no difference in ASD accordingly, although there was no significant difference

between the self and other attribution conditions in the NTP group either. It is interesting,

however, that we measured the biggest distance in ASD and biggest difference between NTP

and ASD participants when they had to maintain eye contact and they had to set a

comfortable distance for themselves. These results might suggest that participants with ASD

are capable of modifying their behaviour to some extent according to others' aspect, in

contrast with previous results [70,71], but to test this hypothesis further studies are needed.

An invention in our experimental design is that we combined interpersonal distance

measurement with heart rate registration. In social behaviour parasympathetic regulation, the

flexibility of vagal tone plays an important role according to Porges' polyvagal theory [72–

74]. Higher resting HRV was found to be associated with cooperative behaviour, using less

disengagement and more socially adaptive emotion regulation strategies among healthy

adults [75,76]. Variables influenced by parasympathetic regulation (e.g. respiratory sinus

arrhythmia) are related to emotion recognition and symptom severity in ASD [77]. In line

with the results of previous studies corroborating altered autonomic nervous system

functioning in ASD [41,44,78], we found reduced baseline heart rate variability in

participants with ASD spectrum disorder. The average heart rate (reflecting sympathetic

activity) was slightly higher in the ASD group than in neurotypicals, but this difference was

not significant.

In social situations, not just the baseline but also the regulatory capacity must be taken

into account. Previous studies showed that the HRV decrease induced by participating in a

social situation was lower, assuming a decreased regulatory capacity in the ASD group [44].

We used the RMSSD method to measure HRV in order to capture the parasympathetic

regulation rather than sympathetic arousal [49]. Psychotropic medication, caffeine intake,

smoking, and exercising habits might influence autonomic regulation [79–82], but study

groups in this study did not differ in actual or regular caffeine intake, smoking and exercising

habits (Table 1). There were no significant differences between participants who took

medication and those who did not, consequently, we argue that the results are attributable to

autism. We found a significant HRV decrease in interpersonal setting compared to the

baseline in NTP, but not in ASD group, which confirms previous research findings of

decreased regulatory capacity of participants with ASD in social situations.

Additionally, we calculated HRV during the social distance regulation task in 10 s

time periods, applying ultra-short-term analysis [56] exactly at the time point when

participants arrived at the reported distance in order to take a closer look at the relationship of

interpersonal distance and autonomic regulation. To test experimentally the more nuanced

aspects of interpersonal distance regulation, we assessed the modulatory effect of eye contact

and attribution. We did not find a significant difference between study conditions regarding

the 10 s HRV metrics, and results did not directly support our hypothesis that HRV predicts

the interpersonal distance. Nevertheless, we can speculate that due to the inherently lower

baseline HRV in ASD, the diminished capacity of reactive decrement might have prevented

the further fine-tuning during the interpersonal task. Reduced regulatory capacity, combined

with elevated amygdala reactivity could lead to early exhaustion even during minimal social

interaction. This can raise the possibility that the larger interpersonal distance is the

consequence of the early exhaustion of regulatory capacity, and by keeping the distance they

might avoid a more severe autonomic disturbance in social situations. To further test this

hypothesis in real-life situations, applying widely available wearable devices might be useful.

This experience gained could later be used, for example, to develop biofeedback tools for

social communication training for people with autism spectrum disorder.

Limitations and further directions

Despite the most careful planning, every study has its limitations. In this study, we

examined adult participants with ASD trying to fill a gap in this research area and measure

interpersonal distance and autonomic regulation simultaneously. We recruited participants

with average or above-average intellectual abilities, which increases the likelihood of

adaptive skill acquisition. To overcome this limitation inclusion of a broader spectrum of

autistic participants is needed in future studies.

We found greater interpersonal distance in ASD measured by the modified version of

the stop distance paradigm, but there was no difference between study groups regarding heart

rate variability during that part of the experiment. In subsequent research, HRV differences

should be measured at fixed distances as well, e.g. closer than the comfortable range for

neurotypicals, farther than the comfortable range for participants with ASD, and at

intermediate distances. Subjective rating of the level of comfort (both by participants and by

the experimenter) might help to gain a better insight of how correctly the experimenter's

perspective can be estimated by the participants.

Unfortunately, we have had to finish the data collection due to the beginning of the

COVID-pandemic. As the introduced social distancing measures deeply affect our senses of

interpersonal distance, we cannot continue with the original study design. We plan to

investigate the effect of this unforeseen but global intervention by repeating our experiment

when the pandemic subsides, and the social aspects of the situation are consolidated.

Additional conditions with and without wearing face masks might be considered, too. These

subsequent studies will be able to show us whether autistic people have been affected

differently than neurotypicals by social distancing measures.

Conclusion

Interpersonal distance regulation is a relevant nonverbal part of social

communication. It reflects the individual needs for personal space and the ability to read

others' intentions. Together with other biomarkers of autonomic functions, this might express

how demanding a simple social interaction can be for people with ASD. In this study, we

introduced a new experimental design to measure these factors together, in a basic social

interaction setting. Participants with ASD preferred larger interpersonal distance. The

difference was most pronounced when they had to approach the experimenter actively with

maintained eye contact. Participants with ASD had lower baseline heart rate variability and

decreased heart rate variability reactivity than neurotypicals; however, their HRV and its

changes in different conditions during the social distance task did not differ significantly.

This raises the possibility that regulatory capacities were exhausted sooner, at a farther

distance in this group, as part of the compensatory avoidant behaviour in ASD. We believe

that applying this experimental design could also be beneficial in studying other psychiatric

conditions, such as borderline personality disorder, social phobia, or psychosis. These results

could further expand our understanding of interpersonal distance regulation in autism

spectrum disorder.

Acknowledgment

This research was supported by the National Brain Research Program (project 2017-1.2.1-NKP- 2017-00002, PI: D. N.); the Hungarian Scientific Research Fund (OTKA K 128016, PI: D. N., OTKA PD 124148, PI: K. J.); the János Bolyai Research Scholarship of the Hungarian Academy of Sciences (to K. J.); the New National Excellence Program of the Ministry for Innovation and Technology (ÚNKP-19-2-I-ELTE-332, A. G.); the IDEXLYON Fellowship of the University of Lyon as part of the Programme Investissements d'Avenir (ANR-16-IDEX-0005 to D. N.) and Higher Education Institutional Excellence Programme of the Ministry of Human Capacities in Hungary, within the framework of the Neurology thematic program of Semmelweis University (to K. F., E. K., B. Sz., J. R.).

References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. Fifth
Edition. American Psychiatric Association; 2013
https://doi.org/10.1176/appi.books.9780890425596

2. Ecker C. The neuroanatomy of autism spectrum disorder: An overview of structural
neuroimaging findings and their translatability to the clinical setting. Autism Int J Res Pract.
2017;21(1):18–28. https://doi.org/10.1177/1362361315627136

3. Herringshaw AJ, Ammons CJ, DeRamus TP, Kana RK. Hemispheric differences in language
processing in autism spectrum disorders: A meta-analysis of neuroimaging studies. Autism Res
Off J Int Soc Autism Res. 2016;9(10):1046–57. https://doi.org/10.1002/aur.1599

4. Hadjikhani N, Åsberg Johnels J, Zürcher NR, Lassalle A, Guillon Q, Hippolyte L, et al. Look
me in the eyes: constraining gaze in the eye-region provokes abnormally high subcortical
activation in autism. Sci Rep. 2017;7(1):3163. https://doi.org/10.1038/s41598-017-03378-5

5. Langen M, Bos D, Noordermeer SDS, Nederveen H, van Engeland H, Durston S. Changes in
the development of striatum are involved in repetitive behavior in autism. Biol Psychiatry.
2014;76(5):405–11. https://doi.org/10.1016/j.biopsych.2013.08.013

6. Gomez IN, Flores JG. Diverse Patterns of Autonomic Nervous System Response to Sensory
Stimuli Among Children with Autism. Curr Dev Disord Rep 2020;7(4):249–57.
https://doi.org/10.1007/s40474-020-00210-0

7. Kushki A, Brian J, Dupuis A, Anagnostou E. Functional autonomic nervous system profile in
children with autism spectrum disorder. Mol Autism. 2014;5(1):39.
https://doi.org/10.1186/2040-2392-5-39

8. Philip RCM, Dauvermann MR, Whalley HC, Baynham K, Lawrie SM, Stanfield AC. A
systematic review and meta-analysis of the fMRI investigation of autism spectrum disorders.
Neurosci Biobehav Rev. 2012;36(2):901–42. https://doi.org/10.1016/j.neubiorev.2011.10.008

9. Thye MD, Bednarz HM, Herringshaw AJ, Sartin EB, Kana RK. The impact of atypical sensory
processing on social impairments in autism spectrum disorder. Dev Cogn Neurosci.
2018;29:151–67. https://doi.org/10.1016/j.dcn.2017.04.010

10. Wood ET, Cummings KK, Jung J, Patterson G, Okada N, Guo J et al. Sensory over-responsivity
is related to GABAergic inhibition in thalamocortical circuits. Transl Psychiatry. 2021;11(1):1–
10. https://doi.org/10.1038/s41398-020-01154-0

11. Demetriou EA, DeMayo MM, Guastella AJ. Executive Function in Autism Spectrum Disorder:
History, Theoretical Models, Empirical Findings, and Potential as an Endophenotype. Front
Psychiatry. 2019;10:753. https://doi.org/10.3389/fpsyt.2019.00753

12. Hill EL. Executive dysfunction in autism. Trends Cogn Sci. 2004;8(1):26–32.
https://doi.org/10.1016/j.tics.2003.11.003

13. Candini M, di Pellegrino G, Frassinetti F. The plasticity of the interpersonal space in autism
spectrum disorder. Neuropsychologia. 2020;147:107589.
https://doi.org/10.1016/j.neuropsychologia.2020.107589

14.   Massaccesi C, Groessing A, Rosenberger LA, Hartmann H, Candini M, di Pellegrino G et al.
604   Neural Correlates of Interpersonal Space Permeability and Flexibility in Autism Spectrum
605   Disorder. Cereb Cortex. 2021; 31(6): 2968–2979 https://doi.org/10.1093/cercor/bhaa404

606   15.   Lough E, Hanley M, Rodgers J, South M, Kirk H, Kennedy DP, Riby DM. Violations of
607   Personal Space in Young People with Autism Spectrum Disorders and Williams Syndrome:
608   Insights from the Social Responsiveness Scale. J Autism Dev Disord. 2015;45(12):4101–8.
609   https://doi.org/10.1007/s10803-015-2536-0

610   16.   Gessaroli E, Santelli E, di Pellegrino G, Frassinetti F. Personal Space Regulation in Childhood
611   Autism Spectrum Disorders. PLoS ONE. 2013;8(9) e74959
612   https://doi.org/10.1371/journal.pone.0074959

613   17.   Perry A, Levy-Gigi E, Richter-Levin G, Shamay-Tsoory SG. Interpersonal distance and social
614   anxiety in autistic spectrum disorders: A behavioral and ERP study. Soc Neurosci.
615   2015;10(4):354–65. https://doi.org/10.1080/17470919.2015.1010740

616   18.   Kennedy DP, Adolphs R. Violations of Personal Space by Individuals with Autism Spectrum
617   Disorder. PLoS ONE. 2014;9(8). https://doi.org/10.1371/journal.pone.0103369

618   19.   Asada K, Tojo Y, Osanai H, Saito A, Hasegawa T, Kumagaya S. Reduced Personal Space in
619   Individuals with Autism Spectrum Disorder. PLoS ONE. 2016;11(1).
620   https://doi.org/10.1371/journal.pone.0146306

621   20.   Ingram DH, Mayes SD, Troxell LB, Calhoun SL. Assessing children with autism, mental
622   retardation, and typical development using the Playground Observation Checklist. Autism Int J
623   Res Pract. 2007;11(4):311–9. https://doi.org/10.1177/1362361307078129

624   21.   Pedersen J, Schelde T. Behavioral aspects of infantile autism: an ethological description. Eur
625   Child Adolesc Psychiatry. 1997;6(2):96–106. https://doi.org/10.1007/BF00566672

626   22.   Candini M, Giuberti V, Manattini A, Grittani S, Pellegrino G di, Frassinetti F. Personal space
627   regulation in childhood autism: Effects of social interaction and person's perspective. Autism
628   Res. 2017;10(1):144–54. https://doi.org/10.1002/aur.1637

629   23.   Mul C, Cardini F, Stagg SD, Sadeghi Esfahlani S, Kiourtsoglou D, Cardellicchio P, Aspell JE.
630   Altered bodily self-consciousness and peripersonal space in autism. Autism [Internet].
631   2019;23(8):2055–67. https://doi.org/10.1177/1362361319838950

632   24.   Simões M, Mouga S, Pereira AC, de Carvalho P, Oliveira G, Castelo-Branco M. Virtual Reality
633   Immersion Rescales Regulation of Interpersonal Distance in Controls but not in Autism
634   Spectrum Disorder. J Autism Dev Disord. 2020;50(12):4317–28.
635   https://doi.org/10.1007/s10803-020-04484-6

636   25.   Kennedy DP, Gläscher J, Tyszka JM, Adolphs R. Personal Space Regulation by the Human
637   Amygdala. Nat Neurosci. 2009;12(10):1226–7. https://doi.org/10.1038/nn.2381

638   26.   Baron-Cohen S. Mindblindness: An Essay on Autism and Theory of Mind. MIT Press; 1997.
639   206 p. https://doi.org/10.7551/mitpress/4635.003.0001

640   27.   Baron-Cohen S, Leslie AM, Frith U. Does the autistic child have a 'theory of mind'? Cognition.
641   1985;21(1):37–46. https://doi.org/10.1016/0010-0277(85)90022-8

642   28.   Black MH, Chen NTM, Iyer KK, Lipp OV, Bölte S, Falkmer M, Tan T, Girdler S. Mechanisms
643   of facial emotion recognition in autism spectrum disorders: Insights from eye tracking and

644   electroencephalography. Neurosci Biobehav Rev. 2017;80:488–515.
645   https://doi.org/10.1016/j.neubiorev.2017.06.016

646   29.   Monteiro R, Simões M, Andrade J, Castelo Branco M. Processing of Facial Expressions in
647   Autism: a Systematic Review of EEG/ERP Evidence. Rev J Autism Dev Disord.
648   2017;4(4):255–76. https://doi.org/10.1007/s40489-017-0112-6

649   30.   Teh EJ, Yap MJ, Rickard Liow SJ. Emotional Processing in Autism Spectrum Disorders:
650   Effects of Age, Emotional Valence, and Social Engagement on Emotional Language Use. J
651   Autism Dev Disord. 2018;48(12):4138–54. https://doi.org/10.1007/s10803-018-3659-x

652   31.   Wieckowski AT, Flynn LT, Richey JA, Gracanin D, White SW. Measuring change in facial
653   emotion recognition in individuals with autism spectrum disorder: A systematic review. Autism
654   Int J Res Pract. 2020;24(7):1607–28. https://doi.org/10.1177/1362361320925334

655   32.   Dalton KM, Nacewicz BM, Johnstone T, Schaefer HS, Gernsbacher MA, Goldsmith HH,
656   Alexander AL, Davidson RJ. Gaze fixation and the neural circuitry of face processing in autism.
657   Nat Neurosci. 2005;8(4):519–26. https://doi.org/10.1038/nn1421

658   33.   Tanaka JW, Sung A. The "eye avoidance" hypothesis of autism face processing. J Autism Dev
659   Disord. 2016;46(5):1538–52. https://doi.org/10.1007/s10803-013-1976-7

660   34.   Tottenham N, Hertzig ME, Gillespie-Lynch K, Gilhooly T, Millner AJ, Casey BJ. Elevated
661   amygdala response to faces and gaze aversion in autism spectrum disorder. Soc Cogn Affect
662   Neurosci. 2014; 9(1):106–17. https://doi.org/10.1093/scan/nst050

663   35.   Joseph RM, Ehrman K, McNally R, Keehn B. Affective response to eye contact and face
664   recognition ability in children with ASD. J Int Neuropsychol Soc JINS. 2008;14(6):947–55.
665   https://doi.org/10.3390/s21155166

666   36.   Madipakkam AR, Rothkirch M, Dziobek I, Sterzer P. Unconscious avoidance of eye contact in
667   autism spectrum disorder. Sci Rep. 2017;7(1):13378. https://doi.org/10.1038/s41598-017-
668   13945-5

669   37.   Frank DW, Dewitt M, Hudgens-Haney M, Schaeffer DJ, Ball BH, Schwarz NF, et al. Emotion
670   regulation: Quantitative meta-analysis of functional activation and deactivation. Neurosci
671   Biobehav Rev. 2014;45:202–11. https://doi.org/10.1016/j.neubiorev.2014.06.010

672   38.   LeDoux J. The amygdala. Curr Biol. 2007;17(20):R868–74.
673   https://doi.org/10.1016/j.cub.2007.08.005

674   39.   Lydon S, Healy O, Reed P, Mulhern T, Hughes BM, Goodwin MS. A systematic review of
675   physiological reactivity to stimuli in autism. Dev Neurorehabilitation. 2016;19(6):335–55.
676   https://doi.org/10.3109/17518423.2014.971975

677   40.   Hutt SJ, Hutt C, Lee D, Ounsted C. A behavioural and electroencephalographic study of autistic
678   children. J Psychiatr Res. 1965;3(3):181–97. https://doi.org/10.1016/0022-3956(65)90028-2

679   41.   Arora I, Bellato A, Ropar D, Hollis C, Groom MJ. Is autonomic function during resting-state
680   atypical in Autism: A systematic review of evidence. Neurosci Biobehav Rev. 2021;125:417–
681   41. https://doi.org/10.1016/j.neubiorev.2021.02.041

682   42.   Laborde S, Mosley E, Thayer JF. Heart Rate Variability and Cardiac Vagal Tone in
683   Psychophysiological Research – Recommendations for Experiment Planning, Data Analysis,
684   and Data Reporting. Front Psychol. 2017;8:213. https://doi.org/10.3389/fpsyg.2017.00213

43.   Colzato LS, Jongkees BJ, de Wit M, van der Molen MJW, Steenbergen L. Variable heart rate
and a flexible mind: Higher resting-state heart rate variability predicts better task-switching.
Cogn Affect Behav Neurosci. 2018;18(4), 730–738 https://doi.org/10.3758/s13415-018-0600-x

44.   Cheng Y-C, Huang Y-C, Huang W-L. Heart rate variability in individuals with autism spectrum
disorders: A meta-analysis. Neurosci Biobehav Rev. 2020;118:463–71.
https://doi.org/10.1016/j.neubiorev.2020.08.007

45.   Lord C, Rutter M, Le Couteur A. Autism Diagnostic Interview-Revised: a revised version of a
diagnostic interview for caregivers of individuals with possible pervasive developmental
disorders. J Autism Dev Disord. 1994;24(5):659–85. https://doi.org/10.1007/BF02172145

46.   Lord C, Rutter M, Goode S, Heemsbergen J, Jordan H, Mawhood L, Schopler E. Autism
diagnostic observation schedule: a standardized observation of communicative and social
behavior. J Autism Dev Disord. 1989;19(2):185–212. https://doi.org/10.1007/BF02211841

47.   Saario T. Polar Electro Oy. Kempele, Finland: Suncom Systems; 2019. www.polar.com

48.   Gilgen-Ammann R, Schweizer T, Wyss T. RR interval signal quality of a heart rate monitor and
an ECG Holter at rest and during exercise. Eur J Appl Physiol. 2019;119(7):1525–32.
https://doi.org/10.1007/s00421-019-04142-5%20

49.   Shaffer F, Ginsberg JP. An Overview of Heart Rate Variability Metrics and Norms. Front Public
Health. 2017;5. https://doi.org/10.3389/fpubh.2017.00258

50.   Veres A. Obimon Prox [Internet]. Obimon Systems Ltd.; 2019. http://obimon.com

51.   Neirynck. An alternative double-sided two-way ranging method. 13th Workshop on Positioning,
Navigation and Communications (WPNC); 2016.

52.   Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D et al. Array
programming with NumPy. Nature. 2020;585(7825):357–62. https://doi.org/10.1038/s41586-
020-2649-2

53.   Reback J, McKinney W, jbrockmendel, Van den Bossche J, Augspurger T, Cloud P et al.
pandas-dev/pandas: Pandas 1.0.3. Zenodo; 2020.
https://zenodo.org/record/3715232#.YKkKa6j7SUm

54.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0:
fundamental algorithms for scientific computing in Python. Nat Methods. 2020;17(3):261–72.
https://doi.org/10.1038/s41592-019-0686-2

55.   Hill LK, Siebenbrock A. Are all measures created equal? Heart rate variability and respiration -
biomed 2009. Biomed Sci Instrum. 2009;45:71–6. PMID: 19369742

56.   Salahuddin L, Cho J, Jeong MG, Kim D. Ultra short term analysis of heart rate variability for
monitoring mental stress in mobile settings. Annu Int Conf IEEE Eng Med Biol Soc IEEE Eng
Med Biol Soc Annu Int Conf. 2007;2007:4656–9. https://doi.org/10.3389/fnins.2020.594880

57.   R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2020. https://www.R-project.org/

58.   RStudio Team. RStudio: Integrated Development for R. 2020. https://rstudio.com/

59.   JASP Team. JASP. 2020. https://jasp-stats.org/

60.   Wickham H, François R, Henry L, Müller K, RStudio. dplyr: A Grammar of Data Manipulation.
725   2020. https://CRAN.R-project.org/package=dplyr

726   61.   Kassambara A. 'ggplot2' Based Publication Ready Plots [R package ggpubr version 0.4.0].
727   Comprehensive R Archive Network (CRAN); 2020. https://CRAN.R-
728   project.org/package=ggpubr

729   62.   Revelle W. Procedures for Psychological, Psychometric, and Personality Research [R package
730   psych version 2.1.3]. Comprehensive R Archive Network (CRAN); 2021. https://CRAN.R-
731   project.org/package=psych

732   63.   Auguie B. Miscellaneous Functions for 'Grid' Graphics [R package gridExtra version 2.3].
733   Comprehensive R Archive Network (CRAN); 2017 https://CRAN.R-
734   project.org/package=gridExtra

735   64.   Kassambara A. ggpubr: 'ggplot2' Based Publication Ready Plots 2020 https://CRAN.R-
736   project.org/package=ggpubr

737   65.   Wickham H, Bryan J, readxl: Read Excel Files 2019. https://CRAN.R-
738   project.org/package=readxl

739   66.   Wei T, Simko V, Levy M, Xie Y, Jin Y, Zemla J. corrplot: Visualization of a Correlation
740   Matrix. 2017. https://CRAN.R-project.org/package=corrplot

741   67.   Harrell FE et al. Hmisc: Harrell Miscellaneous. 2020. https://CRAN.R-
742   project.org/package=Hmisc

743   68.   Wiley J. varian: Variability Analysis in R. Comprehensive R Archive Network (CRAN); 2016.
744   https://CRAN.R-project.org/package=varian

745   69.   Hall ET. The Hidden Dimension. Anchor Books; 1966.

746   70.   Frith U, Morton J, Leslie AM. The cognitive basis of a biological disorder: autism. Trends
747   Neurosci. 1991;14(10):433–8. https://doi.org/10.1016/0166-2236(91)90041-R

748   71.   Livingston LA, Carr B, Shah P. Recent Advances and New Directions in Measuring Theory of
749   Mind in Autistic Adults. J Autism Dev Disord. 2019;49(4):1738–44.
750   https://doi.org/10.1007/s10803-018-3823-3

751   72.   Muscatello RA, Vandekar SN, Corbett BA. Evidence for decreased parasympathetic response to
752   a novel peer interaction in older children with autism spectrum disorder: a case-control study. J
753   Neurodev Disord. 2021;13. https://doi.org/10.1186/s11689-020-09354-x

754   73.   Porges SW. The Polyvagal Theory: phylogenetic contributions to social behavior. Physiol
755   Behav. 2003;79(3):503–13. https://doi.org/10.1016/S0031-9384(03)00156-2

756   74.   Porges SW. The polyvagal theory: New insights into adaptive reactions of the autonomic
757   nervous system. Cleve Clin J Med. 2009;76:S86–90. https://doi.org/10.3949/ccjm.76.s2.17

758   75.   Beffara B, Bret AG, Vermeulen N, Mermillod M. Resting high frequency heart rate variability
759   selectively predicts cooperative behavior. Physiol Behav. 2016; 1;164(Pt A):417–28.
760   https://doi.org/10.1016/j.physbeh.2016.06.011

76.   Geisler FCM, Kubiak T, Siewert K, Weber H. Cardiac vagal tone is associated with social
engagement and self-regulation. Biol Psychol. 2013;93(2):279–86.
https://doi.org/10.1016/j.biopsycho.2013.02.013

77.   Bal E, Harden E, Lamb D, Van Hecke AV, Denver JW, Porges SW. Emotion Recognition in
Children with Autism Spectrum Disorders: Relations to Eye Gaze and Autonomic State. J
Autism Dev Disord. 2010;40(3):358–70. https://doi.org/10.1007/s10803-009-0884-3

78.   Thapa R, Alvares GA, Zaidi TA, Thomas EE, Hickie IB, Park SH, Guastella AJ. Reduced heart
rate variability in adults with autism spectrum disorder. Autism Res. 2019;12(6):922–30.
https://doi.org/10.1002/aur.2104

79.   Alvares GA, Quintana DS, Hickie IB, Guastella AJ. Autonomic nervous system dysfunction in
psychiatric disorders and the impact of psychotropic medications: a systematic review and meta-
analysis. J Psychiatry Neurosci JPN. 2016;41(2):89–104. https://doi.org/10.1503/jpn.140217

80.   Benjamim CJR, Kliszczewicz B, Garner DM, Cavalcante TCF, da Silva AAM, Santana MDR,
Valenti VE. Is Caffeine Recommended Before Exercise? A Systematic Review To Investigate
Its Impact On Cardiac Autonomic Control Via Heart Rate And Its Variability. J Am Coll Nutr.
2020;39(6):563–73. https://doi.org/10.1080/07315724.2019.1705201

81.   Hautala AJ, Kiviniemi AM, Tulppo MP. Individual responses to aerobic exercise: the role of the
autonomic nervous system. Neurosci Biobehav Rev. 2009;33(2):107–15.
https://doi.org/10.1016/j.neubiorev.2008.04.009

82.   Middlekauff HR, Park J, Moheimani RS. Adverse effects of cigarette and noncigarette smoke
exposure on the autonomic nervous system: mechanisms and implications for cardiovascular
risk. J Am Coll Cardiol. 2014; 21;64(16):1740–50. https://doi.org/10.1016/j.jacc.2014.06.1201

Supporting information

S1 Table. Social distance in intentional conditions. Descriptive statistics and group differences.
ASD: Autism Spectrum Disorder, NTP: Neurotypical Participant, N: sample size, SD: standard deviation

S2 Table. HRV in explicit conditions. Descriptive statistics and group differences.
ASD: Autism Spectrum Disorder, NTP: Neurotypical Participant, N: sample size, SD: standard deviation

S1 File. Questionnaires

a

b

c

d

Distance (cm)

Distance (cm)

HRV (RMSSD) transformed

HRV (RMSSD) transformed

Eye contact

Attribution

Eye contact

Attribution

Eye contact

No eye contact

Self

Other

Eye contact

No eye contact

Self

Other

NTP

ASD