MAVS expression in alveolar macrophages is essential for host resistance against *Aspergillus fumigatus*

Xi Wang[1], Cristina Cunha[2,3], Madeleine S. Grau[1], Shelly J. Robertson[4], João F. Lacerda[5,6], António Campos Jr[7], Katrien Lagrou[8,9], Johan Maertens[8,10], Sonja M. Best[4], Agostinho Carvalho[2,3], Joshua J. Obar[1,#]

[1]Geisel School of Medicine at Dartmouth, Department of Microbiology & Immunology, Lebanon, NH, USA

[2]Life and Health Sciences Research Institute (ICVS), School of Medicine, University of Minho, Braga, Portugal

[3]ICVS/3B's – PT Government Associate Laboratory, Braga/Guimarães, Portugal

[4]Laboratory of Virology, Rocky Mountain Laboratories, National Institute of Allergy and Infectious Diseases, NIH, Hamilton MT, USA.

[5]Instituto de Medicina Molecular, Faculdade de Medicina de Lisboa, Universidade de Lisboa, Lisboa, Portugal.

[6]Serviço de Hematologia e Transplantação de Medula, Hospital de Santa Maria, Lisboa, Portugal

[7]Serviço de Transplantação de Medula Óssea (STMO), Instituto Português de Oncologia do Porto, Porto, Portugal

[8]Department of Microbiology, Immunology and Transplantation, KU Leuven, Leuven, Belgium

[9]Clinical Department of Laboratory Medicine and National Reference Center for Medical Mycology, University Hospitals Leuven, Leuven, Belgium

[10]Department of Hematology, University Hospitals Leuven, Leuven, Belgium

Conflict of Interests: The authors have declared that no conflict of interest exists.

[#]Corresponding author:
Joshua J. Obar
Geisel School of Medicine at Dartmouth
Department of Microbiology & Immunology
1 Medical Center Drive
Lebanon, NH 03756
Telephone: +1 (603) 646-5384
Email: joshua.j.obar@dartmouth.edu

ABSTRACT

Our recent data demonstrates a critical role of the RIG-I-like receptor (RLR) family in regulating antifungal immunity against *Aspergillus fumigatus* in a murine model. However, the importance of this pathway in humans and the cell type(s) which utilize this innate immune receptor to detect *A. fumigatus* remains unresolved. Here we demonstrate that a polymorphism in human *MAVS* present in the donor genome was associated with the incidence of invasive pulmonary aspergillosis (IPA) in recipients of hematopoietic stem cell transplantation (HSCT). Moreover, in a separate cohort of confirmed IPA patients, polymorphisms in the *IFIH1* gene alter the inflammatory response, including interferon-responsive chemokines. Returning to our murine model, we now demonstrate that CD11c$^+$ alveolar macrophages require *Mavs* expression to maintain host resistance against *A. fumigatus.* Our data support the role of MAVS signaling in mediating antifungal immunity in both mice and human at least in part through the role of MAVS-dependent signaling in alveolar macrophages.

INTRODUCTION

Aspergillus fumigatus is a ubiquitous environmental mold that humans inhale on a daily basis. Individuals with normal immune systems readily clear *A. fumigatus* conidia from their airways without complications. In contrast, immunocompromised individuals are at significantly greater risk of developing invasive pulmonary aspergillosis (IPA), including those receiving chemotherapy treatments for cancer and patients receiving immunosuppressive regimens to prevent GVHD following hematopoietic stem cell or solid organ transplantation [1-5]. However, only a small proportion of immunosuppressed patients develop invasive fungal infections indicating that additional risk factors must exist. Numerous human studies have identified genetic polymorphisms in key antifungal pattern recognition receptors and inflammatory cytokines associated with fungal infections (reviewed in [6-8]). Thus, it is important to understand how these responses are coordinated in response to fungal infection of the lungs.

Recently, both type I and type III interferons have been shown to be essential for host resistance against pulmonary *A. fumigatus* challenge [9]. Both Dectin 1 (*Clec7a*) and MDA5 (*Ifih1*) are necessary to induce of the type I and type III interferon response following *A. fumigatus* challenge in the mouse model [10, 11]. Engagement of Dectin 1 leads to Syk activation which drive IRF5 activation for the production of IFNβ following *Candida albicans* challenge [12]. MDA5 engagement by dsRNA leads to its interaction with MAVS resulting in the recruitment of IKKε and TBK1 that activate NFκB and IRF3 and IRF7, respectively, for the production of early cytokines and type I interferons [13]. Genetic polymorphisms within *CLEC7A* have been associated with increased risk of developing IPA [14-16], while the role for genetic polymorphisms within *IFIH1* or *MAVS* regarding susceptibility to IPA has not been explored to date.

Herein, our data demonstrates that genetic polymorphisms within *IFIH1* and *MAVS* alter

the production of interferon-dependent chemokines and risk for HSCT patients in developing IPA,

respectively. Interestingly, increased risk for developing IPA was associated with genetic variation

within *MAVS* only in the donor/hematopoietic compartment. Using our murine model, we identify

alveolar macrophages as a key hematopoietic cell in the induction of the MAVS-dependent

interferon response which is necessary for host resistance against *A. fumigatus*. Overall, our study

reveals a critical role for MDA5/MAVS in host anti-fungal immunity in both mice and humans.

RESULTS

Single Nucleotide Polymorphisms (SNPs) in *IFIH1* and *MAVS* influence the risk for developing invasive pulmonary aspergillosis.

Our recent mouse data suggests that MDA5/MAVS signaling is critical in maintaining host resistance against pulmonary challenge with *A. fumigatus* [11]. Therefore, we wanted to explore the role of these molecules in a human cohort comprised of 460 HSCT patients (Supplemental Table 1). Missense SNPs within the coding region of *IFIH1* (rs1990760 and rs3747517) are associated with autoimmune conditions, particularly interferonopathies [17]. Moreover, a missense SNP in *MAVS* (rs17857295) is associated with altered type I interferon regulation [18]. While an individual genotype for each SNP was not associated with risk of developing IPA (Supplemental Figure 1), we further examined these SNPs using a recessive genetic model because the autoimmune risk allele in *IFIH1* rs1990760 has been shown to work in a dominant fashion [17]. Notably, we found a significant association between the CC genotype at rs1990760 in *IFIH1* and the risk of developing IPA in HSCT recipients (Figure 1A). This increased risk occurred when the variant was carried by the recipient in a recessive genetic model, but not when it was carried by the donor. In contrast, no association between the *IFIH1* rs3747517 SNP and the risk of developing IPA was found (Figure 1B). Additionally, we also found a significant association between the *MAVS* rs17857295 SNP and the risk of developing IPA in HSCT recipients (Figure 1C). This increased risk occurred when the variant was carried by the donor in a recessive genetic model, but not when it was carried by the recipient. Moreover, the risk from the donor *MAVS* rs17857295 SNP held up to multivariate analysis accounting for relevant clinical risk factors (Table 1).

To demonstrate a functional effect of the rs1990760 SNP in *IFIH1* and the associated

amino acid substitution on the function of MDA5, we analyzed the inflammatory response in the

bronchoalveolar lavage fluid (BALF) of 23 patients with IPA after stratification by genotypes at

rs1990760 in *IFIH1*. We found that individuals with the CC genotype, which is associated with

increased risk of developing IPA (Figure 1A), had a significant reduction in IP-10 (CXCL10),

RANTES (CCL5), and MIP-1α (CCL3), but not IL-8 and GROα (CXCL1) (Figure 1D). IP-10

(CXCL10), RANTES (CCL5), and MIP-1α (CCL3) are known to be interferon-responsive genes,

while IL-8 and GROα (CXCL1) are not [19]. In the BALF samples from this patient cohort, the

levels of IFN-α were below the detection limit (data not shown). Given the relatively rare

frequency of the rs17857295 genotype in *MAVS*, it was not possible to analyze cytokine production

according to *MAVS* genotype in the BAL samples (data not shown). Overall, these data support an

important role for MDA5/MAVS signaling in regulating the inflammatory response and host

resistance against *Aspergillus* spp. in humans.

Deletion of *Mavs* in CD11c⁺ cells result in increased susceptibility to *Aspergillus fumigatus* in

mice.

Given the observation that patients in the HSCT cohort with the *MAVS* rs17857295

polymorphism within the donor population had altered risk for developing IPA, we next wanted

to assess which hematopoietic cell population(s) in our invasive aspergillosis murine model relies

on *Mavs* to maintain host resistance against *A. fumigatus.* To directly address which cells must

express *Mavs* we utilized the recently developed *Mavs*^(fl/fl) *x Itgax-Cre* conditional knock-out

model, hereafter referred to as *Mavs*^(Cd11c/Cd11c), that will delete *Mavs* in all CD11c-expressing cells

[20], including both dendritic cells and alveolar macrophages. To test of role of *Mavs* expression

in CD11c-expressing cells for the maintenance of host resistance against *A. fumigatus*, we

challenged C57BL/6J, *Mavs*$^{(fl/fl)}$, and *Mavs*$^{(Cd11c/Cd11c)}$ mice with 4×10^7 conidia of the CEA10 strain

and monitored survival over nine days. *Mavs*$^{(Cd11c/Cd11c)}$ mice were more susceptible to pulmonary

challenge with *A. fumigatus* than either C57BL/6J or *Mavs*$^{(fl/fl)}$ mice (Figure 2A; Mantel-Cox log

rank test, $p < 0.0001$). Mice completely lacking either *Ifih1* or *Mavs* have decreased neutrophil

recruitment, which was associated with increased fungal germination after pulmonary challenge

with *A. fumigatus* [11]. Thus, we assessed fungal germination in the lung by histological analysis

at 48 h after conidial instillation. Strikingly, GMS staining of lung tissue from *Mavs*$^{(Cd11c/Cd11c)}$

mice revealed high levels of *A. fumigatus* germination at this time compared with control *Mavs*$^{(fl/fl)}$

mice (Figure 2B). When the percentage of germinated *A. fumigatus* was quantified, *Mavs*$^{(fl/fl)}$ mice

displayed low levels of fungal germination ($23.5\% \pm 9.1$) compared with *Mavs*$^{(Cd11c/Cd11c)}$ mice

($60.1\% \pm 8.2$). Finally, we assessed inflammatory cell accumulation in the airways. The increased

susceptibility of *Mavs*$^{(Cd11c/Cd11c)}$ mice to *A. fumigatus* challenge was associated with decreased

accumulation of neutrophils in the airways compared with *Mavs*$^{(fl/fl)}$ mice (Figure 2C). These data

match our prior observations with mice completely lacking either *Ifih1* or *Mavs* having increased

IPA susceptibility [11], suggesting that *Mavs* expression within CD11c-expressing cells is critical

for host resistance against *A. fumigatus* challenge.

Alveolar macrophage transfer to *Mavs*$^{(Cd11c/Cd11c)}$ conditional knock-out mice reduces their

susceptibility to *Aspergillus fumigatus* in mice.

Multiple cells in the lungs can express *Igtax* (CD11c) including alveolar macrophages,

CD103$^+$ dendritic cells, CD11b$^+$ dendritic cells, monocyte-derived dendritic cells, and

plasmacytoid dendritic cells, all of which have be implicated in the antifungal immune response

against *A. fumigatus* [21-27]. Utilizing the publicly available data assembled by the ImmGen

Consortium [28], in the lungs we see that *Mavs* is highly expressed in Siglec F[+] alveolar

macrophages, but also moderately expressed in both monocytes and dendritic cells (data not

shown). Interestingly, following RSV infection alveolar macrophages have been shown to be the

key cell type for initiating the type I interferon through a MAVS-dependent mechanism [29, 30].

While alveolar macrophages largely populate the lungs during embryogenesis and are maintained

by local proliferation during homeostasis [31, 32] following conditioning for HSC transplantation

alveolar macrophage can be repopulated from the donor HSC compartment [33-36]. Therefore, to

specifically address the role of *Mavs* in alveolar macrophages, we purified Siglec F[+] cells from the

lungs of naïve *Mavs*[(fl/fl)] or *Mavs*[(Cd11c/Cd11c)] mice and transferred 5×10^5 Siglec F[+] cells

intratracheally into *Mavs*[(Cd11c/Cd11c)] mice one day prior to challenging with 4×10^7 conidia of the

CEA10 strain (Figure 3A). Forty-eight hours after challenge, we examined the lungs by GMS

staining of lung tissue from *Mavs*[(Cd11c/Cd11c)] mice receiving the *Mavs*[(Cd11c/Cd11c)] Siglec F[+] cells and

revealed the presence of high levels of germinated *A. fumigatus* (68.1% ± 12.6), while that was

not observed to the same extent in *Mavs*[(Cd11c/Cd11c)] mice receiving the *Mavs*[(fl/fl)] Siglec F[+] cells

(31.4% ± 6.7) (Figure 3B). This is in line with control *Mavs*[(Cd11c/Cd11c)] mice or *Mavs*[(fl/fl)] mice

receiving DMEM (vehicle) administered intratracheally. When we examined the accumulation of

leukocytes in the airways by differential cytospin analysis we found that *Mavs*[(Cd11c/Cd11c)] mice

receiving *Mavs*[(Cd11c/Cd11c)] Siglec F[+] cells had reduced numbers of neutrophils when compared with

Mavs[(Cd11c/Cd11c)] mice receiving *Mavs*[(fl/fl)] Siglec F[+] cells (Figure 3C). Strikingly, when we

examined macrophage numbers in the airways, we found *Mavs*[(Cd11c/Cd11c)] mice receiving *Mavs*[(fl/fl)]

Siglec F[+] cells had many macrophages present, but *Mavs*[(Cd11c/Cd11c)] mice receiving *Mavs*[(Cd11c/Cd11c)]

Siglec F[+] cells had few macrophages in their airways at 48 hours post-inoculation with *A.*

fumigatus (Figure 3D). Importantly, the decrease in macrophages in the airways is not observed in

naïve *Mavs*$^{(Cd11c/Cd11c)}$ mice (Supplemental Figure 2A). Additionally, *Mavs*$^{(Cd11c/Cd11c)}$ mice

receiving an i.t. transfer of *Mavs*$^{(Cd11c/Cd11c)}$ Siglec F$^+$ cells 24h prior only had a moderate decrease

in alveolar macrophage numbers (Supplemental Figure 2B). These data demonstrate that *Mavs*

expression within Siglec F$^+$ alveolar macrophages is critical for host resistance against *A.*

fumigatus.

DISCUSSION

Humans inhale *Aspergillus fumigatus* spores on a daily basis, but individuals with healthy

immune systems readily clear *A. fumigatus* conidia from their airways without problems. One lung

sentinel cell that can be critical in the clearance of *A. fumigatus* is the alveolar macrophage.

Alveolar macrophages are known to phagocytose and destroy *A. fumigatus* [25], but their role in

host resistance has remained controversial [27, 37]. Patients undergoing HSCT are at increased

risk of developing IPA [1-5]. In patients undergoing HSCT, alveolar macrophages are

dysfunctional [35] and found at decreased in numbers [34-36] for at least 50 days post-

transplantation. Thus, it is important to understand the role of alveolar macrophages in the initial

inflammatory response induced by *A. fumigatus.* These lung sentinel alveolar macrophages are

also known to participate in the inflammatory response induced by *A. fumigatus* [38], but their role

in driving the interferon response following *A. fumigatus* challenge was not addressed. During

respiratory syncytial virus infection alveolar macrophages have been demonstrated to be key

drivers of the type I interferon response through a MAVS-dependent mechanism [29, 30]. We

recently describe a critical role for MDA5/MAVS-dependent induction of interferons following *A.*

fumigatus challenge [11], but the cellular localization of MAVS-dependent signaling was not

elucidated. Here we demonstrated that *Mavs* expression in alveolar macrophages was at least

partially necessary for host resistance against *A. fumigatus*, through the regulation of neutrophil

accumulation in the lung for the prevention of fungal growth. While alveolar macrophage transfer

to *Mavs$^{(Cd11c/Cd11c)}$* mice largely restores host resistance, it was not absolute suggesting additional

CD11c$^+$ cells might require *Mavs* expression for resistance. One potential candidate would be

plasmacytoid dendritic cells that are known to produce type I interferons after fungal challenge

[21] and are essential for host resistance *A. fumigatus* [39], but the importance of MAVS signaling

in plasmacytoid dendritic cells in driving the interferon response remains controversial. The role

of MAVS signaling in other CD11c$^+$ cell populations will be explored in future studies. Alveolar

macrophages also participate in the inflammatory response to other fungal pathogens including

Cryptococcus neoformans [40], *Pneumocystis* spp. [41], and *Rhizopus* spp. [42]. Interestingly,

heterogeneity within the alveolar macrophage population has been demonstrated particularly in

regard to CXCL2 expression [40], which could be important in our findings showing that the mice

lacking *Ifih1* or *Mavs* have decreased neutrophil accumulation following *A. fumigatus* challenge

[11].

While our previous studies have shown the importance of MDA5/MAVS signaling [11]

and type I/III interferons [9] in host resistance against *A. fumigatus* in the murine model, the

importance of these pathways in humans has not been described. Herein, we have found that

antifungal role of MDA5/MAVS signaling is also critical in humans. Our data from a cohort of

HSCT patients demonstrate that SNPs in *IFIH* (rs1990760) and *MAVS* (rs17857295) are associated

with an altered risk of developing IPA. Interestingly, SNPs within *IFIH1* have been associated

with numerous autoimmune diseases, particularly those that are highly dependent on type I

interferons for their progression [17, 43]. Recent evidence suggests that the autoimmune risk

genotype of the rs1990760 SNP in *IFIH1* (TT) has functional consequences in human PBMCs,

specifically it is associated with elevated basal levels of *IFNB1* [17]. Knock-in of the *IFIH1*

rs1990760 autoimmune risk genotype (TT) into mice not only recapitulates the susceptibility to

autoimmune diseases, but also results in increased host resistance against encephalomyocarditis

virus (ECMV), a picornavirus specifically recognized by MDA5 [44], and elevated *Ifnb1*

expression [17]. Moreover, in human pancreatic islets it has been shown that the *IFIH1* rs1990760

TC genotype also drove enhanced interferon signaling after Coxsackievirus infection [45]. This

enhanced interferon response during Coxsackievirus infection was associated with greater

signaling from the peroxisomes by the *IFIH1* rs1990760 TC genotype [45], which has previously

been shown to preferentially induce a type III interferon response [46, 47]. Interestingly, our SNP

analysis of a human HSCT cohort found that the T allele of *IFIH1* rs1990760, which as just

mentioned is associated with an increased interferon response [17, 45], was associated with a

reduced risk for the development of IPA. In addition to the *IFIH1* rs1990760 SNP, we also found

a clinical association with the rs17857295 SNP in *MAVS* with the risk of developing IPA after

HSCT. Much less is known about the functional outcome of the rs17857295 SNP in *MAVS*, but

one report suggests overexpression of the GG genotype in 293T cells stably knocked-down for the

endogenous *MAVS* allele resulted in an inability to induce *Ifnb1* expression following PolyI:C

stimulation [18]. Overall, our analysis suggests that HSCT patients harboring these SNPs display

a reduced ability to trigger an interferon response and are at greater risk of developing IPA.

In contrast to our clinical data from HSCT patients who develop IPA, the T allele at

rs1990760 in *IFIH1* is associated with an increased risk for the development of chronic

mucocutaneous candidiasis [48]. Again, the T allele at rs1990760 is associated with an increased

interferon response [17, 45], but the lack of IFNAR-dependent signaling leads to increased

susceptibility to invasive candidiasis in mice [12]. In their study Netea and colleagues did not

observe altered *Ifnb* expression in the absence of MDA5 after stimulation with *Candida albicans*

hyphae [48], but a role for MDA5 in regulating other interferons was not explored. Our data with

A. fumigatus suggests that MDA5/MAVS-signaling are more important in the regulation of type

III interferons (IFNλ/IL-28), rather than type I interferons (IFNα/β). Excessive inflammation and

tissue pathology can be critical in mediating disease during invasive candidiasis [49]. Rivera and

colleagues demonstrated that type III interferon (IFNλ/IL-28) signaling enhances the production

of reactive oxygen species by neutrophils following *A. fumigatus* [9], which could enhance disease

during invasive candidiasis. Thus, there appears to be an interesting dichotomy in MDA5 signaling

that is crucial for tuning host resistance against different fungal pathogens and warrants further

research.

METHODS

Mice and Aspergillus fumigatus challenge model. $Mavs^{(fl/fl)}$ and $Mavs^{(fl/fl)}$ x *Itgax-Cre* mice [20] were bred in-house at Geisel School of Medicine at Dartmouth. C57BL/6J mice were purchased from Jackson Laboratory. All mice were 8-16 weeks of age at the time of challenge.

Preparation of Aspergillus fumigatus conidia and murine challenge model. A. fumigatus CEA10 strains were used for this study. *A. fumigatus* was grown on glucose minimal media (GMM) agar plates for 3 days at 37°C. Conidia were harvested by adding 0.01% Tween 80 to plates and gently scraping conidia from the plates using a cell scraper. Conidia were then filtered through sterile Miracloth, were washed, and resuspended in phosphate buffered saline (PBS), and counted on a hemocytometer.

Mice were challenged with *A. fumigatus* conidia by the intratracheal (i.t.) route. Mice were anesthetized by inhalation of isoflurane; subsequently, mice were challenged i.t. with ~4×10^7 *A. fumigatus* conidia in a volume of 100 µl PBS. At the indicated time after *A. fumigatus* challenge, mice were euthanized using carbon dioxide. Bronchoalveolar lavage fluid (BALF) was collected by washing the lungs with 2 ml of PBS containing $0.05M$ EDTA. BALF was clarified by centrifugation and stored at -20°C until analysis. After centrifugation, the cellular component of the BALF was resuspended in 200 µl of PBS and total BAL cells were determined by hemocytometer count. BALF cells were subsequently spun onto glass slides using a Cytospin4 cytocentrifuge (Thermo Scientific) and stained with the Hema 3™ Stat Pack (Fisher Scientific) stain set for differential counting. For histological analysis lungs were filled with and stored in 10% buffered formalin phosphate for at least 24 hours. Lungs were then embedded in paraffin and sectioned into 5-micron sections. Sections were stained with Grocott-Gomori methenamine silver

(GMS) using standard histological techniques to assess lung inflammatory infiltrates and fungal

germination, respectively. Representative pictures of lung sections were taken using an Olympus

BX50WI microscope with a QImaging Retiga 2000R camera.

Alveolar macrophage isolation and adoptive transfer. Lungs from naïve *Mavs*$^{(fl/fl)}$ and *Mavs*$^{(fl/fl)}$ *x*

Itgax-Cre were perfused with 20-30 ml of PBS. Lungs were then removed, injected with 700 µl of

collagenase buffer [10ml RPMI, 1ml FBS, 25 µl 0.5*M* CaCl$_2$, 25 µl 0.5*M* MgCl$_2$, 50 µl HEPES,

100 µl L-glutamine, 100 µl Penn-Strep, 2 µl gentamycin, 25 µl DNase (VWR, Catalog # 77001-

900), and 2 ml Liberase (Sigma-Aldrich)], and finally placed in 15 ml conical tube containing 2

ml of collagenase buffer. Lungs were digested or 30min at 37°C while shaking at 160-180 rpm.

After which, 5 ml of ice cold stop buffer [25 ml RPMI, 1.25 ml FBS, 50 µl 0.5*M* EDTA] was

added to each tube. Lungs were pushed through a 70 µm filter to generate a single cell suspension.

Single cell lung suspensions were labeled with anti-Siglec-F MicroBeads (Miltenyi Biotec, Cat.

No. 130-118-513) and selected for using LS Columns (Miltenyi Biotec, Cat. No. 130-042-401).

Siglec F$^+$ were eluted from the column, spun down, and resuspended in DMEM at 5x10^6 cells per

ml. Subsequently 5x10^5 Siglec F$^+$ alveolar macrophages were transferred i.t. to *Mavs*$^{(fl/fl)}$ *x Itgax-*

Cre. Twenty-four hours later mice were challenge with *A. fumigatus* as described above.

Human HSCT cohort for SNP analysis. A total of 460 hematologic patients undergoing allogeneic

hematopoietic stem cell transplantation (HSCT) at the Hospital of Santa Maria, Lisbon and

Instituto Português de Oncologia (IPO), Porto, between 2009 and 2015 were enrolled in the study.

Cases of IPA were identified and classified as 'probable' or 'proven' according to the 2008 criteria

from the European Organization for Research and Treatment of Cancer/Mycology Study Group

(EORTC/MSG) [50]. Exclusion criteria included diagnosis of 'possible' IPA, infection with

invasive molds other than *Aspergillus* spp. or history of pre-transplant mold infection. Study

approval was obtained from the institutional review boards (SECVS-125/2014, HSM-632/14 and

CES.26/015) and from the National Data Protection Commission (CNPD, 1950/2015) and was in

compliance with all local relevant ethical regulations.

Genomic DNA was isolated from whole blood of patients using the QIAcube automated

system (Qiagen). SNPs were selected based on their previous association with infection or with

putative functional consequences to the gene [17, 18, 45]. Genotyping was performed using

KASPar assays (LGC Genomics) in an Applied Biosystems 7500 Fast Real-Time PCR system

(Thermo Fisher Scientific), according to the manufacturer's instructions.

Cytokine analysis of BALF from human IPA patients. BALF and blood samples were collected

from hospitalized adult patients (≥18 years of age) at the Leuven University Hospitals, Leuven,

Belgium, as previously described [51]. This study was approved and carried out in accordance

with recommendations of the Ethics Subcommittee for Life and Health Sciences of the University

of Minho, Portugal (SECVS-125/2014), and the Ethics Committee of the University Hospitals of

Leuven, Belgium. Written informed consent was obtained from all subjects in accordance with the

Declaration of Helsinki.

For this cytokine analysis, twenty-three cases of "probable" or "proven" IPA were

identified according to the standard criteria from the European Organization for Research and

Treatment of Cancer/Mycology Study Group (EORTC/MSG) [50] and included for cytokine

analysis. Genomic DNA was isolated from EDTA venous blood and genotyped for the *IFIH1*

rs1990760 polymorphism. Cytokine levels were determined using the Cytokine & Chemokine 34-

Plex Human ProcartaPlex and stratified based on their *IFIH1* rs1990760 genotype.

Statistical analysis. Statistical significance for *in vitro* and *ex vivo* data was determined by a Mann-

Whitney U test, one-way ANOVA using a Bonferroni post-test, or Kruskal-Wallis one-way

ANOVA with Dunn's post-test through the GraphPad Prism 7 software as outlined in the figure

legends. Mouse survival data were analyzed with the Mantel-Cox log rank test using GraphPad

Prism. For the human HSCT cohort, the probability of IPA resulting from *IFIH1* and *MAVS* SNPs

was analyzed using the cumulative incidence method and compared using Gray's test [52].

Cumulative incidences were computed with the *cmprsk* package for R version 2.10.1 [53], with

censoring of data at the date of last follow-up visit and defining relapse and death as competing

events. A period of 24 months after transplant was chosen to include all cases of IPA.

Study approvals. All animal experiments were approved by the Dartmouth College Institutional

Animal Care and Use Committee under protocol number 00002168. For our human studies

approval was obtained from the institutional review boards (SECVS-125/2014, HSM-632/14 and

CES.26/015) and from the National Data Protection Commission (CNPD, 1950/2015) and was in

compliance with all local relevant ethical regulations.

AUTHOR CONTRIBUTIONS

Conceived and designed the experiments: XW, AC, JJO. Performed the experiments: XW, CC, MG. Analyzed the data: XW, CC, MG, AC, JJO. Provided clinical samples: JFL, AC Jr, KL, JM. Wrote the paper: XW, JJO.

ACKNOWLEDGEMENTS

Thank you to Drs. Robert Cramer, Claudia Jakubzick, and David Leib (Geisel School of Medicine at Dartmouth) for helpful discussion on this project. Research in this study was supported in part by institutional startup funds to JJO in part through the Dartmouth Lung Biology Center for Molecular, Cellular, and Translational Research grant P30 GM106394 (PI: Bruce A. Stanton) and Center for Molecular, Cellular and Translational Immunological Research grant P30 GM103415 (PI: William R. Green). JJO was partially supported by a Munck-Pfefferkorn Award from Dartmouth College and NIH R01 AI139133 grant. SMB is supported by the Division of Intramural Research, National Institutes of Health, National Institute of Allergy and Infectious Diseases. AC and CC were supported by the Fundação para a Ciência e Tecnologia (FCT) (PTDC/SAU-SER/29635/2017, PTDC/MED-GEN/28778/2017, UIDB/50026/2020, UIDP/50026/2020, and CEECIND/04058/2018), the Northern Portugal Regional Operational Programme (NORTE 2020), under the Portugal 2020 Partnership Agreement, through the European Regional Development Fund (ERDF) (NORTE-01-0145-FEDER-000039), the European Union's Horizon 2020 research and innovation programme under grant agreement no. 847507, and the "la Caixa" Foundation (ID 100010434) and FCT under the agreement LCF/PR/HR17/52190003. The funders had no role in the preparation or publication of the manuscript.

REFERENCES

1.   Baddley, J.W., et al., *Factors associated with mortality in transplant patients with invasive aspergillosis.* Clin Infect Dis, 2010. **50**(12): p. 1559-67.

2.   Steinbach, W.J., et al., *Clinical epidemiology of 960 patients with invasive aspergillosis from the PATH Alliance registry.* J Infect, 2012. **65**(5): p. 453-64.

3.   Garcia-Vidal, C., et al., *Epidemiology of invasive mold infections in allogeneic stem cell transplant recipients: biological risk factors for infection according to time after transplantation.* Clin Infect Dis, 2008. **47**(8): p. 1041-50.

4.   Upton, A., et al., *Invasive aspergillosis following hematopoietic cell transplantation: outcomes and prognostic factors associated with mortality.* Clin Infect Dis, 2007. **44**(4): p. 531-40.

5.   Thompson, G.R., 3rd and T.F. Patterson, *Pulmonary aspergillosis.* Semin Respir Crit Care Med, 2008. **29**(2): p. 103-10.

6.   Romani, L., *Immunity to fungal infections.* Nat Rev Immunol, 2011. **11**(4): p. 275-88.

7.   Cunha, C., et al., *Human genetic susceptibility to invasive aspergillosis.* PLoS Pathog, 2013. **9**(8): p. e1003434.

8.   Campos, C.F., et al., *Host Genetic Signatures of Susceptibility to Fungal Disease.* Curr Top Microbiol Immunol, 2019. **422**: p. 237-263.

9.   Espinosa, V., et al., *Type III interferon is a critical regulator of innate antifungal immunity* Science Immunology, 2017. **2**(16): p. eaan5357.

10.   Dutta, O., et al., *Dectin-1 Promotes Type I and III Interferon Expression to Support Optimal Antifungal Immunity in the Lung.* Front Cell Infect Microbiol, 2020. **10**: p. 321.

11.   Wang, X., et al., *MDA5 Is an Essential Sensor of a Pathogen-Associated Molecular Pattern Associated with Vitality That Is Necessary for Host Resistance against Aspergillus fumigatus.* J Immunol, 2020. **205**(11): p. 3058-3070.

12.   del Fresno, C., et al., *Interferon-beta production via Dectin-1-Syk-IRF5 signaling in dendritic cells is crucial for immunity to C. albicans.* Immunity, 2013. **38**(6): p. 1176-86.

13.   Loo, Y.M. and M. Gale, Jr., *Immune signaling by RIG-I-like receptors.* Immunity, 2011. **34**(5): p. 680-92.

14.   Cunha, C., et al., *Dectin-1 Y238X polymorphism associates with susceptibility to invasive aspergillosis in hematopoietic transplantation through impairment of both recipient- and donor-dependent mechanisms of antifungal immunity.* Blood, 2010. **116**(24): p. 5394-402.

15.   Sainz, J., et al., *Dectin-1 and DC-SIGN polymorphisms associated with invasive pulmonary*
Aspergillosis infection. PLoS One, 2012. **7**(2): p. e32273.

16.   Fisher, C.E., et al., *Validation of single nucleotide polymorphisms in invasive aspergillosis*
following hematopoietic cell transplantation. Blood, 2017. **129**(19): p. 2693-2701.

17.   Gorman, J.A., et al., *The A946T variant of the RNA sensor IFIH1 mediates an interferon*
program that limits viral infection but increases the risk for autoimmunity. Nat Immunol,
2017. **18**(7): p. 744-752.

18.   Xing, F., et al., *Alteration of Antiviral Signalling by Single Nucleotide Polymorphisms*
(SNPs) of Mitochondrial Antiviral Signalling Protein (MAVS). PLoS One, 2016. **11**(3): p.
e0151173.

19.   Bauer, J.W., et al., *Elevated serum levels of interferon-regulated chemokines are*
biomarkers for active human systemic lupus erythematosus. PLoS Med, 2006. **3**(12): p.
e491.

20.   Dutta, M., et al., *A Systems Approach Reveals MAVS Signaling in Myeloid Cells as Critical*
for Resistance to Ebola Virus in Murine Models of Infection. Cell Rep, 2017. **18**(3): p. 816-
829.

21.   Ramirez-Ortiz, Z.G., et al., *A nonredundant role for plasmacytoid dendritic cells in host*
defense against the human fungal pathogen Aspergillus fumigatus. Cell Host Microbe,
2011. **9**(5): p. 415-24.

22.   Espinosa, V., et al., *Inflammatory monocytes orchestrate innate antifungal immunity in the*
lung. PLoS Pathog, 2014. **10**(2): p. e1003940.

23.   Hohl, T.M., et al., *Inflammatory monocytes facilitate adaptive CD4 T cell responses during*
respiratory fungal infection. Cell Host Microbe, 2009. **6**(5): p. 470-81.

24.   Zelante, T., et al., *CD103(+) Dendritic Cells Control Th17 Cell Function in the Lung.* Cell
Rep, 2015. **12**(11): p. 1789-801.

25.   Ibrahim-Granet, O., et al., *Phagocytosis and intracellular fate of Aspergillus fumigatus*
conidia in alveolar macrophages. Infect Immun, 2003. **71**(2): p. 891-903.

26.   Grimm, M.J., et al., *Monocyte- and macrophage-targeted NADPH oxidase mediates*
antifungal host defense and regulation of acute inflammation in mice. J Immunol, 2013.
190(8): p. 4175-84.

27.   Bhatia, S., et al., *Rapid host defense against Aspergillus fumigatus involves alveolar*
macrophages with a predominance of alternatively activated phenotype. PLoS One, 2011.
6(1): p. e15943.

28.   Heng, T.S., M.W. Painter, and C. Immunological Genome Project, *The Immunological*
Genome Project: networks of gene expression in immune cells. Nat Immunol, 2008. **9**(10):
p. 1091-4.

29.   Goritzka, M., et al., *Alveolar macrophage-derived type I interferons orchestrate innate*
immunity to RSV through recruitment of antiviral monocytes. J Exp Med, 2015. **212**(5): p.
699-714.

30.   Kirsebom, F.C.M., et al., *Neutrophil recruitment and activation are differentially*
dependent on MyD88/TRIF and MAVS signaling during RSV infection. Mucosal Immunol,
2019. **12**(5): p. 1244-1255.

31.   Guilliams, M., et al., *Alveolar macrophages develop from fetal monocytes that differentiate*
into long-lived cells in the first week of life via GM-CSF. J Exp Med, 2013. **210**(10): p.
1977-92.

32.   Svedberg, F.R., et al., *The lung environment controls alveolar macrophage metabolism*
and responsiveness in type 2 inflammation. Nat Immunol, 2019. **20**(5): p. 571-580.

33.   Zinter, M.S. and J.R. Hume, *Effects of Hematopoietic Cell Transplantation on the*
Pulmonary Immune Response to Infection. Front Pediatr, 2021. **9**: p. 634566.

34.   Thomas, E.D., et al., *Direct evidence for a bone marrow origin of the alveolar macrophage*
in man. Science, 1976. **192**(4243): p. 1016-8.

35.   Winston, D.J., et al., *Alveolar macrophage dysfunction in human bone marrow transplant*
recipients. Am J Med, 1982. **73**(6): p. 859-66.

36.   Nakata, K., et al., *Augmented proliferation of human alveolar macrophages after*
allogeneic bone marrow transplantation. Blood, 1999. **93**(2): p. 667-73.

37.   Mircescu, M.M., et al., *Essential role for neutrophils but not alveolar macrophages at early*
time points following Aspergillus fumigatus infection. J Infect Dis, 2009. **200**(4): p. 647-
56.

38.   Dubourdeau, M., et al., *Interaction of Aspergillus fumigatus with the alveolar macrophage.*
Med Mycol, 2006. **44**(Supplement_1): p. S213-S217.

39.   Guo, Y., et al., *During Aspergillus Infection, Monocyte-Derived DCs, Neutrophils, and*
Plasmacytoid DCs Enhance Innate Immune Defense through CXCR3-Dependent
Crosstalk. Cell Host Microbe, 2020: p. doi: 10.1016/j.chom.2020.05.002.

40.   Xu-Vanpala, S., et al., *Functional heterogeneity of alveolar macrophage population based*
on expression of CXCL2. Sci Immunol, 2020. **5**(50).

41. Bhagwat, S.P., et al., *Intrinsic Programming of Alveolar Macrophages for Protective Antifungal Innate Immunity Against Pneumocystis Infection.* Front Immunol, 2018. **9**: p. 2131.

42. Andrianaki, A.M., et al., *Iron restriction inside macrophages regulates pulmonary host defense against Rhizopus species.* Nat Commun, 2018. **9**(1): p. 3333.

43. Cen, H., et al., *Association of IFIH1 rs1990760 polymorphism with susceptibility to autoimmune diseases: a meta-analysis.* Autoimmunity, 2013. **46**(7): p. 455-62.

44. Kato, H., et al., *Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses.* Nature, 2006. **441**(7089): p. 101-5.

45. Domsgen, E., et al., *An IFIH1 gene polymorphism associated with risk for autoimmunity regulates canonical antiviral defence pathways in Coxsackievirus infected human pancreatic islets.* Sci Rep, 2016. **6**: p. 39378.

46. Odendall, C., et al., *Diverse intracellular pathogens activate type III interferon expression from peroxisomes.* Nat Immunol, 2014. **15**(8): p. 717-26.

47. Dixit, E., et al., *Peroxisomes are signaling platforms for antiviral innate immunity.* Cell, 2010. **141**(4): p. 668-81.

48. Jaeger, M., et al., *The RIG-I-like helicase receptor MDA5 (IFIH1) is involved in the host defense against Candida infections.* Eur J Clin Microbiol Infect Dis, 2015. **34**(5): p. 963-74.

49. Pappas, P.G., et al., *Invasive candidiasis.* Nat Rev Dis Primers, 2018. **4**: p. 18026.

50. De Pauw, B., et al., *Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group.* Clin Infect Dis, 2008. **46**(12): p. 1813-21.

51. Goncalves, S.M., et al., *Evaluation of Bronchoalveolar Lavage Fluid Cytokines as Biomarkers for Invasive Pulmonary Aspergillosis in At-Risk Patients.* Front Microbiol, 2017. **8**: p. 2362.

52. Gray, R.J., *A Class of K-Sample Tests for Comparing the Cumulative Incidence of a Competing Risk.* The Annals of Statistics, 1988. **16**(3): p. 1141-1154.

53. Scrucca, L., A. Santucci, and F. Aversa, *Competing risk analysis using R: an easy guide for clinicians.* Bone Marrow Transplant, 2007. **40**(4): p. 381-7.

Table 1. Multivariate analysis of the association of *IFIH1 and MAVS* SNPs with the risk of invasive pulmonary aspergillosis among transplant recipients.

Genetic/clinical variables	Adjusted HR† (95% CI)	P value
Donor rs17857295 in *MAVS*	2.39 (1.07 – 5.32)	0.033
Recipient rs1990760 in *IFIH1*	1.46 (0.89 – 2.40)	0.121
Acute GVHD grades III-IV	1.73 (0.94 – 3.19)	0.047

HR, hazard ratio; CI, confidence interval. Multivariate analyses were based on the subdistribution regression model of Fine and Gray. †Hazard ratios were adjusted for patient age and gender, and clinical variables with a P<0.15 in the univariate analyses. Only the variables remaining significant after adjustment are shown.

Figure 1. *IFIH1* **and** *MAVS* **polymorphisms are associated with invasive pulmonary aspergillosis in human transplant patients using a recessive allele model.** Cumulative incidence analysis of invasive aspergillosis after transplantation according to donor or recipient genotypes at rs1990760 in *IFIH1* **(A)**, rs3747517 in *IFIH1* **(B)**, or rs17857295 in *MAVS* **(C)** over 24 months after HSCT. Data were analyzed by two-sided Gray's test. **(D)** Inflammatory cytokine levels in the bronchoalveolar lavage fluid from 23 patient with invasive pulmonary aspergillosis were measured using a 32-plex ProCarta Luminex assay and plotted based on their *IFIH1* rs1990760 genotype. Data were analyzed using a Mann-Whitney U-test (* $p < 0.05$).

Figure 2. _Mavs_-dependent responses are essential in CD11c[+] cells for host resistance against _Aspergillus fumigatus_. _Mavs_[<Cd11c/Cd11c>], _Mavs_[<fl/fl>] and C57BL/6J mice were challenged i.t. with 4×10^{7} resting conidia of the CEA10 isolate of _Aspergillus fumigatus_. **(A)** Survival analysis in immune-competent wild-type and knock-out mice were tracked over the first 9 days. ****, P < 0.0001 by Mantel-Cox log rank test. **(B)** Forty hours after _A. fumigatus_ challenge mice were euthanized and fungal germination was assessed in the lungs by GMS staining. **, P < 0.01 by Mann-Whitney U-test. **(C)** At the same time point, cell differentials in the lung airways was determined by differential staining of BALF cytospins. Statistical significance was determined using a two-way ANOVA with a Sidak's post-test.

__Figure 3. Alveolar macrophages require a Mavs-dependent response to maintain host resistance against Aspergillus fumigatus.__ **(A)** $Mavs^{<Cd11c/Cd11c>}$ mice were complemented with $5x10^5$ $Mavs^{<fl/fl>}$ or $Mavs^{<Cd11c/Cd11c>}$ alveolar macrophages (Siglec F$^+$) given intranasally. One day later mice were challenged i.t. with $4x10^7$ resting conidia of the CEA10 isolate of *Aspergillus fumigatus*. Forty hours after *A. fumigatus* challenge mice were euthanized. **(B** Fungal germination was assessed in the lungs by GMS staining. Representative 20x GMS images are shown. Statistically significant different were determined using a one-way ANOVA with a Tukey's post-test: *$p<0.05$ (vs. DMEM → $Mavs^{(fl//fl)}$), ****$p<0.0001$ (vs. DMEM → $Mavs^{(fl//fl)}$), and $\tau\tau\tau\tau$ $p<0.0005$ (vs. $Mavs^{(fl//fl)}$ → $Mavs^{(Cd11c)}$). Neutrophils **(C)** and macrophages **(D)** in the lung airways was determined by differential staining of BALF cytospins.

 Supplemental Table 1. Baseline characteristics of transplant recipients enrolled in the study.

Variables	IPA (n=91)	No IPA (n=348)	P value
Age at transplantation, no (%)			
≤20 years	13 (14.3)	69 (19.8)	0.264
21 – 40 years	23 (25.3)	101 (29.0)	
>40 years	55 (60.4)	178 (51.2)	
Gender, no (%)			
Female	38 (41.8)	150 (43.1)	0.859
Male	53 (58.2)	198 (56.9)	
Underlying disease, no. (%)			
Acute leukemia	49 (53.8)	179 (51.5)	0.115
Chronic lymphoproliferative diseases	14 (15.4)	69 (19.8)	
Myelodysplastic/myeloproliferative diseases	13 (14.3)	30 (8.6)	
Chronic myeloproliferative diseases	7 (7.7)	20 (5.7)	
Aplastic anemia	6 (6.6)	17 (4.9)	
Other	2 (2.2)	33 (9.5)	
Transplantation type, no. (%)			
Matched, related	34 (37.4)	169 (48.6)	0.037
Matched, unrelated	33 (36.3)	81 (23.3)	
Mismatched, related	0 (0.0)	7 (2.0)	
Mismatched, unrelated	24 (26.4)	91 (26.2)	
Graft source, no. (%)			
Peripheral blood	80 (87.9)	287 (82.5)	0.506
Bone-marrow	10 (11.0)	53 (15.2)	
Cord blood	1 (1.1)	8 (2.3)	
Disease stage, no. (%)			
First complete remission	49 (53.8)	188 (54.0)	0.800

Second or subsequent remission, or relapse	13 (14.3)	59 (17.0)	
Active disease	29 (31.9)	101 (29.0)	
Conditioning regimen, no (%)			
RIC	68 (74.7)	228 (65.5)	0.091
Myeloablative	23 (25.3)	120 (34.5)	
CMV serostatus of donor and recipient, no. (%)			
D-/R+ or D+/R+	80 (87.9)	313 (89.9)	0.504
D-/R- or D+/R-	11 (12.1)	35 (10.1)	
Duration of neutropenia, mean days (range)†	13.1 (8 – 39)	13.5 (5 – 35)	0.460
Acute GVHD, no. (%)			
No GVHD or grades I – II	63 (69.2)	302 (86.8)	0.0002
Grades III – IV	28 (30.8)	46 (13.2)	
Antifungal prophylaxis, no. (%)‡			
Fluconazole	42 (46.2)	117 (33.6)	0.002
Posaconazole	26 (28.6)	107 (30.8)	
Other	9 (9.9)	14 (4.0)	
None or unknown	14 (15.4)	110 (31.6)	

Twenty-one patients with "possible" IPA were excluded. Chronic lymphoproliferative diseases
included cases of chronic lymphocytic leukemia, multiple myeloma, and B- and T-cell
lymphomas. Chronic myeloproliferative diseases included cases of chronic myelogenous leukemia
and primary myelofibrosis. Other diseases included cases of idiopathic medullar aplasia,
lymphohistiocytosis, hemoglobinopathies and paroxysmal nocturnal hemoglobinuria. RIC,
reduced intensity conditioning; CMV, cytomegalovirus; D, donor; R, recipient; GVHD, graft-
versus-host-disease. †Neutropenia was defined as $\leq 0.5 \times 10^9$ cells/L. ‡Other antifungals used in
prophylaxis included voriconazole, liposomal amphotericin B, itraconazole and caspofungin. P
values were calculated by Fisher's exact probability t-test or Student's t-test for continuous
variables.

Supplemental Figure 1. **Individual genotype of *IFIH1* and *MAVS* polymorphisms in human transplant patients.** Cumulative incidence analysis of invasive aspergillosis after transplantation according to donor or recipient genotypes at rs1990760 in *IFIH1* (**A**), rs3747517 in *IFIH1* (**B**), or rs17857295 in *MAVS* (**C**) over 24 months after HSCT. Data were analyzed by two-sided Gray's test.

560
561

Supplemental Figure 2. **Alveolar macrophage numbers are normal in naïve mice.** BAL cells from naïve *Mavs[Cd11c/Cd11c]* *and Mavs[fl/fl]* mice **(A)** or naïve *Mavs[Cd11c/Cd11c]* mice given $5x10^5$ SiglecF$^+$ lung cells from naïve *Mavs[Cd11c/Cd11c]* *and Mavs[fl/fl]* mice 24h prior **(B)** were collected in PBS/EDTA. Alveolar macrophage numbers were quantified by Cytospin analysis and differential staining. n.s. = not significant; *, $P < 0.01$; by Mann-Whitney U-test.