Neuro-COVID long-haulers exhibit broad dysfunction in T cell memory generation and

responses to vaccination

Lavanya Visvabharathy[1*¶], Barbara Hanson[1,3], Zachary Orban[1], Patrick H. Lim[1], Nicole Palacio[2],

Rishi Jain[1], Eric Michael Liotta[1], Pablo Penaloza-MacMaster[2], Igor J. Koralnik[1*]

[1]Ken and Ruth Davee Department of Neurology, Feinberg School of Medicine, Northwestern

University, Chicago IL 60611 USA

[2]Department of Microbiology-Immunology, Feinberg School of Medicine, Northwestern

University, Chicago IL 60611 USA

[3]Rush Medical College, Chicago IL 60612 USA

*Corresponding authors: Igor J. Koralnik, M.D.: igor.koralnik@northwestern.edu;

Lavanya Visvabharathy, Ph.D: lavanya.visvabharathy@northwestern.edu

¶ Lead contact: Lavanya Visvabharathy, Ph.D

Summary

The high prevalence of post-acute sequelae of SARS-CoV-2 infection (PASC) is a significant health concern. In particular, virus-specific immunity in patients who suffer from chronic neurologic symptoms after mild acute COVID remain poorly understood. Here, we report that neuro-PASC patients have a specific signature composed of humoral and cellular immune responses that are biased towards different structural proteins compared to healthy COVID convalescents. Interestingly, the severity of cognitive deficits or quality of life markers in neuro-PASC patients are associated with reduced effector molecule expressionn in memory T cells. Furthermore, we demonstrate that T cell responses to SARS-CoV-2 mRNA vaccines are aberrantly elevated in longitudinally sampled neuro-PASC patients compared with healthy COVID convalescents. These data provide a framework for the rational design of diagnostics and predictive biomarkers for long-COVID disease, as well as a blueprint for improved therapeutics.

Keywords: COVID-19 immunity, T cell memory, neuro-PASC, long-COVID, vaccine-induced immunity

Introduction

SARS-CoV-2 is the causative agent of a worldwide pandemic that started in Wuhan, China in December, 2019. There have been more than 200 million cases and over 4 million deaths globally attributable to COVID-19 disease (Center, 2021). Although highly effective vaccines are now used to prevent severe disease and death caused by SARS-CoV-2, long-term sequelae of SARS-CoV-2 infection are increasingly becoming an important medical concern.

SARS-CoV-2 infection is associated with presentations ranging from asymptomatic carriage to severe multi-organ dysfunction (Li et al., 2020; Liguori et al., 2020; Syed et al., 2020). Globally, the estimated fatality rate following SARS-CoV-2 infection is approximately 1-2%, but not all patients recover to their baseline states (Higgins et al., 2020). "Long COVID" affects an estimated 10-40% of all infected people and includes symptoms persisting more than 28 days after diagnosis of SARS-CoV-2 infection, termed "post-acute sequelae of SARS-CoV-2 infection" or PASC (Ladds et al., 2020; Vehar et al., 2021). Greater than two-thirds of hospitalized COVID-19 patients experience ongoing fatigue, breathlessness, and psychiatric issues such as post-traumatic stress disorder (PTSD) 4-8 weeks after discharge (Halpin et al., 2021). However, the vast majority of people with COVID experience mild disease not requiring hospitalization, and more than half of these have symptoms persisting more than 4 months after acute infection (Petersen et al., 2020). Many people who survived Middle-East respiratory virus (MERS) and the original SARS-CoV pandemic also experienced PTSD and neurological impairments up to 3.5 years after acute infection (Ahmed et al., 2020; Lam et al., 2009). More recent studies on recovered COVID-19 patients showed significant cognitive deficits in attention, working memory, and emotional processing months after acute infection had resolved (Hampshire, 2021). Here, we focus on a cohort of mostly non-hospitalized long-COVID patients

presenting with neurological symptoms ("neuro-PASC") who exhibit a reduction in quality of

life with regards to cognitive and fatigue parameters (Graham et al., 2021).

Numerous studies suggest that a robust T cell response is necessary for viral clearance in

infected individuals. In particular, CD4[+] T cell responses directed against the Spike protein were

found in 100% of healthy COVID convalescents, while 70% displayed Spike-specific CD8[+] T

cell responses (Grifoni et al., 2020). Another study found that memory T cells cross-reactive to

SARS-CoV-2 nucleocapsid (N) protein persisted in people exposed to SARS-CoV up to 17 years

after the epidemic of 2003 (Le Bert et al., 2020). Post-mortem autopsies of COVID-19 patients

have shown that innate immune cells but not lymphocytes are enriched in lung infiltrate, and that

these patients exhibit impaired germinal center formation in lung draining lymph nodes (Duan et

al., 2020). Additional studies have shown that CD8[+] T cell depletion after SARS-CoV-2

infection of rhesus macaques impairs anamnestic responses, suggesting a role for T cell memory

in long-term immunity to SARS-CoV-2 infection (McMahan et al., 2021).

Most studies on T cell responses to SARS-CoV-2 have focused on acutely symptomatic

individuals. However, adaptive immune responses in patients who remain chronically

symptomatic remain largely unexplored. Our studies indicate that neuro-PASC patients ("CN")

had hypo-functional Ag-specific CD8+ T cell memory responses SARS-CoV-2 compared with

healthy convalescents ("CC") and enhanced reactivity to N and M proteins compared with CC

subjects. The severity of cognitive deficits and quality of life markers were associated with

enhanced polarization in cytolytic granule expression in memory T cell subsets. Finally, we

demonstrate that T cell responses to SARS-CoV-2 vaccination are more robust and display

aberrant kinetics in neuro-PASC patients compared with control groups. Together, these data

demonstrate a wide-scale dysfunction in SARS-CoV-2 T cell memory in neuro-PASC long-

hauler patients, with important implications for both appropriate diagnostic and vaccination

strategies.

Results

Clinical characteristics of neuro-PASC patients

We enrolled a total of 111 participants prior to SARS-CoV-2 vaccination drawn from the neuro-COVID-19 outpatient clinic at Northwestern Memorial Hospital or from the surrounding Chicago area. This included 56 neuro-PASC "CN" patients (confirmed RT-PCR+ or anti-SARS-CoV-2 Spike IgG+) meeting Infectious Disease Society of America clinical criteria for COVID-19 starting after February 2020 and had neurologic symptoms lasting at least 6 weeks post-infection, as previously reported (Graham et al., 2021). Among those, 48 (86%) were never hospitalized for pneumonia or hypoxemia. We additionally recruited 24 healthy COVID convalescents without lasting symptoms (RT-PCR+ or seropositive for anti- SARS-CoV-2 Spike RBD IgG, "CC"); and 31 healthy controls who were RT-PCR- and seronegative for SARS-CoV-2 Spike-IgG ("HC"; study description in Fig. 1A).

CN patients displayed a constellation of neurological symptoms similar to those previously reported by our group and in other studies (Fig. 1B). In addition, we utilized standardized methods to quantify their quality of life and cognitive disturbances relative to healthy convalescents. Results from the patient reported outcomes information system (PROMIS-57) survey (Tang et al., 2019) showed that CN patients scored significantly lower on physical function and higher on anxiety, depression, pain and other metrics compared with CC subjects or the national average (Fig. 1C). NIH toolbox tests administered to CN patients to assess their cognitive function (Weintraub et al., 2013) showed CN patients had significantly lower T scores in the attention module, which was indicative of cognitive dysfunction relative to the national average (Fig. 1D).

Neuro-PASC is associated with a distinct immunodominance hierarchy.

To determine whether neuro-PASC included alterations in the magnitude and/or

specificity of Ag-specific T cell responses in CN patients, we performed ELISPOT for SARS-

CoV-2-specific T cells. PBMC from each subject were cultured in the presence of peptide pools

derived from spike (S), nucleocapsid (N), or membrane (M) proteins of SARS-CoV-2 (Fig. S1).

The magnitude and specificity of both IFN-γ and IL-2 responses to S sub-pools were similar

between CN and CC subjects (Fig. 2A-B). However, CN subjects produced high levels of IFN-γ

against all sub-pools of N and M proteins, while CC subjects had a specificity for the N1 pool

and lower responses against all M pools (Fig. 2C-D), indicative of different immunodominance

hierarchies between groups. No significant differences were found in IL-2 production, and

healthy controls exhibited some response to N pools likely caused by cross-reactivity with

endemic coronaviruses (Fig. 2E-F). Antibody titers against the Spike receptor-binding domain

(RBD) did not differ between CN and CC groups (Fig. 2G). Interestingly, high IgG titers did not

correlate with the magnitude of the IFN-γ T cell response against S2 or S3 peptide pools

comprising the 2 halves of the Spike RBD (Fig. S2).

CD4$^+$ Tfh cells display opposing reactivity to SARS-CoV-2 N- and M-proteins in neuro-PASC vs.

healthy COVID convalescents

RNA-Seq analysis of CD4$^+$ T cells from hospitalized and non-hospitalized COVID-19

patients showed that severe disease is associated with elevated CD4$^+$ T follicular helper (Tfh)

proportions relative to patients with mild disease (Meckiff et al., 2020). We sought to determine

whether Tfh responses (gating in Fig. S3) that prime antibody production from plasma B cells

could similarly differentiate neuro-PASC CN patients from CC individuals. Immunophenotyping

showed that there were no differences in total percentages of T cell subsets, including Tfh cells,

between groups (Fig. S4). The activation-induced marker (AIM) assay has been previously used

to detect TCR-mediated activation of T cells (Grifoni et al., 2020) and we used this method to

investigate Tfh cell activation (gating for Tfh cells in Fig. S3). N protein-specific

$CD134^+CD137^+$ (AIM^+) CD4 Tfh cells were significantly elevated in CN compared with CC

subjects, while the opposite trend was observed in M protein-specific activation (Fig. 3A-B). No

differences were seen in Tfh activation between CN and CC groups across Spike peptide pools

(Fig. 3B). The magnitude of N- and M-specific Tfh cell activation did not correlate with the time

since acute infection in either CN or CC patients (Fig. 3C-D) despite reports showing that

antibody titers against N protein decrease rapidly post-infection (Van Elslande et al., 2021)

which would presumably lead to decreased N-specific Tfh cell activation over time. N-specific

IgG titers were significantly elevated in CN compared with CC subjects (Fig. 3E), consistent

with the enhanced N-specific $CD4^+$ Tfh cell activation shown in 3B. Similar to Tfh cells, there

was no correlation between anti-N IgG titers and time post-symptom onset for either CN or CC

subjects (Fig. S5A). Studies have shown that IgG titers against N protein decline to undetectable

levels in 40% of COVID convalescents within 4 months (Muecksch et al., 2021). We largely did

not observe this decline in CN patients despite collecting their blood samples an average of 193

days post-symptom onset (Fig. 1B).

$CD4^+$ T cell polyfunctionality to Spike protein differs in neuro-PASC vs. healthy COVID

convalescents

Cytotoxic $CD4^+$ T cells acquire the ability to produce cytolytic granules upon activation

with cognate viral antigen (Goubard et al., 2015; Sledzinska et al., 2020). Viral infections

specifically induce expansion of the memory $CD4^+$ T effector memory cells re-expressing

CD45RA (TEMRA) population that secrete copious amounts of cytolytic granules upon antigen

encounter (Tian et al., 2017). We therefore investigated whether Ag-specific production of

multiple granzymes and cytokines in CD4$^+$ T cells were altered between groups. While CN and

HC groups maintained polyfunctionality and polarization, CC subjects had significantly more

polyfunctional CD4$^+$ T cells producing granzymes A, B, and M after Spike pool stimulation

(category 2 in yellow, Fig. 3F). A heatmap quantifying this effect showed that CC had a 2-fold

elevation in category 2 effector molecules over CN subjects (Fig. 3G). This data suggests that

cytotoxic responses to Spike protein in CD4$^+$ T cells from healthy COVID-19 convalescents

significantly differ from those experiencing chronic neurologic symptoms.

CD8$^+$ memory T cell functionality in neuro-PASC patients

CD8$^+$ memory T cells are crucial for protection from recurrent viral infections (Mockus

et al., 2019). Studies have shown that memory CD8$^+$ cells can persist for several years after

patient exposure to SARS-CoV (Chen et al., 2005) and that acute mild disease induces stronger

Ag-specific CD8$^+$ memory responses than severe disease (Peng et al., 2020). However, little is

known about whether CD8$^+$ memory differs in neuro-PASC long-haulers vs. healthy COVID

convalescents months after acute infection resolves. To address this, we probed the dynamics of

CD8$^+$ TEM, TCM, and TEMRA activation in CN, CC, and HC subjects. CD8$^+$ TEM were

significantly activated by various S, N, and M peptide pools in CC subjects while remaining

relatively quiescent after antigen stimulation in CN patients (Fig. 4A-B). CD8$^+$ TEMRA are

terminally differentiated memory cells with distinct transcriptional programs that respond

quickly to viral infections (Tian et al., 2019). Total percentages of CD8$^+$ TEMRA cells were

significantly elevated in CN compared with CC and HC groups (Fig. 4C), but despite their

increased numbers, CD8$^+$ TEMRA cells were less activated by the S3/4 peptide pool in CN

patients and showed a trend towards decreased activation after N pool stimulation compared with

CC subjects (Fig. 4D-E).

Similarly, antigen stimulation resulted in altered cytokine polarization in CD8+ T cell

subsets in CN and CC groups. Effector molecule production in CD8+ T cells was similar across

groups in the unstimulated condition, despite showing some statistically significant differences in

perforin production (Fig. S6B). However, stimulation with the S3/4 pool induced greater TNF-α

production in the CC group while CN patients had enhanced IL-6 production to the same

stimulus (Fig. 4F). S pool stimulation also induced CD8+ TEMRA cells from the CC group to

produce elevated levels of granzymes, which was not seen in CN or HC groups (Fig. 4G, black

lines), demonstrating that CD8+ T cell subsets display decreased functionality in CN vs. CC

patients after stimulation.

Impaired cognition and decreased quality of life metrics correlate with distinct patterns of

polyfunctionality in memory T cell subsets

Having shown that Tfh and T cell memory responses differed between patient groups,

we next sought to probe whether within-group differences in T cell responses correlated with

various clinical measures in CN patients. We found a significant positive correlation between the

magnitude of IFN-γ production to N protein and higher pain interference scores (Fig. 5A). There

was also a trend towards a positive correlation between high scores for depression and the

magnitude of the N-specific T cell response (Fig. 5B). To look at associations between clinical

parameters and T cell activation, we separated T scores from NIH Toolbox or PROMIS-57

measurements (Fig. 1C-D) into quartiles and only the lowest and highest groups (Q1 vs. Q4)

were used for analysis (Fig. S7A, red boxes). CN subjects reporting high (Q4) pain interference

scores produced significantly less granzyme A or B and more IL-6 than those with low scores

(Fig. 5C-D) from CD8$^+$ T cells after S pool stimulation. Further, patients reporting low

depression scores had highly polyfunctional Spike-specific CD8$^+$ TEM, while those reporting

high levels of depression had TEM producing ~3-fold higher granzymes A, B, and M alone

(category 2; Fig. 5E, F). The severity of cognitive impairment could also be significantly

correlated with T cell responses. Q1 patients scoring low on executive function tests had CD8$^+$

TCM polarized towards granzymes A, B, M and perforin production (category 1), while those in

Q4 were biased to produce granzymes A, B, and perforin (Fig. 5G-H). Similar analyses were

performed for other CD8$^+$ and CD4$^+$ memory T cell subsets, and significant differences by

quartile were also found in correlations with depression, processing speed, working memory, and

global pain (Fig. S7B-K). These data show that the severity of cognitive deficits or quality of life

measures can be significantly correlated with differences in SARS-CoV-2-specific memory T

cell function.

SARS-CoV-2 vaccination induces sustained elevation in Spike-specific IFN-γ responses in neuro-

PASC but not healthy COVID convalescents

SARS-CoV-2 mRNA vaccines have been proven to be highly effective against

contracting infection in the short-term in both clinical trials and in real-world settings (Haas et

al., 2021; Polack et al., 2020). However, neither clinical trials nor field studies specifically

characterized the vaccine-induced immune response in long-COVID patients, and it is similarly

unknown whether vaccination can ameliorate long-haul symptoms. One prospective study

showed that vaccination did not significantly worsen symptoms in 22 patients with long-COVID

(Arnold et al., 2021), but also did not provide evidence to show any decrease in symptoms. To

address how vaccine-induced immunity evolves over time in neuro-PASC long-haulers, we

conducted a longitudinal vaccine study assessing Spike-specific antibody and T cell responses

over the course of 6 months post-vaccination (study design and patient demographics in Fig. 6A-B). While this study is ongoing, we report data from the first 4 months comparing responses in CN, CC, and HC subjects (n for each group per study visit shown in Fig. S8). Vaccination with either Pfizer or Moderna mRNA vaccines induced robust SARS-CoV-2 Spike RBD IgG titers in all groups tested by 3 weeks post-2nd dose, with the highest titers found in CN patients. Antibody titers were at similar levels in all groups by 3 months post-1st dose, but trended higher in CN than in HC subjects at visit (V)4 (Fig. 6C). In parallel, we measured Spike-specific T cell responses in study subjects. The magnitude and kinetics of the T cell response in CC and HC subjects were similar and peaked at V3. However, CN patients had significantly elevated Spike-specific T cell responses compared with either HC or CC groups at V2 after the vaccination boost (Fig. 6D; individual pool data shown in Fig. S9). IFN-γ-specific T cell responses remained high at 4 months post-vaccination in CN patients while not significantly differing from pre-vaccination levels in both CC and HC subjects. To our knowledge, these are the first data that longitudinally compare the T cell response to vaccination in neuro-PASC patients with healthy COVID convalescents and unexposed individuals.

Overall, our study demonstrates that neuro-PASC patients have elevated IFN-γ responses to N and M proteins, enhanced Ag-specific activation of Tfh cells, sustained anti-N IgG production, and deficient activation of multiple CD8$^+$ memory T cell subsets compared with healthy COVID-19 convalescents. There were also correlations between the severity of cognitive deficits or quality of life impairments and Ag-specific cytokine signatures (Fig. 5, S7). Importantly, vaccination resulted in sustained enhancements in the magnitude of Spike-specific T cell responses in neuro-PASC patients vs. all other groups, regardless of prior COVID exposure. Together, we show that CN patients exhibit a large-scale dysfunction in multiple

235   aspects of the memory T cell response which may inform treatment and/or vaccination options

236   down the line.

Discussion

COVID-19 is well-recognized as a multi-organ disease with long-term sequelae

associated with neurological disease. PASC has been reported in up to 87% of those hospitalized

with SARS-CoV-2 pneumonia and in up to 40% of those with mild disease who do not require

hospitalization (Havervall et al., 2021; Hirschtick et al., 2021; Mahase, 2020). SARS-CoV-2

vaccines have decreased the rates of SARS-CoV-2 infection and mortality in unexposed

individuals, but it is not clear that vaccines help to ameliorate long-COVID symptoms in

previously infected individuals (Arnold et al., 2021) and thus further study is needed to inform

treatment options. Long-term sequelae after other coronavirus infections can persist for years

(Ahmed et al., 2020); therefore, it is important to specifically characterize SARS-CoV-2-specific

immune responses in long-COVID patients. Most studies on effector and memory T cell

responses to SARS-CoV-2 have focused on acute infection or healthy convalescents as opposed

to those with long-COVID (Rodda et al., 2021; Sekine et al., 2020; Weiskopf et al., 2020). We

aimed to fill this knowledge gap and determine whether and how T cell phenotype and function

differ in patients with neuro-PASC and healthy COVID-19 convalescents.

Clinically, neuro-PASC resembles myalgic encephalomyelitis/chronic fatigue syndrome

(ME/CFS), which many patients report as a post-viral infectious complication (Rasa et al., 2018).

The causes of ME/CFS remain elusive, and the underlying mechanisms of neuro-PASC remain

similarly unknown. It is possible that direct infection of the brain is responsible for neuro-PASC

symptoms. It is believed that SARS-CoV-2 gains entry into the central nervous system through

the olfactory bulb and has been shown to infect neurons in vitro, which is supported by findings

of viral protein expression in cortical neurons from post-mortem autopsies (Klingenstein et al.,

2020; Song et al., 2021). However, another study was unable to find any evidence of SARS-

CoV-2 in the brains of 4 patients with neurological symptoms during acute infection (Kantonen

et al., 2020), suggesting that infection of the nervous system may be transient. Another study of

acutely infected neuro-COVID patients did not find any SARS-CoV-2 RNA present in the CSF,

though they did identify enhanced presence of exhausted T cells and dedifferentiated monocytes

(Heming et al., 2021). However, all of these prior studies were conducted on patients with acute

SARS-CoV-2 infection as opposed to neuro-PASC, and thus mechanisms for neurological

dysfunction may differ. As lumbar punctures or brain biopsies are not indicated in neuro-PASC

outpatients, reproducing the above study results in outpatient populations will not be possible.

Additional hypotheses for neuro-PASC pathogenesis include a contribution of autoimmune

mechanisms which is suggested by the skewed ratio of females to males affected, similar to that

seen in rheumatoid arthritis or systemic lupus erythematosus (Chakravarty et al., 2007;

Myasoedova et al., 2010), as well as the possibility of persistent SARS-CoV-2 infection in the

periphery (Al-Aly et al., 2021). Much of our findings on the SARS-CoV-2 T cell response in

neuro-PASC patients supports this latter hypothesis.

Ag-specific IFN-γ and IL-2 production was similar between CN and CC groups in

response to stimulation by different SARS-CoV-2 Spike peptide pools (Fig. 2A-B). However, T

cells from CN patients retained high IFN-γ responses to all N- and M-peptide pools tested, while

CC subjects had a reactivity limited to N1 and N2 peptide pools with little IFN-γ production in

response to M pools (Fig. 2C-D). The presence of an immunodominance hierarchy in CC

subjects is the expected outcome of an effective memory response to SARS-CoV-2 infection.

Studies have shown that while a primary CD4[+] or CD8[+] T cell response to influenza A or LCMV

infection is highly diverse and contains T cells reactive to many viral proteins and epitopes, the

memory response preferentially contains T cells responding to specific immunodominant viral

antigens (Sant et al., 2018; Tebo et al., 2005). This effect is more pronounced in the CD8$^+$ T cell

compartment where memory responses are dependent on the ability of dendritic cells to present

viral antigen (Crowe et al., 2003) or on antigen availability (Henrickson et al., 2013). The

evident lack of a narrow and targeted IFN-γ response to N- and M-peptides in CN patients thus

suggests that they are unable to effectively generate a memory response to SARS-CoV-2.

Though there were no differences observed in the magnitude or specificity of the IFN-γ response

to SARS-CoV-2 Spike peptides, this is somewhat expected as memory T cell responses to Spike

protein remain diverse after both infection (Grifoni et al., 2020) and vaccination (Alter et al.,

2021). It is additionally not surprising that differences in reactivity were observed for IFN-γ and

not IL-2 T cell responses between CN and CC groups (Fig. 2B, E-F). Studies have demonstrated

that while Ag-specific T cells produce IL-2 as a proliferation factor, IFN-γ production is largely

confined to the memory T cell population (Anthony et al., 2012), which is largely where we see

differences in activation and reactivity between groups (Fig. 4).

Antibody production is a key outcome measure after SARS-CoV-2 infection, Tfh cell

activation can inform the effectiveness of an antibody response. Stimulation with SARS-CoV-2

N peptides activated Tfh cells in CN but not CC subjects, while the opposite trend was found

with M pool stimulation (Fig. 3A-B). Tfh cell activation is crucial to the establishment of

germinal centers in secondary lymphoid organs, ultimately resulting in B cell maturation into

long-lived plasma cells that can continuously produce class-switched, high-affinity antibodies

(Good-Jacobson and Shlomchik, 2010). Studies have demonstrated that the magnitude of the Tfh

cell response is dependent on the amount of antigen available and directly correlates with the

magnitude of the B cell response (Baumjohann et al., 2013). Indeed, the same CN patients with

high N-specific Tfh cell activation also displayed anti-N IgG titers significantly greater than CC

subjects (Fig. 3E) despite the fact that we obtained their samples more than 6 months on average

after acute infection. Altogether, these findings are consistent with the presence of a persisting

nucleocapsid antigen reservoir resulting in enhanced N-specific Tfh cell activation in CN

patients. Clinical reports have identified cases of persistent SARS-CoV-2 infection in the

nasopharynx lasting up to 63 days (Bennasrallah et al., 2020) and many patients re-tested RT-

PCR+ for SARS-CoV-2 up to 38 days post-discharge (Dao et al., 2021). Though many of our

CN patients have not re-tested RT-PCR+ by nasal swab (data not shown), others have yet to be

tested. The nasopharynx is not the only possible testing site, however, as there are numerous

studies showing that infectious SARS-CoV-2 particles can be detected in fecal matter and can

persist for up to 70 days post-symptom onset (Tian et al., 2020; van Doorn et al., 2020).

Gastrointestinal symptoms also did not have to be present in order to test RT-PCR+ for SARS-

CoV-2 in stool samples (Chen et al., 2020). Thus, it is possible that the gut may also act as a

reservoir for persistent SARS-CoV-2 infection which leads to aberrant T cell responses and the

development of neuro-PASC.

Effective generation of T cell memory responses is crucial to protect against future

infections with the same pathogen. $CD8^+$ TEM cells from CC subjects retained high levels of

activation to all S, N, and M pools assayed while CN patients displayed very little TEM

activation (Fig. 4A-B), suggestive of aberrant memory T cell function. Studies on convalescents

from the original SARS-CoV epidemic of 2002-03 found that the majority of Ag-specific $CD8^+$

memory T cells were TEM, and these persisted up to 4 years after infection (Channappanavar et

al., 2014). Sterilizing immunity can be a precondition for the development and maintenance of

memory T cells because high levels of foreign antigen favor the generation of short-lived

effector cells (Mueller et al., 2013), which is why chronic infections often limit the formation

and/or function of memory T cells (Wherry, 2011). In fact, chronic LCMV infection in mice

induced aberrantly functioning Ag-specific memory CD8[+] T cells requiring the presence of viral

peptide rather than simply the homeostatic cytokines IL-7 and IL-15 to proliferate (Shin et al.,

2007; Wherry et al., 2004). The inability of CD8[+] TEM to become activated by antigenic

stimulation in CN patients is thus suggestive of a chronic infection state wherein viral antigen

can persist but limits the formation of CD8[+] T cell memory. It is also possible that chronic

SARS-CoV-2 infection in CN patients leads to memory T cell exhaustion, as shown in chronic

HIV and HBV infections in humans (McLane et al., 2019). However, we did not find any

differences in the inhibitory marker PD-1 expression on CD8[+] memory T cell subsets between

CN and CC patients (data not shown).

Further providing support to the chronic infection hypothesis in neuro-PASC patients,

there was a significant elevation in the CD8[+] TEMRA cell population in CN patients over CC or

HC groups (Fig. 4C). CD8[+] TEMRA cells are a terminally differentiated memory subset that do

not traffic through secondary lymphoid organs, and their induction during viral infection can be

protective (Tian et al., 2019). Yet, they have also been shown to accumulate during persistent

viral infections and contribute to immunosenescence (Derhovanessian et al., 2011). CD8[+]

TEMRA cell reactivity to SARS-CoV-2 peptides was also decreased in CN patients over CC

subjects. Functionally, the polarization in granzyme production in S3/4-specific CD8[+] TEMRA

cells from CC patients (Fig. 4G) suggests higher cytotoxic capacity compared with CN patients

and coincides with their higher activation state in Fig. 4E. Ag-specific CD8[+] TEMRA cells are

expanded and functionally active in people with significant anti-Dengue virus immunity, and this

phenotype is seen as a goal for vaccine-induced protection (Tian et al., 2019). We propose that

CD8[+] TEMRA cells are more expanded but less functionally active in CN subjects over CC as a

consequence of inappropriate CD8[+] T cell memory formation, which is lacking in multiple T cell

subsets for neuro-PASC patients.

We also showed that CD8[+] T cells from CC subjects skewed towards TNF-α production

after S pool stimulation compared with CN patients. Comparatively, CN patients displayed

significantly more IL-6 production from S-specific CD8[+] T cells (Fig. 4F). Vaccination against

the intracellular pathogen *Shigella flexneri* showed that polyfunctional TNF-α-producing CD8[+]

TEM were an important correlate of protection (Toapanta et al., 2018). Despite multiple reports

indicating that enhanced TNF-α production is correlated with worse COVID-19 outcomes (Karki

et al., 2020, 2021; Pedersen and Ho, 2020), we speculate that Ag-specific TNF-α production is

protective in our model system because we are looking at chronically and not acutely infected

CN patients. In contrast, IL-6 is known to suppress T_H1 differentiation (Diehl et al., 2000) and

can promote pathogen survival and exacerbate clinical disease during the original SARS-CoV

infection (Channappanavar and Perlman, 2017). Indeed, studies in severely ill, hospitalized

COVID-19 patients demonstrated that high serum levels of IL-6 significantly correlated with

poor clinical outcome (Weiskopf et al., 2020). These data suggest a role for enhanced IL-6

production by CD8[+] T cells in the pathogenesis of neuro-PASC, and open new avenues of

research for the treatment of long-COVID through limiting IL-6 activity.

Clinically, neuro-PASC patients reported significantly elevated levels of anxiety,

depression, fatigue, sleep disturbance and pain as well as decreased physical function compared

with healthy convalescents (Fig. 1C). The severity of these deficits was highly correlated with

Ag-specific enhancements in polyfunctionality and decreases in polarization of various memory

T cell subsets (Fig. 5, S7). It is possible that T cell function contributes to the genesis and

persistence of some of these symptoms. Studies in rodents have shown that T cell activation and

function can affect the severity of pain and analgesia (Rosen et al., 2019; Sorge et al., 2015).

Indeed, pain is a common hallmark of chronic viral infection (Addis et al., 2020) and recognized

among post-COVID sequelae (Kemp et al., 2020); it might follow then that aberrant T cell

activation can associate with high pain scores. Additionally, reports have shown that

transcriptional programs in immunity and inflammation were differentially regulated in $CD4^+$ T

cells from patients with depression compared with healthy controls (Wang et al., 2015). T_{reg} cells

may also decrease depressive behavior through negative regulation of inflammation (Miller,

2010), and CN patients do display elevated T_H1-type cytokine production to S pool stimulation

(Fig. 2, 6) while not displaying any compensatory upregulation in T_{reg} total numbers or function

(data not shown). T cell-derived cytokines can also impact learning and memory. Studies in

mouse models of West Nile and Zika viral encephalitis have demonstrated that IFN-γ production

from $CD8^+$ T cells in the brain is responsible for neuronal apoptosis and spatial learning deficits

(Garber et al., 2019). Thus, there is a precedent for correlating T cell function with cognitive

deficits, pain, or depression. Ag-specific cytokine signatures associated with the severity of

cognitive and quality of life deficits in neuro-PASC patients may therefore provide some

predictive value in terms of clinical outcomes.

Preliminary reports showed that the Pfizer mRNA vaccine could elicit a T cell response 7

days after completion of the full prime-boost protocol (Sahin et al., 2020), but until our studies

there has been no data on longitudinal T cell responses primed by vaccination and how these

vary between groups with different types of prior SARS-CoV-2 exposure. Our results

demonstrate for the first time that vaccine-elicited immune responses are significantly divergent

in neuro-PASC versus healthy COVID convalescents (Fig. 6C-D). CN patients consistently had

higher antibody titers after receiving the second dose of the vaccine compared with CC and HC

groups, though titers were similar in all groups at 11-15 weeks post-1st dose (Fig. 6C). In

contrast, the magnitude of Spike-specific IFN-γ production by T cells remained high in CN

patients out to 4 months post-vaccination while not significantly higher than pre-vaccination

levels in CC and HC subjects along the same timeline (Fig. 6D). These data suggest that the

mRNA vaccines do not induce robust long-term T cell responses in many individuals, regardless

of prior COVID exposure, if they are not neuro-PASC patients. Yet, despite vaccines enhancing

Spike-specific IFN-γ production in CN T cells, the fact that these responses continue to increase

at 15 weeks post-vaccination suggests that CN patients may still have an active SARS-CoV-2

infection or a persistent antigen reservoir rather than developing a robust T cell memory

response. These results suggest that alternate SARS-CoV-2 vaccines that induce long-lasting

memory T cell responses in previously unexposed individuals as well as healthy COVID

convalescents are needed in order to mediate long-term protection from infection. Conversely,

vaccination may not be indicated for long-COVID patients who might have a persistent infection

as it may be ineffective in the absence of viral clearance. Indeed, current clinical guidance from

the CDC recommends that vaccination be delayed until 3 months after acute infection in

unvaccinated COVID convalescents. Thus, vaccination strategies that include the memory T cell

response as a marker for efficacy should be carefully considered. Together, these data show that

irregular Tfh responses, broad scale dysfunction in CD8^{+} T cell memory generation and aberrant

T cell responses to vaccination are hallmarks of neuro-PASC and require further study to inform

treatment and vaccination strategies across the population.

Limitations of study

One limitation of our study is the relatively small sample size of unvaccinated neuro-

PASC patients. This was due to the wide implementation of SARS-CoV-2 vaccines in the

Chicago area soon after beginning study enrollment. Another limitation was not being able to

control for time of sample collection with respect to date of COVID-19 symptom onset. As it is

possible that neuro-PASC could be the result of a persistent infection, further investigations

would require testing of potential cryptic reservoirs, including stool samples from CN patients.

Materials and Methods

Ethics Statement

This study was approved by the Northwestern University Institutional Review Board (Koralnik Lab, IRB STU00212583). Informed consent was obtained from all enrolled participants. Samples were de-identified before banking.

Study participants, NIH Toolbox, and PROMIS-57 data collection

We enrolled consenting unvaccinated adult outpatients seen in the Neuro-PASC-19 clinic at Northwestern Memorial Hospital from September 2020-June 2021, including 56 neuro-PASC "long-hauler" patients with documented PCR+ or seropositive IgG results for SARS-CoV-2 (CN). In parallel, we recruited 24 unvaccinated healthy COVID convalescents from the surrounding community who tested PCR+ for SARS-CoV-2 but had no lingering neurological symptoms (CC) and 31 healthy controls who tested PCR- for SARS-CoV-2 and were also seronegative for IgG against SARS-CoV-2 Spike RBD. All study subjects remained living throughout the period of observation. Heparinized blood samples were collected one time from each subject at an average of 155-315 days post-symptom onset (as in Fig. 1B). Other demographic information is contained in Fig. 1B. CN patients completed a cognitive function evaluation in the clinic coincident or near the date of their blood sample acquisition with the National Institutes of Health (NIH) Toolbox v2.1 instrument, including assessments of: processing speed (pattern comparison processing speed test); attention and executive memory (inhibitory control and attention test); executive function (dimensional change card sort test); and working memory (list sorting working memory test) (Lai et al., 2011; Weintraub et al., 2013). PROMIS-57 was administered to CN and CC patients an average of 72 days post-sample

collection. Both PROMIS-57 and NIH Toolbox results are expressed as T-scores adjusted for

age, education, gender, and race/ethnicity with a score of 50 representing the normative

mean/median of the US reference population with a standard deviation of 10. Lower cognition T-

scores indicate worse performance while higher fatigue, depression, anxiety, or pain interference

T-scores indicate greater symptom severity.

PBMC and plasma collection

30mL of venous blood from study volunteers was collected in blood collection tubes containing

sodium heparin from BD Biosciences. Whole blood was layered on top of 15mL of Histopaque

1077 (Sigma-Aldrich) in 50mL Leucosep blood separation tubes (Greiner Bio-One) and spun at

1000g for 18min at RT. Plasma was collected and stored at -80°C. The PBMC layer was

collected and washed 2x in sterile PBS before red blood cell lysis with ACK buffer (Quality

Biologicals). PBMCs were used in assays either immediately or frozen down for use in the near

term.

SARS-CoV-2 peptide antigens

All S, N and M peptide arrays used in ELISPOT and flow cytometry studies were obtained from

BEI Resources, NIAID, NIH: Peptide Array, SARS-Related Coronavirus 2 Spike (S) Protein;

NR-52402, Nucleocapsid (N) Protein, NR-52404; Membrane (M) Protein, NR-52403. The S

peptide array consisted of 181 peptides of 13-17aa in length and split into 6 sub-pools (S1-S6)

containing 30-31 peptides each. The N peptide array consisted of 59 peptides of 13-17aa each

split into 3 sub-pools containing 29-30 peptides each. The M peptide array consisted of 31

peptides of 12-17aa and split into 3 sub-pools of 10-11 peptides each (Fig. S1). All peptides were

dissolved in either sterile H_2O or 50% sterile H_2O-DMSO up to 1mL for a universal 1mg/mL

stock concentration. Peptides were used at a final concentration at 2μg/mL in all assays.

IgG Spike RBD and Nucleocapsid ELISA

Antigen-specific total antibody titers were measured by ELISA as described previously (Dangi et

al., 2020; Palacio et al., 2020). In brief, 96-well flat-bottom MaxiSorp plates (Thermo Scientific)

were coated with 1 μg/ml of Spike RBD for 48 hr at 4°C. Plates were washed three times with

wash buffer (PBS + 0.05% Tween 20). Blocking was performed with blocking solution (PBS +

0.05% Tween 20 + 2% bovine serum albumin), for 4 hr at room temperature. 6 μl of sera was

added to 144 μl of blocking solution in the first column of the plate, 1:3 serial dilutions were

performed until row 12 for each sample, and plates were incubated for 60 min at room

temperature. Plates were washed three times with wash buffer followed by addition of secondary

antibody conjugated to horseradish peroxidase, goat anti-human IgG (H + L) (Jackson

ImmunoResearch) diluted in blocking solution (1:1000) and 100 μl/well was added and

incubated for 60 min at room temperature. After washing plates three times with wash buffer,

100 μl/well of Sure Blue substrate (SeraCare) was added for 1 min. Reaction was stopped using

100 μl/well of KPL TMB Stop Solution (SeraCare). Absorbance was measured at 450 nm using

a Spectramax Plus 384 (Molecular Devices). SARS-CoV-2 RBD and N proteins used for ELISA

were produced at the Northwestern Recombinant Protein Production Core by Dr. Sergii

Pshenychnyi using plasmids that were produced under HHSN272201400008C and obtained from

BEI Resources, NIAID, NIH: Vector pCAGGS containing the SARS-related coronavirus 2,

Wuhan-Hu-1 spike glycoprotein gene (soluble, stabilized), NR-52394 and receptor binding

domain (RBD), NR-52309, nucleocapsid gene NR-53507.

Cell stimulation and IFN-γ/IL-2 ELISPOT

Multiscreen-IP plates (Millipore-Sigma) were coated overnight at 4°C with 2μg/mL anti-IFN-γ

(clone 1-D1K, Mabtech) or 5μg/mL anti-IL-2 (clone MT2A91/2C95, Mabtech), washed with

sterile PBS, and blocked with complete RPMI-10% FBS. PBMC isolated from CN, CC, and HC

subjects were used either freshly isolated or after thawing and resting overnight in media

containing 10ng/μL recombinant human IL-15 (Peprotech) at 37°C, 5% CO_2. Cells were then

plated at a concentration of $2.5x10^5$ cells/well in 100μL of media and stimulated with the

indicated antigen mixtures from SARS-CoV-2 at a concentration of 2μg/mL in complete RPMI

medium containing 5% human AB serum (Sigma-Aldrich) and 5ng/mL IL-15. Plates were

incubated at 37°C, 5% CO_2 for 20 h and washed 5x with dH_2O and PBS-0.05% Tween-20 (PBS-

T). 2μg/mL biotinylated IFN-γ (clone 7-B6-1, Mabtech) or 5μg/mL IL-2 (clone MT8G10,

Mabtech) diluted in PBS-10% FBS (PBS-F) was added to the respective wells and plates were

incubated for 1.5h at RT. Plates were subsequently incubated for 40 minutes at RT in

streptavidin-alkaline phosphatase in PBS-F (Jackson ImmunoResearch) was added after washing

plates 5x in PBS-T. ELISPOT plates were developed using an Alkaline Phosphatase Conjugate

Substrate Kit according to manufacturer's instructions (Bio-Rad Laboratories, Carlsbad, CA).

IFN-γ or IL-2-producing cells were quantified using an ImmunoSpot reader (Cellular

Technologies, Ltd., Shaker Heights, OH).

Antibodies and Flow Cytometry

Fresh or frozen PBMCs isolated from the indicated patient groups were stimulated with antigen

mixtures as above for 20-22h at 37°C, 5% CO_2. For intracellular staining and cytokine detection,

the Brefeldin-A Golgi plug (Biolegend) was added at a 1:1000 concentration 2 hours after

antigenic stimulation commenced. Cells were washed with PBS-1% BSA after incubation and

incubated with the indicated antibodies for surface phenotyping by AIM assay or for intracellular

cytokine staining (ICS). Cells from each subject were left unstimulated in medium containing

5ng/mL IL-15 ("background") or stimulated in the presence of the indicated antigens. Fixation

and permeabilization was performed using Cytofix/Cytoperm (BD Biosciences). Surface staining

was done in the dark at 4°C for 30 minutes, while ICS was done in the dark at RT for 45

minutes. Flow cytometry was conducted on $2\text{-}5\text{x}10^5$ cells per condition. Data was acquired on a

BD FACSymphony Spectral analyzer and analyzed using FlowJo v10 (BD Biosciences) and

SPICE-Pestle (Roederer et al., 2011).

Quantification and Statistical Analysis

Statistical tests to determine significance are described in figure legends and conducted largely in

Prism (GraphPad). For pie graphs or heatmaps generated using SPICE analysis, statistics were

determined by Permutation test following unstimulated background subtraction, with additional

thresholding of 0.03% to account for noise, using SPICE-Pestle. *P*-values lower than 0.05 were

considered statistically significant. Quantile stratification was performed within group for CN

cohort. Clinical data were collected and managed using REDCap electronic data capture tools

hosted at Northwestern University Feinberg School of Medicine (Harris et al., 2009). REDCap

(Research Electronic Data Capture) is a secure, web-based software platform designed to support

data capture for research studies, providing 1) an intuitive interface for validated data capture; 2)

audit trails for tracking data manipulation and export procedures; 3) automated export procedures

for seamless data downloads to common statistical packages; and 4) procedures for data

integration and interoperability with external sources.

Acknowledgements

We would like to thank the Flow Cytometry Core Facility at the Robert H. Lurie Comprehensive Cancer Center at Northwestern University supported by Cancer Center Support Grant (NCI CA060553) for their assistance in optimizing antibody panels and help with flow cytometry instrumentation. L.V. was supported by a T32 grant T32AR007611 from the Department of Rheumatology, Northwestern University Feinberg School of Medicine. P.P.M. is supported by a grant from the Emerging and Re-Emerging Pathogens Program (EREPP) at Northwestern University, and a grant from the National Institute on Drug Abuse (NIDA, DP2DA051912).

Author Contributions

Conceptualization L.V. P.P.M. and I.K; Investigation L.V., B.H., Z.O., P.H.L, and N.P.; Formal Analysis L.V., B.H., E.M.L., P.P.M. and N.P.; Resources L.V., P.P.M., I.K., Data Curation L.V.; Writing L.V. with feedback from all authors; Supervision P.P.M and I.K.; Project Administration L.V.; Funding Acquisition L.V., P.P.M, and I.K.

Declaration of Interests

The authors declare no competing interests.

References

Addis, D.R., DeBerry, J.J., and Aggarwal, S. (2020). Chronic Pain in HIV. Mol Pain *16*,

1744806920927276.

Ahmed, H., Patel, K., Greenwood, D.C., Halpin, S., Lewthwaite, P., Salawu, A., Eyre, L., Breen,

A., O'Connor, R., Jones, A*., et al.* (2020). Long-term clinical outcomes in survivors of severe

acute respiratory syndrome and Middle East respiratory syndrome coronavirus outbreaks after

hospitalisation or ICU admission: A systematic review and meta-analysis. J Rehabil Med *52*,

jrm00063.

Al-Aly, Z., Xie, Y., and Bowe, B. (2021). High-dimensional characterization of post-acute

sequelae of COVID-19. Nature *594*, 259-264.

Alter, G., Yu, J., Liu, J., Chandrashekar, A., Borducchi, E.N., Tostanoski, L.H., McMahan, K.,

Jacob-Dolan, C., Martinez, D.R., Chang, A*., et al.* (2021). Immunogenicity of Ad26.COV2.S

vaccine against SARS-CoV-2 variants in humans. Nature.

Anthony, D.D., Milkovich, K.A., Zhang, W., Rodriguez, B., Yonkers, N.L., Tary-Lehmann, M.,

and Lehmann, P.V. (2012). Dissecting the T Cell Response: Proliferation Assays vs. Cytokine

Signatures by ELISPOT. Cells *1*, 127-140.

Arnold, D.T., Milne, A., Samms, E., Stadon, L., Maskell, N.A., and Hamilton, F.W. (2021).

Symptoms After COVID-19 Vaccination in Patients With Persistent Symptoms After Acute

Infection: A Case Series. Ann Intern Med.

Baumjohann, D., Preite, S., Reboldi, A., Ronchi, F., Ansel, K.M., Lanzavecchia, A., and

Sallusto, F. (2013). Persistent antigen and germinal center B cells sustain T follicular helper cell

responses and phenotype. Immunity *38*, 596-605.

Bennasrallah, C., Bannour, R., Jlassi, O., Kacem, M., Fredj, M.B., Abroug, H., Zemni, I.,

Garrach, B., Bahri, R., Charfeddine, N.*, et al.* (2020). Three COVID-19 cases with a long-term

viral shedding period in Tunisia. Pan Afr Med J *35*, 117.

Center, J.C.R. (2021). Cumulative worldwide Covid-19 cases (Baltimore, MD: Johns Hopkins

University of Medicine).

Chakravarty, E.F., Bush, T.M., Manzi, S., Clarke, A.E., and Ward, M.M. (2007). Prevalence of

adult systemic lupus erythematosus in California and Pennsylvania in 2000: estimates obtained

using hospitalization data. Arthritis Rheum *56*, 2092-2094.

Channappanavar, R., and Perlman, S. (2017). Pathogenic human coronavirus infections: causes

and consequences of cytokine storm and immunopathology. Semin Immunopathol *39*, 529-539.

Channappanavar, R., Zhao, J., and Perlman, S. (2014). T cell-mediated immune response to

respiratory coronaviruses. Immunol Res *59*, 118-128.

Chen, H., Hou, J., Jiang, X., Ma, S., Meng, M., Wang, B., Zhang, M., Zhang, M., Tang, X.,

Zhang, F.*, et al.* (2005). Response of memory CD8+ T cells to severe acute respiratory syndrome

(SARS) coronavirus in recovered SARS patients and healthy individuals. J Immunol *175*, 591-

598.

Chen, Y., Chen, L., Deng, Q., Zhang, G., Wu, K., Ni, L., Yang, Y., Liu, B., Wang, W., Wei, C.*,

et al.* (2020). The presence of SARS-CoV-2 RNA in the feces of COVID-19 patients. J Med

Virol *92*, 833-840.

Crowe, S.R., Turner, S.J., Miller, S.C., Roberts, A.D., Rappolo, R.A., Doherty, P.C., Ely, K.H.,

and Woodland, D.L. (2003). Differential antigen presentation regulates the changing patterns of

CD8+ T cell immunodominance in primary and secondary influenza virus infections. J Exp Med

198, 399-410.

Dao, T.L., Hoang, V.T., and Gautret, P. (2021). Recurrence of SARS-CoV-2 viral RNA in

recovered COVID-19 patients: a narrative review. Eur J Clin Microbiol Infect Dis *40*, 13-25.

Derhovanessian, E., Maier, A.B., Hahnel, K., Beck, R., de Craen, A.J.M., Slagboom, E.P.,

Westendorp, R.G.J., and Pawelec, G. (2011). Infection with cytomegalovirus but not herpes

simplex virus induces the accumulation of late-differentiated CD4+ and CD8+ T-cells in

humans. J Gen Virol *92*, 2746-2756.

Diehl, S., Anguita, J., Hoffmeyer, A., Zapton, T., Ihle, J.N., Fikrig, E., and Rincon, M. (2000).

Inhibition of Th1 differentiation by IL-6 is mediated by SOCS1. Immunity *13*, 805-815.

Duan, Y.Q., Xia, M.H., Ren, L., Zhang, Y.F., Ao, Q.L., Xu, S.P., Kuang, D., Liu, Q., Yan, B.,

Zhou, Y.W.*, et al.* (2020). Deficiency of Tfh Cells and Germinal Center in Deceased COVID-19

Patients. Curr Med Sci *40*, 618-624.

Garber, C., Soung, A., Vollmer, L.L., Kanmogne, M., Last, A., Brown, J., and Klein, R.S.

(2019). T cells promote microglia-mediated synaptic elimination and cognitive dysfunction

during recovery from neuropathogenic flaviviruses. Nat Neurosci *22*, 1276-1288.

Good-Jacobson, K.L., and Shlomchik, M.J. (2010). Plasticity and heterogeneity in the generation

of memory B cells and long-lived plasma cells: the influence of germinal center interactions and

dynamics. J Immunol *185*, 3117-3125.

Goubard, A., Loiez, C., Abe, J., Fichel, C., Herwegh, S., Faveeuw, C., Porte, R., Cayet, D.,

Sebbane, F., Penet, S.*, et al.* (2015). Superantigenic Yersinia pseudotuberculosis induces the

expression of granzymes and perforin by CD4+ T cells. Infect Immun *83*, 2053-2064.

Graham, E.L., Clark, J.R., Orban, Z.S., Lim, P.H., Szymanski, A.L., Taylor, C., DiBiase, R.M.,

Jia, D.T., Balabanov, R., Ho, S.U.*, et al.* (2021). Persistent neurologic symptoms and cognitive

dysfunction in non-hospitalized Covid-19 "long haulers". Ann Clin Transl Neurol *8*, 1073-1085.

Grifoni, A., Weiskopf, D., Ramirez, S.I., Mateus, J., Dan, J.M., Moderbacher, C.R., Rawlings,

S.A., Sutherland, A., Premkumar, L., Jadi, R.S.*, et al.* (2020). Targets of T Cell Responses to

SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease and Unexposed Individuals. Cell

181, 1489-1501 e1415.

Haas, E.J., Angulo, F.J., McLaughlin, J.M., Anis, E., Singer, S.R., Khan, F., Brooks, N., Smaja,

M., Mircus, G., Pan, K.*, et al.* (2021). Impact and effectiveness of mRNA BNT162b2 vaccine

against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a

nationwide vaccination campaign in Israel: an observational study using national surveillance

data. Lancet *397*, 1819-1829.

Halpin, S.J., McIvor, C., Whyatt, G., Adams, A., Harvey, O., McLean, L., Walshaw, C., Kemp,

S., Corrado, J., Singh, R.*, et al.* (2021). Postdischarge symptoms and rehabilitation needs in

survivors of COVID-19 infection: A cross-sectional evaluation. J Med Virol *93*, 1013-1022.

Hampshire, A., Trender W., Chamberlain SR, Jolly AE, Grant JE, Patrick F, Mazibuko N,

Williams S, Barnaby JM, Hellyer H, Mehta MA. (2021). Cognitive deficits in people who have

recovered from COVID-19. EClinicalMedicine.

Harris, P.A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., and Conde, J.G. (2009). Research

electronic data capture (REDCap)--a metadata-driven methodology and workflow process for

providing translational research informatics support. J Biomed Inform *42*, 377-381.

Havervall, S., Rosell, A., Phillipson, M., Mangsbo, S.M., Nilsson, P., Hober, S., and Thalin, C.

(2021). Symptoms and Functional Impairment Assessed 8 Months After Mild COVID-19

Among Health Care Workers. JAMA *325*, 2015-2016.

Heming, M., Li, X., Rauber, S., Mausberg, A.K., Borsch, A.L., Hartlehnert, M., Singhal, A., Lu,

I.N., Fleischer, M., Szepanowski, F.*, et al.* (2021). Neurological Manifestations of COVID-19

Feature T Cell Exhaustion and Dedifferentiated Monocytes in Cerebrospinal Fluid. Immunity *54*,

164-175 e166.

Henrickson, S.E., Perro, M., Loughhead, S.M., Senman, B., Stutte, S., Quigley, M., Alexe, G.,

Iannacone, M., Flynn, M.P., Omid, S.*, et al.* (2013). Antigen availability determines CD8(+) T

cell-dendritic cell interaction kinetics and memory fate decisions. Immunity *39*, 496-507.

Higgins, V., Sohaei, D., Diamandis, E.P., and Prassas, I. (2020). COVID-19: from an acute to

chronic disease? Potential long-term health consequences. Crit Rev Clin Lab Sci, 1-23.

Hirschtick, J.L., Titus, A.R., Slocum, E., Power, L.E., Hirschtick, R.E., Elliott, M.R., McKane,

P., and Fleischer, N.L. (2021). Population-based estimates of post-acute sequelae of SARS-CoV-

2 infection (PASC) prevalence and characteristics. Clin Infect Dis.

Kantonen, J., Mahzabin, S., Mayranpaa, M.I., Tynninen, O., Paetau, A., Andersson, N., Sajantila,

A., Vapalahti, O., Carpen, O., Kekalainen, E.*, et al.* (2020). Neuropathologic features of four

autopsied COVID-19 patients. Brain Pathol *30*, 1012-1016.

Karki, R., Sharma, B.R., Tuladhar, S., Williams, E.P., Zalduondo, L., Samir, P., Zheng, M.,

Sundaram, B., Banoth, B., Malireddi, R.K.S.*, et al.* (2020). COVID-19 cytokines and the

hyperactive immune response: Synergism of TNF-alpha and IFN-gamma in triggering

inflammation, tissue damage, and death. bioRxiv.

Karki, R., Sharma, B.R., Tuladhar, S., Williams, E.P., Zalduondo, L., Samir, P., Zheng, M.,

Sundaram, B., Banoth, B., Malireddi, R.K.S.*, et al.* (2021). Synergism of TNF-alpha and IFN-

gamma Triggers Inflammatory Cell Death, Tissue Damage, and Mortality in SARS-CoV-2

Infection and Cytokine Shock Syndromes. Cell *184*, 149-168 e117.

Kemp, H.I., Corner, E., and Colvin, L.A. (2020). Chronic pain after COVID-19: implications for

rehabilitation. Br J Anaesth *125*, 436-440.

Klingenstein, M., Klingenstein, S., Neckel, P.H., Mack, A.F., Wagner, A.P., Kleger, A., Liebau,

S., and Milazzo, A. (2020). Evidence of SARS-CoV2 Entry Protein ACE2 in the Human Nose

and Olfactory Bulb. Cells Tissues Organs *209*, 155-164.

Ladds, E., Rushforth, A., Wieringa, S., Taylor, S., Rayner, C., Husain, L., and Greenhalgh, T.

(2020). Persistent symptoms after Covid-19: qualitative study of 114 "long Covid" patients and

draft quality principles for services. BMC Health Serv Res *20*, 1144.

Lai, J.S., Cella, D., Choi, S., Junghaenel, D.U., Christodoulou, C., Gershon, R., and Stone, A.

(2011). How item banks and their application can influence measurement practice in

rehabilitation medicine: a PROMIS fatigue item bank example. Arch Phys Med Rehabil *92*, S20-

27.

Lam, M.H., Wing, Y.K., Yu, M.W., Leung, C.M., Ma, R.C., Kong, A.P., So, W.Y., Fong, S.Y.,

and Lam, S.P. (2009). Mental morbidities and chronic fatigue in severe acute respiratory

syndrome survivors: long-term follow-up. Arch Intern Med *169*, 2142-2147.

Le Bert, N., Tan, A.T., Kunasegaran, K., Tham, C.Y.L., Hafezi, M., Chia, A., Chng, M.H.Y.,

Lin, M., Tan, N., Linster, M.*, et al.* (2020). SARS-CoV-2-specific T cell immunity in cases of

COVID-19 and SARS, and uninfected controls. Nature *584*, 457-462.

Li, Y.C., Bai, W.Z., and Hashikawa, T. (2020). The neuroinvasive potential of SARS-CoV2 may

play a role in the respiratory failure of COVID-19 patients. J Med Virol *92*, 552-555.

Liguori, C., Pierantozzi, M., Spanetta, M., Sarmati, L., Cesta, N., Iannetta, M., Ora, J., Mina,

G.G., Puxeddu, E., Balbi, O.*, et al.* (2020). Subjective neurological symptoms frequently occur

in patients with SARS-CoV2 infection. Brain Behav Immun *88*, 11-16.

Mahase, E. (2020). Covid-19: What do we know about "long covid"? BMJ *370*, m2815.

McLane, L.M., Abdel-Hakeem, M.S., and Wherry, E.J. (2019). CD8 T Cell Exhaustion During

Chronic Viral Infection and Cancer. Annu Rev Immunol *37*, 457-495.

McMahan, K., Yu, J., Mercado, N.B., Loos, C., Tostanoski, L.H., Chandrashekar, A., Liu, J.,

Peter, L., Atyeo, C., Zhu, A.*, et al.* (2021). Correlates of protection against SARS-CoV-2 in

rhesus macaques. Nature *590*, 630-634.

Meckiff, B.J., Ramirez-Suastegui, C., Fajardo, V., Chee, S.J., Kusnadi, A., Simon, H.,

Eschweiler, S., Grifoni, A., Pelosi, E., Weiskopf, D.*, et al.* (2020). Imbalance of Regulatory and

Cytotoxic SARS-CoV-2-Reactive CD4(+) T Cells in COVID-19. Cell *183*, 1340-1353 e1316.

Miller, A.H. (2010). Depression and immunity: a role for T cells? Brain Behav Immun *24*, 1-8.

Mockus, T.E., Ren, H.M., Shwetank, and Lukacher, A.E. (2019). To Go or Stay: The

Development, Benefit, and Detriment of Tissue-Resident Memory CD8 T Cells during Central

Nervous System Viral Infections. Viruses *11*.

Muecksch, F., Wise, H., Batchelor, B., Squires, M., Semple, E., Richardson, C., McGuire, J.,

Clearly, S., Furrie, E., Greig, N.*, et al.* (2021). Longitudinal Serological Analysis and

Neutralizing Antibody Levels in Coronavirus Disease 2019 Convalescent Patients. J Infect Dis

223, 389-398.

Mueller, S.N., Gebhardt, T., Carbone, F.R., and Heath, W.R. (2013). Memory T cell subsets,

migration patterns, and tissue residence. Annu Rev Immunol *31*, 137-161.

Myasoedova, E., Crowson, C.S., Kremers, H.M., Therneau, T.M., and Gabriel, S.E. (2010). Is

the incidence of rheumatoid arthritis rising?: results from Olmsted County, Minnesota, 1955-

2007. Arthritis Rheum *62*, 1576-1582.

Pedersen, S.F., and Ho, Y.C. (2020). SARS-CoV-2: a storm is raging. J Clin Invest *130*, 2202-

2205.

Peng, Y., Mentzer, A.J., Liu, G., Yao, X., Yin, Z., Dong, D., Dejnirattisai, W., Rostron, T.,

Supasa, P., Liu, C.*, et al.* (2020). Broad and strong memory CD4(+) and CD8(+) T cells induced

by SARS-CoV-2 in UK convalescent individuals following COVID-19. Nat Immunol *21*, 1336-

1345.

Petersen, M.S., Kristiansen, M.F., Hanusson, K.D., Danielsen, M.E., B, A.S., Gaini, S., Strom,

M., and Weihe, P. (2020). Long COVID in the Faroe Islands - a longitudinal study among non-

hospitalized patients. Clin Infect Dis.

Polack, F.P., Thomas, S.J., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Perez, J.L., Perez

Marc, G., Moreira, E.D., Zerbini, C.*, et al.* (2020). Safety and Efficacy of the BNT162b2 mRNA

Covid-19 Vaccine. N Engl J Med *383*, 2603-2615.

Rasa, S., Nora-Krukle, Z., Henning, N., Eliassen, E., Shikova, E., Harrer, T., Scheibenbogen, C.,

Murovska, M., Prusty, B.K., and European Network on, M.C. (2018). Chronic viral infections in

myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). J Transl Med *16*, 268.

Rodda, L.B., Netland, J., Shehata, L., Pruner, K.B., Morawski, P.A., Thouvenel, C.D., Takehara,

K.K., Eggenberger, J., Hemann, E.A., Waterman, H.R.*, et al.* (2021). Functional SARS-CoV-2-

Specific Immune Memory Persists after Mild COVID-19. Cell *184*, 169-183 e117.

Roederer, M., Nozzi, J.L., and Nason, M.C. (2011). SPICE: exploration and analysis of post-

cytometric complex multivariate datasets. Cytometry A *79*, 167-174.

Rosen, S.F., Ham, B., Haichin, M., Walters, I.C., Tohyama, S., Sotocinal, S.G., and Mogil, J.S.

(2019). Increased pain sensitivity and decreased opioid analgesia in T-cell-deficient mice and

implications for sex differences. Pain *160*, 358-366.

Sahin, U., Muik, A., Derhovanessian, E., Vogler, I., Kranz, L.M., Vormehr, M., Baum, A.,

Pascal, K., Quandt, J., Maurus, D., *et al.* (2020). COVID-19 vaccine BNT162b1 elicits human

antibody and TH1 T cell responses. Nature *586*, 594-599.

Sant, A.J., DiPiazza, A.T., Nayak, J.L., Rattan, A., and Richards, K.A. (2018). CD4 T cells in

protection from influenza virus: Viral antigen specificity and functional potential. Immunol Rev

284, 91-105.

Sekine, T., Perez-Potti, A., Rivera-Ballesteros, O., Stralin, K., Gorin, J.B., Olsson, A.,

Llewellyn-Lacey, S., Kamal, H., Bogdanovic, G., Muschiol, S., *et al.* (2020). Robust T Cell

Immunity in Convalescent Individuals with Asymptomatic or Mild COVID-19. Cell *183*, 158-

168 e114.

Shin, H., Blackburn, S.D., Blattman, J.N., and Wherry, E.J. (2007). Viral antigen and extensive

division maintain virus-specific CD8 T cells during chronic infection. J Exp Med *204*, 941-949.

Sledzinska, A., Vila de Mucha, M., Bergerhoff, K., Hotblack, A., Demane, D.F., Ghorani, E.,

Akarca, A.U., Marzolini, M.A.V., Solomon, I., Vargas, F.A., *et al.* (2020). Regulatory T Cells

Restrain Interleukin-2- and Blimp-1-Dependent Acquisition of Cytotoxic Function by CD4(+) T

Cells. Immunity *52*, 151-166 e156.

Song, E., Zhang, C., Israelow, B., Lu-Culligan, A., Prado, A.V., Skriabine, S., Lu, P., Weizman,

O.E., Liu, F., Dai, Y., *et al.* (2021). Neuroinvasion of SARS-CoV-2 in human and mouse brain. J

Exp Med *218*.

Sorge, R.E., Mapplebeck, J.C., Rosen, S., Beggs, S., Taves, S., Alexander, J.K., Martin, L.J.,

Austin, J.S., Sotocinal, S.G., Chen, D., *et al.* (2015). Different immune cells mediate mechanical

pain hypersensitivity in male and female mice. Nat Neurosci *18*, 1081-1083.

Syed, A., Khan, A., Gosai, F., Asif, A., and Dhillon, S. (2020). Gastrointestinal pathophysiology

of SARS-CoV2 - a literature review. J Community Hosp Intern Med Perspect *10*, 523-528.

Tang, E., Ekundayo, O., Peipert, J.D., Edwards, N., Bansal, A., Richardson, C., Bartlett, S.J.,

Howell, D., Li, M., Cella, D*., et al.* (2019). Validation of the Patient-Reported Outcomes

Measurement Information System (PROMIS)-57 and -29 item short forms among kidney

transplant recipients. Qual Life Res *28*, 815-827.

Tebo, A.E., Fuller, M.J., Gaddis, D.E., Kojima, K., Rehani, K., and Zajac, A.J. (2005). Rapid

recruitment of virus-specific CD8 T cells restructures immunodominance during protective

secondary responses. J Virol *79*, 12703-12713.

Tian, Y., Babor, M., Lane, J., Schulten, V., Patil, V.S., Seumois, G., Rosales, S.L., Fu, Z.,

Picarda, G., Burel, J*., et al.* (2017). Unique phenotypes and clonal expansions of human CD4

effector memory T cells re-expressing CD45RA. Nat Commun *8*, 1473.

Tian, Y., Babor, M., Lane, J., Seumois, G., Liang, S., Goonawardhana, N.D.S., De Silva, A.D.,

Phillips, E.J., Mallal, S.A., da Silva Antunes, R*., et al.* (2019). Dengue-specific CD8+ T cell

subsets display specialized transcriptomic and TCR profiles. J Clin Invest *129*, 1727-1741.

Tian, Y., Rong, L., Nian, W., and He, Y. (2020). Review article: gastrointestinal features in

COVID-19 and the possibility of faecal transmission. Aliment Pharmacol Ther *51*, 843-851.

Toapanta, F.R., Bernal, P.J., Kotloff, K.L., Levine, M.M., and Sztein, M.B. (2018). T cell

mediated immunity induced by the live-attenuated Shigella flexneri 2a vaccine candidate CVD

1208S in humans. J Transl Med *16*, 61.

van Doorn, A.S., Meijer, B., Frampton, C.M.A., Barclay, M.L., and de Boer, N.K.H. (2020).

Systematic review with meta-analysis: SARS-CoV-2 stool testing and the potential for faecal-

oral transmission. Aliment Pharmacol Ther *52*, 1276-1288.

Van Elslande, J., Oyaert, M., Ailliet, S., Van Ranst, M., Lorent, N., Vande Weygaerde, Y.,

Andre, E., Lagrou, K., Vandendriessche, S., and Vermeersch, P. (2021). Longitudinal follow-up

of IgG anti-nucleocapsid antibodies in SARS-CoV-2 infected patients up to eight months after

infection. J Clin Virol *136*, 104765.

Vehar, S., Boushra, M., Ntiamoah, P., and Biehl, M. (2021). Post-acute sequelae of SARS-CoV-

2 infection: Caring for the 'long-haulers'. Cleve Clin J Med *88*, 267-272.

Wang, T., Ji, Y.L., Yang, Y.Y., Xiong, X.Y., Wang, I.M., Sandford, A.J., Liang, Z.A., and He,

J.Q. (2015). Transcriptomic profiling of peripheral blood CD4(+) T-cells in asthmatics with and

without depression. Gene *565*, 282-287.

Weintraub, S., Dikmen, S.S., Heaton, R.K., Tulsky, D.S., Zelazo, P.D., Bauer, P.J., Carlozzi,

N.E., Slotkin, J., Blitz, D., Wallner-Allen, K.*, et al.* (2013). Cognition assessment using the NIH

Toolbox. Neurology *80*, S54-64.

Weiskopf, D., Schmitz, K.S., Raadsen, M.P., Grifoni, A., Okba, N.M.A., Endeman, H., van den

Akker, J.P.C., Molenkamp, R., Koopmans, M.P.G., van Gorp, E.C.M.*, et al.* (2020). Phenotype

and kinetics of SARS-CoV-2-specific T cells in COVID-19 patients with acute respiratory

distress syndrome. Sci Immunol *5*.

Wherry, E.J. (2011). T cell exhaustion. Nat Immunol *12*, 492-499.

Wherry, E.J., Barber, D.L., Kaech, S.M., Blattman, J.N., and Ahmed, R. (2004). Antigen-

independent memory CD8 T cells do not develop during chronic viral infection. Proc Natl Acad

Sci U S A *101*, 16004-16009.

Figure Legends

Figure 1: Study design and clinical data

A.) Flow chart describing patient populations and experimental assays for each sample. B.) Table

showing patient demographics and neurologic symptoms. C.) PROMIS-57 patient reported

survey T scores for CN (n=36) and CC (n=13) groups. D.) NIH Toolbox cognitive T scores for

CN (n=43). Horizontal black line represents the U.S. national average T score of 50; p values

relative to US national average by one sample Wilcoxon signed rank test. *p<0.05, ***p<0.005,

****p<0.0001 by two-tailed Student's t test.

Figure 2: Neuro-PASC long-haulers display altered reactivity to SARS-CoV-2 N and M

peptides compared to healthy convalescents.

A-B.) CN (PCR+ or seropositive neuro-PASC) and CC (PCR+ or seropositive healthy COVID

convalescent) groups display similar IFN-γ and IL-2 responses to peptides from SARS-CoV-2

Spike protein by ELISPOT. C-D.) CN samples show significantly enhanced IFN-γ responses to

the N3 peptide pool (C) and to the M1 and M3 peptide pools (D) compared with CC or healthy

controls. E-F.) N- and M- specific IL-2 production did not significantly differ between patient

subgroups. G.) Spike RBD IgG endpoint titer quantification for CN, CC, and HC groups. LOD is

limit of detection. Data representative of 10 experiments with all conditions plated in duplicate.

*p<0.05, **p<0.01, ***p<0.005, ****p<0.0001 by two-way ANOVA with Tukey's posttest.

Figure 3: N-specific CD4$^+$ Tfh cells are more activated and S-specific CD4$^+$ T cells display

enhanced polyfunctional granzyme production in neuro-PASC long-haulers.

A.) FACS plots show that Ag-specific CD4 Tfh from CN patients are more highly activated in

response to N peptide pool stimulation compared with CC or HC, but less activated by M

peptides. B). Quantification of AIM$^+$ Tfh cell activation. C-D.) N- and M-specific AIM$^+$ Tfh cell

percentages plotted by time sample obtained post-infection for CN and CC. No correlation was

found between these parameters. E.) Anti-SARS-CoV-2 nucleocapsid IgG endpoint titers for

CN, CC, and HC subjects shown in 3D. CN patients display significantly elevated anti-N IgG

titers compared with CC subjects. F.) Pie graphs show CD4$^+$ T cells from CC patients have

enhanced production of granzymes A, B, & M compared with CN or HC subjects in response to

SARS-CoV-2 Spike protein. G.) Heatmap quantifying polyfunctionality in different categories of

cytokine production between groups. CC subjects produced 2-fold more granzymes A/B/M than

CN patients. Data combined from 6 independent experiments with CN n=35, CC n=11, and HC

n=9. *p<0.05 using one-way ANOVA with Bonferroni's posttest (B) or a Permutation test (E).

All pie graphs are showing data after subtracting background (unstimulated condition).

Figure 4: Neuro-PASC long-haulers have decreased CD8$^+$ memory T cell activation and

function compared with healthy convalescents.

A.) Selected FACS plots showing CD8 TEM cells are less activated in CN vs. CC groups after S

and N peptide pool stimulation. B.) Quantification of AIM+ CD8 TEM cells after SARS-CoV-2

peptide stimulation. C.) CD8$^+$ TEMRA cells accumulate significantly in PBMC from CN

subjects compared with CC or HC. D.) CD8$^+$ TEMRA cells from CN patients are less activated

by S3/4 and N pools compared with those from CC subjects. E.) Quantification of AIM$^+$ CD8$^+$

TEMRA cells from D. F.) Pie charts demonstrating that S3/4-specific CD8$^+$ TEM are polarized

to produce more TNF-α in CC group while those from CN patients produce significantly more

IL-6. G.) CD8$^+$ TEMRA from CC patients are polarized towards category 2 cytokine production

in response to S3/4 stimulation compared with CN subjects. Black lines demonstrate boundaries

of category 2 cytokine production in each group. Data combined from 5 independent

experiments with n=35 CN, n=11 CC. *p<0.05, **p<0.01 using two-tailed Student's t test with

Welch's correction (B, C, E) or permutation test (F, G). All pie graphs are showing data after

subtracting background (unstimulated condition).

Figure 5: Clinical measures of cognitive dysfunction and depression correlate with altered

CD8[+] memory T cell function in neuro-PASC long-haulers.

A.) Higher T cell responses to N pool stimulation are positively correlated with high pain scores

in CN patients. B.) A positive trend exists between elevated N-specific IFN-γ production and

higher depression scores in CN patients. C.) CN patients scoring high on pain interference have

Spike-specific CD8[+] T cells producing more IL-6 than those scoring low. D.) Heatmap showing

CN with low pain scores have Spike-specific CD8[+] T cells producing significantly more

granzymes A or B than those reporting high pain levels. E.) Pie graph to demonstrate that CN

patients in Q4 for depression have significantly enhanced production of granzymes A/B/M

compared with patients with low depression scores. F.) Heatmap demonstrating that CN patients

with high depression scores had Spike-specific CD8[+] TEM producing 3-fold higher levels of

granzymes A, B, and M or perforin than those scoring low on depression. G.) Spike-specific

CD8[+] TCM from CN patients with high scores for executive function are more polarized to

produce granzymes A/B/M compared with CN patients from Q1. H.) Heatmap showing Spike-

specific CD8[+] TCM in Q1 CN patients are more polyfunctional and produce 8-10x more

granzymes A/B/M and perforin than those in Q4. Data representative of 5 independent

experiments with n=8-9/quartile. All pie graphs showing Ag-specific cytokine production are

background subtracted (unstimulated conditions). *$p<0.05$, **$p<0.01$, ****$p<0.001$ using

Permutation tests.

Figure 6: Spike-specific T cell responses wane within 4 months after vaccination in CC and

HC subjects while remaining high in CN patients over time.

A.) Vaccine study visit timeline. V0 was obtained before the first dose of either Pfizer or

Moderna mRNA vaccines. V1 and V2 were conducted 3 weeks after the first and second doses,

respectively. B.) Vaccine study subject demographics. C.) Longitudinal anti-Spike RBD IgG

responses from V0-V4 across groups. Antibody titers are highest in CN patients and wane most

quickly in HC subjects. D.) IFN-γ production from Spike-specific T cells do not significantly

increase in CC and HC groups while remaining high in CN up to V4 post-vaccination. Total S-

specific IFN-γ SFU calculated by averaging responses from each sub-pool for each participant

(data in Fig. S9). Data combined from 10 individual experiments with all ELISA conditions done

in triplicate and all ELISPOT conditions plated in duplicate. *$p<0.05$, **$p<0.01$, ***$p<0.005$,

****$p<0.0001$ by two-way ANOVA with Tukey's posttest or by two-tailed Student's t test.

A.

B.

Patient Demographics	CN	CC	HC
Participants	56	24	31
Sex	38F, 18M	16F, 8M	19F, 12M
Race/Ethnicity	39 White 9 Hispanic 4 Asian 4 Black	15 White 6 Hispanic 3 Asian	11 White 5 Hispanic 13 Asian 2 Black
Hospitalization (%)	8 (14.3)	1 (4.2)	0 (0)
Days post-symptom onset (range)	193.4 (20-407)	155.0 (44-458)	NA
Neurologic symptoms (%)	56 (100)	0 (0)	0 (0)
Brain Fog	49 (87.5		
Fatigue	46 (82.1)		
Headache	41 (73.2)		
Dizziness	33 (58.9)		
Anosmia	35 (62.5)		
Dysgeusia	33 (58.9)		
Myalgias	34 (60.7)		

C.

D.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A.

B.

Vaccine Study Demographics	CN	CC	HC
Participants	19	12	20
Mean Age (range)	50.8 (33-70)	29.2 (22-63)	29.9 (22-59)
Sex	13F, 6M	7F, 5M	13F, 7M
Race/Ethnicity	12 White 5 Hispanic 1 Asian 1 Black	8 White 1 Hispanic 3 Asian	8 White 4 Hispanic 7 Asian 1 Black
Vaccine Brand	13 Pfizer 6 Moderna	10 Pfizer 2 Moderna	16 Pfizer 4 Moderna

C.

D.

Figure 6