1   **Full vaccination against COVID-19 suppresses SARS-CoV-2 delta variant and**

2   **spike gene mutation frequencies and generates purifying selection pressure**

4   **Ting-Yu Yeh and Gregory P. Contreras**

6   Agricultural Biotechnology Laboratory, Auxergen Inc., Columbus Center, 701 E Pratt

7   Street, Baltimore, MD 21202, USA

8
9   *Ting-Yu Yeh is Corresponding author
10
11   E-mail: yehty@auxergen.com

12   Tel: 1-443-762-1974

20   This paper is dedicated to Professor Austin Hughes (1949-2015), an evolution

21   biologist at University of South Carolina, for his lifetime achievement in neural

22   theory and adaptive molecular evolution.

23
24

Abstract

COVID-19 vaccination resistance has become a major challenge to prevent global SARS-CoV-2 transmission. Here we report that the vaccination coverage rate is inversely correlated to the mutation frequency of the full genome (R^2=0.878) and spike gene (R^2=0.829) of SARS-CoV-2 delta variants in 16 countries, suggesting that full vaccination against COVID-19, with other mitigation strategies, is critical to suppress emergent mutations. Neutrality analysis of DH and Zeng's E tests suggested that directional selection was the major driving force of delta variant evolution. To eliminate the homogenous effects (population expansion, selective sweep etc.), the synonymous (D_{syn}) and nonsynonymous (D_{nonsyn}) polymorphisms of the delta variant spike gene were estimated with Tajima's D statistic. Both D ratio (D_{nonsyn}/D_{syn}) and ΔD (D_{syn}-D_{nonsyn}) have positive correlation with the full vaccination rate (R^2= 0.723 and 0.505, respectively) in 19 countries, indicating that purifying selection pressure of SARS-CoV-2 spike gene increased as the vaccination coverage rate increased. Taken together, our data suggests that vaccination plays an important role in the purifying selection force of spike protein of SARS-CoV-2 delta variants.

Introduction

Legally required vaccination against various infectious diseases is essential to public health policy in many countries. However, resistance to voluntary COVID-19 vaccination has emerged worldwide, including in the United States. In mid-June 2021, 33% of Americans said they were not willing to be vaccinated (Kirzinger et al, 2021). Public distrust has undermined COVID-19 vaccine acceptance, in association with a belief that the vaccine is ineffective, dangerous or compromises individual freedom (Schmelz et al, 2021). Therefore, overcoming COVID-19 vaccination resistance has become a major challenge to prevent global SARS-CoV-2 transmission.

Mutations drive genome variability, generating many different SARS-CoV-2 variants as the virus evolves to escape vaccine-mediated immunity and thereby, develop drug or vaccine resistance (Roy et al., 2020). The delta (lineage B1.617.2) variants were first documented in October 2020 in India and were designated variants of concern by the World Health Organization in May of 2021 due to its increased transmissibility and virulence (https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/). All SARS-CoV-2 mutations result from two major mechanisms: (1) spontaneous substitution/deletion of nucleotides, and (2) RNA recombination (Roy et al, 2020, Yeh and Contreras, 2020, Ignatieva et al, 2021, Yeh and Contreras, 2021). Until now, it remained unclear exactly how human vaccinations affected virus selective pressure.

Is there correlation between mutation frequency and vaccination?

To explore this question, we first analyzed the correlation between the rates of full

vaccination and the point mutation frequency (*Mf*) of COVID-19 delta variants'

genome in 20 countries. Complete SARS-CoV-2 genome sequences with high

coverage from June 20 to July 3 2021 in 20 countries: Australia (N=121), France

(N=788), Germany (N=955), Indonesia (N=97), India (N=171), Ireland (N=617),

Israel (N=333), Italy (N=642), Japan (N=105), Mexico (N=368), Netherland (N=456),

Norway (N=142), Portugal (N=782), Singapore (N=131), Spain (N=689),

Switzerland (N=131), Sweden (N=786), Turkey (N=428), United States (N=537), and

United Kingdom (UK, N=3534) were collected from the Global Initiative on Sharing

All Influenza Data (GISAID) (https://www.gisaid.org/). *Mf* was calculated as Pi / (Ln

× Ns). Pi is the number of instances of polymorphism detected within the

genome/locus. Ln is the nucleotide length of the genome/locus. Ns is the number of

sequenced entities present in the dataset (Roy et al., 2020). The information of fully

vaccinated rate on 3 July 2021 in individual counties were obtained in Our World in

Data (https://ourworldindata.org/covid-vaccinations).

We found that *Mf* was logarithmically reduced as the full vaccination rate increased

in 16 of 20 countries (R^2=0.878, P<10^{-7}, Figure 1A). To our knowledge, this is the

first evidence suggesting that vaccinations could successfully suppress viral

mutations. Since the spike protein is the target of the vaccination program, we

further examined mutations of delta variant spike gene. Likewise, we found that

the *Mf* of the spike gene was also logarithmically reduced as the full vaccination rate

increased this time in 17 of the 20 countries (R^2=0.829, $P<10^{-6}$, Figure 1B). With a

10.8% vaccinated rate, *Mf* was exceptionally low in Australia, likely as a result of the

strict state lockdown restrictions. In contrast, the *Mfs* are higher in Japan,

Switzerland, and United States, suggesting that their social mitigation strategies

have been less successful. This also suggests that he mutation frequency may be not

only affected by vaccination only, but may also reflect overall control of the virus.

How does vaccination affect selection pressure?

SARS-CoV-2 is subject to many powerful selection pressures which may contribute

to vaccine escape. Like other viruses, the delta variant appears to have been selected

for its increased transmissibility regardless of vaccination. Previously we reported

on the mutations and selection pressure of SARS-CoV-2 among passengers in close-

quarters quarantine on the Diamond Princess cruise ship between January and

March 2020 (Yeh and Contreras, 2021). Until then, it had been unclear whether

COVID-19 variants were evolving randomly or under selection pressures generated

by vaccination and/or other mitigation efforts.

To understand how selection pressure operates within COVID-19 virus populations,

we analyzed the polymorphism of the full SARS-CoV-2 genome using neutrality tests

based on the site-frequency spectrum: (1) Tajima's *D* test, (2) normalized Wu and

Fay's *DH* test, and (3) Zeng's *E* test, using DNASP6 software with the bat coronavirus

RaTG13 as the outgroup sequence (Rozas, et al., 2017). The Tajima *D* test compares

pairwise nucleotide diversity (π, the average number of nucleotide differences per

site between two sequences) and total polymorphism to infer selection and/or

demographic events (Tajima, 1989). *DH* test is sensitive to the changes in high-

frequency variants and primarily affected by directional selection and no other

driving forces, such as demographic expansion (Zeng et al., 2006). Zeng's *E* test

contrasts high and low frequency regions of frequency spectrum, so this test can

detect population growth after a sweep (Zeng et al., 2006). Statistical significance of

observed values of all tests was obtained by coalescent simulation with free

recombination after 10,000 replicates. The probability for each test statistic was

calculated as the frequency of replicates with a value lower than the observed

statistic (two-tailed test) by DNASP6 (Rozas, et al., 2017).

We observed that Tajima's *D* values were negative and significantly deviated from

zero (-2.199 to -2.878, *P* < 0.05) among 19 countries, indicating an excess of variants

of low frequency (Table 1). Because the negative Tajima's *D* values could result from

demographic expansion and/or purifying/positive selection, we further employed

normalized *DH* (insensitive to population growth) and Zeng's *E* test (Zeng et al.,

2006). Because fixation causes quick loss of high-frequency alleles by drift, and

excess of low-frequency mutations are generated, Zeng's *E* test is very sensitive to

population growth. Genomic polymorphism of these viral populations revealed

significantly negative *DH* values (-1.177 to -2.791, *P*<0.05), but *E* values were not

significantly different from zero (-0.04 to 0.482, *P*>0.05) (Table 1). Similar results

were obtained after confining our analysis to the spike gene of SARS-CoV-2 delta

variants using Tajima' D, DH, and Zeng's E tests (Table 1). Taken together, these data

suggest that directional selection was operating in these countries, and

demographic expansion was less likely the major driving force.

Selection pressure has been observed in unvaccinated SARS-CoV-2 populations,

such as in the aforementioned Diamond Princess shipboard lockdown (Yeh and

Contreras, 2021). Our current results lead to the hypothesis that vaccination could

contribute to a part of the selection pressures on SARS-CoV-2, especially on the

spike protein. It has been shown that, within populations, the frequency distribution

of non-synonymous polymorphisms is negatively skewed relative to the distribution

of synonymous polymorphisms under purifying selection (Hahn et al, 2002; Hughes,

et al. 2005). Therefore, this hypothesis can be examined using a modified Tajima's D

statistics, which takes more negative values for non-synonymous (D_{nonsyn}) than for

synonymous sites (D_{syn}) of spike gene under purifying selection (Hahn et al, 2002,

Hughes, et al., 2005). The average number of pairwise synonymous differences (kS)

and the average number of nonsynonymous pairwise differences (kN), the number

of synonymous segregating sites (S_S), and the number of nonsynonymous

segregating sites (S_N) were computed with equation as Austin Hughes described (1)

S*$_S$=S$_S$/a1, (2) S*$_N$=S$_N$/a1, where a1 is as $a_1 = \sum_{i=1}^{n-1} \frac{1}{i}$ (Tajima, 1989, Hughes, et al.,

2005). D_{syn} was defined as kS − S*$_S$, divided by the standard error of that difference,

and D_{nonsyn} was defined as kN − S*$_N$, divided by the standard error of that difference

(Tajima, 1989, Hughes, et al., 2005). The values of K_a/K_s (the ratio of the number of

nonsynonymous substitutions per non-synonymous site (K_a), to the number of

synonymous substitutions per synonymous site (K_s), in a given period of time), and

the neutrality index of the McDonald-Kreitman (MK) test of the spike protein

(RaTG13 as the outgroup sequence) were analyzed by DNASP6 software (Rozas, et

al., 2017).

Unlike Tajima's D test, it has been demonstrated that D_{nonsyn} and D_{syn} analysis is

independent of sample size, so the value of D_{nonsyn} or D_{syn} can be compared between

data sets of different sizes (Hughes et al., 2008). The same excess of low-frequency

alleles in non-synonymous polymorphism was also observed (D_{nonsyn} = -1.482 to

-2.714; $P < 0.01$). In addition, the K_a/K_s values were less than 1 and the neutrality

index values of the MK test were significantly more than 1 ($P<0.05$) in all countries

(Table 1). These results confirmed that purifying selection led to constraint on the

available neutral mutations at non-synonymous sites of the spike gene of delta

variants, consistent to previous reports of SARS-CoV-2 variants in 2020 (Chaw et al,

2020).

Because non-synonymous and synonymous mutations are affected unequally by

purifying selection, D_{nonsyn} is expected to be disproportionally lower than D_{syn} (Hahn

et al., 2002). It has been proposed that the value of ΔD (D_{syn}-D_{nonsyn}) increased as

purifying selection became stronger, and ΔD (D_{syn}-D_{nonsyn}) is also able to rule out the

homogenous effects (population expansion, selective sweep etc.) in neutrality tests

(Hahn et al, 2002). We observed that ΔD of the spike gene of SARS-CoV-2 delta

variants is positively proportional to fully vaccination coverage rate (R^2= 0.505,

P=0.001). When using D ratio (D_{nonsyn}/D_{syn}), the standard error terms in both

equations of D_{nonsyn} and D_{syn} cancel out the sample size effect (Hughes, 2008). The

positive correlation of D ratio and fully vaccination rate became even more evident

(R^2= 0.723, P<10^{-4}) (Figure 2). Therefore, purifying selection pressure of SARS-CoV-

2 spike gene increased as the vaccination coverage rate increased. This suggests that

vaccination, in combination with other mitigation strategies, plays an important role

in the purifying selection force of SARS-CoV-2 spike protein.

Application of neutrality tests and their limitations

Neutrality tests are very powerful tools to study population genetics and viral

evolution. However, these tests require careful interpretation due to two known

limitations in this study: (1) sampling across countries and the representation of

sequences in public databases, (2) the timing of introductions and vaccination

campaigns. Previously we have applied various neutrality tests to analyze SARS-

CoV-2 transmission and evolution in Diamond Princess cruise (Yeh and Contreras,

2021). In contrast to the shipboard quarantine, where the virus spread in a closed

environment, many more complicated factors (mitigation efforts, travel, herd

immunity, etc.) could influence viral evolution during pandemics among different

countries.

Bhatt et al. have shown that the Tajima D test performs very well with low type I

error rate (less than 5%) when the θ value (the substitutions per site of a sequence

alignment) is between 0.1 to 10^{-4} in different RNA viruses (Bhatt et al., 2010).

Because θ values in this study ranged between 0.0009 to 0.032, the type I error rate

is low. The Tajima D test is also much less influenced by RNA recombination, which

we have reported previously (Yeh and Contreras, 2020, Yeh and Contreras, 2021).

Tajima D test results depend on the interplay of many parameters: positive and

negative selection, population size dynamics, spatial structure and migration,

random genetic drift, etc. A key limitation of the Tajima D test is the difficulty to

separate the effects of each parameter. However, the selection pressure and

demographic expansion can be verified in combination with other neutrality tests.

Our results of DH and Zeng's E test suggests that purifying selection, not

demographic expansion, was the acting force. Tajima D may show negative values

due to a recent bottleneck event rather than selection, but the bottleneck effect

should impact all types of polymorphism equally (Tajima, 1989). Significant

differences between D_{nonsyn} and D_{syn} (either ΔD or D ratio (D_{nonsyn}/D_{syn})), which

efficiently eliminated the homogenous effects, strongly indicates that purifying

selection was at play and is positively corelated with vaccination coverage rate

within these samples.

The inter-species divergence K_a/K_s test has been shown to be poorly equipped to

detect positive selection in short time scales for the comparison of young

evolutionary lineages like SARS-CoV-2 (Mugal, et al., 2014). Consistent with other

analytical attempts, our K_a/K_s data was inconclusive (Table 1). This has been the

case with other applications of the K_a/K_s test in SARS-CoV-2 spike gene, which

resulted in extreme values and controversial results (Kang et al., 2021),

In contrast to K_a/K_s, the MK test cannot be used to locate specific sites under

selection, although this test enables us to estimate the adaptive substitution rate of a

gene (Smith and Eyre-Walker, 2002). Variants of the MK test have been used to

analyze the evolution of RNA viruses (Bhatt et al., 2010). Since the MK test relies on

the comparison of the ratio of polymorphism to fixed differences of both

synonymous and nonsynonymous mutations, selection of an outgroup sequence is

critical for this test. The number of fixed synonymous mutations may be

underestimated in a rapidly evolving virus like SARS-CoV-2 when a distant outgroup

is applied (Baudry and Depaulis, 2003). Since the K_a/K_s or the MK tests may not be

the appropriate tests for SARS-CoV-2 delta variants, we only included them to

confirm the observation of purifying selection of SARS-CoV-2 spike gene here as

well. We also did not include neutrality tests based on haplotype distribution or

linkage disequilibrium, because they are expected to be strongly affected by RNA

recombination that commonly occurred in SARS-CoV-2 populations (Yeh and

Contreras, 2020, Ignatieva et al, 2021, Yeh and Contreras, 2021).

Perspectives

More virulent strains of SARS-CoV-2 have emerged with enhanced transmissibility

and immune evasion properties. For example, multiple variants have escaped

neutralizing antibodies developed to target the spike protein receptor-binding or N-

terminal domain (Harvey, et al., 2021). What's more, the case numbers of

breakthrough infection caused by the delta variant have increased drastically

worldwide (Farinholt et al., 2021).

Of all countries studied, our observations show that higher vaccination rates

corresponded with lower mutation frequencies and higher D_{nonsyn}/D_{syn} values for

the whole genome and the spike gene. We conclude then, that as the vaccination

coverage rate increases, purifying selection forces of nonsynonymous mutations

also increase.

Thus, we recommend that: 1) universal vaccination should be administered as soon

as possible to suppress the generation of deadly mutations; 2) mitigations strategies

such as personal protection equipment, social distancing, etc., should be continued

until full vaccination is reached to prevent viral transmission; 3) genomic

surveillance should be undertaken to monitor for new mutations; and 4) more

sequence summary statistics of RNA virus evolution are required to facilitate

understanding new COVID-19 outbreaks.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Author contributions

All authors contributed to study concept, rationale, and initial manuscript drafts,

interpretation of data, and final manuscript preparation. All authors have read and

approved the final version of the manuscript.

Funding

No funding

Ethical approval

None declared.

References

Bhatt, B., Katzourakis, A., Pybus, O.G., (2010) Detecting natural selection in RNA

virus populations using sequence summary statistics. Infect Genet Evol., *10*, 421-30.

Baudry, E., Depaulis, F. (2003) Effect of misoriented sites on neutrality tests with

outgroup. Genetics, *165*, 1619-1622

Chaw, S.M., Tai, J.H., Chen, S.L., Hsieh, C.H., Chang, S.Y., Yeh, S.H., Yang, W.S., Chen, P.J.,

Wang, H.Y. (2020) The origin and underlying driving forces of the SARS-CoV-2

outbreak. J Biomed Sci., *27*, 73.

Farinholt, T., Doddapaneni, H., Qin, X., Menon, V., Meng, Q., Metcalf, G., Chao, H.,

Gingras M., Farinholt, P., Agrawal, C., Muzny, D.M., Piedra, P.A., Gibbs, R.A.,

Petrosino, J. (2021) Transmission event of SARS-CoV-2 Delta variant reveals

multiple vaccine breakthrough infections. medRxiv doi:

10.1101/2021.06.28.21258780.

Harvey, W.T., Carabelli, A.M., Jackson, B., Gupta, R.K., Thomson, E.C., Harrison, E.M.,

Ludden, C., Reeve, R., Rambaut, A., COVID-19 Genomics UK (COG-UK) Consortium,

Peacock S.H., Robertson D.L. (2021) SARS-CoV-2 variants, spike mutations and

immune escape. Nat Rev Microbiol. ,*19*, 409-424.

Hahn, M.W., Rausher, M.D., Cunningham, C.W. (2002) Distinguishing between

selection and population expansion in an experimental lineage of bacteriophage T7.

Genetics, *161*, 11-20.

Hughes, A.L., (2005) Evidence for Abundant Slightly Deleterious Polymorphisms in

Bacterial Populations. Genetics, 169, 533-538.

Hughes, A.L., Friedman, R., Rivailler, P., French, J.O., (2008) Synonymous and

nonsynonymous polymorphisms versus divergences in bacterial genomes. Mol Biol

Evol. ,*25*, 2199–2209.

Ignatieva, A., Hein, J., Jenkins, P.A, (2021) Evidence of ongoing recombination in

SARS-CoV-2 through genealogical reconstruction. bioRxiv,

https://doi.org/10.1101/2021.01.21.427579

Kang, L., He, G., Sharp, A.K., Wang, X., Brown, A.M., Michalak. P., Weger-Lucarelli, J.

(2021) A selective sweep in the Spike gene has driven SARS-CoV-2 human

adaptation. Cell, *184*, 4392-4400.

Kirzinger, A., Sparks, G., Brodie, M. (2021) KFF COVID-19 vaccine monitor-Rural

America. https://www.kff.org/coronavirus-covid-19/poll-finding/kff-covid-19-

vaccine-monitor-in-their-own-words-six-months-later/ Accessed 13 July 2021.

Mugal, C.F., Wolf, J.B., Kaj, I. (2013) Why time matters: codon evolution and the

temporal dynamics of dN/dS. Mol Biol Evol., *31*,212–231.

Rozas, J., Ferrer-Mata, A., Sánchez-DelBarrio, J.C., Guirao-Rico, S., Librado, P., Ramos-

Onsins, S.E., Sánchez-Gracia, A. (2017) DnaSP 6: DNA Sequence Polymorphism

Analysis of Large Data Sets. Mol Biol Evol., *34*, 3299–3302.

Roy, C., Mandal, S.M., Mondal, S.K., Mukherjee, S., Mapder, T., Ghosh, W., Chakraborty,

R. (2020) Trends of mutation accumulation across global SARS-CoV-2 genomes:

Implications for the evolution of the novel coronavirus. Genomics ,*112*, 5331-5342.

Schmelz, K., Bowles, S., (2021) Overcoming COVID-19 vaccination resistance when

alternative policies affect the dynamics of conformism, social norms, and crowding

out. Proc Natl Acad Sci U S A ,*118*, e2104912118.

Smith, N.G., Eyre-Walker, A., (2002) Adaptive protein evolution in *Drosophila*.

Nature, *415*, 1022–1024.

Tajima, F. (1989) Statistical method for testing the neutral mutation hypothesis by

DNA polymorphism. Genetics, *123*, 585–595.

Yeh, T.Y., Contreras, G.P. (2020) Emerging viral mutants in Australia suggest RNA

recombination event in the SARS-CoV-2 genome. Med J Aust ,*213*, 44-44.e1.

Yeh, T.Y., Contreras, G.P. (2021) Viral transmission and evolution dynamics of SARS-

CoV-2 in shipboard quarantine. Bull World Health Organ., *99*, 486–495.

Zeng, K., Fu, Y.X., Shi, S., Wu, C.I. (2006) Statistical tests for detecting positive

selection by utilizing high-frequency variants. Genetics, *174*, 1431-1439.

Figure Legend

Figure 1. Correlation between full vaccinated rate

(https://ourworldindata.org/covid-vaccinations) and mutation frequency (*Mf*) of

the full genome (A) or the spike gene (B) of SARS-CoV-2 delta variants from June 20

to July 3 2021 in 20 countries: Australia (AUS), France (FRA), Germany (GER),

Indonesia (IDA), India (IND), Ireland (IRL), Israel (ISR), Italy (ITA), Japan (JPN),

Mexico (MEX), Netherland (NED), Norway (NOR), Portugal (POR), Singapore (SGP),

Spain (ESP), Switzerland (SUI), Sweden (SWE), Turkey (TUR), United States (USA),

and UK. Logarithmic regression (solid) line was draw based on 16 countries (pink)

with a calculated 95% confidence interval (dashed lines). Japan, Switzerland, USA,

and Australia are labeled in different colors as outliers.

Figure 2. Correlation between full vaccinated rate and D ratio (D_{nonsyn} /D_{syn}) of the

spike gene of SARS-CoV-2 delta variants. Exponential regression (solid) line was

draw based on 19 countries (pink square) with a calculated 95% confidence interval

(dashed lines). Australia (AUS) is labeled as outliers.

Figure 1.

Figure 2.
361

362
363

365 Table 1. Neutrality analysis of SARS-CoV-2 delta variants of the full genome and the viral spike gene
366 from June 20 to July 3, 2021 in 20 countries. The statistical significance was estimated using 10,000
367 coalescent simulations in DNASP6. Not significant values (*P*>0.05) are indicated in italic.
368

Country	Tajima's D	DH	Zeng's E	DH	Zeng's E	Tajima's D	ΔD (D_{Nonsyn} -D_{syn})	D ratio (D_{Nonsyn} /D_{syn})	Ka /Ks	Neutrality index
Israel	-2.401	-2.539	*-0.040*	-4.468	*1.792*	-2.065	0.611	1.424	0.055	13.209
France	-2.518	-2.482	*-0.278*	-2.973	*0.176*	-2.482	0.307	1.144	0.055	15.612
Germany	-2.800	-2.703	*-0.370*	-3.831	*0.655*	-2.661	0.065	1.026	0.034	10.938
Portugal	-2.645	-2.269	*-0.558*	-3.16	*0.402*	-2.395	0.593	1.314	0.058	8.393
Spain	-2.660	-2.61	*-0.283*	*-1.606*	*-0.943*	-2.530	0.306	1.143	0.065	12.986
Australia	-2.578	-1.914	*-0.626*	*-1.949*	*-0.078*	-2.064	0.091	1.056	0.052	10.609
USA	-2.657	-2.207	*-0.594*	-4.551	*1.306*	-2.580	0.290	1.124	0.015	14.467
Ireland	-2.833	-2.069	*-0.868*	-2.337	*-0.464*	-2.659	0.289	1.126	0.055	12.385
Italy	-2.533	-2.341	*-0.388*	-3.691	*0.73*	-2.51	0.095	1.042	0.045	17.699
Turkey	-2.451	-2.638	*-0.252*	-3.748	*0.916*	-2.398	-0.09	0.961	0.044	16.292
Indonesia	-2.339	-2.116	*-0.211*	*-1.35*	*-0.342*	*-1.773*	-0.381	0.796	0.055	11.236
Japan	-2.523	-1.801	*-0.675*	-3.697	*1.095*	-2.459	-0.582	0.776	0.052	5.068
Mexico	-2.522	-2.532	*-0.166*	-3.111	*0.444*	-2.405	-0.307	0.869	0.073	11.340
Norway	*-1.566*	*-1.177*	*-0.370*	*-0.682*	*-0.732*	*-1.562*	0.359	1.307	0.054	9.059
Switzerland	-2.560	-2.319	*-0.279*	-4.591	*1.756*	-2.494	0.333	1.158	0.045	10.81
Sweden	-2.473	-2.295	*-0.383*	-4.275	*1.101*	-2.565	0.182	1.08	0.021	11.19
Netherlands	-2.372	-2.327	*-0.231*	-3.811	*0.893*	-2.48	-0.107	0.956	0.049	10.64
Singapore	-2.199	-2.791	*0.482*	-2.538	*0.507*	*-1.909*	0.500	1.364	0.053	9.059
India	-2.878	-2.470	*-0.469*	-3.961	*0.884*	-2.826	-0.073	0.793	0.051	7.102
UK	-2.748	-2.546	*-0.540*	-3.299	*0.168*	-2.663	0.355	1.162	0.013	32.537

Table 1. Neutrality analysis of SARS-CoV-2 delta variants of the full genome and the viral spike gene from June 20 to July 3, 2021 in 20 countries. The statistic significance was estimated using 10,000 coalescent simulations in DNASP6. Not significant values (*P*>0.05) are indicated in italic.

	Full genome			Spike gene						
Country	Tajima's D	DH	Zeng's E	DH	Zeng's E	Tajima's D	ΔD (D_{Nonsyn} -D_{syn})	D ratio (D_{Nonsyn} /D_{syn})	Ka /Ks	Neutrality index
Israel	-2.401	-2.539	*-0.040*	-4.468	*1.792*	-2.065	0.611	1.424	0.055	13.209
France	-2.518	-2.482	*-0.278*	-2.973	*0.176*	-2.482	0.307	1.144	0.055	15.612
Germany	-2.800	-2.703	*-0.370*	-3.831	*0.655*	-2.661	0.065	1.026	0.034	10.938
Portugal	-2.645	-2.269	*-0.558*	-3.16	*0.402*	-2.395	0.593	1.314	0.058	8.393
Spain	-2.660	-2.61	*-0.283*	*-1.606*	*-0.943*	-2.530	0.306	1.143	0.065	12.986
Australia	-2.578	-1.914	*-0.626*	*-1.949*	*-0.078*	-2.064	0.091	1.056	0.052	10.609
USA	-2.657	-2.207	*-0.594*	-4.551	*1.306*	-2.580	0.290	1.124	0.015	14.467
Ireland	-2.833	-2.069	*-0.868*	-2.337	*-0.464*	-2.659	0.289	1.126	0.055	12.385
Italy	-2.533	-2.341	*-0.388*	-3.691	*0.73*	-2.51	0.095	1.042	0.045	17.699
Turkey	-2.451	-2.638	*-0.252*	-3.748	*0.916*	-2.398	-0.09	0.961	0.044	16.292
Indonesia	-2.339	-2.116	*-0.211*	*-1.35*	*-0.342*	*-1.773*	-0.381	0.796	0.055	11.236
Japan	-2.523	-1.801	*-0.675*	-3.697	*1.095*	-2.459	-0.582	0.776	0.052	5.068
Mexico	-2.522	-2.532	*-0.166*	-3.111	*0.444*	-2.405	-0.307	0.869	0.073	11.340
Norway	*-1.566*	*-1.177*	*-0.370*	*-0.682*	*-0.732*	*-1.562*	0.359	1.307	0.054	9.059
Switzerland	-2.560	-2.319	*-0.279*	-4.591	*1.756*	-2.494	0.333	1.158	0.045	10.81
Sweden	-2.473	-2.295	*-0.383*	-4.275	*1.101*	-2.565	0.182	1.08	0.021	11.19
Netherlands	-2.372	-2.327	*-0.231*	-3.811	*0.893*	-2.48	-0.107	0.956	0.049	10.64
Singapore	-2.199	-2.791	*0.482*	-2.538	*0.507*	*-1.909*	0.500	1.364	0.053	9.059
India	-2.878	-2.470	*-0.469*	-3.961	*0.884*	-2.826	-0.073	0.793	0.051	7.102
UK	-2.748	-2.546	*-0.540*	-3.299	*0.168*	-2.663	0.355	1.162	0.013	32.537