A randomised controlled trial to test the effects of fish aggregating devices and SBC activities promoting fish consumption in Timor-Leste: A study protocol

Alexander Tilley[1], Kendra A Byrd[1], Lauren Pincus[1], Katherine Klumpyan[2], Katherine Dobson[2], Joctan do Reis Lopes[3], Kelvin Mashisia Shikuku[1]

[1]WorldFish, Bayan Lepas, Penang, Malaysia, [2] Mercy Corps Timor-Leste, [3]WorldFish Timor-Leste, Dili, Timor-Leste

ABSTRACT

Timor-Leste is one of the world's most malnourished nations where micronutrient-deficient diets are a contributing factor to the prevalence of child stunting, currently estimated to be 45.6% of children under five. Fish are an important source of micronutrients and one that may assist the country's predominantly rural population of agriculturalists to exit poverty and malnutrition. However, a small national fishing fleet producing low catch volumes place fish out of reach of most inland and upland populations where it is needed most. Fish consumption is very low in rural areas compared to coastal, regional, and global averages. This trial is a one-year, cluster-randomized, partially masked, controlled trial among families living in rural, inland Timor-Leste. This trial aims to test and compare the effects of two treatments, alone and in combination, on the frequency and volume of household fish consumption in upland areas as a proxy for improved dietary diversity and micronutrient intake. Treatment 1 is the installation of nearshore, moored fish aggregating devices (FADs) to improve catch rates with existing fishing gears. Treatment 2 is social and behaviour change (SBC) activities to promote fish consumption. Villages in inland communities will be randomized to receive treatment 1, treatment 2, both treatments, or neither treatment. Some households with one child under five will be recruited, and data will be collected at baseline (prior to the rollout of the treatments) and endline. Our study will determine the impact of an improved supply of fish, along with nutrition-oriented SBC activities, on the fish purchasing and consumption practices of rural, inland households. Findings from this study are urgently needed by small island developing states in order to make policy and investment decisions on how best to improve households' diets using locally available, nutrient-dense foods such as fish. Investments such as these are needed to break the cycle of malnutrition.

Keywords: FAD, Dietary diversity, VSLA, RCT, nutrition, aquatic foods

1. Background and rationale

Timor-Leste is a country afflicted by multiple burdens of malnutrition, and sustains one of the world's highest rates of stunting (low height for age), afflicting 47.1% of children under five [1]. Stunted growth is multifactorial, but in Timor-Leste, a contributing cause is likely to be a monotonous diet [2]. There is mounting evidence that fish is an underutilised resource that has the potential to improve dietary quality, but consumption of fish is low at ~6 kg/person/yr [3] compared to neighbouring countries in Asia and the global average of ~20.5 kg/person/yr [4]. However, this 6 kg average figure masks the large variation in fish consumption; coastal communities consume ~17 kg/person/year, while inland communities consume fish only seasonally, resulting in a consumption rate of ~4 kg/person/yr [3].

Fish consumption in inland and rural regions of Timor-Leste is constrained in part due to the low catch volumes landed by the small national fleet of mostly non-motorized canoes [5]. Additional barriers, such as limited road and market infrastructures in inland areas, also reduce inland consumers' access to fish. Decades of low access to marine fish in inland areas [3] has decreased consumers' expectations of consuming this food source and made them unlikely to demand it from local traders.

Fish aggregating devices (FADs) are a technology to concentrate fish to make them easier to find and catch, and there is evidence of their improving catch rates in both inshore [6,7] and offshore settings [8,9]. Offshore FADs can be anchored or drifting, and are used to target tuna and other highly valued pelagic fish [10,11]. Anchored nearshore FADs are utilized by shore-based fishers to catch small pelagic fishes such as mackerels and scads [6]. By enabling access to a previously inaccessible fish stock, FADs can increase the climate adaptive capacity of fishing communities by diversifying the species caught and by reducing the dependence on production from vulnerable reef habitat [13].

Investigating the barriers for fish from small-scale fishers to reach the populations in need is important for understanding the contribution of fisheries to nutrition security. Proximity to inland fisheries is associated with an increase in fish consumption in children in rural Sub-Saharan Africa [14]; however, more information is needed to determine if this relationship holds in other contexts. Fish aggregating devices can increase supply, but increased availability alone may not be adequate to increase consumption, especially when it comes to using animal-source foods for infant and child feeding [15]. Building consumers' awareness and demand for high-quality marine fish must go hand-in-hand with increasing fish supply to improve the contribution of fisheries to nutrition security in inland areas.

Beyond increasing supply and awareness on benefits of fish consumption, a woman's decision-making power, sociocultural practices and seasonality have strong impacts on a household's ability to consume fish. A Gender Equality and Social Inclusion Analysis (GESIA) conducted in 2017 [15,16] found that a major gender equity issue in nutrition relates to the constraints that women face in decision-making and their lack of control over food, which impacts significantly on nutrition choices and the prevention of malnutrition. Research has also found that mothers

(younger mothers in particular) were not confident to spend money on small quantities of meat or
canned fish without permission from their husband [17,18].

Promoting a healthy diet and optimal infant and child feeding practices through contextualized
SBC interventions is another avenue to improve dietary quality and eventual health outcomes.
The influence of targeted SBC approaches on diets and expenditures is mixed [19], however,
some programs have shown success [20]. A key component of successful interventions is a deep
understanding of the community in which the SBC intervention is taking place, and inclusion of
local perspectives and beliefs [21]. Mercy Corps, an international NGO, has been working in
Timor-Leste since 2007, and has provided interventions that are compatible with its ethnography
[22]. Mercy Corps provides rural households with support through a sequenced layering
approach. The sequence begins with financial services through Village Savings and Loans
Associations (VSLAs). With access to secure savings, a source of credit, and an opportunity to
earn interest on their investment, VSLAs provide a foundation for the promotion of key nutrition
practices through participatory discussions within established VSLA members [23]. Thus, our
study design will allow us to evaluate the feasibility of using VSLAs with contextually-appropriate
nutrition discussions to improve dietary quality. Furthermore, this study will provide robust
evidence to determine the impact of both supply and demand interventions, alone and in
combination, to increase fish consumption and improve dietary quality in rural, inland Timor-Leste.

2. Design and methods

Figure 1. *A CONSORT flow diagram of the progress through the phases of the parallel randomised trial.*

2.1 Objectives

93 Our trial aims to evaluate the effectiveness of increased fish production and targeted promotion
94 of the nutritional benefits of fish consumption through social behaviour change (SBC) activities on
95 the frequency and quantity of fish consumption in rural households in Timor-Leste. Findings from
96 our study will inform governments, practitioners, and donors on the implementation of effective
97 interventions to improve household diets through increased fish consumption.

2.2 Hypotheses

99 H_0 Deployment of FADs has no effect on volume and frequency of fish consumed

100 H_0 Providing SBC has no effect on volume and frequency of fish consumed

101 H_0 Combined deployment of FADs and SBC has no effect on volume and frequency of fish
102 consumed

103   H_0 Deployment of FADs and SBC have no differential impacts on consumption

2.3 Trial design

Our study will implement a one year-long, cluster randomised controlled trial (RCT) among rural
inland households. We will test the effects of a bundled intervention addressing both supply-side
and demand-side constraints to consumption of fish.3Intervention

On the supply side, the experiment will involve the deployment of two fish aggregating devices in
the nearshore fishing grounds of a coastal fishing village with already established trading links to
inland study sites. Randomisation of the interventions will occur at the site level. Using WorldFish
and government data on fisheries landing sites in six municipalities, rural coastal fishing villages
with over 10 active fishing vessels will be listed. Villages that already have or utilise FADs will be
excluded. Six villages (one per municipality) will then be randomly assigned to one of two RCT
arms, namely FAD (4 villages) and non-FAD (2 villages) (Figure 2). This treatment arm is
imbalanced 2:1 because although the efficacy of nearshore FADs at increasing catch rates of fish
has been established [6], it is also shown to be dependent upon local ecological and bathymetric
conditions. These coastal villages are geographically dispersed, hence minimizing the risk of
contamination between treatment and control sites. In addition, by mapping the value chains of
traders in baseline focus groups, we will minimise contamination of non-FAD sites with fish caught
at a FAD.

On the demand side, 3-6 inland villages within a 30 km radius of the coastal treatment sites that
have a VSLA established by Mercy Corps, will be further randomised into an SBC arm (treatment)
and a non-SBC arm (control). A further inclusion criterion for the villages is that fish traders from
the coastal landing sites confirm through interview that they currently sell fish products to those
respective villages. Villages will be randomly allocated to treatment or control arms using an Excel
random number table to a 1:1 allocation ratio. The principal researcher will generate the allocation
sequence.

Within each SBC inland village, Mercy Corps will layer nutrition content and learning through
VSLAs. The VSLA members will participate in skill-building activities that promote five key
behaviours[1] about fish nutrition and household decision making on fish consumption as part of
facilitated discussions, interactive learning sessions, and a video-based facilitated dialogue. At
the village level (N = 12), Mercy Corps will promote increased fish consumption through SBC
mass attendance events through marketing of fish vendors, interactive learning, and *edu-tainment*
competitions promoting fish consumption. SBC will raise awareness of the benefits of eating fish

---

[1]Key promoted behaviours include: 1) Incorporate fish into family meals at least twice a week; 2) Parents pick
bones out of fish for small children and start offering fish to infants at 6 months of age; 3) VSLA members save with
a purpose of protein consumption for their families, with a focus on fish consumption; 4) Couples initiate
conversations on a weekly basis on allocation of resources for weekly protein purchase, inclusive of fish; 5)
Households use resources to purchase fish two times a week.

as part of a balanced diet, with a particular focus on pregnant and lactating mothers, and children under 5 years of age. The non-SBC communities will not receive the nutrition campaign package.

Figure 2. A sketch representation of the two treatment levels of the randomised controlled trial. 1. Coastal nearshore fish aggregating devices and 2. Social and behaviour change activities in rural inland communities in Timor-Leste. This diagram is a visualization only and does not represent the location of villages in the study.

Table 1. Description of study arms and their exposures

Arm	Exposures
FAD + SBC	- Mercy Corps VSLA site - Within 30 km of a FAD - SBC activities (nutrition messaging and skill building on fish consumption to improve dietary diversity) - One or more retailers who provides the village with fish
FAD only	- Mercy Corps VSLA site - Within 30 km of a FAD - No SBC activities - One or more retailers who provides the village with fish
SBC only	- Mercy Corps VSLA site - SBC activities (nutrition messaging and skill building on fish consumption to improve dietary diversity) - One or more retailers who provides the village with fish
No FAD, No SBC (control)	- Mercy Corps VSLA site - No FAD within 30 km - No SBC activities - One or more retailers who provides the village with fish

2.4 Study setting

This study will be carried out in six municipalities of Timor-Leste (Bobonaro, Dili, Liquica, Manufahi, Manatuto and Covalima) (Figure 3). The rural, upland areas of these municipalities exhibit some of the highest rates of child stunting in the world at 47.1% [26].

Figure 3. Map of Timor-Leste showing municipalities. Red shading shows the municipalities represented by coastal producer sites. The black dots indicate where the FADS will be placed offshore.

2.5 Sample size and sampling procedure

The sampling frame comprises rural, inland households in Timor Leste. We conducted a power calculation using the command *clustersampsi* in STATA based on secondary data, and a primary outcome of fish consumption per household. We assumed a minimum detectable effect of 15% and an intra-cluster correlation coefficient equal to 0.05. In order to achieve at least 80% statistical power, we require a sample size of 575 inland community households selected from 25 communities (that is, we sample 23 households in each community). Adjusting for 25% attrition, we estimate that a sample size of 700 is required—about 30 households per site.

Sampling follows a multi-stage procedure. In the first stage, 6 coastal municipalities were purposively selected based on the coverage of Mercy Corps programming of VSLA groups. The second step involves compiling a list of rural coastal fishing villages with over 10 active fishing vessels and randomly selecting one village within each municipality. In the third step, a list of inland villages within a 30 km radius of each coastal villages and with active presence of a VSLA was compiled. Although we include only villages participating in VSLAs, the typical size of a VSLA (about 15 members) is less than the required number of households per village. In the fourth step, therefore, we generate a list of all households in the village, both VSLA members and non-members. All VSLA participating households are selected to participate in the study. In order to ensure the required sample size is satisfied (in accordance with the power analysis) we randomly

167 select additional households from the list of those not participating in VSLAs. The chosen
168 households will be visited and asked if they would be willing to participate in the study. Where
169 households are not willing, the closest neighbouring household will be asked, and so on.

2.6 Recruitment and Consent

Communities and study households will be informed of the study and sensitized by the research
team to its activities and purpose through community meetings and meetings with local leaders.
Treatment and control groups will not be informed of their assigned group to avoid the possibility
of introducing Hawthorne or John Henry effects. An informed consent including maintenance of
the confidentiality of personal data, and the possibility to refuse the consent without having to
justify the refusal, will be obtained from all community leaders and study participants prior to
responding to any surveys.

2.7 Data collection

2.7.1 FAD fisheries catch and effort data

At each of the six coastal sites, a baseline of fishing activities will be collected from participating
fishers including trip frequency, duration, method, location and substrate type, along with the
catch volume. Data will be collected from fishers as they return to shore by on-site enumerators,
consistent with the national fisheries monitoring system in Timor-Leste, PeskAAS (following [24]).
This is a tablet-based survey where government enumerators collect and upload near-real-time
landings data to the central database. All participation in vessel tracking and contribution of data
is voluntary.

Information collected includes the number of fishers (sex-disaggregated), boat and gear types,
individuals sizes and quantities of fish captured, trip duration, the primary habitat fished, and the
proportion and price of fish sold. These data are also paired with geospatial vessel tracks for
boats that are fitted with a GPS tracker. Fishing activities and catch monitoring will be established
prior to installation of any FADs, to evaluate any change in catch rates or production volume
following the FAD deployment. Enumerators at each of the six coastal sites will be responsible
for daily data collection at landing sites of small-scale fishers, and will collect data on fish catch
by family group.

2.7.2 Household surveys

Following enrolment and informed consent, a baseline survey will be conducted with one member
per household. The baseline survey will consist of household demographic information, asset
ownership, and food consumption questions to assess relative wealth and standard of living,
access to fish and other animal source foods, food insecurity, and the frequency and volume of
fish consumed at the household level. Qualitative 24-hour recalls will be done with the household
member responsible for food preparation. If there are children under five years old in the
household and the respondent is not the primary caregiver, the caregiver will be asked 24-hour
recall questions related to the diet of the child(ren). The 24-hour recalls will be used to calculate

the minimum dietary diversity for both the woman and child [25,26]. We will also ascertain if there
are any cultural or traditional beliefs that influence fish consumption for specific people or at
certain times.

Interviews will be conducted by enumerators extensively trained on the tools, methods, research
ethics and confidentiality. Questionnaires will be translated into Tetum and carried out in Tetum
or a combination of Tetum and other local languages. After enumerator training and survey pre-
testing, the survey tools and procedures will be adapted as necessary to account for errors in
interpretation by enumerators and respondents. Staff will inform the village chief and VSLA
members in advance of the baseline and endline data collection sessions, and the latter will
arrange for participants to be present.

214 2.7 Data management

All data will be kept confidential. Data will be submitted daily via 3G for review by in-country
researchers and only stored on password protected mobile phone tablets and laptops temporarily.
The household survey questionnaire will include skip logics and field specifications and delimiters
to minimise input errors, but also supervising researchers will flag any unusual data for checking
with enumerators. All variables will be visualized and checked for outliers. Outliers will be closely
inspected, and sensitivity analyses will be conducted on high leverage data points. Outcome
variables will be visually checked for normality and transformed if necessary. Participants will be
assigned randomly-generated ID numbers, and names and identifying information will be de-
linked from the dataset. Data will be analysed for groups and no individual response will be
identifiable. Survey data will be archived and accessible to only the research team and those
approved by the entire research team. Anonymised datasets will be securely maintained
indefinitely to facilitate future research for example pooled analysis with other studies for purposes
of meta-analysis or multi-site analysis.

228 2.8 Statistical analysis

2.8.1 Main outcome variables

The main outcomes of interest are:

1) *Per capita household fish consumption* measured as total quantity (g) of fish consumed
by the household in the previous seven days divided by the number adult equivalents in
the household.

2) *Frequency of fish consumed at household level* measured as the number of days fish were
consumed out of the previous seven days.

2.8.2 Other outcome variables:

1) *Frequency of fish consumption in women* measured as the number of days fish were
consumed out of the previous seven days.

2) *Frequency of fish consumption in children* measured as the number of days fish were
240   consumed out of the previous seven days.

241   3) *Average catch rates* calculated as the increase in fish availability by comparing average
242   catch rates prior to FAD deployment with catch rates post-deployment. Catch rate and
243   total production values for different coastal community sites will be calculated and
244   extrapolated following Tilley et al. [24].

245   4) *Average price and supply of fish* calculated as the change in price and volume of fish
246   available in upland village sites by comparing price and volume of fish proffered by traders
247   at baseline with volumes at midline and endline.

248   5) *Knowledge and practices about fish purchase and handling.* A series of questions
249   assessing knowledge of nutritional benefits of fish and handling practices will be
250   administered to respondents at baseline, midline and endline.

2.8.3 Treatment variable

The main treatment variable is a dummy variable that indicates to which of the four treatment
arms a village was assigned. Our primary model is an unadjusted model.

However, if we find that after unmasking, there are variables in the dataset that appear different
by treatment arm, we may consider adjusting for confounders in the model.

2.9 Mediation analysis

We will assess whether the effect of SBC is mediated by an exogenous exposure of coastal
communities to the FADs. Therefore, we will construct an interaction of the SBC treatment dummy
with a dummy variable equal to one if an inland community is linked to a coastal community with
access to a FAD and zero otherwise.

Additionally, we will assess whether increased fish consumption is mediated by a change in self-
reported knowledge. We will do this by constructing a dummy composite variable of all knowledge
questions, and looking for an interaction by knowledge and treatment arm on per capita fish
intakes.

2.10 Checking for spillover effects

In addition to our research design feature for controlling spillovers, we will formally test for
evidence of spillovers across neighbouring villages using Global Positioning System (GPS)
coordinates of the households. We test whether the presence of an individual from another
experimental arm in a neighbouring village affects fish consumption status of the neighbour. To
check whether fish consumption status of the control group neighbours were affected by
spillovers, we will compare outcomes of control group neighbours who are close to a treated
neighbour and control group neighbours further away from treated units. Using a border-to-
treatment dummy variable, a *t*-test will be conducted to check that control group neighbours' fish

consumption status were not significantly affected by the presence of a neighbour from another
experimental arm.

2.11 Dealing with attrition

Our study design incorporates strategies to minimize attrition including working with existing VSLA
groups. However, we cannot rule out that attrition may still occur. Therefore, in case of substantial
attrition, the reasons will be carefully recorded. This will help us to assess whether potential
causes of attrition are systematically linked to specific treatments. We will check whether attrition
rates are equal across the three experimental arms or if concentrated in one arm. High attrition is
potentially problematic, as it could introduce selection bias in our randomised experiment. In case
of high attrition, we will examine the implication of attrition for our results in several ways. First,
we will test whether attrition is affected by treatment assignment. If this is the case, we will
implement Lee bounds to estimate lower and upper bounds of treatment effects [28,29]. Second,
we will test whether our remaining sample is (still) balanced along key observable dimensions,
and whether attrition is driven by different characteristics across treatment groups. To do this, a
probit regression will be used where we regress attrition status on the treatment dummies,
household characteristics, and the interaction of these. The hypothesis that attrition is not
differentially determined by treatment status will be assessed through an F-test for the joint
significance of the interaction terms in this regression. If differential determinants of attrition are
observed, we will attempt to control for this through a weighting procedure as a robustness
analysis [30]. Specifically, we will follow a two-stage procedure. In the first stage, a logit regression
is used to estimate the predicted probability of having non-missing measures for our outcomes
given treatment assignment and a vector of observable covariates. In the second stage, we will
weight each observation using the inverse of the thus estimated probability of having a non-
missing measure of our outcomes. We will check whether our main results remain robust to all
these robustness tests.

2.12 Multiple hypothesis testing

Because we are making inference on a large number of hypotheses, it is possible that significant
results emerge from our analysis due to chance rather than actual treatment effects. We will follow
[31] and adjust the *p*-values using a number of different methods. We will calculate Romano-Wolf
adjusted *p*-values (following [32]) to correct for the familywise error rate (FWER), the probability
of making at least one false discovery among a family of comparisons. We will also calculate
sharpened *q*-values (as in [33]) to correct for the false discovery rate (FDR), the probability of
making at least one false discovery among the discoveries already made.

2.13 Balance at baseline

The goal of the randomization is to ensure that the treatment and control groups are similar in
terms of average characteristics at baseline. This will be tested formally through an F-test of joint
orthogonality using a multinomial logit regression, which tests whether the observable
characteristics in table 1 are jointly unrelated to treatment status [34,35]. Failure to reject this null
hypothesis indicates that the randomization succeeded in achieving balance across the

experimental arms. In addition, we will calculate the standardized difference in means [35,36].
We will check whether the standardized difference in means are below the threshold of 0.25 as
recommended in literature [37], indicating balance.

Table 1. Baseline characteristics by experiment arm

Variable	SBC only	FAD only	SBC + FAD	Control (no SBC & no FAD)
Panel A: *Outcomes*				
Per capita household fish consumption				
Frequency of fish consumed at household level				
Frequency of fish consumption in women				
Frequency of fish consumption in children				
Knowledge about nutrition benefits of fish				
Dietary diversity				
Panel B: *Other variables*				
Age of respondent				
Sex of respondent				
Education of respondent				
Household size				
Number of women of reproductive age (15-49)				
Number of infants				
Per capita household income				
Food insecurity				
Asset score				
Participation in other community projects				
Number of nutrition information exchange links				

Empirical estimation

Main estimation approach

Intent-to-treat (ITT) estimates will be presented following equation 1:

(1) $y_{iv} = \alpha + \sum_{k=1}^{3} \beta_k treat_v^k + \gamma_i W_{iv} + C_c + \varepsilon_{ic}$

where y_{iv} is the outcome variable for household i in village v (Per capita household fish
consumption, frequency of fish consumed at household level, frequency of fish consumption in
women, frequency of fish consumption in children, knowledge about nutrition benefits of fish). The
variable $treat_v^k$ denotes the treatment dummy variables. W_{ijm} is a vector of control variables
including household characteristics and baseline levels of outcome variables, C_c captures
municipality fixed effects, and ε_{ijm} is the usual idiosyncratic error term, clustered at the village
level (the unit of randomization). Here the parameter of interest is β_k, the average treatment effect.
Equation will be estimated using OLS. We will first estimate a parsimonious model with only the
treatment dummies and the municipality fixed effects. Then we will control for additional baseline

covariates. The control group in equation 1 is the no-FAD and no-SBC group. Equation 1 will be estimated separately for midline and endline subsamples. However, we will compare the magnitude of impacts at midline and endline.

2.14 Ethical and safety considerations

Ethics approval was received from the Timor-Leste Instituto Nasionau do Saude (National Institute of Health) in December 2020 (1934MS-INS/DE/ /XII/2020). Written informed consent will be obtained from all VSLA members and respondents before recruitment into the study. Based on the experience of similar surveys and trials, no harm is expected from trial participation.

2.15 COVID19

Fieldwork will be conducting according to Government of Timor-Leste directives regarding the Standard Operating Procedure to minimise transmission of COVID19. Precautions taken will include conducting interviews outside, at a distance not less than 2 metres from respondents and others, and with interviewer and respondent wearing masks.

3. Discussion

The development and comparison of new technologies and approaches for reducing malnutrition in Timor-Leste responds to an urgent need captured in the country's current ranking at the very bottom of the global hunger index [38]. The Timor-Leste Strategic Development Plan [39], stated the goal of doubling capture fisheries productivity (an increase of 6,500 t) by 2020 to harness more animal-source foods from fisheries and aquaculture, yet to do so the government must have robust scientific evidence on which to base investment decisions. Fisheries play an important role in nutrition and livelihoods for coastal dwellers in Timor-Leste [8,43], who consume almost three times more fish than the national average [40]. Yet, there remains a lack of economic incentives to invest in the sector due to poor infrastructure and low economic returns [41–43].

There is convincing evidence that since their adoption in the Pacific in the late 1970s, FADs have in many instances substantially increased SSF catch rates [8,44]. As such many Pacific nations have integrated FAD programs into their national fisheries action plans and policy [8,44]. However, there is very little empirical research on the contribution of FADs to improved small-scale fisheries livelihoods and nutrition. Tilley et al. (2019) [6] established that nearshore FADs effect on catch rates was highly dependent on site ecology, but on average FADs provided a return on investment in five months or less. Furthermore, the oceanographic characteristics of Timor-Leste (steep slopes and strong currents) limit the longevity of deployed FADs, so one year is sufficient time to see any changes in catch rates.

Our study seeks to understand if increased fish volumes in coastal areas is absorbed by current markets, or is sufficient surplus to drive greater supply into inland areas. It is designed to approach the push and pull market dynamic from either end, without directly incentivising actors in the market chain. By increasing the volume of fish available for sale, we are pushing surplus supply

into new and smaller markets; and by encouraging fish consumption through a contextualized SBC intervention, we aim to increase the demand for fish to pull more into rural market. We do not anticipate large effects on the market price of fish in the study timeline, but we will record fisher sale price as part of monitoring. The reason for not incentivising traders to go to specific study villages is the lack of a reliable mechanism through which this could be sustained beyond the project life cycle. Individual variation between traders and their extant connections with inland communities is likely to be an important factor in decisions over where to direct greater fish production. We aim to mitigate this as far as possible through selection of traders for baseline and endline focus groups that already trade in the study inland communities.

While many people are aware of the importance of animal-source foods in Timor-Leste, and in particular the importance of fish [3], many women and children suffer from in adequate diets [45]. This study, utilizing robust evidence from interventions both alone and in combination, will generate important knowledge on household barriers to consuming more fish. In particular, our study will determine if additional encouragement to consume fish, when combined with the resources to purchase fish (via a Village Savings and Loan Association) is effective in increasing fish consumption, even without increased supply. As previously stated, there are other factors that can impact fish consumption, including seasonality, fish price, and decision-making power over household income expenditure and cultural practices. Alternatively, we may see that increased supply alone, in combination is additional household spending money is enough to increase fish consumption, even absent of an SBC campaign. These sets of interventions have not been tested in this context before, and the evidence will thus provide important information to policy-makers and stakeholders working to improved nutrition in small island developing states.

Limitations

There are two limitations of this trial. First, the dietary data on fish consumption is based on recall and self-reporting, and therefore risks response bias. It is expected that response bias would favour over-reporting of consumption, which may suggest lower effectiveness of the interventions. Sharing of foods with family members would likely be under-reported, again leading to underestimation of effectiveness. Collection of data on potential confounders / modifiers of effect.

While gender and power dynamics are recognised as important influencers of dietary choice and diversity, the intersection of these factors with increased supply and knowledge of nutrition benefits of fish will not be explored in this study. However, VSLAs are composed of both men and women members and SBC activities will involve all members and encourage collaborative food purchasing decision-making.

 The price of fish is a potential confounder in the study; however, we will collect market price and spending data on fish and other animal source foods. The market price variation of fish across sites (e.g. based on the geographical distance between producer and consumer) may affect access and thereby influence dietary preference. Information on prices and spending at each of the sites will be routinely collected as part of this study, and if variation occurs, will be included as a covariate in our adjusted models. In the same way, variable fish quality and safety between sites (fish deteriorating further with longer exposure to heat in reaching further sites) may affect consumer dietary preference. In baseline and endline surveys, we will ask if fish quality is a

407 concern or constraint to purchasing behaviour at each site. Travel time will be established to each
408 site under normal conditions, and temperature loggers will also be used to track the temperature
409 inside coolers carried by traders from producer to consumer sites.

4. Conclusion

From a programmatic point of view, if FADs and SBC prove successful at increasing rates and
volume of fish consumption, not only would it provide an acceptable and scalable program, but it
might also highlight where other technological innovations could be paired with SBC to drive
sustainable market effects without the need for costly incentives.

This study will contribute broadly to the body of knowledge on fish consumption patterns and
dietary choice by the rural poor in Timor-Leste, and due to close collaboration by WorldFish and
Mercy Corps with the Ministry of Agriculture and Fisheries and the FAO, our findings will be used
to inform decision making around new interventions, policies and investments in Timor-Leste. The
effects of FADs on the supply of fish to inland areas will further highlight possible opportunities
for using this technology to achieve national development priority objectives around nutrition and
rural development. Furthermore, new knowledge generated about the effects of SBC on dietary
choice with specific relevance to fish are of broad interest to those studying food systems, and
the dynamics of poverty in small-island developing states and beyond.

References

1. Timor-Leste: Peskiza Ai-Han No Nutrisaun 2020. Ministry of Health, GOTL; 2020.

2. Provo A, Atwood S, Sullivan EB, Mbuya N. Malnutrition in Timor-Leste:A review of the burden,
drivers, and potential response. World Bank; 2017. Available:
https://pdfs.semanticscholar.org/6622/1a1d1e2cdfe63edb1f356abb7669e4562dd8.pdf

3. AMSAT International,. Fish and animal protein consumption and availability in Timor-Leste.
Regional Fisheries Livelihoods Programme for South and Southeast Asia (GCP/RAS/237/SPA);
2011.

4. FAO. The State of World Fisheries and Aquaculture 2020. Sustainability in action. Rome: FAO;
2020. doi:10.4060/ca9229en

5. López-Angarita J, Hunnam KJ, Pereira M, Mills DJ, Pant J, Teoh SJ, et al. Fisheries and aquaculture
of Timor-Leste in 2019: Current knowledge and opportunities. Penang, Malaysia: WorldFish; 2019.
Available:
https://worldfish.archive.knowledgearc.net/bitstream/handle/20.500.12348/3737/Program-Report-
2019-15-Timor-Leste-Fisheries.pdf?sequence=2&isAllowed=y

6. Tilley A, Wilkinson SP, Kolding J, López-Angarita J, Pereira M, Mills DJ. Nearshore Fish
Aggregating Devices Show Positive Outcomes for Sustainable Fisheries Development in Timor-
Leste. Frontiers in Marine Science. 2019;6: 487.

7. Albert JA, Beare D, Schwarz A-M, Albert S, Warren R, Teri J, et al. The contribution of nearshore

fish aggregating devices (FADs) to food security and livelihoods in Solomon Islands. PLoS One.
2014;9: e115386.

8.   Sharp M. The benefits of fish aggregating devices in the Pacific. SPC Fisheries Newsletter.
2011;135: 28–36.

9.   Sharp M. Positive results of a FAD monitoring programme in Yap. SPC Fisheries Newsletter.
2014;143: 34–38.

10.   Girard C, Benhamou S, Dagorn L. FAD: Fish Aggregating Device or Fish Attracting Device? A new
analysis of yellowfin tuna movements around floating objects. Anim Behav. 2004;67: 319–326.

11.   Hilborn R, Medley P. Tuna Purse-Seine Fishing with Fish-Aggregating Devices (FAD): Models of
Tuna FAD Interactions. Can J Fish Aquat Sci. 1989;46: 28–32.

12.   Duffy JE, Lefcheck JS, Stuart-Smith RD, Navarrete SA, Edgar GJ. Biodiversity enhances reef fish
biomass and resistance to climate change. Proc Natl Acad Sci U S A. 2016;113: 6230–6235.

13.   O'Meara L, Cohen PJ, Simmance F, Marinda P, Nagoli J, Teoh SJ, et al. Inland fisheries critical for
the diet quality of young children in sub-Saharan Africa. Global Food Security. 2021;28: 100483.

14.   Robert RC, Bartolini RM, Creed-Kanashiro HM, Verney Sward A. Using formative research to
design context-specific animal source food and multiple micronutrient powder interventions to
improve the consumption of micronutrients by infants and young children in Tanzania, Kenya,
Bangladesh and Pakistan. Matern Child Nutr. 2020; e13084.

15.   TOMAK. Gender Equality and Social Inclusion Analysis. TOMAK; 2016. Available:
https://beamexchange.org/resources/1112/

16.   Wong JT, Bagnol B, Grieve H, da Costa Jong JB, Li M, Alders RG. Factors influencing animal-
source food consumption in Timor-Leste. Food Security. 2018;10: 741–762.

17.   Byrd K, Dentz HN, Williams A, Kiprotich M, Pickering AJ, Omondi R, et al. A behaviour change
intervention with lipid-based nutrient supplements had little impact on young child feeding
indicators in rural Kenya. Matern Child Nutr. 2019;15: e12660.

18.   Lindsey LLM. The influence of persuasive messages on healthy eating habits: a test of the Theory of
Reasoned Action when attitudes and subjective norm are targeted for change. J Vet Behav. 2017;22:
e12106.

19.   Menon P, Ruel MT, Nguyen PH, Kim SS, Lapping K, Frongillo EA, et al. Lessons from using
cluster-randomized evaluations to build evidence on large-scale nutrition behavior change
interventions. World Dev. 2020;127: 104816.

20.   Tumilowicz A, Neufeld LM, Pelto GH. Using ethnography in implementation research to improve
nutrition interventions in populations. Matern Child Nutr. 2015;11 Suppl 3: 55–72.

21.   Mercy Corps Timor-Leste Country Strategy 2019-2024. Mercy Corps; 2019.

22.   Australian Aid, TOMAK, Mercy Corps. The Impact of Savings Groups on Food Security &
Nutrition: What does the evidence show us? Australian Aid, TOMAK, Mercy Corps; 2018.
Available: http://tomak.org/wp-
content/uploads/2018/09/Think_Piece_2_Savings_Groups_English.pdf

23.   Timor-Leste Food and Nutrition Survey 2020 preliminary results: steady progress made, yet
challenges remain. [cited 28 Dec 2020]. Available: https://www.unicef.org/timorleste/press-
releases/timor-leste-food-and-nutrition-survey-2020-preliminary-results-steady-progress-made

24.   Tilley A, Dos Reis Lopes J, Wilkinson SP. PeskAAS: A near-real-time, open-source monitoring and
analytics system for small-scale fisheries. PLoS One. 2020;15: e0234760.

25.   FAO, FHI. Minimum Dietary Diversity for Women: A Guide for Measurement. FAO and FHI 360;
2016.

26.   WHO. Global Nutrition Monitoring Framework: operational guidance for tracking progress in
meeting targets for 2025. Geneva: World Health Organization; 2017.

27.   Lee DS. Training, wages, and sample selection: Estimating sharp bounds on treatment effects. Rev
Econ Stud. 2009;76: 1071–1102.

28.   Gerber AS, Green DP. Field Experiments: Design, Analysis, and Interpretation. W. W. Norton;
2012.

29.   Bulte E, Beekman G, Di Falco S, Hella J, Lei P. Behavioral Responses and the Impact of New
Agricultural Technologies: Evidence from a Double-blind Field Experiment in Tanzania. American
Journal of Agricultural Economics. 2014. pp. 813–830. doi:10.1093/ajae/aau015

30.   Arouna A, Michler JD, Yergo WG, Saito K. One Size Fits All? Experimental Evidence on the
Digital Delivery of Personalized Extension Advice in Nigeria. Am J Agric Econ. 2020;n/a.
doi:10.1111/ajae.12151

31.   Clarke D, Romano JP, Wolf M. The Romano-Wolf Multiple Hypothesis Correction in Stata. 2020.
Available: https://papers.ssrn.com/abstract=3513687

32.   Anderson ML. Multiple Inference and Gender Differences in the Effects of Early Intervention: A
Reevaluation of the Abecedarian, Perry Preschool, and Early Training Projects. J Am Stat Assoc.
2008;103: 1481–1495.

33.   McKenzie D. Identifying and spurring high-growth entrepreneurship: Experimental evidence from a
business plan competition. The World Bank; 2015.

34.   Deaton A, Cartwright N. Understanding and Misunderstanding Randomized Controlled Trials.
National Bureau of Economic Research; 2016.

35.   Canavari M, Drichoutis AC, Lusk JL, Nayga RM. How to run an experimental auction: a review of
recent advances. Eur Rev Agric Econ. 2019;46: 862–922.

36.   Cochran WG, Rubin DB. Controlling Bias in Observational Studies: A Review. Sankhyā: The Indian
Journal of Statistics, Series A (1961-2002). 1973;35: 417–446.

37.   von Grebmer K, Bernstein J, Alders R, Dar O, Kock R, Rampa F, et al. 2020 Global Hunger Index:
One Decade to Zero Hunger: Linking Health and Sustainable Food Systems. Bonn: Welthungerhilfe;
and Dublin: Concern Worldwide; 2020. Available:
https://www.globalhungerindex.org/pdf/en/2020.pdf

38.   GOTL. Timor-Leste Strategic Development Plan 2011-2030. Dili: Government of Timor-Leste;
2011.

39. Mills DJ, Tilley A, Pereira M, Hellebrandt D. Livelihood diversity and dynamism in Timor-Leste;
insights for coastal resource governance and livelihood development. Mar Policy. 2017. Available:
https://www.sciencedirect.com/science/article/pii/S0308597X17302695

40. Hartmann WD. Consultancy to Verify National RFLP Work Plan Activities, Collaborators and
Indicators - Final Report on Timor-Leste. Regional Fisheries Livelihoods Programme for South and
Southeast Asia; 2010.

41. Matsumoto WM, Kazama TK, Aasted DC. Anchored Fish Aggregating Devices in Hawaiian Waters.
Mar Fish Rev. 1981;43: 1–13.

42. Dempster T, Taquet M. Fish aggregation device (FAD) research: Gaps in current knowledge and
future directions for ecological studies. Rev Fish Biol Fish. 2004;14: 21–42.

43. Désurmont A, Chapman L. The use of anchored FADs in the area served by the Secretariat of the
Pacific Community (SPC): regional synthesis. Pêche thonière et dispositifs de concentration de
poissons, Caribbean-Martinique, 15-19 Oct 1999 15-19 octobre 1999. 2000. Available:
https://archimer.ifremer.fr/doc/00042/15283/

44. Campbell B, Hanich Q, Delisle A. Not just a passing FAD: Insights from the use of artisanal fish
aggregating devices for food security in Kiribati. Ocean Coast Manag. 2016;119: 38–44.

45. Bonis-Profumo G, Stacey N, Brimblecombe J. Maternal diets matter for children's dietary quality:
Seasonal dietary diversity and animal-source foods consumption in rural Timor-Leste. Matern Child
Nutr. 2020; e13071.