Original Article

Title: Treatment interruption is a risk factor for sickness presenteeism: a large-scale cross-sectional

study during the COVID-19 pandemic

Makoto Okawara[1], M.D., Tomohiro Ishimaru[1], M.D., M.P.H., Ph.D., Seiichiro Tateishi[2], M.D., Ph.D.,

Ayako Hino[3], M.D., Ph.D., Mayumi Tsuji[4], M.D., Ph.D., Akira Ogami[5], M.D., Ph.D., Tomohisa

Nagata[6], M.D., Ph.D., Shinya Matsuda[7], M.D., Ph.D., Yoshihisa Fujino[1], M.D., M.P.H., Ph.D., for

the CORoNaWork project

Author contributions: Y.F. planned the overall survey; all authors designed the research protocol

and questionnaire; M.O. handled the data and conducted statistical analysis with Y.F.; M.O. drafted

the initial manuscript; all authors revised and approved the final manuscript.

Authors' affiliations:

1 Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,

University of Occupational and Environmental Health, Japan

2 Department of Occupational Medicine, School of Medicine, University of Occupational and

Environmental Health, Japan

3 Department of Mental Health, Institute of Industrial Ecological Sciences, University of

Occupational and Environmental Health, Japan

4 Department of Environmental Health, School of Medicine, University of Occupational and

Environmental Health, Japan

5 Department of Work Systems and Health, Institute of Industrial Ecological Sciences, University of

Occupational and Environmental Health, Japan

6 Department of Occupational Health Practice and Management, Institute of Industrial Ecological

Sciences, University of Occupational and Environmental Health, Japan

7 Department of Preventive Medicine and Community Health, School of Medicine, University of

Occupational and Environmental Health, Japan

Corresponding author:

Makoto Okawara, M.D.

Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences, University

of Occupational and Environmental Health, Japan

1-1, Iseigaoka, Yahatanishiku, Kitakyushu, 807-8555, Japan

Tel: +81-93-691-7401

E-mail: makotouoeh@med.uoeh-u.ac.jp

Abstract

Objectives: This study examined the relationship between interruption to routine medical care

during the coronavirus disease 2019 pandemic and sickness presenteeism among workers in Japan.

Methods: A cross-sectional study using data obtained from an internet monitor questionnaire was

conducted. Interruption to medical care was defined based on the response "I have not been able to

go to the hospital or receive treatment as scheduled." The fraction of sickness presenteeism days in

the past 30 days was employed as the primary outcome. A fractional logit model was used for

analysis to treat bounded data.

Results: Of the 27,036 participants, 17,526 (65%) were workers who did not require routine medical

care, 8,451 (31%) were using medical care as scheduled, and 1,059 (4%) experienced interrupted

medical care. The adjusted odds ratio (aOR) of sickness presenteeism was significantly higher

among workers who experienced interrupted medical care (3.44; 95% confidence interval [CI]:

3.04–3.89) than those who did not require routine medical care. In terms of symptoms, the highest

aOR was observed among workers with mental health symptoms (aOR: 5.59, 95%CI: 5.04–6.20).

Conclusions: This study suggests the importance of continuing necessary treatment during a

pandemic to prevent presenteeism.

Keywords: occupational health; patient dropouts; presenteeism; COVID-19; regression analysis

Introduction

Sickness presenteeism is an increasingly important issue in occupational health. Aronsson defined sickness presenteeism as "people, despite complaints and ill health that should prompt rest and absence from work, still turning up at their jobs"[1]. Sickness presenteeism is the result of a choice made by a worker with ill-health, disease, or capacity loss between sickness presenteeism and sickness absence[2]. This decision is influenced by the individual's personality, values, economic status, workplace "demands for presence" and support for adaptation, and national culture and employment customs[2]. Evidence suggests that sickness presenteeism can lead to sickness absence and future worsening of physical and mental health conditions[3–9]. In addition, the impact of working while ill on productivity is also gaining attention, especially in North America[10,11]. A variety of diseases and health conditions have been found to be associated with sickness presenteeism, suggesting the importance of managing disease and maintaining good condition[4,12].

Under the coronavirus disease 2019 (COVID-19) pandemic, there is concern that both organizational and individual factors will increase sickness presenteeism above that observed under normal conditions[13–15]. Increased workload on workers can lead to a negative work culture around taking sick leave, such as where workers who choose to work while ill are valued for their loyalty to the company and motivation to work, thus promoting sickness presenteeism[16]. Examples of

individual factors that may increase sickness presenteeism include worsening economic situations

and job insecurity; increased telecommuting, which can make it easier for workers to work even

while sick; the impact of the pandemic on anxiety and mental health; worsening of health conditions

and diseases due to lifestyle changes; and worsening of chronic diseases due to the inability to access

medical resources. All of these factors are expected to lead to an increase in sickness presenteeism.

Interruption to medical care is an important problem in the COVID-19 pandemic. Access to

necessary routine medical care and medical resources is reportedly being affected in many countries

around the world[17–19]. In Japan, there is data showing that the number of prescriptions issued has

decreased[20]. Interruption to medical care can adversely affect management of chronic diseases and

delay the detection and treatment of new diseases[22]. In fact, excess deaths unrelated to COVID-19

have been reported[23]. Thus, interruption to medical care during the COVID-19 pandemic may lead to

worsening of non-COVID-19 diseases and health conditions.

During the COVID-19 pandemic, the number of people working while ill may increase as a

result of worsening health conditions arising from treatment interruptions and delays. This may

result in an increase in sickness presenteeism. However, few studies have examined the effect of

medical care interruption on sickness presenteeism during the COVID-19 pandemic. We

hypothesized that sickness presenteeism has increased among workers who experienced

interruptions to their medical care during the COVID-19 pandemic. The purpose of this study was to

examine the association between medical care interruption and sickness presenteeism in Japanese

workers during the COVID-19 pandemic.

Materials and Methods

We performed a cross-sectional study based on baseline survey data obtained in the

Collaborative Online Research on the Novel- Coronavirus and Work (CORoNaWork) project, a

prospective cohort study that performed a questionnaire-based survey of Internet monitors to

determine the effect of the COVID-19 pandemic on workers' health. Details of the study protocol are

published elsewhere[23]. The practical aspects of the survey, namely recruitment, data sampling, initial

data clean-up, and management of respondents IDs for tracking cohort data, were conducted by

Cross Marketing Inc. (Tokyo, Japan), which has 4.7 million registered monitors. Before completing

the online survey, participants read a description of the survey's aims and details about the handling

of their information. Only participants who agreed with the contents of the description were allowed

to participate. Participants' IDs were stored by the survey company. To participate in surveys offered

by the survey company, users had to register in advance. Once registered, participants were given an

ID to use when completing online surveys. Thus, all surveys were conducted anonymously. The

survey company generated cohort data after linking participants' IDs. Participants' personal data

were anonymized prior to receipt by the researchers, and were protected based on the survey

company's privacy policy. This study was approved by the Ethics Committee of the University

of Occupational and Environmental Health, Japan (Approval No. R2-079 and R3-006).

The baseline survey was conducted from December 22 to 26, 2020. A total of 33,302

participants aged from 20 and 65 years who indicated they were working when completing the

survey were included. A total of 605,381 registered monitors were emailed invitations to participate.

Sampling was designed such that sex and occupation (office and non-office workers) were

approximately equal among the five regions of residence. Specifically, we predetermined a total of

20 units comprising five regions of residence, two sexes, and two occupations. Sampling was

continued until each unit reached 1500 responses plus a margin of 10%, after which the unit was

closed to participation. We devised the five regions of residence according to infection and

geographic status: first, Japan's 47 prefectures were divided into four categories based on the

cumulative COVID-19 infection rate; second, the category with the highest infection rate was further

divided into Kanto and non-Kanto regions. A total of 55,045 participants answered the initial

screening questions, of whom 33,302 matched the survey's criteria (worker status, region of

residence, sex, and age). Participants answered one questionnaire item per page, with the overall

questionnaire containing 55 pages. Participants could review and change their responses using the

back button. Participants who provided fraudulent responses (n=6,266) according to the survey

company or a predefined definition of a fraudulent response were excluded. Fraudulent responses

included an unusually short response time (below 6 minutes), unusually short height (below 140 cm),

unusually low weight (below 30 kg), varying answers to similar questions in the survey (e.g.,

varying answers to questions about marital status or area of residence), and incorrect answers to

tiered questions used to identify inappropriate responses (e.g., choose the third highest number from

the following five numbers). After exclusion, responses from 27,036 participants were included in

the analysis.

Assessment of treatment status

We assessed the presence of disease and use of medical care using the following question:

"Do you have a disease that requires regular visits to the hospital or treatment?" Responses were "I

do not have any such disease"; "I am able to go to the hospital or receive treatment as scheduled"; "I

have not been able to go to the hospital or receive treatment as scheduled."

Those who answered "I do not have any such disease" were defined as workers who did not

require routine medical care, and thus did not have any disease that requires hospital visits or

treatment. Those who answered "I am able to go to the hospital or receive treatment as scheduled"

141 were defined as workers who used medical care. Those who answered "I have not been able to go to

142 the hospital or receive treatment as scheduled" were defined as workers who experienced interrupted

143 medical care.

145 Assessment of sickness presenteeism and other covariates

146  Respondents' number of sickness presenteeism days was ascertained based on the following

147 question and used as the primary outcome: "In the last 30 days, how many days have you worked

148 (including work from home) despite feeling that you really should have taken sick leave due to your

149 state of health?"[2] According to a previous study, the target period for sickness presenteeism can

150 range from four weeks to 12 months.[24] We chose a target period of 30 days to reduce the effects of

151 recall bias and changes in the COVID-19 pandemic situation in Japan.

152  Socioeconomic and work-related factors included sex, age, job type (mainly desk work,

153 mainly interpersonal communication, mainly physical work), marital status (married,

154 divorced/deceased, never married), equivalent income (household income divided by the square root

155 of the number of people in the household), education (junior high school, high school, vocational

156 school/junior college/college of technology, university/graduate school), frequency of working from

157 home (at least one day a month, at least one day a week, at least two days a week, at least four days a

week), company size (total number of employees in the respondent's main place of work [1 for

self-employed]), presence of a policy by the employer requesting that employees refrain from

attending work while ill (yes or no), and the number of days worked per week.

The cumulative infection rate of COVID-19 in the province of residence was employed as a

community-level variable.

To control for potential confounders, we also asked participants to indicate their main

symptoms using the following question: "Which of the following conditions or body parts give you

the most trouble during your work?" The options were "No problem"; "pain"; "movement";

"tightness, loss of energy, appetite, fever, dizziness, or feeling poor"; "toileting or elimination";

"mental health"; "skin, hair, or beauty"; "sleep"; "eyes"; "nose"; "ears"; and "other."

Statistical analysis

Age was treated as a continuous variable and reported as mean and standard deviation (SD).

The number of sickness presenteeism days was treated as a discrete variable and converted to a

fractional response variable by dividing by the maximum value of 30 days. Categorical variables

were reported as number and percentage. Equivalent income was categorized into quartiles.

We compared the results of Poisson regression, Zero-inflated Poisson regression (ZIP),

negative binomial regression, Zero-inflated Negative Binomial regression (ZINB), and fractional

logistic regression as statistical models. To handle data with excess zeros, which indicates a

population at low risk of sickness presenteeism, we considered a zero-inflated model[24]. Fractional

logistic regression[26] was also considered because the maximum possible number of days of sickness

presenteeism was 30, allowing the data to be treated as bounded data. As a measure of model fitness,

we compared the Akaike's Information Criterion (AIC), and ultimately adopted the fractional logit

model.

Fractional logistic regression analysis was conducted with the fraction of sickness

presenteeism days in last 30 days (0 indicates 0 days of sickness presenteeism; 1 indicates 30 days of

sickness presenteeism) as the dependent variable and the respondents' category of treatment status as

an independent variable.

We adjusted for the following potential confounders: sex, age, job type, marital status,

equivalent income, education, frequency of working from home, company size, cumulative infection

rate by prefecture, main symptoms, presence of a policy by the employer requesting that employees

refrain from attending work while ill, and the number of days worked per week.

In further analysis, we estimated the margins of sickness presenteeism days for each

treatment status and symptom. First, we used the same statistical model as that in the main analysis.

Second, we calculated the predictive margins of sickness presenteeism days, substituting measured

values for other covariates, dividing the data into 36 groups (3 treatment statuses and 12

symptoms)[27]. In the fractional logit model, because predictive margins were calculated as fractions,

we multiplied margins and standard errors by 30 to obtain predictive margins for sickness

presenteeism days in the last 30 days. Preliminarily, we confirmed the simple main effects for each

treatment status compared to workers who did not require routine medical care by adding the

interaction term between treatment status and symptoms to the model used for the main analysis. For

all analyses, the Bonferroni method was used to adjust for multiple comparisons.

All comparisons were performed in Stata (Stata Statistical Software: Release 16; StataCorp

LLC, TX, USA), with $p<0.05$ indicating statistical significance.

Results

The demographic and sociological characteristics of the analyzed population are shown in

Table 1. A total of 13,814 (51%) were men, and the mean age was 47.0 years (SD: 10.5). Of the total

population, 17,526 (65%) were workers who did not require routine medical care, 8,451 (31%) were

using medical care as scheduled, and 1,059 (4%) experienced interrupted medical care. The

distribution of sickness presenteeism is shown for the three treatment statuses in a histogram in

Figure 1. While the majority of respondents reported zero days of sickness presenteeism, a large

number also selected the maximum of 30 days. There were also small clusters at 5, 10, and 20 days,

which may be due to digit preference.

The association between treatment status and the fraction of sickness presenteeism days is

shown in Table 2. There was a significant association between the fraction of sickness presenteeism

days and treatment status. After adjusting for other covariates in the multivariate model, the odds

ratio (aOR) of sickness presenteeism days was significantly higher among workers who used

medical care (aOR: 1.36, 95%CI: 1.26–1.46, $p<0.001$) and workers who experienced interrupted

medical care (aOR: 3.28, 95%CI: 2.93–3.67, $p<0.001$) compared to workers who did not require

routine medical care.

The association between participants' main symptoms and the fraction of sickness

presenteeism days is shown in Table 3. There were significant associations between the fraction of

sickness presenteeism days and some symptoms using the model presented in Table 2. The highest

OR of sickness presenteeism days was observed for mental health symptoms (aOR: 5.35, 95%CI:

4.85–5.91, $p<0.001$).

The predictive margins of sickness presenteeism days for each treatment status and

symptom are shown in Table 4 and Figure 2. When the analysis was performed based on the three

treatment statuses, irrespective of symptom, the predictive margin of sickness presenteeism days

among workers who experienced interrupted medical care was 6.6 days (standard error [SE]=0.25),

while that among workers who did not require routine medical care was 1.4 days (SE=0.03). When

the analysis was performed based on the 36 treatment-symptom groups (3 treatment statuses and 12

symptoms), the largest predictive margin of sickness presenteeism days was observed for mental

health symptoms AND interrupted medical care (predictive margin: 9.9 days, SE=0.38). The simple

effect comparisons test, which included the interaction term between treatment status and symptoms,

showed that there were significant differences between workers with the same symptoms who did

and did not require routine medical care, and between workers with the same symptoms who

experienced interruption to medical care and who did not require routine medical care. For example,

the fraction of sickness presenteeism days significantly differed between those with mental health

symptoms who used medical care and those with mental health symptoms who did not require

routine medical care ($p<0.001$).

Discussion

This study demonstrated an association between treatment interruption and sickness

presenteeism among Japanese workers during the country's third wave of the COVID-19 pandemic.

Compared to workers who did not require routine medical care, workers who had diseases that

required routine medical care reported more days of sickness presenteeism, and those who

experienced interrupted medical care reported even more such days. Furthermore, our findings

revealed that there may be some symptoms that are more likely to lead to sickness presenteeism.

We found that workers who experienced interrupted medical care had increased sickness

presenteeism. This is because appropriate treatment can improve work function and productivity by

improving workers' health and subjective symptoms[10,28]. Employees who experience interrupted

treatment for chronic diseases may be forced to return to work due to fear of being laid off,

depending on the financial situation of their workplace during the pandemic. It is important to

continue regular treatments to manage disease and maintain health[29] to prevent inappropriate

presence at work while ill.

We found that the proportion of sickness presenteeism days varies by the type of symptoms

experienced by workers. While ORs among workers with symptoms related to mental health

problems, loss of energy/fever, body movement, and sleep were high, those among workers with

nose and skin, hair, beauty symptoms were moderate. Sickness presenteeism is the result of a

worker's choice to attend work despite being unwell rather than being absent from work. Many

previous studies conducted mainly in Europe have evaluated "sickness presenteeism" as a health

behavior based on whether or not workers "worked one or more days in a certain period of time with

a health condition for which they think they really should be absent"[2,30]. This is in contrast to studies

of "presenteeism" in North America, where the concept is evaluated as productivity loss due to

illness or a health condition using the Work Limitations Questionnaire or other tools[10,31]. One reason

as to why sickness presenteeism is more likely to be reported by workers experiencing mental health

problems may be that these symptoms are severe enough to cause "health conditions for which they

think they really should be absent." In contrast, physical symptoms such as those of allergic rhinitis

may not be considered "health conditions for which they think they really should be absent"

regardless of severity. If workers themselves do not consider their symptoms to be "health conditions

for which they think they really should be absent," they may not report experiencing sickness

presenteeism. However, there may be cases in which workers do not deem it necessary to be absent

from work, despite having a symptom that reduces productivity. Allergies with nose-related

symptoms are one example that causes loss of productivity due to presenteeism[11,32]. Workers may

not link these symptoms to sickness presenteeism due to differences in interpretation of "health

conditions that require absence from work." This is an important point when evaluating "sickness

presenteeism" as a health behavior.

We also found that the impact of continuing treatment on the prevention of sickness

presenteeism varied by symptom. Sickness presenteeism was more frequent in workers with

treatment interruption and symptoms related to pain, tightness and loss of energy, toileting and

elimination, mental health, sleep, and eyes than those who did not require routine medical care.

Among those who reported toileting and elimination, sleep, and eye symptoms, there was no

difference in sickness presenteeism between workers who used medical care and those who did not

require routine medical care. This suggests the importance of continuing necessary routine medical

care for preventing sickness presenteeism due to these symptoms. For those with pain, tightness and

loss of energy, and mental health symptoms, sickness presenteeism remained high even with

continued treatment, indicating the need to identify appropriate treatment and manage one's daily

health condition in addition to continuing treatment. Meanwhile, the prevalence of sickness

presenteeism was comparable among treatment statuses for participants with movement and mobility,

nose, and ear symptoms. This may be because individuals may not consider these symptoms

sufficiently adverse to engage in sickness presenteeism, or may experience chronic symptoms for

which support and adaptive behaviors have already been put into place such as movement symptoms.

The impact of continuing treatment on sickness presenteeism may be related to whether an

individual considers their symptoms to be sufficiently adverse to require an absence from work, or

whether or not the symptoms can be improved with treatment.

294   There are several limitations to this study. First, since this study is a survey of Internet

295 monitors, limitations regarding selection bias and generalizability are unavoidable. To reduce

296 potential bias, sampling and recruitment were conducted according to occupation and sex in each

297 region and the COVID-19 infection rate. To understand the characteristics of the target population of

298 this survey, we compared the results with those of national surveys and occupational surveys using

299 various batteries[23]. Second, we did not obtain detailed information related to treatment interruptions,

300 including the type of disease, duration, and reasons for interruption. We were thus unable to

301 determine whether the reason for interruption to treatment was due to patient-related reasons (e.g.,

302 economic situation and anxiety) or hospital-related reasons (e.g., schedule adjustment). Additionally,

303 the questions we used to identify exposure factors were related to the presence or absence of diseases

304 that require hospital visits and use of medical care. However, sickness presenteeism is not just

305 related to disease and medical status, but also a wide range of health conditions or concerns. Third,

306 interruptions to treatment may be the result of better disease control and improved health. It is

307 unclear how these factors would affect the occurrence of sickness presenteeism. Fourth, because this

308 study examined sickness presenteeism among workers during the third wave of the COVID-19

309 pandemic in Japan, caution is needed when interpreting the results. Further, causality is unclear. For

310 example, because most workers were told to remain home if they were experiencing COVID-like

311 symptoms during the pandemic, sickness presenteeism may have been reduced during this period.

However, it is unclear how such instructions affected overall sickness presenteeism, including that

associated with symptoms unrelated to COVID-19. Finally, we did not consider all possible

confounders affecting sickness presenteeism because we did not obtain information on some

confounders, such as job insecurity, annual leave rights, and the culture around employment and sick

leave in each company.

Conclusion

Interruption to medical care was associated with the occurrence of sickness presenteeism

during Japan's third wave of the COVID-19 pandemic. While the occurrence of sickness

presenteeism largely differed according to symptoms, it may be possible that there were important

cases of sickness presenteeism that were undetectable using the questionnaire. This study

demonstrates the importance of maintaining one's health condition and continuing necessary

treatment even during an infectious disease pandemic.

Acknowledgements:

This study was supported and partly funded by research grants from the University of

Occupational and Environmental Health, Japan (no grant number); the Japanese Ministry of Health,

Labour and Welfare (grant numbers H30-josei-ippan-002, H30-roudou-ippan-007, 19JA1004,

20JA1006, 210301-1, and 20HB1004); Anshin Zaidan (no grant number); the Collabo-Health Study

Group (no grant number); and Hitachi Systems, Ltd. (no grant number); and scholarship donations

from Chugai Pharmaceutical Co., Ltd. (no grant number).

The current members of the CORoNaWork Project, in alphabetical order, are as follows:

Dr. Yoshihisa Fujino (present chairperson of the study group), Dr. Akira Ogami, Dr. Arisa Harada,

Dr. Ayako Hino, Dr. Hajime Ando, Dr. Hisashi Eguchi, Dr. Kazunori Ikegami, Dr. Kei Tokutsu, Dr.

Keiji Muramatsu, Dr. Koji Mori, Dr. Kosuke Mafune, Dr. Kyoko Kitagawa, Dr. Masako Nagata, Dr.

Mayumi Tsuji, Ms. Ning Liu, Dr. Rie Tanaka, Dr. Ryutaro Matsugaki, Dr. Seiichiro Tateishi, Dr.

Shinya Matsuda, Dr. Tomohiro Ishimaru, and Dr. Tomohisa Nagata. All members are affiliated with

the University of Occupational and Environmental Health, Japan.

Disclosure:

Ethics Approval: This study was approved by the ethics committee of the University of Occupational

and Environmental Health, Japan (reference No. R2-079 and R3-006).

Informed Consent: Informed consent was obtained using a form on the survey website.

Registry and the Registration No. of the study/Trial: N/A

Animal Studies: N/A

Conflict of Interest: None declared.

References

1. Aronsson G, Gustafsson K, Dallner M. Sick but yet at work. An empirical study of sickness
presenteeism. *J Epidemiol Community Health*. 2000;54(7):502-509.

2. Aronsson G, Gustafsson K. Sickness presenteeism: prevalence, attendance-pressure factors, and
an outline of a model for research. *J Occup Environ Med*. 2005;47(9):958-966.

3. Skagen K, Collins AM. The consequences of sickness presenteeism on health and wellbeing
over time: A systematic review. *Soc Sci Med*. 2016;161:169-177.

4. Johns G. Presenteeism in the workplace: A review and research agenda. *J Organ Behav*.
2010;31(4):519-542.

5. Kivimäki M, Head J, Ferrie JE, et al. Working while ill as a risk factor for serious coronary
events: the Whitehall II study. *Am J Public Health*. 2005;95(1):98-102.

6. Janssens H, Clays E, De Clercq B, De Bacquer D, Braeckman L. The relation between
presenteeism and different types of future sickness absence. *J Occup Health*.
2013;55(3):132-141.

7. Bergström G, Bodin L, Hagberg J, Aronsson G, Josephson M. Sickness presenteeism today,
sickness absenteeism tomorrow? A prospective study on sickness presenteeism and future
sickness absenteeism. *J Occup Environ Med*. 2009;51(6):629-638.

8. Gustafsson K, Marklund S. Associations between health and combinations of sickness presence
and absence. *Occup Med* . 2014;64(1):49-55.

9. Gustafsson K, Marklund S. Consequences of sickness presence and sickness absence on health
and work ability: a Swedish prospective cohort study. *Int J Occup Med Environ Health*.
2011;24(2):153-165.

10. Schultz AB, Chen C-Y, Edington DW. The cost and impact of health conditions on
presenteeism to employers: a review of the literature. *Pharmacoeconomics*.
2009;27(5):365-378.

11. Nagata T, Mori K, Ohtani M, et al. Total Health-Related Costs Due to Absenteeism,
Presenteeism, and Medical and Pharmaceutical Expenses in Japanese Employers. *J Occup
Environ Med*. 2018;60(5):e273-e280.

12. Burton WN, Pransky G, Conti DJ, Chen C-Y, Edington DW. The association of medical

conditions and presenteeism. *J Occup Environ Med*. 2004;46(6 Suppl):S38-45.

13.   Reuter M, Dragano N, Wahrendorf M. Working while sick in context of regional
unemployment: a Europe-wide cross-sectional study. *J Epidemiol Community Health*.
2021;75(6):574-580.

14.   Tilchin C, Dayton L, Latkin CA. Socioeconomic Factors Associated With an Intention to Work
While Sick From COVID-19. *J Occup Environ Med*. 2021;63(5):363-368.

15.   Kinman G, Grant C. Presenteeism during the COVID-19 pandemic: risks and solutions. *Occup*
Med . Published online November 18, 2020. doi:10.1093/occmed/kqaa193

16.   Kinman G. Sickness presenteeism at work: prevalence, costs and management. *Br Med Bull*.
2019;129(1):69-78.

17.   Das AK, Mishra DK, Gopalan SS. Reduced access to care among older American adults during
CoVID-19 pandemic: results from a prospective cohort study. *bioRxiv*. Published online
November 30, 2020. doi:10.1101/2020.11.29.20240317

18.   Czeisler MÉ, Mpp KM, Clarke KEN, et al. Delay or avoidance of medical care because of
COVID-19–related concerns —. doi:10.1101/2020.04.22.20076141v1

19.   van Ballegooijen H, Goossens L, Bruin RH, Michels R, Krol M. Concerns, quality of life,
access to care and productivity of the general population during the first 8 weeks of the
coronavirus lockdown in Belgium and the Netherlands. *BMC Health Serv Res*. 2021;21(1):227.

20.   Lopresti M Seo T Sato. Pandemics and access to care: use of real-world DATA to examine the

IMPACT of COVID - 19 on pharmacy visits in Japan. *Value Health*.
doi:10.1016/j.jval.2020.08.1707

21.   Erol MK, Kayıkçıoğlu M, Kılıçkap M, et al. Treatment delays and in-hospital outcomes in acute
myocardial infarction during the COVID-19 pandemic: A nationwide study. *Anatol J Cardiol*.
2020;24(5):334-342.

22.   Woolf SH, Chapman DA, Sabo RT, Weinberger DM, Hill L. Excess Deaths From COVID-19
and Other Causes, March-April 2020. *JAMA*. 2020;324(5):510-513.

23.   Fujino Y, Ishimaru T, Eguchi H, et al. Protocol for a nationwide Internet-based health survey in
workers during the COVID-19 pandemic in 2020. *bioRxiv*. Published online February 5, 2021.

doi:10.1101/2021.02.02.21249309

24. Navarro A, Salas-Nicás S, Llorens C, Moncada S, Molinero-Ruíz E, Moriña D. Sickness
presenteeism: Are we sure about what we are studying? A research based on a literature review
and an empirical illustration. *Am J Ind Med*. 2019;62(7):580-589.

25. Papke LE, Wooldridge JM. Econometric methods for fractional response variables with an
application to 401(k) plan participation rates. *J Appl Econ*. 1996;11(6):619-632.

26. Williams R. Using the Margins Command to Estimate and Interpret Adjusted Predictions and
Marginal Effects. *Stata J*. 2012;12(2):308-331.

27. Burton WN, Morrison A, Wertheimer AI. Pharmaceuticals and worker productivity loss: a
critical review of the literature. *J Occup Environ Med*. 2003;45(6):610-621.

28. Chudasama YV, Gillies CL, Zaccardi F, et al. Impact of COVID-19 on routine care for chronic
diseases: A global survey of views from healthcare professionals. *Diabetes Metab Syndr*.
2020;14(5):965-967.

29. Hansen CD, Andersen JH. Going ill to work--what personal circumstances, attitudes and
work-related factors are associated with sickness presenteeism? *Soc Sci Med*.
2008;67(6):956-964.

30. Ishimaru T, Mine Y, Fujino Y. Two definitions of presenteeism: sickness presenteeism and
impaired work function. *Occup Med* . 2020;70(2):95-100.

Table 1. Basic characteristics of the study subjects

	Total (n=27,036)	Workers who did not require routine medical care (n= 17,526)	Workers who used medical care (n=8,451)	Workers who experienced interrupted medical care (n=1,059)
	n (%)	n (%)	n (%)	n (%)
Age, mean (SD)	47.0 (10.5)	45.4 (10.5)	50.4 (9.8)	46.0 (10.2)
Sex, men	13814 (51.1%)	8422 (48.1%)	4885 (57.8%)	507 (47.9%)
Marital status, married	15029 (55.6%)	9627 (54.9%)	4894 (57.9%)	508 (48.0%)
Job type				
Mainly desk work	13468 (49.8%)	8545 (48.8%)	4440 (52.5%)	483 (45.6%)
Mainly interpersonal communication	6927 (25.6%)	4621 (26.4%)	2032 (24.0%)	274 (25.9%)
Mainly physical work	6641 (24.6%)	4360 (24.9%)	1979 (23.4%)	302 (28.5%)
Equivalent income (Japanese Yen)				
≤2500000	6859 (25.4%)	4,374 (25.0%)	2,147 (25.4%)	338 (31.9%)
2500001-3800000	7148 (26.4%)	4,777 (27.3%)	2,119 (25.1%)	252 (23.8%)
3800001-5250000	7002 (25.9%)	4,594 (26.2%)	2,167 (25.6%)	241 (22.8%)
>5250000	6027 (22.3%)	3,781 (21.6%)	2,018 (23.9%)	228 (21.5%)
Education				
Junior high school	368 (1.4%)	244 (1.4%)	107 (1.3%)	17 (1.6%)
High school	6953 (25.7%)	4381 (25.0%)	2284 (27.0%)	288 (27.2%)
Vocational school, junior college, or technical college	6544 (24.2%)	4378 (25.0%)	1913 (22.6%)	253 (23.9%)

University or graduate school	13171 (48.7%)	8523 (48.6%)	4147 (49.1%)	501 (47.3%)
Number of employees				
1	2,556 (9.5%)	1,625 (9.3%)	848 (10.0%)	83 (7.8%)
2-49	7,999 (29.6%)	5,378 (30.7%)	2,320 (27.5%)	301 (28.4%)
50-999	9,703 (35.9%)	6,262 (35.7%)	3,036 (35.9%)	405 (38.2%)
1000-9999	4,719 (17.5%)	2,986 (17.0%)	1,552 (18.4%)	181 (17.1%)
≥10000	2,059 (7.6%)	1,275 (7.3%)	695 (8.2%)	89 (8.4%)
Frequency of working from home				
Almost zero	21,276 (78.7%)	13,944 (79.6%)	6,484 (76.7%)	848 (80.1%)
At least 1 day a month	615 (2.3%)	370 (2.1%)	217 (2.6%)	28 (2.6%)
At least 1 day a week	878 (3.2%)	571 (3.3%)	277 (3.3%)	30 (2.8%)
At least 2 days a week	1,477 (5.5%)	915 (5.2%)	501 (5.9%)	61 (5.8%)
At least 4 days a week	2,790 (10.3%)	1,726 (9.8%)	972 (11.5%)	92 (8.7%)
Presence of a policy by the employer requesting that employees refrain from attending work while ill, Yes	20,230 (74.8%)	13,057 (74.5%)	6,484 (76.7%)	689 (65.1%)

SD: standard deviation

Table 2. Association between treatment status and sickness presenteeism

Treatment status	Univariate			Multivariate*		
	OR	95% CI	*p*	Adjusted OR	95% CI	*p*
Workers who did not require routine medical care	Reference			Reference		
Workers who used medical care	1.57	1.47 1.69	<0.001	1.36	1.26 1.46	<0.001
Workers who experienced interrupted medical care	5.75	5.16 6.40	<0.001	3.28	2.93 3.67	<0.001

*Adjusted for sex, age, marital status, equivalent income, education, company size, job type, frequency of working from home, number of days worked per week, presence of a policy by the employer requesting that employees refrain from attending work while ill, cumulative infection rate for COVID-19 in the region of residence, and main symptom.

OR: odds ratio, CI: confidence interval

Table 3. Association between main symptoms and sickness presenteeism

Main symptom	Adjusted* OR	95% CI		p
No problem	Reference			
Pain	2.76	2.46	3.10	<0.001
Movement	3.02	2.57	3.55	<0.001
Tightness, loss of energy, appetite, fever, dizziness, or feeling poor	4.53	4.05	5.06	<0.001
Toileting or elimination	2.62	2.14	3.21	<0.001
Mental health	5.35	4.85	5.91	<0.001
Skin, hair, or beauty	1.96	1.60	2.39	<0.001
Sleep	3.07	2.73	3.45	<0.001
Eyes	2.01	1.74	2.31	<0.001
Nose	1.54	1.09	2.17	0.014
Ears	2.35	1.74	3.18	<0.001
Other	2.58	2.22	3.00	<0.001

*Adjusted for sex, age, marital status, equivalent income, education, company size, job type, frequency of working from home, number of days worked per week, presence of a policy by the employer requesting that employees refrain from attending work while ill, cumulative infection rate for COVID-19 in the region of residence, and treatment status.

OR: odds ratio, CI: confidence interval

Table 4. Predictive margins of sickness presenteeism days for each symptom and comparison between each treatment status

	Workers who did not require routine medical care (n=17,526)				Workers who used medical care (n=8,451)					Workers who experienced interrupted medical care (n=1,059)				
	n	%	Predictive margins*	Standard error	n	%	Predictive margins*	Standard error	p †	n	%	Predictive margins*	Standard error	p †
Total	17526	100	1.4	0.03	8451	100	2.2	0.05	<0.001	1059	100	6.6	0.25	<0.001
Main symptom														
No problem	10938	62.4	0.8	0.03	3642	43.1	0.9	0.03	1.000	160	15.1	2.6	0.15	<0.001
Pain	849	4.8	2.0	0.10	842	10.0	2.5	0.12	<0.001	144	13.6	6.1	0.33	<0.001
Movement	481	2.7	2.3	0.16	344	4.1	2.8	0.20	1.000	54	5.1	6.5	0.44	0.428
Tightness, loss of energy, appetite, fever, dizziness, or feeling poor	715	4.1	3.4	0.15	583	6.9	4.1	0.17	0.027	133	12.6	8.9	0.39	<0.001
Toileting or elimination	263	1.5	2.0	0.18	201	2.4	2.5	0.23	1.000	40	3.8	5.5	0.50	<0.001
Mental health	1143	6.5	3.8	0.14	909	10.8	4.7	0.17	<0.001	219	20.7	9.9	0.38	<0.001
Skin, hair, or beauty	389	2.2	1.5	0.14	160	1.9	1.9	0.18	0.006	29	2.7	4.9	0.44	0.513
Sleep	997	5.7	2.3	0.11	637	7.5	2.8	0.14	0.127	116	11.0	6.5	0.35	<0.001
Eyes	795	4.5	1.5	0.09	512	6.1	1.8	0.11	0.319	65	6.1	4.3	0.30	<0.001
Nose	98	0.6	1.2	0.20	71	0.8	1.5	0.25	1.000	9	0.9	3.4	0.54	1.000

Ears	80	0.5	1.7	0.25	84	1.0	2.2	0.30	1.000	12	1.1	5.7	0.73	0.056
Other	778	4.4	2.0	0.13	466	5.5	2.5	0.17	0.031	78	7.4	6.3	0.41	<0.001

*predictive margins: mean predicted number of sickness presenteeism days for each symptom and treatment status using the model in Table 2 and 3 with substitution of measured values for other covariates (adjusted using the Bonferroni method); calculated using the formula: mean predicted fraction × 30 (days)

† p-value for simple main effects for each treatment status compared to workers who did not require routine medical care using a model that included the interaction term for treatment status and main symptoms (adjusted using the Bonferroni method)

%: proportion of the total number of respondents for each treatment status

Figure 1. Number of days of sickness presenteeism among workers with each treatment status

(Figure 1)

Figure 2. Predictive margins with 95% confidence intervals for each treatment status and symptom

(Figure 2)

*others include loss of energy, appetite, fever, dizziness, or feeling poor

†Error bars indicate 95% confidence intervals

Predicted mean percent of sickness presenteeism days