Macular vascular complexity analysis of diabetes mellitus by swept-source optical coherence tomographic angiography

Jiahui Liu, [1] Shuhui Chen, [1] Zhiyi Xu, [1] Wei Wang, [2] Dingqiao Wang, [2] Yongyue Su, [2] Shulun Zhao, [1] Meichan Li, [1] Fengling Chen, [1] Chengguo Zuo, [2*] Minyu Chen[1*]

[1] Affiliated Dongguan People's Hospital, Southern Medical University, Dongguan, China

[2] State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangzhou, China

* Corresponding authors:

Minyu Chen, Ophthalmology department, Affiliated Dongguan People's Hospital, Southern Medical University, Dongguan, China, Email:

chenminyu99@tom.com

Chengguo Zuo, Zhongshan Ophthalmic Center, State Key Laboratory of Ophthalmology, Sun Yat-sen University, 54S.Xianlie Road, Guangzhou, China, Email:

chengguozuo@163.com

Word count: 3025

Funding information: This research was supported by the National Natural Science Foundation of China (82000901, 82102993), the Fundamental

32 Abstract

Purpose To evaluate the associations of retinal vascular complexity features,

including fractal dimension (FD) and blood vessel tortuosity (BVT), with the

severity of diabetic retinopathy (DR) by using noninvasive optical coherence

tomographic angiography (OCTA).

Methods This prospective cross-sectional study recruited ocular-treatment-

naïve patients with type 2 diabetes mellitus (DM) registered in the community

of Guangzhou, China. OCTA was used to obtain the measurements of FD and

BVT in the superficial capillary plexus (SCP) and the deep capillary plexus

(DCP). Univariate and multivariate linear regression was used to analyse the

correlation of FD and BVT in different layers with DR severity.

Results A total of 1282 patients with DM (1282 eyes), with a mean age of

64.2±7.8 years, were included. FD in the DCP decreased and BVT in the DCP

increased in DR patients compared to non-DR patients, even after adjusting

for confounding factors (P< 0.05). Trend analysis showed a significant

decrease in the FD values as the DR progressed, while the BVT progressively

increased with worsening DR severity (P< 0.01). The FD in DCP had a

statistically significant positive correlation with FD in SCP and a negative

correlation with BVT in SCP and BVT in DCP in all of the participants,

including the non-DR group, moderate DR group and severe DR group (P<

0.01).

Conclusion FD and BVT determined using OCTA might be useful parameters

for objectively distinguishing DR from non-DR and indicating DR progression.

Keywords: fractal dimension; blood vessel tortuosity; diabetic retinopathy;

OCTA

Introduction

Diabetic retinopathy (DR) is the leading cause of blindness in the

working-age population around the world [1-3]. Patients with diabetes mellitus

(DM) are usually symptomless until the advanced stages of DR, leading to a

poor visual prognosis after treatment. Thus, early detection and prompt

treatment of DR play a pivotal role in protecting DM patients from vision

impairment. Although the exact pathogenesis of DR remains unclear, it is

universally acknowledged that damage to the retinal microvasculature,

especially in the macula, might contribute to the occurrence and development

of DR[4, 5] . Capillary drop-out has been reported to precede proliferative

disease [6]. However, there are no available clinically quantitative parameters to

identify the degree of microvascular damage, especially in the early stage of

DR.

As optical coherence tomography angiography (OCTA) has become more

widely available, detailed microvascular networks can now be quantified

conveniently and noninvasively with high resolution. With OCTA, some

abnormal changes in the retinal capillary network in DM patients without DR

have been recognized, demonstrating that OCTA metrics could be early

indicators of subtle, subclinical alterations in DM before the first clinically

observable signs of DR [7-9]. Fractal dimension (FD) and blood vessel tortuosity

(BVT) have been used to evaluate the overall structural and geometric

complexity of retinal vasculatures. Recent studies revealed that FD and BVT

were altered in DR and therefore were DR biomarker candidates [2, 3, 10, 11].

However, these findings did not consider confounding factors, such as age,

sex, blood pressure, HbA1c or OCTA signal strength intensity. In addition, the

correlation of FD and BVT has not been elucidated to date. We are unaware

of any studies about vascular complexity features in diabetes patients from

Chinese communities. Therefore, the objective of this study was to assess FD

and BVT to identify localized complexity distortions across various stages of

DR with OCTA and to determine the relationship between them in a large

sample of Chinese patients with DM.

Methods

Participants

This prospective cross-sectional study was conducted in Zhongshan

Ophthalmic Center (ZOC) with the permission of the Ethics Committee of the

ZOC (2017KYPJ094). All investigations followed the tenets of the Declaration

of Helsinki. Subjects in the study were enrolled from November 01, 2017, to

December 30, 2020, in the community of Guangzhou, China. Patients aged

35–80 years with type 2 diabetes and without any history of eye treatments or

surgeries (ocular treatment naïve) were included in this study. The exclusion

criteria were as follows: (1) eyes with other significant ocular pathology

(including hypertensive retinopathy, retinal vascular occlusions, and uveitis);

(2) history of other systemic diseases, such as malignant tumours, ischaemic

heart disease, stroke, cancer, or kidney disease; (3) a history of heart bypass,

thrombolysis, or kidney transplantation; cognitive impairments or mental

illnesses that made them unable to complete questionnaires or cooperate with

the examiner; (4) spherical equivalent (SE) > −12.0 D, astigmatism > +4.0 D,

axial length (AL) > 26.0 mm; (5) a best corrected visual acuity (BCVA) less

than 20/200; (6) abnormal refractive media (such as moderate to severe

cataract, corneal ulcer, or severe pterygium), poor fixation or other causes

that lead to poor quality of fundus, OCT or OCTA images. When both eyes

were eligible for the study, the worse eye graded by DR from each patient was

selected as the study eye. When both eyes had an equal grade of DR, the

right eye was selected as the study eye.

General and laboratory parameters

Demographic and systemic information, including age, sex, and medical

history, was obtained using standardized questionnaires, and the information

was confirmed after further inspection of outpatient history. Height, weight,

systolic blood pressure (SBP), and diastolic blood pressure (DBP) were

measured using standardized processes. Haemoglobin A1c (HbA1c), total

cholesterol and triglycerides were measured in a laboratory certified by the

Chinese government.

Ocular examination

Each participant underwent a full clinical examination, including slit-lamp

bioscopy, ophthalmoscopy, intraocular pressure (IOP), visual acuity,

refraction, ocular biometry, retinal photography, and OCTA imaging. Ocular

biometric parameters were measured with a Lenstar LS900 (Haag-Streit AG,

Koeniz, Switzerland), including AL, corneal curvature, lens thickness and

anterior chamber depth. Pupil dilation was performed with instillation of 0.5%

tropicamide plus 0.5% phenylephrine in eye drops. Once the pupils were fully

dilated, standardized 7-field colour retinal photographs adhering to the Early

Treatment of Diabetic Retinopathy Study (ETDRS) protocol were taken using a digital fundus camera (Canon CR-2, Tokyo, Japan). The clinician retinopathy severity grading was used for patient recruitment. For the analysis, a masked grading of the colour photographs assigned eyes into the non-DR group (no retinopathy), mild DR group (mild nonproliferative diabetic retinopathy), moderate DR group (moderate nonproliferative diabetic retinopathy), and severe DR group (including severe nonproliferative diabetic retinopathy and proliferative DR) for final analysis by two trained ophthalmologists and if there was any disagreement about the grade, it was confirmed by a senior retina specialist [12].

OCTA imaging and measurements

A commercial swept-source OCTA instrument (DRI OCT-2 Triton; Topcon, Tokyo, Japan) was used for retinal microvasculature imaging. It had a central wavelength of 1050 nm with an axial resolution of 8 μm, a transverse resolution of 20 μm, and a speed of 100,000 A-scans per second. We used 3 mm × 3 mm scans that provided a more detailed description of the fovea and its contour. Automated segmentation and image exportation generated SS-OCTA images of the superficial capillary plexus (SCP) and the deep capillary plexus (DCP). Once moderate segmentation errors appeared, we corrected the segmentation manually. All OCTA images were acquired by a well-trained examiner with no knowledge of the study protocol. The image quality of each

scan was automatically graded by the built-in software, with a signal strength

index (SSI) ranging from 0 to 100 [13]. Images with SSI≤50 were included.

We measured two complexity quantitative features, FD and BVT, in the

OCTA images. The OCTA images were standardized, cropped, and binarized

using Fiji (free downloadable software, https://fiji.sc) [14]. Before measuring the

OCTA features, a binary vessel map and skeleton map were extracted from

the input images. FD can quantify the degree of branching complexity (i.e.,

blood vessels). In our study, the box counting technique was used to measure

the FD values in both SCP and DCP[15] . The FD analysis procedure has been

described in a previous study [15]. The average BVT was measured as the ratio

of the sum of the actual branch lengths to the sum of the straight lengths

between the branch nodes[16] . The FD and BVT were measured using

segmented branches within the skeleton map in the SCP and DCP (Figure 1).

As the FD and BVT values were too small, we enlarged their values by 1000

times for statistical analysis.

Statistical analysis

The Shapiro–Wilk test was used to verify the normal distribution, and

analysis of variance (ANOVA) was performed to compare the differences in

the demographic, systemic, and ocular parameters among the groups with

different stages of DR. Categorical variables were analysed by χ2. Linear

regression analysis was used to evaluate the association between FD in DCP

and the other OCTA features among the different groups. Univariate and

multivariate linear regressions were used to compare FD and BVT in both

SCP and DCP between the control group and the various DR groups. Model 1

of the multivariable linear regression was adjusted for age, sex and OCTA

signal strength intensity. Model 2 was performed by adjusting for more

confounding factors, including age, sex, systolic blood pressure, diastolic

blood pressure, HbA1c, serum creatinine, blood cholesterol, triglycerides,

axial length, and OCTA signal strength intensity. All analyses were performed

using Stata Version 16.0 (Stata Corporation, College Station, TX, USA). A P

value < 0.05 was considered statistically significant.

Results

Demographics

A total of 1282 eyes from 1282 subjects were included in the study. The

demographic and clinical characteristics of the study participants are shown in

Table 1. The average age was 64.2±7.8 years, and 756 (58.97%) patients

were male. Of the 1059 patients without DR, the mean age was 64.3 ± 7.5

years, and 644 (60.81%) were male. Of the 54 patients with mild DR, the

mean age was 62.8 ± 6.9 years, and 30 (55.56%) were male. Of the 133

patients with moderate DR, the mean age was 63.8 ± 9.3 years, and 64

(48.12%) were male. Of the 36 patients with severe DR, the mean age was

62.1 ± 8.8 years, and 18 (50%) were male. The parameters relating to sex,

SBP, HbA1c, serum creatinine and OCTA macular signal strength intensity

differed among the groups (P < 0.05). No statistically significant differences

were present with respect to the other parameters.

Distribution of FD and BVT by DR severity

A summary of the average and detailed univariate analysis of FD and

BVT in SCP and DCP between the control group and the different DR groups

is shown in Table 2 and Figure 2.

A post hoc study after ANOVA showed that all of the features were able to

distinguish controls versus moderate DR patients (P< 0.001) and controls

versus severe DR patients (P< 0.05), but none of the features showed

statistically significant differences between moderate DR eyes and severe DR

eyes. Comparison of FD and BVT in DCP between non-DR patients and mild

DR patients showed a significant difference (P< 0.05). Furthermore, there was

a significant increase in BVT in SCP and in DCP of mild DR eyes compared

with moderate DR eyes (P ≤ 0.01). Only FD in SCP was able to distinguish

between the mild DR group and the severe DR group (P=0.038). The FD in

DCP was higher in non-DR eyes than in DR eyes at each stage (P ≤ 0.001),

but there was no significant difference among the different stages of DR.

Consequently, we considered that FD in DCP was an important feature to

differentiate DR and non-DR.

Association between Vascular Complexity and Diabetic Retinopathy

A univariate linear regression analysis was performed to observe the relationship between FD in DCP and other OCTA features (Table 3). FD in DCP had a statistically significant positive correlation with FD in SCP and a negative correlation with BVT in SCP and BVT in DCP in all of the participants, non-DR group, moderate DR group and severe DR group ($P<0.01$). Only BVT in DCP showed a negative trend with FD in DCP in mild DR eyes ($P<0.001$).

To reduce the factors known to be relevant to OCTA features, we performed two multivariate linear regression analyses to compare FD and BVT in both SCP and DCP between the control group and the various DR groups, which was adjusted for confounding factors. The results of the two models were consistent with the univariate analysis in Table 2, but nearly all of the differences between groups were of a greater magnitude.

Discussion

This OCTA study based on a large sample demonstrated that FD in DCP decreased and BVT in DCP increased in various grades of DR compared to non-DR. The differences were of a greater magnitude after adjusting for potential confounding factors. In addition, our results showed that FD in DCP was positively correlated with FD in SCP and negatively correlated with BVT in SCP and BVT in DCP in all groups except the mild DR group. These

findings confirmed a close relationship between alterations in the retinal

microcirculation, particularly in DCP, and the pathophysiology of DR. To the

best of our knowledge, this study is the first to use SS-OCTA to investigate the

association of vascular complexity features, including FD and BVT, and

different stages of DR in patients with diabetes from communities in China.

Fractal analysis has been employed as a useful tool to study the retinal

vascular pattern and measure the complexity of retinal vasculature branching.

As DR develops, the high rate of vascular dropouts results in a lower vessel

density, which is represented by a lower average number of FD values[3] [9].

Chen et al. reported that the value of FD in the four quadrants of the deep

retinal capillary layer was lower in diabetic patients with minimal DR than in

those with no DR, and FD in the deep retinal capillary layer was much lower in

DM eyes without DR than in nondiabetic control subjects[17] . Suruchi et al.

showed that FD in both the superficial and deep capillary plexuses of diabetic

eyes was significantly lower than that in control eyes[10] . Consistent with

previous studies, our findings illustrated that retinal vascular FD in both SCP

and DCP showed a significant reduction in DR patients. Moreover, the

superiority of our research was that we adjusted for a large number of

confounding factors associated with vascular complexity. However, no

statistically significant changes were observed in FD in either the superficial or

deep vascular networks among the different stages of DR, apart from FD at

SCP in the severe DR group compared with the moderate DR group. These

results suggested that FD in DCP could be an important indicator that

distinguishes DR from non-DR, indicating that the early changes of retinal

capillaries associated with DR may first occur in the deep retinal capillary

layer before extending to the superficial plexus.

There was no significant difference in FD of the superficial capillary

plexus when comparing mild DR and non-DR, suggesting that the superficial

capillary plexus may be damaged until DR progresses to a greater degree.

This is further supported by our analysis of BVT, which did not yield a

significant difference in the superficial capillary plexus between control eyes

and mild DR eyes. Similar to the FD in DCP, the BVT in DCP in non-DR was

lower than in any other stages of DR, even after adjusting for all known

covariates. This result is consistent with a previous study by Minhaj et al. [3].

Consequently, BVT in DCP could be used for non-DR vs. DR classification.

Retinal vessel tortuosity has been considered a characteristic of

haemodynamic changes and grading of DR [18-20]. Dysfunction of the

endothelium and pericytes induces unstable vessel walls, which can easily

become tortuous[21, 22] . Neovascularization often occurs in DR patients and

leads to an increase in BVT[20] . However, neither FD nor BVT in DCP

distinguishes between different severities of DR, which is inconsistent with

Suruchi's[10]   and Hyungwoo's research[16] . This may be due to the small

sample size of the enrolled DR populations or heterogeneous study

populations.

As FD in DCP plays a vital role in the diagnosis and risk assessment of

DR, we performed univariate linear regression analyses between FD in DCP

and other OCTA features. Few studies have analysed the association

between FD and BVT. It was shown in our correlation analysis that the

negative associations of FD in DCP with both BVT in SCP and DCP are

plausible because all of these parameters are related to the abnormal

vascular architecture with DR progression. As DR progresses, avascular

regions are assigned small FD values, and interrupted blood flow results in

short branches. Short capillaries are prone to become tortuous rather than

long branches. The cerebral microvascular meshwork demonstrated that

vessels closer to the capillary network have higher tortuosity and more

curvature[23] , which supports our deduction. Additional studies are necessary

to estimate the cause of high tortuosity by short branches.

An advantage of our study is that all participants were treatment-naïve

patients with type 2 DM from a single Chinese community, consisting of a

homogenous and large-scale cohort. In addition, the model was adjusted for

several potential confounding factors, making our results more accurate and

reliable. There are also some limitations to this study. First, our observations

were only of a very small area of the retina because the 3 × 3 mm area

showed more clearly demarcated vessels, ensuring more precise binarization

of vessels and, subsequently, more accurate quantification of the parameters.

A larger scan area could be performed to quantify OCTA features in more

regions of the retina. Second, projection and motion artefacts in OCTA and

blood flow velocity below the threshold value can affect quantitative feature

analysis. We strictly controlled the image quality by a quality index. Third, we

did not have enough patients in each stage of DR because the participants in

our study were recruited from communities. Further work is warranted with a

larger sample size of DR patients. Finally, this study, as a cross-sectional

study, cannot illustrate the cause-effect relationship between vascular

alterations and DR occurrence and development. Longitudinal studies are

required in the future.

Conclusions

In summary, we found that FD and BVT in DCP by OCTA imaging may be

new quantitative biomarkers for the diagnosis and risk assessment of DR.

Future studies are necessary to determine whether intensive investigations of

retinal vascular geometry can benefit patients and improve clinical decision-

making in the risk evaluation and management of DR.

Disclosures

Financial Disclosures: None.

Figures

Figure 1. Illustration of fractal dimension (FD) and blood vessel tortuosity

(BVT) measurements in the superficial capillary plexus (SCP) and deep

capillary plexus (DCP) by OCTA. Column 1: OCTA image. Column 2:

Segmented blood vessel map including large blood vessels and small

capillaries. Column 3: Skeletonized blood vessel map. Column 4: Image from

Column 3 magnified by a thousand times.

Figure 2. Boxplots showing the distribution of fractal dimension (FD) and

blood vessel tortuosity (BVT) in the superficial retinal layer (SRL) and the

deep retinal layer (DRL) by swept-source optical coherence tomographic

angiography stratified by various stages of DR. (A) Average FD in the SRL of

various stages of DR; (B) Average FD in the DRL of various stages of DR; (C)

Average BVT in the SRL of various stages of DR; (B) Average BVT in the DRL

of various stages of DR.

Tables

Table 1. Demographics and clinical characteristics of the study participants.

Characteristics	ALL	Non-DR	Mild DR	Moderate DR	Severe DR	p Value
No. of subjects	1282	1059	54	133	36	
Male (%)	756(58.97%)	644(60.81%)	30(55.56%)	64(48.12%)	18(50%)	0.024
Mean age, year	64.2±7.8	64.3±7.5	62.8±6.9	63.8±9.3	62.1±8.8	0.182
Systolic blood pressure, mm Hg	135.5±19.0	134.9±18.8	133.3±16.0	138.8±19.3	145.3±24.2	0.002
Diastolic blood pressure, mm Hg	70.9±10.3	71.1±10.1	70.9±12.0	69.0±10.1	70.8±12.7	0.161
HbA1c, %	7.0±1.5	6.8±1.3	7.9±1.9	7.9±1.7	8.6±1.8	< 0.001
Triglycerides, mmol/L	2.3±1.6	2.3±1.6	2.2±1.6	2.3±1.9	2.2±1.3	0.919
Total cholesterol, mmol/L	4.8±1.0	4.9±1.0	4.8±1.1	4.8±1.1	4.9±1.1	0.691
Serum creatinine, μmol/L	70.9±21.1	69.3±19.1	72.6±23.7	79.4±28.9	82.3±28.0	< 0.001
Axial length, mm	23.4±0.9	23.4±0.9	23.2±0.9	23.2±0.8	23.2±0.9	0.136
OCTA signal strength intensity	59.2±11.5	60.0±11.1	59.5±12.3	55.7±12.3	49.5±14.1	< 0.001

358   **Table 2.** Comparisons of FD and BVT in different stages of diabetic retinopathy.

Features	Non-DR (I)	Mild DR (II)	Moderate DR (III)	Severe DR (IV)	P values						
					I vs. II	I vs. III	I vs. IV	II vs. III	II vs. IV	III vs. IV	ANOVA
FD(SCP)	1714.98±1.97	1714.24±15.00	1709.50±18.59	1706.33±24.30	1.000	< 0.001	0.001	0.179	0.038	1.000	< 0.001
FD(DCP)	1697.15±12.37	1689.91±20.28	1687.09±17.89	1682.73±25.76	0.001	< 0.001	< 0.001	1.000	0.102	0.582	< 0.001
BVT(SCP)	1023.32±10.08	1022.52±10.24	1028.14±13.82	1028.46±13.71	1.000	< 0.001	0.027	0.007	0.060	1.000	< 0.001
BVT(DCP)	1018.94±7.27	1022.27±8.81	1026.33±11.06	1026.25±12.71	0.017	< 0.001	< 0.001	0.010	0.126	1.000	< 0.001

359   FD=fractal dimension, BVT=blood vessel tortuosity, SCP=superficial capillary plexus, DCP=deep capillary plexus.

361   **Table 3.** Univariate linear regression analysis of the relationship between features of vascular complexity and FD in DCP.

Features	ALL		Non-DR (I)		Mild DR (II)		Moderate DR (III)		Severe DR (IV)	
	Coefficient (95% CI)	P value	Coefficient (95% CI)	P value	Coefficient (95% CI)	P value	Coefficient (95% CI)	P value	Coefficient (95% CI)	P value
FD(SCP)	0.359 (0.312, 0.406)	< 0.001	0.314 (0.259, 0.370)	< 0.001	0.173(-0.276, 0.373)	0.089	0.382 (0.215, 0.549)	< 0.001	0.673 (0.442, 0.903)	< 0.001
BVT(SCP)	-0.156 (-0.196, -0.116)	< 0.001	-0.117 (-0.166, -0.682)	< 0.001	-0.005 (-0.146, 0.136)	0.941	-0.239 (-0.366, -0.112)	< 0.001	-0.239 (-0.405, -0.730)	0.006
BVT(DCP)	-0.281 (-0.308, -0.252)	< 0.001	-0.244 (-0.276, -0.212)	< 0.001	-0.278 (-0.371, -0.186)	< 0.001	-0.273 (-0.372, -0.174)	< 0.001	-0.255 (-0.402, -0.109)	0.001

362   FD=fractal dimension, BVT=blood vessel tortuosity, SCP=superficial capillary plexus, DCP=deep capillary plexus.

Table 4. Multivariable linear regression of OCTA parameters in non-DR eyes compared with various stages of DR eyes after adjusting for confounding factors.

Features	Non-DR vs. Mild DR Coefficient (95% CI)	P value	Non-DR vs. Moderate DR Coefficient (95% CI)	P value	Non-DR vs. Severe DR Coefficient (95% CI)	P value
Model 1						
FD(SCP)	-0.43(-3.59, 3.09)	0.810	-6.72(-9.05, -4.40)	< 0.001	-13.63(-17.98, -9.27)	< 0.001
FD(DCP)	-7.96(-11.91, -4.01)	< 0.001	-10.19(-12.79, -7.59)	< 0.001	-16.26(-21.15, -11.37)	< 0.001
BVT(SCP)	-1.01(-4.04, 2.03)	0.516	4.65(2.65, 6.64)	< 0.001	6.48(2.72, 10.24)	0.001
BVT(DCP)	3.58(1.29, 5.86)	0.002	7.06(5.55, 8.57)	< 0.001	7.90(5.08, 10.73)	< 0.001
Model 2						
FD(SCP)	-0.89(-4.49, 2.72)	0.630	-6.72(-9.15, -4.28)	< 0.001	-13.74(-18.31, -9.17)	< 0.001
FD(DCP)	-6.75(-10.72, -2.78)	0.001	-8.93(-11.62, -6.25)	< 0.001	-13.37(-18.40, -8.34)	< 0.001
BVT(SCP)	-0.61(-3.71, 2.49)	0.700	4.62(2.53, 6.72)	< 0.001	6.63(2.70, 10.55)	0.001
BVT(DCP)	3.55(1.23, 5.87)	0.003	6.66(5.09, 8.23)	< 0.001	7.43(4.49, 10.37)	< 0.001

Model 1 was adjusted for age, sex and OCTA signal strength intensity. Model 2 was further adjusted for confounding factors, including age, sex, systolic blood pressure, diastolic blood pressure, HbA1c, serum creatinine, blood cholesterol, triglycerides, axial length, and OCTA signal strength intensity. FD=fractal dimension, BVT=blood vessel tortuosity, SCP=superficial capillary plexus, DCP=deep capillary plexus.

References

1.   Wong TY, Cheung CM, Larsen M, Sharma S, Simó R. Diabetic retinopathy. *Nat Rev Dis*
Primers 2016;2:16012.
2.   Alam M, Zhang Y, Lim JI, Chan RVP, Yang M, Yao X. Quantitative optical coherence
tomography angiography features for objective classification and staging of diabetic retinopathy.
Retina 2020;40:322-332.
3.   Alam M, Le D, Lim JI, Yao X. Vascular complexity analysis in optical coherence tomography
angiography of diabetic retinopathy. *Retina* 2021;41:538-545.
4.   Stitt AW, Curtis TM, Chen M, et al. The progress in understanding and treatment of diabetic
retinopathy. *Progress in retinal and eye research* 2016;51:156-186.
5.   Cheung CY, Ikram MK, Klein R, Wong TY. The clinical implications of recent studies on the
structure and function of the retinal microvasculature in diabetes. *Diabetologia* 2015;58:871-885.
6.   Golubovic-Arsovska M. Correlation of diabetic maculopathy and level of diabetic
retinopathy. *Prilozi* 2006;27:139-150.
7.   Hwang TS, Zhang M, Bhavsar K, et al. Visualization of 3 Distinct Retinal Plexuses by
Projection-Resolved Optical Coherence Tomography Angiography in Diabetic Retinopathy.
JAMA Ophthalmol 2016;134:1411-1419.
8.   Hwang TS, Gao SS, Liu L, et al. Automated Quantification of Capillary Nonperfusion Using
Optical Coherence Tomography Angiography in Diabetic Retinopathy. *JAMA Ophthalmol*
2016;134:367-373.
9.   Rosen RB, Andrade Romo JS, Krawitz BD, et al. Earliest Evidence of Preclinical Diabetic
Retinopathy Revealed Using Optical Coherence Tomography Angiography Perfused Capillary
Density. *Am J Ophthalmol* 2019;203:103-115.
10.   Bhardwaj S, Tsui E, Zahid S, et al. Value of fractal analysis of optical coherence tomography
angiography in various stages of diabetic retinopathy. *Retina* 2018;38:1816-1823.
11.   Ting DSW, Tan GSW, Agrawal R, et al. Optical Coherence Tomographic Angiography in
Type 2 Diabetes and Diabetic Retinopathy. *JAMA Ophthalmol* 2017;135:306-312.
12.   Wilkinson CP, Ferris FL, 3rd, Klein RE, et al. Proposed international clinical diabetic
retinopathy and diabetic macular edema disease severity scales. *Ophthalmology* 2003;110:1677-
1682.
13.   Tang F, Sun Z, Wong R, et al. Relationship of intercapillary area with visual acuity in diabetes
mellitus: an optical coherence tomography angiography study. *Br J Ophthalmol* 2018.
14.   Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for biological-
image analysis. *Nat Methods* 2012;9:676-682.
15.   Alam M, Thapa D, Lim JI, Cao D, Yao X. Quantitative characteristics of sickle cell retinopathy
in optical coherence tomography angiography. *Biomed Opt Express* 2017;8:1741-1753.
16.   Lee H, Lee M, Chung H, Kim HC. Quantification of retinal vessel tortuosity in diabetic
retinopathy using optical coherence tomography angiography. *Retina* 2018;38:976-985.
17.   Chen Q, Ma Q, Wu C, et al. Macular Vascular Fractal Dimension in the Deep Capillary Layer
as an Early Indicator of Microvascular Loss for Retinopathy in Type 2 Diabetic Patients. *Invest*
Ophthalmol Vis Sci 2017;58:3785-3794.

18.   Kristinsson JK, Gottfredsdóttir MS, Stefánsson E. Retinal vessel dilatation and elongation
precedes diabetic macular oedema. *Br J Ophthalmol* 1997;81:274-278.
19.   Witt N, Wong TY, Hughes AD, et al. Abnormalities of retinal microvascular structure and risk
of mortality from ischemic heart disease and stroke. *Hypertension* 2006;47:975-981.
20.   Hartnett ME, Martiniuk D, Byfield G, Geisen P, Zeng G, Bautch VL. Neutralizing VEGF
decreases tortuosity and alters endothelial cell division orientation in arterioles and veins in a rat
model of ROP: relevance to plus disease. *Invest Ophthalmol Vis Sci* 2008;49:3107-3114.
21.   Curtis TM, Gardiner TA, Stitt AW. Microvascular lesions of diabetic retinopathy: clues
towards understanding pathogenesis? *Eye (Lond)* 2009;23:1496-1508.
22.   Dobrin PB, Schwarcz TH, Baker WH. Mechanisms of arterial and aneurysmal tortuosity.
Surgery 1988;104:568-571.
23.   Lorthois S, Lauwers F, Cassot F. Tortuosity and other vessel attributes for arterioles and
venules of the human cerebral cortex. *Microvasc Res* 2014;91:99-109.