TITLE PAGE

Manuscript title

Association of COVID-19 vaccines ChAdOx1 and BNT162b2 with major venous,
arterial, or thrombocytopenic events: whole population cohort study in 46 million adults
in England

Short Title

COVID-19 vaccines and with major venous, arterial, and thrombocytopenic events

Authors

CVD-COVID-UK consortium

Writing committee

William N Whiteley FRCP (Chair)

Centre for Clinical Brain Sciences, University of Edinburgh

Nuffield Department of Population Health University of Oxford

Samantha Ip PhD

Department of Public Health and Primary Care, University of Cambridge

Jennifer A Cooper PhD

Bristol Population Health Science Institute, University of Bristol

Thomas Bolton PhD

Department of Public Health and Primary Care, University of Cambridge

Spencer Keene MSc

Department of Public Health and Primary Care, University of Cambridge

Venexia Walker PhD

Bristol Medical School: Population Health Sciences, University of Bristol

Rachel Denholm

Bristol Population Health Science Institute, University of Bristol

Ashley Akbari MSc

Population Data Science, Health Data Research UK, Swansea University

Efosa Omigie BEng MSc

NHS Digital

Sam Hollings PhD

NHS Digital

Emanuele Di Angelantonio FRCP

Department of Public Health and Primary Care, University of Cambridge

NIHR Blood and Transplant Research Unit in Donor Health and Genomics, University
of Cambridge

Spiros Denaxas PhD

Institute of Health Informatics, University College London

Angela Wood PhD*

Department of Public Health and Primary Care, University of Cambridge

Jonathan A C Sterne PhD*

Department of Population Health Sciences, University of Bristol

NIHR Bristol Biomedical Research Centre

HDR UK South West

Cathie Sudlow FRSE*

BHF Data Science Centre, Health Data Research UK

*Drs Wood, Sudlow and Sterne contributed equally

Word count:

Abstract: 349

Manuscript: 3637

Corresponding author

Email: william.whiteley@ed.ac.uk

Telephone: 07811113610

ABSTRACT

Background:

Thromboses in unusual locations after the COVID-19 vaccine ChAdOx1-S have been reported. Better understanding of population-level thrombotic risks after COVID-19 vaccination is needed.

Methods:

We analysed linked electronic health records from adults living in England, from 8th December 2020 to 18th March 2021. We estimated incidence rates and hazard ratios (HRs) for major arterial, venous and thrombocytopenic outcomes 1-28 and >28 days after first vaccination dose for ChAdOx1-S and BNT162b2 vaccines. Analyses were performed separately for ages <70 and ≥70 years, and adjusted for age, sex, comorbidities, and social and demographic factors.

Results:

Of 46,162,942 adults, 21,193,814 (46%) had their first vaccination during follow-up. Adjusted HRs 1-28 days after ChAdOx1-S, compared with unvaccinated rates, at ages <70 and ≥70 respectively, were 0.97 (95% CI: 0.90-1.05) and 0.58 (0.53–0.63) for venous thromboses, and 0.90 (0.86-0.95) and 0.76 (0.73-0.79) for arterial thromboses. Corresponding HRs for BNT162b2 were 0.81 (0.74–0.88) and 0.57 (0.53–0.62) for venous thromboses, and 0.94 (0.90-0.99) and 0.72 (0.70-0.75) for arterial thromboses. HRs for thrombotic events were higher at younger ages for venous thromboses after ChAdOx1-S, and for arterial thromboses after both vaccines.

Rates of intracranial venous thrombosis (ICVT) and thrombocytopenia in adults aged <70 years were higher 1-28 days after ChAdOx1-S (adjusted HRs 2.27, 95% CI:1.33–3.88 and 1.71, 1.35–2.16 respectively), but not after BNT162b2 (0.59, 0.24–1.45 and 1.00, 0.75–1.34) compared with unvaccinated. The corresponding absolute excess risks of ICVT 1-28 days after ChAdOx1-S were 0.9–3 per million, varying by age and sex.

Conclusions:

Increases in ICVT and thrombocytopenia after ChAdOx1-S vaccination in adults aged <70 years were small compared with its effect in reducing COVID-19 morbidity and mortality, although more precise estimates for adults <40 years are needed. For people aged ≥70 years, rates of arterial or venous thrombotic, events were generally lower after either vaccine.

INTRODUCTION

In late February 2021 several groups reported a rare syndrome of thrombosis and thrombocytopenia after vaccination against severe acute respiratory syndrome coronavirus-2 disease 2019 (COVID-19) with ChAdOx1-S (developed by Oxford-AstraZeneca).[1-3] Thromboses were found in unusual sites such as the cerebral venous sinuses, or mesenteric or portal veins. This syndrome, named 'vaccine-induced immune thrombotic thrombocytopenia' (VITT), is probably due to an autoimmune response to platelet factor 4 (PF4) in the absence of exposure to heparin.[1] Whether the risk of common thrombotic illnesses is also increased post-vaccination is uncertain, although there have been reports of post-vaccination ischaemic stroke and intracerebral haemorrhage.[1,3,4]

Estimates of vaccine-associated excess risk based on case reports and routine reporting may be biased if diagnostic thresholds vary between vaccinated and unvaccinated individuals or there is less than population-wide coverage. In England, the National Health Service (NHS) provides almost all healthcare; therefore, linked health care records provide comprehensive, population-wide data on outcomes before and after vaccination.[5] Vaccination in England began on December 8th 2020: the first groups vaccinated were care home residents, then those aged ≥80 years and frontline health and social care workers. Those clinically extremely vulnerable were invited alongside people aged ≥70 years, and those with other co-morbidities aged <65 alongside people aged >65 years. Consequently, individuals vaccinated earlier are expected to have higher rates of venous and arterial events.

We conducted a cohort study using routinely collected, linked health data from multiple sources, covering almost the entire English adult population (>46 million). Associations of ChAdOx1-S and BNT162b2 (developed by Pfizer–BioNTech) with

115   major venous and arterial events and thrombocytopenic haematological outcomes

116   were estimated from the start of the vaccination program to mid-March 2021, before

117   VITT surveillance was widespread.

METHODS

Population

The study population was adults (aged ≥18 years), alive and registered with an English NHS general practice on December 8th 2020. The data resource[5] includes primary care data (General practice extraction service (GPES) data for Pandemic Planning and Research, GDPPR) from 98% of general practices linked at individual-level to nationwide secondary care data including all NHS hospital admissions (Hospital Episode Statistics (HES) and Secondary Uses Service (SUS) data from 1997 onwards), COVID-19 laboratory testing data, COVID-19 vaccination data (NHS England immunisation management system), national community drug dispensing data (NHS BSA Dispensed Medicines from 2018) and death registrations. We accessed and analysed pseudonymised data within NHS Digital's secure, privacy protecting Trusted Research Environment.[6]

Covariates

Covariates were defined from primary care, hospital admissions, community drug dispensing and COVID-19 laboratory testing data, using phenotyping algorithms verified by specialist physicians (see study protocol). Phenotypes for co-morbidities, risk factors and other covariates used Systematized Nomenclature of Medicine Clinical Terms (SNOMED-CT) concepts for primary care data, and ICD-10 codes for hospital admission data. Sex, age, region, deprivation and smoking status were defined as the latest recorded in primary care records before 8th December 2020, and the most recently recorded ethnicity in either primary care or hospital admissions records was used. History of diabetes, depression, obesity, cancer, chronic obstructive pulmonary disease, chronic kidney disease, stroke, MI, DVT or PE, cancer,

thrombophilia, liver disease or dementia were defined as any record in primary care

and/or hospital admission data before December 8th 2020. The number of unique

diseases in SNOMED-CT for the year before December 8[th] 2020 were derived from

primary care records and major surgery in the previous year from hospital admission

records (using the Office of Population Censuses and Surveys classification of

surgical procedures).

A history of coronavirus (SARS-CoV2) infection was defined as either a positive

COVID-19 antigen test in national laboratory data covering swab tests performed in

the general population and hospitals or a confirmed COVID-19 diagnosis in primary

care or hospital admission records. Prior medication was defined using community

dispensing data on all prescriptions dispensed by community pharmacists, appliance

contractors, and doctors in England. British National Formulary (BNF) codes were

used to define the total number of types of medication prescribed and dispensed in

the following groups: antiplatelets, antihypertensives, lipid lowering agents, oral

anticoagulants, combined oral contraceptives, and hormone replacement therapy.

Outcomes

Outcomes were derived from primary care data, hospital admission data (SUS

dataset) and the national death registry (see supplementary material).[7] Specialist

clinician-verified SNOMED-CT and ICD-10 rule-based phenotyping algorithms were

used to define fatal or non-fatal (i) arterial thrombotic events: myocardial infarction

(MI), ischaemic stroke (ischaemic or unclassified stroke, spinal stroke or retinal

infarction), other non-stroke non-MI arterial thrombo-embolism; (ii) venous thrombo-

embolic events: pulmonary embolism (PE), lower limb deep venous thrombosis (DVT),

intracranial venous thrombosis (ICVT), portal vein thrombosis, and venous thrombosis

at other sites; (iii) thrombocytopenic haematological events: any thrombocytopenia

(idiopathic, primary, secondary or unspecified), disseminated intravascular

coagulation (DIC), and thrombotic thrombocytopenic purpura (TTP); (iv) death from

any cause; (v) other vascular outcomes: haemorrhagic stroke (intracerebral or

subarachnoid), and mesenteric thrombus for which available codes did not distinguish

between arterial or venous causes; and (vi) lower limb fracture as a control condition

unlikely to be affected by vaccination. The event date was the earliest of: start of

hospital episode, death, or recorded date of primary care event or consultation. Events

within the death registry were identified based on underlying cause of death and in

hospital admission data based on the primary cause of the care episode.

Statistical analyses

Data were analysed according to a pre-specified plan, published on GitHub on April

27th 2021.[8] Follow-up was from December 8th 2020 to March 18th 2021, the date the

European Medicine Agency (EMA) Pharmacovigilance Risk Assessment Committee

(PRAC) discussed the first reported complications of vaccination, after which

diagnostic effort is expected to be concentrated in people receiving ChAdOx1-S.[9] Any

cell numbers <5 and any potentially disclosive numbers <10 are not reported exactly

but as <5 and <10 respectively.

Follow up time for each person was split into periods before, and 1-28 and >28 days

after, first vaccination. Censoring was at the earliest of the outcome, death, March 18th

2021 and, in analyses of specific vaccines, receipt of the other vaccine. Incidence

rates per 100,000 people per year were estimated for each outcome, before and 1-28

and >28 days after vaccination.

Cox models with time zero December 8th 2020 were fitted, ensuring that all analyses

accounted for changes in rates of outcome events with calendar time. Hazard ratios

comparing the periods 1-28 and >28 days after vaccination with unvaccinated or pre-

vaccination person-time (reference) were estimated. Cox models were fitted

separately by age group (<70 and ≥70 years), both overall and separately for males

and females. All Cox models were stratified by geographic region. For rare outcomes,

sex-specific hazard ratios were estimated from vaccine-sex interaction terms. For

computational efficiency, each model included data from all people with, and a 10%

random sample of people without, the outcome of interest; analyses incorporated

inverse probability weights to account for this sampling. Confidence intervals were

derived using robust standard errors.

Region-; age-sex-region-; and fully-adjusted hazard ratios (HR) for associations of

ChAdOx1-S and BNT162b2 with outcome events were estimated. The following

covariates were included in all fully-adjusted models: (a) sex, age and age^2, ethnicity,

postcode-derived deprivation level; (b) risk factors for venous thromboses

(anticoagulant medication, combined oral contraceptive medication, hormone

replacement therapy medication, history of PE or DVT, and history of coronavirus

infection); (c) risk factors for arterial thromboses (diabetes, hypertension, smoking,

antiplatelet medication, blood pressure lowering medication, lipid lowering medication,

anticoagulant medication, history of stroke, and history of MI); (d) further covariates

selected using a backwards stepwise procedure with p value threshold 0.2, from

models using MI as outcome (separately in the four groups defined by age and sex).

The covariates had few missing values (apart from ethnicity, for which the 5.9%

'missing' were included as a separate category); hence all analyses used "complete-

cases".

Sensitivity analyses examined outcome events recorded as the primary or secondary

reason for admission or death in hospital admissions or death records; fatal outcome

events (those followed by death from any cause within 28-days); and all venous and

all arterial thrombotic outcome events associated with thrombocytopenia. Effect

modification by age, sex, ethnicity, medication, diabetes, deprivation and medical

history was examined for all venous and all arterial thromboses.

Absolute excess risks of ICVT were estimated by applying estimated hazard ratios to

incidence rates of first fatal or non-fatal events in decades of age and sex in 2019,

derived from linked hospital and death records.

The study was approved by the Newcastle & North Tyneside 2 Research Ethics

Committee (20/NE/0161), the NHS Digital Data Access Request Service (DARS-NIC-

381078-Y9C5K) and the British Heart Foundation Data Science Centre CVD-COVID-

UK Approvals and Oversight Board.

Data manipulation and analyses used SQL and Python in Databricks and RStudio

(Professional) Version 1.3.1093.1 driven by R Version 4.0.3. All code and phenotypes

are available at github.com/BHFDSC/CCU002_02.

RESULTS

On December 8[th] 2020, 46,162,942 eligible adults were registered with an English

general practice. By March 18[th] 2021, 21,193,814 (46%) had received their first

vaccination (8,712,477 BNT162b2; 12,481,337 ChAdOx1-S) (supplementary figure 1).

Post-vaccination person-years of follow up for ages <40, 40-69 and ≥70 years

respectively were, for ChAdOx1-S: 91,206, 417,160 and 401,343 person years; for

BNT162b2: 159,856, 427,874, and 583, 890 person years.

The risks per 100,000 persons from December 8[th] 2020 to March 18[th] 2021 were: any

venous thrombosis 45.3, any arterial thrombosis 189; thrombocytopenia 4.2 (Table 1).

Risks of venous and arterial thromboses were higher in people with co-morbidities and

with increasing age and deprivation, and varied substantially by ethnicity. Absolute

and relative increases in risk with increasing age and deprivation were greater for

arterial than for venous thromboses. The risk of venous thromboses was higher in

people with prior DVT or PE, thrombophilia, or oral anticoagulant medication, while

the risk of arterial thromboses was higher in people with prior stroke, MI or antiplatelet

medication. The risk of thrombocytopenia increased with age and co-morbidities but

varied less markedly with ethnicity and deprivation than the risk of venous or arterial

thromboses.

Incidence rates of all events varied substantially in unvaccinated and post-vaccination

(Table 2). In the unvaccinated, the crude incidence rates per 100,000 person years

were: all venous thromboses 141 (mostly PE and DVT), ICVT 1.97; portal vein

thrombosis 1.15; any arterial thrombosis 549 (with slightly more ischaemic stroke

(273) than MI (268)); any thrombocytopenia 13, with very low rates of DIC (0.12) and

TTP (0.67). For each vaccine, unadjusted incidence rates of venous and arterial

thromboses, haemorrhagic stroke, mesenteric thrombus and thrombocytopenia were

higher after than before vaccination in the under 70s, but lower after than before

vaccination in the over 70s.

The additional confounding factors selected using backward selection were: for both

sexes and age strata, previous diagnosis of cancer, number of unique diseases in the

last year, surgery in the last year, obesity, liver disease; additionally for women aged

<70, history of depression and total number of types of medication (by BNF chapters);

additionally for men aged <70, history of SARS-CoV2 infection and thrombocytopenia;

and additionally for men and women aged ≥70, chronic kidney disease and dementia.

Crude hazard ratios were substantially attenuated after adjusting for age, and further

attenuated after adjusting for confounding factors. For example, in <70 year-olds the

HRs 1-28 days post- versus unvaccinated were attenuated from 2.81 to 2.27 for ICVT;

from 4.85 to 1.71 for thrombocytopenia; and from 3.21 to 0.90 for ischaemic stroke

(supplementary table 1).

In people aged <70 years, the fully adjusted HR for any venous thrombosis was 0.97

(95% CI:0.90-1.05) 1–28 days after ChAdOx1-S compared with before vaccination,

with similar adjusted HRs for PE and DVT. Compared with before vaccination, the

hazard of any venous thrombosis was lower 1-28 days after ChAdOx1-S in people

aged ≥70 (adjusted HR 0.58, 95%CI:0.53–0.63) and 1–28 days after BNT162b2 (<70:

HR 0.81, 0.74–0.88; ≥70 HR: 0.57, 0.53–0.62), with similar adjusted HRs for PE and

DVT. (Figure 1, supplementary table 1).

In people aged <70 years, adjusted HRs for ICVT were 2.27 (95%CI: 1.32–3.88) 1-28

days after ChAdOx1-S and 0.51 (0.16–1.68) 1-28 days after BNT162b2. HRs were

similar in men and women. In a post-hoc analyses, in people aged <40 years, the HRs

1-28 days after vaccination were 3.73 (1.58–8.83) and 1.07 (0.33–3.48) for ChAdOx1-

S and BNT162b2 respectively. The same estimates for people 40-69 years were 1.93

(1.19–2.13) and 0.74 (0.29–1.87). In people aged ≥70 years, associations of

vaccination with ICVT were estimated imprecisely (<10 events for ChAdOx1-S and 8

for BNT162b2 (Figure 1, supplementary table 1).

Adjusted HRs for any arterial thrombosis 1-28 days after ChAdOx1-S were 0.90

(95%CI: 0.86-0.95) and 0.76 (0.73–0.79) in people aged <70 and ≥70 years

respectively. Corresponding HRs 1-28 days after BNT162b2 were 0.94 (0.90-0.99)

and 0.72 (0.70–0.75). HRs were similar for MI and ischaemic stroke, and >28 days

after vaccination (Figure 2, supplementary table 1).

In people aged <70 years, adjusted HRs for thrombocytopenia were 1.71

(95%CI:1.35–2.16) 1–28 days and 1.69 (1.16–2.46) >28 days after ChAdOx1-S.

Corresponding HRs after BNT162b2 were 1.00 (0.75–1.34) and 0.97 (0.66–1.41). For

those aged ≥70 years, adjusted HRs for thrombocytopenia 1-28 days after vaccination

were similar for ChAdOx1 (0.79, 95%CI: 0.56–1.10) and BNT162b2 (0.68, 0.51-0.90)

(Figure 3, supplementary table 1).

Adjusted HRs for lower limb fracture 1–28 days after ChAdOx1-S were 0.85 (95% CI:

0.77–0.93), and 0.83 (0.78–0.89) in people aged <70 and ≥70 years respectively.

Corresponding HRs after BNT162b2 were 0.93 (0.84–1.02) and 0.69 (0.66–0.73).

(Figure 3, supplementary table 1). Adjusted HRs for all-cause mortality 1–28 days after

ChAdOx1-S were 0.37 (95%CI: 0.35–0.39), and 0.28 (0.28–0.29) in people aged <70

and ≥70 years respectively. Corresponding HRs after BNT162b2 were 0.24 (0.22–

0.26) and 0.19 (0.19–0.20) (supplementary figure 2, supplementary table 1).

Subgroup analyses

HRs for all venous and for all arterial thrombotic events 1-28 days after vaccination

were lower in older age groups for each vaccine (all interaction p values <0.0001)

(supplementary table 2). HRs for venous thromboses were >1 1-28 days after

compared with before vaccination in people aged <50 years for ChAdOx1-S but not

BNT162b2. HRs for arterial thromboses were >1 1-28 days after compared with before

vaccination in people aged <50 years for both ChAdOx1-S and BNT162b2. A higher

proportion of venous thromboses in those aged <40 years compared with those aged

≥40 years were due to ICVT. A higher proportion of arterial thromboses at age <40

was due to stroke than at ages 40–59 and 60–69 years. Less marked trends for lower

HRs in older individuals were observed for HRs >28 days after vaccination. Despite

modest differences between the magnitude of HRs in subgroups defined by sex,

ethnicity, prior COVID, COCP, diabetes, deprivation or history of MI, DVT or PE,

anticoagulation or antiplatelet medication, HRs in all of these subgroups were

consistent with lower rates of arterial or venous thromboses after vaccination with

ChAdOx1-S or BNT162b2.

Sensitivity analyses

When using a less restrictive outcome definition (outcomes recorded as primary or

secondary reason for admission), estimated HRs were consistent with those from

analyses of outcomes in the primary position (supplementary figure 3). Similarly,

estimated HRs for fatal outcomes, were similar to HRs for outcomes in the primary

position (supplementary figure 4). There were very few people for whom outcomes in

the primary position were recorded as well as thrombocytopenia in any position in the

same hospital admission or death record (after ChAdOx1-S and BNT162b

respectively: ICVT 7 and 0; all arterial thromboses 47 and 58; all venous thromboses

37 and 26).

Intracranial venous thrombosis

Characteristics were similar among patients who developed ICVT while unvaccinated,

after ChAdOx1 and after BNT162b. Most were women (66% unvaccinated, 67%

ChAdOx1, 79% BNT162b), of white ethnicity (75% unvaccinated, 92% ChAdOx1,

100% BNT162b), and had no prior co-morbidities (71% unvaccinated, 50% ChAdOx1,

50% BNT162b). Patients with post vaccination ICVT were older than those with

unvaccinated ICVT. Post-vaccination, no patients had a recorded history of

thrombophilia, and fewer than 5% had a recorded dispensed prescription of oral

contraceptive or HRT (supplementary Table 3).

Applying the HR for ICVT 1-28 days after ChAdOx1 to the monthly incidence of ICVT

in 2019, the excess risk of ICVT in the month after vaccination with ChAdOx1 among

those aged 19–30, 31–39, 40–49, 50–59 and 60–69-years was estimated to be,

respectively: 0.9, 1.1, 1.5, 1.5 and 1.6 per million vaccinated in men and 3.0, 2.7, 2.0,

1.6 and 1.5 per million vaccinated in women.

DISCUSSION

In this cohort study, which included almost all adults alive in England at the start of the

national COVID-19 vaccination programme, vaccination with ChAdOx1-S, but not

BNT162b2, was associated with approximately 2-fold higher rates of ICVT or

hospitalisations due to thrombocytopenia in people aged under 70 years, after

adjusting for a comprehensive range of demographic characteristics and

comorbidities. The corresponding absolute increases in the risk of these events were

very small. ChAdOx1-S and BNT162b2 were associated with lower hazards of major

vascular events after vaccination in those aged ≥70 years. Post-hoc analyses stratified

by age suggest that in people aged <50 years, rates of venous thromboses were

higher 1-28 days after vaccination with ChAdOx1-S, and that rates of arterial

thrombotic events were higher 1-28 days after vaccination with ChAdOx1-S or

BNT162b2, compared with rates in unvaccinated people.

The large population analysed (including almost 10 times as many vaccinated people

as any other published study) adds substantial precision to estimated associations of

thromboses with vaccination, which is of particular importance for very rare events and

associations in subgroups. The extensive data linkages across different healthcare

settings enabled adjustment for a wide range of confounding variables. The analysed

population is slightly larger than the mid-year estimate of the 2020 English population

aged ≥18 (44,456,850) that excluded short-term residents and students, and modelled

international migrant numbers.[10]

Smaller Danish and Norwegian electronic health record studies reported a small

excess of intracranial haemorrhage (2 per 100,000 1–28 days after ChAdOx1-S),

cerebral venous thrombosis (3 per 100,000) and other venous thrombosis (2 per

100,000).[11] The randomised trials of ChAdOx1-S, reported no venous thrombotic

events among 8597 participants who received the vaccine versus two among

8581who received placebo.[12] A Scottish study found a risk of of 1.13 (0.62–1.63)

cases of idiopathic thrombocytopenic purpura per 100,000 after ChAdOx1-S. There

was no clear evidence on other outcomes, although the number of post-vaccine events

was small.[13] Reporting of ICVT cases to the UK Medicines and Healthcare products

Regulatory Agency (MHRA) identified more cases of ICVT with thrombocytopenia

after ChAdOx1-S than the present study (44 up to 31st March 2021).[14]

Healthcare systems planning to use ChAdOx1-S should balance the very small harms

against the known benefits of the vaccine. For older populations, who are most

vulnerable to COVID-19, we found no evidence of increased risk of any event with

ChAdOx1-S. In younger populations, who have a lower morbidity and mortality due to

COVID-19, other available vaccines might be prioritise, especially when the risk of

COVID-19 is otherwise low.

Our study has several limitations. First, identification of exposures, covariates and

outcomes relies on the accuracy of data collected during routine healthcare. Additional

data on results of laboratory and radiology investigations would have improved

diagnostic coding, particularly for ascertainment of thrombocytopenia. Second, people

who were not registered with an NHS GP (for example the homeless, recent

immigrants, those using only private healthcare and those not eligible for NHS care)

or who opted out of their data being provided to NHS Digital were excluded. Third,

follow up ended on March 18th 2021, but a small number of events that occurred before

this date may have been excluded because they had yet to be coded or the people

affected were still in hospital. Analyses after this date will likely lead to overestimation

of associations, because speciality societies recommended further investigations of

mild symptoms in vaccinated populations. Fourth, our primary outcome used the

primary reason for death or hospital admission, which improves the positive predictive

value but may lead to an underestimation of incidence. This is necessary because

historical non-incident events are frequently recorded in secondary positions.

Analyses of events recorded in any position as fatal within 28 days were consistent

with the primary analyses. Fifth, we did not address time-varying confounding, which

can occur when factors which vary during follow up, such as admission to hospital,

predict both vaccination and outcomes of interest. Sixth, comparison of adjusted and

unadjusted associations suggests that studies that do not adjust for a comprehensive

range of potential confounders will overestimate the thrombotic effects of vaccination.

However, adjusted associations in this study may still be biased by unmeasured

confounding by patient characteristics that predict both vaccination and thromboses

and that are difficult to ascertain in electronic health records. Examples include general

health at the time of vaccination, and people at higher risk of thromboses (e.g. with

end-stage diseases).[15] There were slightly lower post-vaccination rates of the

'negative control' outcome lower limb fracture in those aged ≥70 years, which implies

that this degree of unmeasured confounding may affect results for thrombotic

outcomes in that age group.

Associations of vaccination with thromboses varied with age. This may be because in

older people small increases in the risk of major thrombotic events after vaccination

with ChAdOx1-S were more than offset by reductions in major thrombotic events

(particularly PE) subsequent to COVID-19. By contrast, in younger people any

increase in risk associated with ChAdOx1-S vaccine is less likely to be offset by a

lower risk of COVID-associated thrombosis, because the chance of severe COVID

disease is lower in younger than older individuals.

Further analyses assessing the effects of other vaccines and the effects of second

doses of these vaccines on thrombotic, neurological and cardiac complications will be

important to inform vaccination programs. Access to data from radiology or laboratory

systems (which in the UK will rely on regional rather than national data collection

systems) will allow more comprehensive case ascertainment and more granular

phenotyping. Such efforts are currently underway across the UK.

DATA SHARING

The de-identified data used in this study is available via the CVD-COVID-UK consortium coordinated by BHF Data Science Centre (https://www.hdruk.ac.uk/projects/cvd-covid-uk-project/) for accredited researchers working on approved projects in the NHS Digital trusted research environment. The study protocol, analytic and phenotyping code is available at: https://github.com/BHFDSC/CCU002_02

FUNDING

The British Heart Foundation Data Science Centre (grant No SP/19/3/34678, awarded to Health Data Research (HDR) UK) funded co-development (with NHS Digital) of the trusted research environment, provision of linked datasets, data access, user software licences, computational usage, and data management and wrangling support. Support was also provided through the Data and Connectivity and Longitudinal Health and Wellbeing National Core Studies, which were established through the UK Government's Chief Scientific Adviser's National Core Studies programme to coordinate COVID-19 priority research. Consortium partner organisations funded the time of contributing data analysts, biostatisticians, epidemiologists, and clinicians.

WW is supported by the Chief Scientist's Office (CAF/01/17). CS, AW and WW are supported by Stroke Association (SA CV 20\100018). WW has given expert testimony to UK courts.

AMW is supported by the BHF-Turing Cardiovascular Data Science Award (BCDSA\100005) and by core funding from UK MRC (MR/L003120/1), BHF (RG/13/13/30194; RG/18/13/33946), and NIHR Cambridge Biomedical Research Centre (BRC-1215-20014). AMW is part of the BigData@Heart Consortium, funded by the Innovative Medicines Initiative-2 Joint Undertaking under grant agreement No 116074. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

JAC, JS, and RD are supported by the Health Data Research (HDR) UK South West Better Care Partnership, and the NIHR Bristol Biomedical Research Centre at University Hospitals Bristol, and Weston NHS Foundation Trust and the University of Bristol.

VMW is supported by the Medical Research Council Integrative Epidemiology Unit, which receives its funding from the Medical Research Council and the University of Bristol (MC_UU_00011/4).

AA, SD is supported by Health Data Research UK (grant number: HDR-9006), which receives its funding from the UK Medical Research Council, Engineering and Physical Sciences Research Council, Economic and Social Research Council, Department of Health and Social Care (England), Chief Scientist Office of the Scottish Government Health and Social Care Directorates, Health and Social Care Research and Development Division (Welsh Government), Public Health Agency (Northern Ireland), British Heart Foundation (BHF) and the Wellcome Trust; and Administrative Data Research UK, which is funded by the Economic and Social Research Council (grant number: ES/S007393/1).

This work uses data provided by patients and collected by the NHS as part of their
care and support. We would also like to acknowledge all data providers who make
anonymised data available for research.

Funders had no role in the study design, collection, analysis, and interpretation of data;
in the writing of the report; and in the decision to submit the article for publication

Conflicts of interest

None

CONTRIBITIONS

WW chaired the writing committee. WW, CS, JS and AW conceived the idea, and
prepared the protocol and drafted the manuscript. SI, JAC, SK, VW, TB, RD and JS
wrote the analytic code. VW, JAC, SK and SI prepared analytic code for open access
publication. SI, VW and TB prepared the tables and figures. EO and SH prepared data
from NHS Digital. SD, SK and WW prepared phenotype code. AW and JS designed
and supervised the statistical analyses. CS is the Director of the BHF Data Science
Centre and coordinated approvals for and access to data within the NHS Digital TRE.
All authors commented on the protocol, analyses, and manuscript, and approved the
final version before submission. All authors had full access to the study data.

REFERENCES

Schultz NH, Sørvoll IH, Michelsen AE, *et al.* Thrombosis and Thrombocytopenia after ChAdOx1 nCoV-19 Vaccination. *N Engl J Med* 2021; published online April 9. DOI:10.1056/nejmoa2104882.

Greinacher A, Thiele T, Warkentin TE, Weisser K, Kyrle PA, Eichinger S. Thrombotic Thrombocytopenia after ChAdOx1 nCov-19 Vaccination. *N Engl J Med* 2021; : NEJMoa2104840.

Scully M, Singh D, Lown R, *et al.* Pathologic Antibodies to Platelet Factor 4 after ChAdOx1 nCoV-19 Vaccination. *N Engl J Med* 2021; : NEJMoa2105385.

Blauenfeldt RA, Kristensen SR, Ernstsen SL, Kristensen CCH, Simonsen CZ, Hvas A-M. Thrombocytopenia with acute ischemic stroke and bleeding in a patient newly vaccinated with an adenoviral vector-based COVID-19 vaccine. *J Thromb Haemost* 2021; published online April 20. DOI:10.1111/jth.15347.

Wood A, Denholm R, Hollings S, *et al.* Linked electronic health records for research on a nationwide cohort of more than 54 million people in England: Data resource. *BMJ* 2021; **373**. DOI:10.1136/bmj.n826.

Trusted research environment. https://digital.nhs.uk/coronavirus/coronavirus-data-services-updates/trusted-research-environment-service-for-england.

Herrett E, Shah AD, Boggon R, *et al.* Completeness and diagnostic validity of recording acute myocardial infarction events in primary care, hospital care, disease registry, and national mortality records: cohort study. *BMJ* 2013; **346**: f2350.

BHF Data Science Centre. Study protocol. 2021. https://github.com/BHFDSC/Protocols/blob/main/COVID vaccination and risk of arterial and venous event BHF DSC v1.2.pdf.

COVID-19 Vaccine AstraZeneca: PRAC investigating cases of thromboembolic events - vaccine's benefits currently still outweigh risks - Update | European Medicines Agency. https://www.ema.europa.eu/en/news/covid-19-vaccine-astrazeneca-prac-investigating-cases-thromboembolic-events-vaccines-benefits (accessed May 10, 2021).

Office for National Statistics. Estimates of the population for the UK, England and Wales, Scotland and Northern Ireland. 2021. https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/datasets/populationestimatesforukenglandandwalesscotlandandnorthernireland.

Pottegård A, Lund LC, Karlstad Ø, *et al.* Arterial events, venous thromboembolism, thrombocytopenia, and bleeding after vaccination with Oxford-AstraZeneca ChAdOx1-S in Denmark and Norway: population based cohort study. *BMJ* 2021; **373**: n1114.

Voysey M, Costa Clemens SA, Madhi SA, *et al.* Single-dose administration and the influence of the timing of the booster dose on immunogenicity and efficacy of ChAdOx1 nCoV-19 (AZD1222) vaccine: a pooled analysis of four randomised trials. *Lancet* 2021; **397**: 881–91.

13   Simpson CR, Shi T, Vasileiou E, *et al.* First-dose ChAdOx1 and BNT162b2
COVID-19 vaccines and thrombocytopenic, thromboembolic and hemorrhagic
events in Scotland. *Nat Med* 2021; published online June 9.
DOI:10.1038/s41591-021-01408-4.

14   Hunter PR. Thrombosis after covid-19 vaccination. *BMJ* 2021; **373**.
DOI:10.1136/BMJ.N958.

15   OpenSAFELY Collaborative T, Curtis HJ, Inglesby P, *et al.* Trends and clinical
characteristics of COVID-19 vaccine recipients: a federated analysis of 57.9
million patients' primary care records in situ using OpenSAFELY. *medRxiv*
2021; : 2021.01.25.21250356.

TABLES

Table 1 Numbers of patients analysed and (in parentheses) risk per 100,000 of venous and arterial events and thrombocytopenia (TCP) during follow up, overall and according to age.

Table 2 Numbers of events and incidence rates pre and post first vaccination. Incidence rate per 100,000 per annum (a) ChAdOx1-S (b) BNT162b2 vaccine

FIGURE LEGENDS

Figure 1 Adjusted Hazard ratios for all venous thromboses, intracranial venous thromboses, pulmonary embolism and deep vein thromboses after ChAdOx1-S or BNT162b2 vaccine.

Figure 2 Adjusted hazard ratios for all arterial thromboses, myocardial infarction, ischaemic stroke and other arterial thromboses after ChAdOx1-S or BNT162b2 vaccine.

Figure 3 Adjusted hazard ratios for thrombocytopenia haemorrhagic stroke, mesenteric thrombosis, lower limb fracture after ChAdOx1-S or BNT162b2 vaccine

SUPPLEMENTARY MATERIAL

Protocol

Supplementary table 1 Unadjusted, age-, sex- and region- adjusted, and fully adjusted hazard ratios for (a) ChAdOx1-S (b) BNT162b2 vaccine

Supplementary table 2 Stratum specific estimates for (a) venous and (b) arterial events.

Supplementary table 3. Characteristics of patient who had an ICVT event before and after vaccination. Only percentage are presented, as disclosure control does not allow numbers <5 to be released

Supplementary figure 1: Cumulative frequency of vaccines of different types during follow up

Supplementary figure 2: Adjusted hazard ratios for portal vein thrombosis, other venous events, and death after ChAdOx1-S or BNT162b2 vaccine

Supplementary Figure 3 Hazard ratios for major (A) arterial and (B) venous thrombotic events, and (C) haematological events, other events and lower limb fractures recorded in any position in EHR

Supplementary Figure 4: Hazard ratios for major (A) arterial and (B) venous thrombotic events, (C) haematological events, other events and fractures recorded in death record or in hospital record in first position followed by death <28 days

Table 1. Numbers of patients analysed and (in parentheses) risk per 100,000 people of venous and arterial events and thrombocytopenia (TCP), overall and according to age.

		Overall				Age, <70				Age, ≥70			
		All	Venous	Arterial	TCP	All	Venous	Arterial	TCP	All	Venous	Arterial	TCP
N		46,162,942	20,903 (45.3)	87,251 (189)	1,926 (4.2)	38,744,171	11,835 (30.5)	38,283 (98.8)	1,167 (3.0)	7,418,771	9,068 (122)	48,968 (660)	759 (10.2)
Sex	Male	22,765,779	10,358 (45.5)	51,851 (228)	986 (4.3)	19,425,454	6,406 (33.0)	26,549 (137)	555 (2.9)	3,340,325	3,952 (118)	25,302 (757)	431 (12.9)
	Female	23,397,163	10,545 (45.1)	35,400 (151)	940 (4.0)	19,318,717	5,429 (28.1)	11,734 (60.7)	612 (3.2)	4,078,446	5,116 (125)	23,666 (580)	328 (8.0)
Age	18 - 29	8,821,973	781 (8.9)	400 (4.5)	143 (1.6)	8,821,973	781 (8.9)	400 (4.5)	143 (1.6)				
	30 - 49	16,131,025	3,544 (22.0)	6,424 (39.8)	338 (2.1)	16,131,025	3,544 (22.0)	6,424 (39.8)	338 (2.1)				
	50 - 69	13,791,173	7,510 (54.5)	31,459 (228)	686 (5.0)	13,791,173	7,510 (54.5)	31,459 (228)	686 (5.0)				
	70 - 79	4,674,973	4,820 (103)	22,249 (476)	418 (8.9)					4,674,973	4,820 (103)	22,249 (476)	418 (8.9)
	80+	2,743,798	4,248 (155)	26,719 (974)	341 (12.4)					2,743,798	4,248 (155)	26,719 (974)	341 (12.4)
Ethnicity	Asian or Asian British	3,718,442	599 (16.1)	5,325 (143)	152 (4.1)	3,467,223	475 (13.7)	3,378 (97.4)	125 (3.6)	251,219	124 (49.4)	1,947 (775)	27 (10.7)
	Black or Black British	1,634,470	654 (40.0)	2,086 (128)	58 (3.5)	1,525,603	513 (33.6)	1,384 (90.7)	45 (2.9)	108,867	141 (130)	702 (645)	13 (11.9)
	Mixed	712,534	182 (25.5)	642 (90.1)	22 (3.1)	682,037	153 (22.4)	426 (62.5)	19 (2.8)	30,497	29 (95.1)	216 (708)	<5 (9.8)
	Other ethnic groups	1,497,133	217 (14.5)	1,072 (71.6)	39 (2.6)	1,427,187	162 (11.4)	678 (47.5)	33 (2.3)	69,946	55 (78.6)	394 (563)	<10 (8.6)
	White	36,446,855	18,995 (52.1)	77,118 (212)	1,643 (4.5)	29,619,868	10,362 (35.0)	31,821 (107)	939 (3.2)	6,826,987	8,633 (126)	45,297 (663)	704 (10.3)
	Unknown or missing	2,153,508	256 (11.9)	1,008 (46.8)	12 (0.6)	2,022,253	170 (8.4)	596 (29.5)	6 (0.3)	131,255	86 (65.5)	412 (314)	<10 (4.6)
Deprivation[1]	1 - 2	9,118,746	4,565 (50.1)	18,830 (206)	365 (4.0)	8,069,275	3,064 (38.0)	10,493 (130)	272 (3.4)	1,049,471	1,501 (143)	8,337 (794)	93 (8.9)
	3 - 4	9,567,633	4,105 (42.9)	17,436 (182)	370 (3.9)	8,273,028	2,454 (29.7)	8,370 (101)	234 (2.8)	1,294,605	1,651 (128)	9,066 (700)	136 (10.5)
	5 - 6	9,355,549	4,091 (43.7)	17,642 (189)	392 (4.2)	7,784,840	2,235 (28.7)	7,263 (93.3)	221 (2.8)	1,570,709	1,856 (118)	10,379 (661)	171 (10.9)
	7 - 8	9,076,313	4,146 (45.7)	16,939 (187)	381 (4.2)	7,362,375	2,128 (28.9)	6,334 (86.0)	221 (3.0)	1,713,938	2,018 (118)	10,605 (619)	160 (9.3)
	9 - 10	8,772,902	3,926 (44.8)	16,024 (183)	406 (4.6)	7,000,538	1,908 (27.3)	5,602 (80.0)	210 (3.0)	1,772,364	2,018 (114)	10,422 (588)	196 (11.1)
Smoking status	Current	7,814,045	3,079 (39.4)	16,115 (206)	260 (3.3)	7,287,225	2,492 (34.2)	11,811 (162)	219 (3.0)	526,820	587 (111)	4,304 (817)	41 (7.8)
	Former	10,623,072	6,930 (65.2)	31,768 (299)	581 (5.5)	7,826,062	3,233 (41.3)	11,703 (150)	277 (3.5)	2,797,010	3,697 (132)	20,065 (717)	304 (10.9)
	Never	25,722,494	10,667 (41.5)	38,701 (150)	1,053 (4.1)	21,674,974	5,929 (27.4)	14,346 (66.2)	643 (3.0)	4,047,520	4,738 (117)	24,355 (602)	410 (10.1)
Medical history	Stroke	727,218	1,020 (140)	21,489 (2,955)	104 (14.3)	286,650	323 (113)	8,147 (2,842)	40 (14.0)	440,568	697 (158)	13,342 (3,028)	64 (14.5)
	Myocardial infarction	1,189,182	1,436 (121)	24,212 (2,036)	158 (13.3)	485,506	436 (89.8)	10,486 (2,160)	46 (9.5)	703,676	1,000 (142)	13,726 (1,951)	112 (15.9)
	DVT or PE	603,351	3,263 (541)	4,050 (671)	142 (23.5)	341,788	1,919 (561)	1,377 (403)	76 (22.2)	261,563	1,344 (514)	2,673 (1,022)	66 (25.2)
	Thrombophilia	44,593	146 (327)	202 (453)	23 (51.6)	38,136	126 (330)	144 (378)	21 (55.1)	6,457	20 (310)	58 (898)	<5 (31.0)
	Coronavirus infection[2]	1,284,984	1,131 (88.0)	4,167 (324)	101 (7.9)	1,124,135	576 (51.2)	1,642 (146)	63 (5.6)	160,849	555 (345)	2,525 (1,570)	38 (23.6)
	Diabetes	4,069,412	3,654 (89.8)	25,595 (629)	423 (10.4)	2,460,649	1,509 (61.3)	9,949 (404)	205 (8.3)	1,608,763	2,145 (133)	15,646 (973)	218 (13.6)
	Depression	9,516,020	6,272 (65.9)	24,146 (254)	500 (5.3)	8,040,590	4,077 (50.7)	12,859 (160)	348 (4.3)	1,475,430	2,195 (149)	11,287 (765)	152 (10.3)
	Obesity	10,918,274	8,863 (81.2)	31,194 (286)	679 (6.2)	8,757,004	5,430 (62.0)	15,848 (181)	411 (4.7)	2,161,270	3,433 (159)	15,346 (710)	268 (12.4)
	Cancer	6,609,691	5,674 (85.8)	17,827 (270)	786 (11.9)	4,854,822	2,583 (53.2)	4,462 (91.9)	412 (8.5)	1,754,869	3,091 (176)	13,365 (762)	374 (21.3)
	COPD	1,582,366	2,497 (158)	12,323 (779)	182 (11.5)	714,880	877 (123)	3,839 (537)	76 (10.6)	867,486	1,620 (187)	8,484 (978)	106 (12.2)
	Liver disease	230,243	284 (123)	1,069 (464)	97 (42.1)	173,423	208 (120)	547 (315)	75 (43.2)	56,820	76 (134)	522 (919)	22 (38.7)
	CKD	2,934,107	4,467 (152)	25,816 (880)	526 (17.9)	978,886	1,179 (120)	5,164 (528)	200 (20.4)	1,955,221	3,288 (168)	20,652 (1,056)	326 (16.7)
	Major surgery[3]	3,995,870	5,407 (135)	24,164 (605)	763 (19.1)	2,773,174	2,735 (98.6)	9,898 (357)	440 (15.9)	1,222,696	2,672 (219)	14,266 (1,167)	323 (26.4)
	Dementia	524,293	1,127 (215)	6,073 (1,158)	45 (8.6)	48,821	81 (166)	415 (850)	10 (20.5)	475,472	1,046 (220)	5,658 (1,190)	35 (7.4)
Medications	Antiplatelet	2,510,382	2,814 (112)	36,415 (1,451)	218 (8.7)	1,000,010	747 (74.7)	15,556 (1,556)	65 (6.5)	1,510,372	2,067 (137)	20,859 (1,381)	153 (10.1)
	BP lowering	8,589,860	7,969 (92.8)	55,462 (646)	777 (9.0)	4,286,561	2,779 (64.8)	20,579 (480)	305 (7.1)	4,303,299	5,190 (121)	34,883 (811)	472 (11.0)
	Lipid lowering	6,808,408	5,796 (85.1)	47,859 (703)	564 (8.3)	3,212,600	1,878 (58.5)	19,543 (608)	194 (6.0)	3,595,808	3,918 (109)	28,316 (787)	370 (10.3)
	Anticoagulant	1,338,585	2,278 (170)	12,758 (953)	205 (15.3)	381,588	1,144 (300)	2,773 (727)	57 (14.9)	956,997	1,134 (118)	9,985 (1043)	148 (15.5)
	Oral contraceptive	622,529	138 (22.2)	42 (6.7)	8 (1.3)	622,529	138 (22.2)	42 (6.7)	8 (1.3)				
	HRT	540,722	213 (39.4)	516 (95.4)	16 (3.0)	501,957	185 (36.9)	368 (73.3)	14 (2.8)	38,765	28 (72.2)	148 (382)	<5 (5.2)
Number of diagnoses	0	36,455,307	12,200 (33.5)	38,617 (106)	980 (2.7)	31,819,272	7,612 (23.9)	18,316 (57.6)	643 (2.0)	4,636,035	4,588 (99.0)	20,301 (438)	337 (7.3)
	1 - 5	9,573,068	8,422 (88.0)	46,012 (481)	925 (9.7)	6,872,479	4,137 (60.2)	19,183 (279)	515 (7.5)	2,700,589	4,285 (159)	26,829 (993)	410 (15.2)
	6+	134,567	281 (209)	2,622 (1948)	21 (15.6)	52,420	86 (164)	784 (1496)	9 (17.2)	82,147	195 (237)	1,838 (2,237)	12 (14.6)
Number of medications	0	22,970,920	3,679 (16.0)	9,629 (41.9)	321 (1.4)	22,122,043	3,143 (14.2)	7,258 (32.8)	290 (1.3)	848,877	536 (63.1)	2,371 (279)	31 (3.7)
	1 - 5	20,875,516	12,872 (61.7)	56,255 (269)	1,210 (5.8)	15,618,756	7,203 (46.1)	25,416 (163)	713 (4.6)	5,256,760	5,669 (108)	30,839 (587)	497 (9.5)
	6+	2,316,506	4,352 (188)	21,367 (922)	395 (17.1)	1,003,372	1,489 (148)	5,609 (559)	164 (16.3)	1,313,134	2,863 (218)	15,758 (1,200)	231 (17.6)

[1] Index of Multiple Deprivation deciles where 10 indicates least deprived and 1 indicates most deprived; [2] After 31/12/2019 and prior to 08/12/2020; [3] In the last year.

Table 2(a) Numbers and incidence rates pre-, post-first ChAdOx1-S. Incidence rate per 100,000 person years. Disclosure control prevents presentation of n>5

Outcome	Age	Pre vaccination or unvaccinated		Post vaccination ≤28 days		Post vaccination >28 days	
		N (events)	Incidence rate (95%	N (events)	Incidence rate (95% CI)	N (events)	Incidence rate (95%
All venous	All	14769	140.7 (138.5–143.0)	2006	309.0 (295.8–322.9)	995	382.6 (359.6–407.2)
	<70	9892	104.3 (102.2–106.3)	902	219.5 (205.7–234.3)	234	240.6 (211.7–273.5)
	≥70	4877	484.5 (471.1–498.3)	1104	463.4 (436.9–491.6)	761	467.5 (435.4–501.9)
Intracranial venous thrombosis	All	207	1.97 (1.72–2.26)	27	4.16 (2.85–6.06)	9	3.46 (1.80–6.65)
	<70	188	1.98 (1.72–2.29)	22	5.35 (3.52–8.13)	<10	5.14 (2.14–12.35)
	≥70	19	1.89 (1.20–2.96)	5	2.10 (0.87–5.04)	<5	2.45 (0.92–6.54)
Portal vein thrombosis	All	121	1.15 (0.96–1.38)	14	2.16 (1.28–3.64)	9	3.46 (1.80–6.65)
	<70	98	1.03 (0.85–1.26)	9	2.19 (1.14–4.21)	<5	2.06 (0.51–8.22)
	≥70	23	2.28 (1.52–3.44)	5	2.10 (0.87–5.04)	<10	4.30 (2.05–9.01)
Pulmonary embolism	All	8473	80.7 (79.0–82.5)	1203	185.3 (175.1–196.1)	587	225.7 (208.1–244.7)
	<70	5399	56.9 (55.4–58.4)	515	125.3 (114.9–136.6)	134	137.8 (116.3–163.2)
	≥70	3074	305.3 (294.7–316.3)	688	288.7 (267.9–311.1)	453	278.2 (253.7–305.0)
Deep vein thrombosis	All	5664	54.0 (52.6–55.4)	739	113.8 (105.9–122.3)	374	143.8 (129.9–159.1)
	<70	3971	41.9 (40.6–43.2)	345	83.9 (75.5–93.3)	90	92.5 (75.3–113.8)
	≥70	1693	168.1 (160.3–176.3)	394	165.3 (149.8–182.5)	284	174.4 (155.2–195.9)
Other	All	459	4.37 (3.99–4.79)	51	7.85 (5.97–10.33)	30	11.5 (8.1–16.5)
	<70	359	3.78 (3.41–4.20)	34	8.27 (5.91–11.58)	9	9.25 (4.81–17.78)
	≥70	100	9.93 (8.16–12.08)	17	7.13 (4.43–11.47)	21	12.9 (8.4–19.8)
All arterial	All	57602	549.1 (544.7–553.6)	8256	1274.4 (1247.2–1302.1)	4634	1787.0 (1736.3–1839.2)
	<70	31396	331.0 (327.4–334.7)	2993	729.4 (703.8–756.0)	755	777.4 (723.9–834.9)
	≥70	26206	2607.7 (2576.3–2639.5)	5263	2215.7 (2156.7–2276.4)	3879	2391.5 (2317.4–2467.9)
Myocardial infarction	All	28134	268.1 (265.0–271.3)	3814	587.9 (569.6–606.9)	2050	788.9 (755.5–823.8)
	<70	17341	182.8 (180.1–185.5)	1628	396.4 (377.6–416.2)	363	373.4 (336.9–413.9)
	≥70	10793	1072.7 (1052.6–1093.1)	2186	918.3 (880.6–957.6)	1687	1037.3 (988.9–1088.0)
Ischaemic stroke (ischaemic, unknown, retinal and spinal)	All	28639	272.9 (269.8–276.1)	4334	668.2 (648.6–688.4)	2539	977.6 (940.3–1016.4)
	<70	13505	142.4 (140.0–144.8)	1309	318.7 (301.9–336.4)	377	387.8 (350.6–429.0)
	≥70	15134	1504.7 (1480.9–1528.9)	3025	1271.6 (1227.1–1317.7)	2162	1330.4 (1275.5–1387.7)
Other arterial	All	1308	12.5 (11.8–13.2)	189	29.1 (25.2–33.6)	98	37.7 (30.9–45.9)
	<70	775	8.17 (7.61–8.76)	82	19.9 (16.1–24.8)	22	22.6 (14.9–34.3)
	≥70	533	52.9 (48.6–57.6)	107	44.9 (37.1–54.2)	76	46.6 (37.3–58.4)
Other							
Disseminated intravascular coagulation	All	13	0.12 (0.07–0.21)	<5	0.46 (0.15–1.43)	<5	-
	<70	5	0.05 (0.02–0.13)	<5	0.49 (0.12–1.95)	<5	-
	≥70	8	0.79 (0.40–1.59)	<5	0.42 (0.06–2.98)	<5	-
Thrombotic thrombocytopenic purpura	All	70	0.67 (0.53–0.84)	<5	0.62 (0.23–1.64)	<5	0.77 (0.19–3.07)
	<70	64	0.67 (0.53–0.86)	<5	0.73 (0.24–2.26)	<5	2.06 (0.51–8.22)
	≥70	6	0.60 (0.27–1.33)	<5	0.42 (0.06–2.98)	<5	-
Any thrombocytopenia	All	1357	12.9 (12.3–13.6)	200	30.8 (26.8–35.4)	89	34.2 (27.8–42.1)
	<70	915	9.64 (9.04–10.29)	124	30.2 (25.3–36.0)	41	42.1 (31.0–57.2)
	≥70	442	43.9 (40.0–48.2)	76	31.9 (25.5–39.9)	48	29.5 (22.2–39.1)
Haemorrhagic stroke (intra-cerebral or subarachnoid)	All	3747	35.7 (34.6–36.9)	487	75.0 (68.6–82.0)	289	111.1 (99.0–124.6)
	<70	2028	21.4 (20.5–22.3)	157	38.2 (32.7–44.7)	45	46.3 (34.5–62.0)
	≥70	1719	170.7 (162.8–179.0)	330	138.4 (124.3–154.2)	244	149.8 (132.1–169.8)
Mesenteric thrombosis	All	1170	11.1 (10.5–11.8)	171	26.3 (22.7–30.6)	109	41.9 (34.7–50.5)
	<70	531	5.60 (5.14–6.09)	56	13.6 (10.5–17.7)	20	20.6 (13.3–31.9)
	≥70	639	63.5 (58.7–68.6)	115	48.2 (40.2–57.9)	89	54.6 (44.4–67.2)
Lower limb fracture	All	18347	174.9 (172.3–177.4)	2657	409.4 (394.2–425.3)	1632	628.0 (598.3–659.2)
	<70	8732	92.0 (90.1–94.0)	539	131.2 (120.5–142.7)	155	159.4 (136.2–186.5)
	≥70	9615	955.8 (936.9–975.1)	2118	890.0 (852.9–928.7)	1477	908.3 (863.2–955.9)
Death	All	123580	1177.5 (1170.9–1184.1)	16192	2493.1 (2455.0–2531.8)	11738	4510.6 (4429.7–4592.9)
	<70	29802	314.1 (310.5–317.7)	1940	471.9 (451.4–493.4)	897	922.0 (863.6–984.3)
	≥70	93778	9312.7 (9253.3–9372.5)	14252	5978.2 (5880.8–6077.1)	10841	6653.3 (6529.2–6779.7)

Table 2(b) Numbers and incidence rates pre-, post-first BNT162b2. Incidence rate per 100,000 person years. Disclosure control prevents presentation of n>5

Outcome	Age	Pre vaccination or unvaccinated		Post vaccination ≤28 days		Post vaccination >28 days	
		N (events)	Incidence rate (95%	N (events)	Incidence rate (95%	N (events)	Incidence rate (95%
All venous	All	14769	140.7 (138.5–143.0)	1546	250.1 (237.9–262.9)	1587	287.0 (273.2–301.5)
	<70	9892	104.3 (102.2–106.3)	548	154.9 (142.4–168.4)	259	110.8 (98.1–125.2)
	≥70	4877	484.5 (471.1–498.3)	998	377.7 (355.0–401.9)	1328	415.9 (394.1–438.9)
Intracranial venous thrombosis	All	207	1.97 (1.72–2.26)	13	2.10 (1.22–3.62)	6	1.08 (0.49–2.41)
	<70	188	1.98 (1.72–2.29)	5	1.41 (0.59–3.39)	<5	1.28 (0.41–3.98)
	≥70	19	1.89 (1.20–2.96)	8	3.03 (1.51–6.05)	<5	0.94 (0.30–2.91)
Portal vein thrombosis	All	121	1.15 (0.96–1.38)	5	0.81 (0.34–1.94)	12	2.17 (1.23–3.82)
	<70	98	1.03 (0.85–1.26)	<5	0.56 (0.14–2.26)	6	2.57 (1.15–5.71)
	≥70	23	2.28 (1.52–3.44)	<5	1.13 (0.37–3.52)	6	1.88 (0.84–4.18)
Pulmonary embolism	All	8473	80.7 (79.0–82.5)	928	150.1 (140.8–160.1)	955	172.7 (162.1–184.0)
	<70	5399	56.9 (55.4–58.4)	306	86.5 (77.3–96.7)	141	60.3 (51.2–71.2)
	≥70	3074	305.3 (294.7–316.3)	622	235.3 (217.6–254.6)	814	254.8 (237.9–273.0)
Deep vein thrombosis	All	5664	54.0 (52.6–55.4)	555	89.8 (82.6–97.6)	590	106.7 (98.4–115.6)
	<70	3971	41.9 (40.6–43.2)	213	60.2 (52.6–68.8)	103	44.1 (36.3–53.5)
	≥70	1693	168.1 (160.3–176.3)	342	129.4 (116.4–143.9)	487	152.4 (139.5–166.6)
Other	All	459	4.37 (3.99–4.79)	55	8.89 (6.83–11.59)	39	7.05 (5.15–9.65)
	<70	359	3.78 (3.41–4.20)	28	7.91 (5.46–11.46)	13	5.56 (3.23–9.58)
	≥70	100	9.93 (8.16–12.08)	27	10.2 (7.0–14.9)	26	8.14 (5.54–11.95)
All arterial	All	57602	549.1 (544.7–553.6)	7960	1289.8 (1261.7–1318.4)	8799	1594.8 (1561.8–1628.5)
	<70	31396	331.0 (327.4–334.7)	2263	640.2 (614.4–667.2)	876	375.1 (351.1–400.8)
	≥70	26206	2607.7 (2576.3–2639.5)	5697	2160.4 (2105.0–2217.2)	7923	2490 (2435.7–2545.4)
Myocardial infarction	All	28134	268.1 (265.0–271.3)	3722	602.5 (583.4–622.2)	4023	728.0 (705.9–750.9)
	<70	17341	182.8 (180.1–185.5)	1285	363.3 (344.0–383.7)	489	209.3 (191.6–228.7)
	≥70	10793	1072.7 (1052.6–1093.1)	2437	922.9 (886.9–960.2)	3534	1107.9 (1072.0–1145.1)
Ischaemic stroke (ischaemic, unknown, retinal and spinal)	All	28639	272.9 (269.8–276.1)	4143	670.6 (650.5–691.4)	4702	851.2 (827.2–875.9)
	<70	13505	142.4 (140.0–144.8)	949	268.2 (251.7–285.9)	369	157.9 (142.6–174.9)
	≥70	15134	1504.7 (1480.9–1528.9)	3194	1209.9 (1168.6–1252.6)	4333	1359.3 (1319.4–1400.4)
Other arterial	All	1308	12.5 (11.8–13.2)	156	25.2 (21.6–29.5)	167	30.2 (25.9–35.1)
	<70	775	8.17 (7.61–8.76)	46	13.0 (9.7–17.3)	26	11.1 (7.6–16.3)
	≥70	533	52.9 (48.6–57.6)	110	41.6 (34.5–50.2)	141	44.1 (37.4–52.0)
Other							
Disseminated intravascular coagulation	All	13	0.12 (0.07–0.21)	<5	0.32 (0.08–1.29)	<5	0.36 (0.09–1.45)
	<70	5	0.05 (0.02–0.13)	<5	0.28 (0.04–2.01)	<5	-
	≥70	8	0.79 (0.40–1.59)	<5	0.38 (0.05–2.69)	<5	0.63 (0.16–2.50)
Thrombotic thrombocytopenic purpura	All	70	0.67 (0.53–0.84)	<5	0.65 (0.24–1.72)	<5	0.36 (0.09–1.45)
	<70	64	0.67 (0.53–0.86)	<5	0.85 (0.27–2.63)	<5	0.86 (0.21–3.42)
	≥70	6	0.60 (0.27–1.33)	<5	0.38 (0.05–2.69)	<5	-
Any thrombocytopenia	All	1357	12.9 (12.3–13.6)	136	22.0 (18.6–26.0)	144	26.0 (22.1–30.6)
	<70	915	9.64 (9.04–10.29)	55	15.5 (11.9–20.2)	32	13.7 (9.7–19.4)
	≥70	442	43.9 (40.0–48.2)	81	30.6 (24.6–38.1)	112	35.1 (29.1–42.2)
Haemorrhagic stroke (intra-cerebral or subarachnoid)	All	3747	35.7 (34.6–36.9)	440	71.2 (64.8–78.1)	579	104.7 (96.5–113.5)
	<70	2028	21.4 (20.5–22.3)	93	26.3 (21.4–32.2)	55	23.5 (18.1–30.7)
	≥70	1719	170.7 (162.8–179.0)	347	131.3 (118.2–145.8)	524	164.0 (150.6–178.7)
Mesenteric thrombosis	All	1170	11.1 (10.5–11.8)	161	26.0 (22.3–30.4)	217	39.2 (34.3–44.8)
	<70	531	5.60 (5.14–6.09)	28	7.91 (5.46–11.46)	18	7.70 (4.85–12.22)
	≥70	639	63.5 (58.7–68.6)	133	50.3 (42.4–59.6)	199	62.3 (54.2–71.6)
Lower limb fracture	All	18347	174.9 (172.3–177.4)	2573	416.2 (400.5–432.6)	3176	574.6 (554.9–594.9)
	<70	8732	92.0 (90.1–94.0)	443	125.2 (114.0–137.4)	228	97.6 (85.7–111.1)
	≥70	9615	955.8 (936.9–975.1)	2130	806.2 (772.7–841.2)	2948	923.9 (891.2–957.9)
Death	All	123580	1177.5 (1170.9–1184.1)	9117	1474.4 (1444.0–1504.9)	13935	2518.7 (2477.3–2560.9)
	<70	29802	314.1 (310.5–317.7)	775	218.9 (204.0–234.9)	531	227.2 (208.7–247.4)
	≥70	93778	9312.7 (9253.3–9372.5)	8342	3155.6 (3088.6–3224.1)	13404	4194.8 (4124.4–4266.4)

Figure 1 Adjusted Hazard ratios for all venous thromboses, intracranial venous thromboses, pulmonary embolism and deep vein thromboses after ChAdOx1-S or BNT162b2 vaccine

Figure 2 Adjusted hazard ratios for all arterial thromboses, myocardial infarction, ischemic stroke and other arterial thromboses after ChAdOx1-S or BNT162b2 vaccine

Figure 3 Adjusted hazard ratios for thrombocytopenia hemorrhagic stroke, mesenteric thrombosis, lower limb fracture after ChAdOx1-S or BNT162b2 vaccine

SUPPLEMENTARY TABLES AND FIGURES

Supplementary Table 1: Hazard ratios (95% CIs) for thrombotic and other outcomes 1-28 and >28 days post-vaccination, compared with pre-vaccination rates. All analyses were stratified on geographical region (A) ChAdOx1-S vaccine (B) BNT162b2 vaccine

Outcome	Age	Sex	1-28 days post-vaccination			>28 days post vaccination s		
			Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]	Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]
(A) ChAdOx1-S vaccine								
All venous events	<70	Overall	2.33 (2.16–2.51)	1.32 (1.22–1.43)	0.97 (0.90–1.05)	2.74 (2.39–3.14)	1.60 (1.38–1.84)	0.94 (0.81–1.08)
		Men	2.68 (2.41–2.97)	1.33 (1.19–1.49)	1.04 (0.93–1.16)	3.13 (2.54–3.84)	1.52 (1.23–1.89)	0.97 (0.78–1.20)
		Women	2.08 (1.87–2.31)	1.31 (1.17–1.47)	0.96 (0.85–1.07)	2.56 (2.13–3.07)	1.67 (1.38–2.02)	1.02 (0.84–1.23)
	≥70	Overall	0.65 (0.59–0.72)	0.64 (0.59–0.70)	0.58 (0.53–0.63)	0.67 (0.59–0.76)	0.58 (0.52–0.65)	0.49 (0.44–0.55)
		Men	0.65 (0.56–0.76)	0.65 (0.56–0.74)	0.59 (0.51–0.68)	0.70 (0.58–0.84)	0.63 (0.53–0.75)	0.55 (0.46–0.65)
		Women	0.65 (0.58–0.74)	0.64 (0.56–0.71)	0.57 (0.50–0.64)	0.64 (0.54–0.76)	0.54 (0.47–0.64)	0.45 (0.39–0.53)
Intracranial venous thrombosis	<70	Overall	2.81 (1.70–4.64)	2.99 (1.75–5.09)	2.27 (1.33–3.88)	2.37 (0.91–6.12)	2.47 (0.95–6.42)	1.72 (0.66–4.49)
		Men	2.72 (1.15–6.39)	2.58 (0.99–6.74)	2.06 (0.79–5.37)	3.43 (0.74–15.87)	3.37 (0.71–15.90)	2.40 (0.53–10.86)
		Women	2.69 (1.45–5.00)	3.27 (1.74–6.17)	2.24 (1.17–4.31)	1.75 (0.52–5.93)	2.14 (0.63–7.22)	1.47 (0.43–5.04)
	≥70	Overall	0.62 (0.17–2.23)	0.65 (0.19–2.21)	0.67 (0.20–2.18)	1.35 (0.13–13.46)	1.13 (0.14–9.20)	1.11 (0.14–9.02)
		Men	10.8 (3.3–35.8)	8.99 (2.75–29.42)	10.1 (2.7–38.5)	-	-	-
		Women	0.27 (0.05–1.42)	0.29 (0.06–1.43)	0.28 (0.06–1.34)	0.87 (0.10–7.32)	0.82 (0.11–6.24)	0.76 (0.10–5.70)
Portal vein thrombosis	<70	Overall	2.49 (1.20–5.14)	1.56 (0.72–3.40)	1.00 (0.24–4.15)	2.76 (0.73–10.51)	1.99 (0.53–7.52)	1.00 (0.06–18.18)
		Men	3.38 (1.40–8.17)	1.95 (0.76–5.02)	1.22 (0.48–3.08)	3.55 (0.52–24.13)	2.16 (0.32–14.52)	0.98 (0.14–6.75)
		Women	1.74 (0.48–6.26)	1.06 (0.28–4.07)	0.72 (0.18–2.86)	2.39 (0.37–15.53)	1.57 (0.24–10.12)	0.86 (0.12–6.28)
	≥70	Overall	0.66 (0.11–4.06)	0.67 (0.10–4.24)	0.61 (0.10–3.71)	2.72 (0.49–15.02)	3.00 (0.48–18.99)	2.61 (0.41–16.52)
		Men	0.43 (0.09–2.09)	0.42 (0.08–2.15)	0.37 (0.07–1.90)	8.60 (2.33–31.70)	7.99 (1.93–33.03)	7.99 (2.03–31.38)
		Women	0.92 (0.06–13.23)	0.92 (0.06–13.99)	0.82 (0.06–11.05)	0.87 (0.12–6.02)	0.95 (0.12–7.39)	0.86 (0.11–6.92)
Pulmonary embolism	<70	Overall	2.54 (2.30–2.80)	1.28 (1.15–1.43)	0.95 (0.85–1.05)	3.12 (2.61–3.73)	1.63 (1.34–1.96)	0.95 (0.79–1.15)
		Men	2.83 (2.46–3.24)	1.27 (1.09–1.48)	1.01 (0.87–1.17)	3.32 (2.52–4.38)	1.43 (1.07–1.91)	0.92 (0.69–1.23)
		Women	2.35 (2.04–2.70)	1.30 (1.12–1.52)	0.95 (0.82–1.10)	3.09 (2.44–3.92)	1.81 (1.41–2.32)	1.09 (0.85–1.41)
	≥70	Overall	0.62 (0.55–0.70)	0.60 (0.54–0.68)	0.54 (0.48–0.61)	0.61 (0.52–0.72)	0.54 (0.46–0.62)	0.45 (0.39–0.53)
		Men	0.63 (0.52–0.75)	0.62 (0.52–0.74)	0.57 (0.47–0.67)	0.63 (0.50–0.79)	0.57 (0.46–0.70)	0.49 (0.40–0.62)
		Women	0.61 (0.52–0.72)	0.59 (0.51–0.69)	0.53 (0.45–0.61)	0.60 (0.48–0.75)	0.51 (0.42–0.63)	0.43 (0.35–0.53)
Deep vein thrombosis	<70	Overall	2.08 (1.85–2.34)	1.34 (1.18–1.52)	0.99 (0.87–1.12)	2.37 (1.90–2.96)	1.58 (1.25–1.99)	0.97 (0.77–1.22)
		Men	2.61 (2.22–3.07)	1.48 (1.25–1.76)	1.15 (0.97–1.36)	2.85 (2.04–3.98)	1.62 (1.14–2.29)	1.06 (0.75–1.50)
		Women	1.67 (1.40–2.00)	1.21 (1.00–1.46)	0.90 (0.75–1.09)	2.12 (1.57–2.86)	1.57 (1.15–2.13)	1.02 (0.75–1.40)
	≥70	Overall	0.73 (0.62–0.86)	0.71 (0.61–0.83)	0.63 (0.54–0.74)	0.71 (0.57–0.88)	0.62 (0.51–0.75)	0.51 (0.42–0.62)
		Men	0.72 (0.56–0.94)	0.71 (0.55–0.91)	0.63 (0.49–0.81)	0.77 (0.56–1.06)	0.70 (0.51–0.95)	0.58 (0.42–0.79)
		Women	0.73 (0.59–0.90)	0.72 (0.59–0.87)	0.64 (0.53–0.77)	0.67 (0.51–0.88)	0.57 (0.45–0.73)	0.47 (0.37–0.60)
Other	<70	Overall	2.48 (1.70–3.62)	1.60 (1.06–2.40)	1.02 (0.69–1.52)	3.04 (1.54–6.00)	2.04 (1.01–4.09)	0.94 (0.47–1.91)
		Men	2.61 (1.51–4.53)	1.45 (0.79–2.64)	1.02 (0.57–1.83)	3.17 (1.13–8.90)	1.74 (0.60–5.05)	0.89 (0.31–2.58)
		Women	2.38 (1.42–4.00)	1.74 (1.01–2.98)	1.12 (0.65–1.91)	3.15 (1.29–7.73)	2.40 (0.97–5.96)	1.26 (0.50–3.17)
	≥70	Overall	0.46 (0.21–1.01)	0.46 (0.21–0.99)	0.43 (0.20–0.92)	1.13 (0.50–2.54)	1.03 (0.48–2.21)	0.93 (0.42–2.03)
		Men	0.38 (0.11–1.30)	0.39 (0.12–1.31)	0.33 (0.10–1.08)	1.10 (0.36–3.39)	1.02 (0.34–3.04)	0.79 (0.26–2.36)
		Women	0.52 (0.19–1.44)	0.51 (0.20–1.36)	0.50 (0.19–1.32)	1.16 (0.37–3.63)	1.03 (0.36–2.98)	0.98 (0.33–2.94)

Outcome	Age	Sex	1-28 days post-vaccination			>28 days post vaccination s		
			Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]	Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]
All arterial events	<70	Overall	3.02 (2.90–3.14)	1.25 (1.20–1.31)	0.90 (0.86–0.95)	3.55 (3.29–3.83)	1.57 (1.45–1.70)	0.91 (0.84–1.00)
		Men	3.42 (3.26–3.60)	1.26 (1.20–1.34)	0.91 (0.86–0.96)	4.70 (4.27–5.17)	1.66 (1.50–1.84)	0.96 (0.86–1.07)
		Women	2.96 (2.76–3.18)	1.24 (1.14–1.34)	0.93 (0.86–1.00)	3.30 (2.90–3.75)	1.43 (1.25–1.64)	0.88 (0.77–1.02)
	≥70	Overall	0.89 (0.85–0.93)	0.84 (0.80–0.87)	0.76 (0.73–0.79)	1.03 (0.97–1.10)	0.83 (0.78–0.87)	0.72 (0.68–0.77)
		Men	0.90 (0.85–0.96)	0.87 (0.82–0.92)	0.78 (0.74–0.83)	1.06 (0.97–1.16)	0.89 (0.82–0.96)	0.78 (0.71–0.85)
		Women	0.88 (0.82–0.94)	0.80 (0.76–0.85)	0.73 (0.69–0.78)	1.02 (0.93–1.12)	0.77 (0.71–0.83)	0.68 (0.62–0.74)
Myocardial infarction	<70	Overall	2.85 (2.70–3.01)	1.24 (1.17–1.32)	0.88 (0.83–0.94)	2.93 (2.63–3.26)	1.40 (1.24–1.56)	0.83 (0.73–0.93)
		Men	3.23 (3.02–3.44)	1.25 (1.17–1.35)	0.89 (0.83–0.95)	3.52 (3.08–4.03)	1.33 (1.15–1.53)	0.78 (0.68–0.90)
		Women	2.93 (2.63–3.26)	1.22 (1.09–1.38)	0.92 (0.82–1.04)	3.56 (2.95–4.29)	1.58 (1.29–1.92)	1.01 (0.82–1.24)
	≥70	Overall	0.86 (0.80–0.92)	0.83 (0.78–0.88)	0.76 (0.71–0.81)	0.97 (0.89–1.07)	0.83 (0.76–0.90)	0.74 (0.68–0.81)
		Men	0.88 (0.80–0.96)	0.85 (0.78–0.93)	0.77 (0.71–0.84)	0.96 (0.84–1.09)	0.84 (0.74–0.94)	0.73 (0.64–0.82)
		Women	0.85 (0.76–0.94)	0.80 (0.72–0.88)	0.76 (0.69–0.84)	1.03 (0.90–1.19)	0.82 (0.72–0.93)	0.78 (0.68–0.88)
Ischaemic stroke[4]	<70	Overall	3.21 (3.02–3.41)	1.25 (1.17–1.34)	0.90 (0.84–0.96)	4.36 (3.91–4.86)	1.76 (1.56–1.98)	0.94 (0.84–1.07)
		Men	3.73 (3.44–4.05)	1.28 (1.17–1.40)	0.91 (0.83–0.99)	6.58 (5.73–7.57)	2.14 (1.84–2.49)	1.12 (0.96–1.31)
		Women	2.91 (2.64–3.20)	1.21 (1.09–1.35)	0.90 (0.81–1.00)	3.10 (2.59–3.71)	1.32 (1.09–1.59)	0.77 (0.63–0.94)
	≥70	Overall	0.92 (0.87–0.98)	0.85 (0.81–0.90)	0.77 (0.73–0.82)	1.08 (0.99–1.18)	0.83 (0.77–0.90)	0.72 (0.67–0.78)
		Men	0.94 (0.85–1.03)	0.89 (0.82–0.96)	0.81 (0.74–0.88)	1.14 (1.01–1.30)	0.93 (0.83–1.04)	0.82 (0.73–0.92)
		Women	0.92 (0.84–1.00)	0.82 (0.77–0.89)	0.75 (0.69–0.80)	1.04 (0.92–1.17)	0.76 (0.68–0.84)	0.66 (0.59–0.73)
Other arterial	<70	Overall	3.18 (2.48–4.07)	1.24 (0.93–1.64)	0.82 (0.63–1.07)	4.20 (2.62–6.73)	1.68 (0.99–2.86)	0.77 (0.46–1.30)
		Men	3.17 (2.29–4.40)	1.05 (0.73–1.51)	0.75 (0.53–1.07)	5.56 (3.07–10.06)	1.66 (0.86–3.20)	0.89 (0.47–1.69)
		Women	3.67 (2.51–5.37)	1.59 (1.03–2.46)	1.04 (0.68–1.59)	3.60 (1.64–7.91)	1.74 (0.73–4.18)	0.83 (0.35–2.01)
	≥70	Overall	0.44 (0.33–0.59)	0.45 (0.34–0.59)	0.41 (0.31–0.53)	0.55 (0.38–0.79)	0.49 (0.35–0.70)	0.44 (0.31–0.63)
		Men	0.46 (0.31–0.67)	0.45 (0.31–0.66)	0.42 (0.29–0.61)	0.61 (0.35–1.05)	0.56 (0.33–0.95)	0.51 (0.30–0.87)
		Women	0.43 (0.28–0.66)	0.44 (0.30–0.67)	0.39 (0.26–0.59)	0.51 (0.30–0.86)	0.44 (0.27–0.71)	0.38 (0.23–0.62)
Haematological events								
Any thrombocytopenia	<70	Overall	4.85 (3.87–6.07)	3.00 (2.34–3.85)	1.71 (1.35–2.16)	7.48 (5.31–10.53)	4.68 (3.24–6.77)	1.69 (1.16–2.46)
		Men	6.47 (4.62–9.06)	3.52 (2.42–5.13)	1.85 (1.30–2.63)	13.2 (8.1–21.4)	6.84 (3.99–11.73)	2.02 (1.18–3.44)
		Women	3.81 (2.80–5.17)	2.65 (1.89–3.71)	1.95 (1.40–2.72)	4.59 (2.78–7.58)	3.34 (1.98–5.64)	2.08 (1.22–3.54)
	≥70	Overall	0.92 (0.65–1.29)	0.87 (0.63–1.22)	0.79 (0.56–1.10)	1.14 (0.66–1.95)	0.99 (0.59–1.65)	0.84 (0.50–1.41)
		Men	0.84 (0.53–1.32)	0.79 (0.50–1.22)	0.68 (0.43–1.06)	1.37 (0.64–2.91)	1.17 (0.57–2.38)	0.92 (0.45–1.90)
		Women	1.04 (0.62–1.75)	0.99 (0.61–1.63)	0.93 (0.56–1.54)	1.02 (0.46–2.24)	0.88 (0.42–1.85)	0.80 (0.38–1.70)
Other events								
Haemorrhagic stroke[5]	<70	Overall	2.25 (1.88–2.69)	1.05 (0.86–1.27)	1.00 (0.73–1.37)	2.87 (2.10–3.92)	1.39 (1.00–1.93)	1.00 (0.51–1.95)
		Men	2.68 (2.11–3.41)	1.15 (0.89–1.49)	1.01 (0.78–1.30)	2.86 (1.74–4.71)	1.21 (0.72–2.04)	0.87 (0.52–1.46)
		Women	1.93 (1.48–2.52)	0.93 (0.70–1.24)	0.87 (0.66–1.16)	2.94 (1.95–4.42)	1.47 (0.95–2.27)	1.23 (0.80–1.91)
	≥70	Overall	0.83 (0.69–1.00)	0.78 (0.66–0.92)	0.73 (0.62–0.87)	1.09 (0.84–1.40)	0.84 (0.67–1.04)	0.76 (0.61–0.95)
		Men	0.86 (0.65–1.14)	0.82 (0.64–1.05)	0.79 (0.62–1.01)	1.34 (0.93–1.94)	1.03 (0.75–1.43)	0.97 (0.70–1.34)
		Women	0.81 (0.63–1.03)	0.75 (0.60–0.94)	0.70 (0.56–0.87)	0.92 (0.65–1.30)	0.71 (0.52–0.96)	0.64 (0.47–0.86)

Outcome	Age	Sex	1-28 days post-vaccination			>28 days post vaccination s		
			Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]	Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]
Mesenteric thrombosis	<70	Overall	3.25 (2.42–4.37)	1.24 (0.89–1.73)	0.84 (0.61–1.16)	5.47 (3.33–8.99)	2.16 (1.24–3.77)	1.04 (0.60–1.80)
		Men	2.87 (1.72–4.77)	0.98 (0.57–1.71)	0.72 (0.43–1.22)	4.34 (1.59–11.83)	1.42 (0.48–4.25)	0.75 (0.26–2.17)
		Women	3.39 (2.36–4.87)	1.43 (0.94–2.17)	1.01 (0.67–1.52)	5.66 (3.21–9.99)	2.59 (1.36–4.94)	1.40 (0.73–2.69)
	≥70	Overall	0.66 (0.49–0.89)	0.60 (0.46–0.79)	0.53 (0.41–0.70)	0.95 (0.64–1.43)	0.74 (0.52–1.05)	0.62 (0.43–0.89)
		Men	0.65 (0.39–1.09)	0.61 (0.38–0.98)	0.52 (0.32–0.83)	0.81 (0.44–1.49)	0.69 (0.40–1.18)	0.54 (0.31–0.94)
		Women	0.66 (0.46–0.95)	0.60 (0.44–0.84)	0.55 (0.40–0.77)	1.00 (0.60–1.68)	0.76 (0.48–1.20)	0.67 (0.42–1.08)
Lower limb fracture	<70	Overall	1.71 (1.55–1.87)	1.01 (0.91–1.11)	0.85 (0.77–0.93)	2.26 (1.91–2.67)	1.35 (1.13–1.60)	0.98 (0.83–1.17)
		Men	1.35 (1.15–1.58)	0.98 (0.83–1.16)	0.86 (0.73–1.02)	2.22 (1.67–2.93)	1.60 (1.20–2.14)	1.24 (0.93–1.66)
		Women	1.91 (1.70–2.15)	1.03 (0.91–1.16)	0.88 (0.78–0.99)	2.21 (1.80–2.72)	1.25 (1.01–1.55)	0.95 (0.77–1.18)
	≥70	Overall	1.11 (1.02–1.21)	0.94 (0.88–1.01)	0.83 (0.78–0.89)	1.56 (1.39–1.76)	0.95 (0.87–1.04)	0.78 (0.71–0.85)
		Men	1.06 (0.90–1.24)	0.94 (0.83–1.08)	0.84 (0.73–0.95)	1.50 (1.19–1.89)	0.97 (0.81–1.17)	0.81 (0.67–0.97)
		Women	1.11 (1.01–1.22)	0.95 (0.88–1.03)	0.84 (0.77–0.91)	1.51 (1.32–1.73)	0.95 (0.85–1.06)	0.78 (0.70–0.87)
Death	<70	Overall	1.96 (1.87–2.06)	0.60 (0.57–0.63)	0.37 (0.35–0.39)	4.51 (4.20–4.85)	1.41 (1.30–1.53)	0.51 (0.47–0.56)
		Men	2.35 (2.20–2.51)	0.63 (0.58–0.67)	0.44 (0.41–0.47)	6.47 (5.87–7.13)	1.57 (1.40–1.75)	0.75 (0.67–0.85)
		Women	1.69 (1.56–1.82)	0.56 (0.52–0.61)	0.36 (0.33–0.39)	3.48 (3.13–3.89)	1.21 (1.07–1.37)	0.50 (0.44–0.57)
	≥70	Overall	0.35 (0.34–0.36)	0.34 (0.33–0.35)	0.28 (0.28–0.29)	0.49 (0.47–0.51)	0.38 (0.37–0.39)	0.27 (0.26–0.28)
		Men	0.31 (0.30–0.32)	0.32 (0.31–0.33)	0.26 (0.25–0.27)	0.44 (0.42–0.47)	0.37 (0.35–0.38)	0.26 (0.24–0.27)
		Women	0.39 (0.37–0.40)	0.36 (0.35–0.37)	0.30 (0.29–0.31)	0.54 (0.51–0.56)	0.39 (0.38–0.41)	0.29 (0.27–0.30)

Outcome	Age	Sex	1-28 days post-vaccination			>28 days post vaccination s		
			Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]	Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]
(B) BNT162b2 vaccine **All venous events**	<70	Overall	1.47 (1.35–1.61)	1.01 (0.93–1.11)	0.81 (0.74–0.88)	1.14 (1.01–1.29)	0.88 (0.77–1.00)	0.73 (0.64–0.83)
		Men	1.98 (1.75–2.23)	1.15 (1.01–1.30)	0.92 (0.81–1.05)	1.56 (1.28–1.89)	1.02 (0.83–1.25)	0.83 (0.68–1.02)
		Women	1.20 (1.06–1.37)	0.91 (0.80–1.03)	0.74 (0.65–0.84)	1.01 (0.86–1.20)	0.81 (0.69–0.96)	0.70 (0.59–0.83)
	≥70	Overall	0.65 (0.60–0.71)	0.56 (0.52–0.61)	0.57 (0.53–0.62)	0.68 (0.62–0.75)	0.50 (0.46–0.55)	0.50 (0.45–0.55)
		Men	0.67 (0.59–0.76)	0.59 (0.52–0.66)	0.60 (0.53–0.67)	0.75 (0.65–0.87)	0.59 (0.51–0.68)	0.58 (0.50–0.67)
		Women	0.64 (0.58–0.72)	0.54 (0.49–0.60)	0.56 (0.50–0.63)	0.63 (0.56–0.72)	0.45 (0.40–0.51)	0.45 (0.40–0.52)
Intracranial venous thrombosis	<70	Overall	0.76 (0.31–1.85)	0.72 (0.29–1.78)	0.59 (0.24–1.45)	0.65 (0.20–2.13)	0.59 (0.18–1.93)	0.51 (0.16–1.68)
		Men	1.60 (0.50–5.16)	1.39 (0.43–4.49)	1.09 (0.33–3.61)	1.04 (0.14–7.77)	0.92 (0.12–7.05)	0.74 (0.10–5.62)
		Women	0.39 (0.10–1.60)	0.44 (0.11–1.78)	0.37 (0.09–1.52)	0.48 (0.11–2.07)	0.51 (0.12–2.20)	0.45 (0.10–1.95)
	≥70	Overall	1.65 (0.64–4.29)	1.25 (0.48–3.26)	1.43 (0.55–3.75)	1.13 (0.13–9.75)	0.71 (0.08–6.56)	0.88 (0.10–7.52)
		Men	-	-	-	-	-	-
		Women	2.01 (0.69–5.88)	1.45 (0.49–4.26)	1.57 (0.54–4.54)	1.26 (0.13–11.94)	0.73 (0.07–7.29)	0.82 (0.09–7.61)
Portal vein thrombosis	<70	Overall	0.58 (0.15–2.30)	0.44 (0.11–1.72)	0.29 (0.07–1.17)	3.06 (1.29–7.24)	2.84 (1.19–6.74)	2.19 (0.89–5.37)
		Men	1.28 (0.33–4.95)	0.81 (0.21–3.09)	0.51 (0.13–2.04)	-	-	-
		Women	-	-	-	5.30 (1.98–14.22)	4.28 (1.55–11.78)	3.95 (1.39–11.20)
	≥70	Overall	0.49 (0.10–2.28)	0.43 (0.10–1.87)	0.42 (0.09–1.91)	0.67 (0.23–1.97)	0.54 (0.19–1.56)	0.48 (0.17–1.38)
		Men	0.85 (0.17–4.38)	0.55 (0.11–2.66)	0.49 (0.10–2.46)	1.26 (0.36–4.41)	0.70 (0.23–2.17)	0.48 (0.17–1.38)
		Women	0.22 (0.01–3.38)	0.24 (0.01–3.80)	0.26 (0.02–4.47)	0.37 (0.09–1.52)	0.44 (0.10–2.01)	0.46 (0.11–2.04)
Pulmonary embolism	<70	Overall	1.52 (1.36–1.71)	0.99 (0.87–1.11)	0.78 (0.69–0.88)	1.20 (1.02–1.43)	0.89 (0.75–1.06)	0.71 (0.60–0.85)
		Men	1.93 (1.63–2.28)	1.05 (0.88–1.25)	0.85 (0.71–1.01)	1.56 (1.19–2.05)	0.97 (0.74–1.29)	0.77 (0.58–1.03)
		Women	1.32 (1.13–1.56)	0.94 (0.79–1.10)	0.75 (0.64–0.89)	1.10 (0.88–1.37)	0.84 (0.67–1.05)	0.71 (0.57–0.89)
	≥70	Overall	0.61 (0.55–0.68)	0.53 (0.48–0.59)	0.54 (0.49–0.60)	0.61 (0.54–0.69)	0.45 (0.40–0.51)	0.45 (0.39–0.50)
		Men	0.64 (0.55–0.75)	0.56 (0.48–0.65)	0.57 (0.49–0.66)	0.65 (0.54–0.77)	0.50 (0.42–0.59)	0.49 (0.41–0.59)
		Women	0.59 (0.52–0.68)	0.51 (0.44–0.58)	0.52 (0.45–0.60)	0.58 (0.49–0.68)	0.42 (0.36–0.49)	0.42 (0.36–0.49)
Deep vein thrombosis	<70	Overall	1.38 (1.21–1.59)	1.03 (0.89–1.18)	0.82 (0.71–0.95)	1.05 (0.86–1.28)	0.85 (0.69–1.04)	0.73 (0.59–0.89)
		Men	1.93 (1.60–2.33)	1.23 (1.01–1.49)	0.99 (0.81–1.20)	1.55 (1.15–2.10)	1.10 (0.81–1.48)	0.94 (0.69–1.28)
		Women	1.08 (0.88–1.33)	0.87 (0.71–1.08)	0.72 (0.58–0.89)	0.88 (0.67–1.15)	0.74 (0.56–0.97)	0.65 (0.50–0.86)
	≥70	Overall	0.69 (0.60–0.79)	0.58 (0.51–0.67)	0.61 (0.53–0.70)	0.81 (0.69–0.96)	0.59 (0.50–0.70)	0.61 (0.52–0.73)
		Men	0.66 (0.53–0.82)	0.59 (0.48–0.73)	0.61 (0.49–0.76)	0.95 (0.74–1.23)	0.78 (0.60–1.01)	0.80 (0.61–1.05)
		Women	0.71 (0.59–0.85)	0.59 (0.49–0.70)	0.62 (0.52–0.74)	0.72 (0.58–0.90)	0.49 (0.40–0.61)	0.52 (0.41–0.64)
Other	<70	Overall	2.16 (1.47–3.19)	1.60 (1.07–2.37)	1.18 (0.79–1.77)	1.64 (0.93–2.90)	1.34 (0.75–2.40)	1.03 (0.57–1.87)
		Men	3.25 (1.93–5.50)	2.03 (1.16–3.54)	1.46 (0.83–2.57)	2.14 (0.86–5.34)	1.51 (0.59–3.86)	1.03 (0.39–2.70)
		Women	1.48 (0.82–2.65)	1.20 (0.67–2.16)	0.98 (0.54–1.77)	1.55 (0.75–3.21)	1.31 (0.63–2.74)	1.21 (0.57–2.55)
	≥70	Overall	1.13 (0.66–1.94)	0.97 (0.58–1.64)	1.01 (0.60–1.72)	0.86 (0.39–1.88)	0.66 (0.31–1.41)	0.67 (0.30–1.47)
		Men	1.50 (0.77–2.95)	1.25 (0.65–2.42)	1.27 (0.64–2.49)	1.51 (0.50–4.51)	1.09 (0.36–3.34)	1.07 (0.33–3.44)
		Women	0.79 (0.31–2.04)	0.72 (0.30–1.75)	0.75 (0.30–1.85)	0.42 (0.17–1.06)	0.36 (0.15–0.88)	0.36 (0.14–0.91)
All arterial events	<70	Overall	2.21 (2.12–2.31)	1.31 (1.25–1.37)	0.94 (0.90–0.99)	1.45 (1.35–1.55)	1.08 (1.01–1.16)	0.88 (0.82–0.95)
		Men	3.06 (2.90–3.22)	1.44 (1.36–1.52)	0.99 (0.93–1.05)	2.17 (1.99–2.38)	1.19 (1.08–1.31)	0.91 (0.82–1.00)
		Women	1.85 (1.71–1.99)	1.11 (1.02–1.20)	0.92 (0.84–0.99)	1.44 (1.30–1.60)	0.96 (0.86–1.07)	0.91 (0.81–1.02)
	≥70	Overall	0.90 (0.87–0.93)	0.69 (0.66–0.71)	0.72 (0.70–0.75)	1.15 (1.10–1.20)	0.69 (0.66–0.72)	0.71 (0.68–0.74)
		Men	0.96 (0.91–1.00)	0.75 (0.71–0.78)	0.77 (0.73–0.81)	1.17 (1.11–1.25)	0.74 (0.70–0.79)	0.74 (0.70–0.79)
		Women	0.84 (0.80–0.89)	0.63 (0.60–0.66)	0.68 (0.65–0.71)	1.13 (1.06–1.20)	0.64 (0.61–0.68)	0.68 (0.64–0.73)

Outcome	Age	Sex	1-28 days post-vaccination			>28 days post vaccination s		
			Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]	Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]
Myocardial infarction	<70	Overall	2.24 (2.12–2.37)	1.39 (1.30–1.47)	0.94 (0.88–1.00)	1.42 (1.29–1.55)	1.14 (1.04–1.26)	0.88 (0.80–0.97)
		Men	3.07 (2.87–3.28)	1.51 (1.40–1.62)	0.97 (0.90–1.05)	1.98 (1.76–2.23)	1.13 (1.00–1.28)	0.81 (0.71–0.93)
		Women	1.94 (1.74–2.17)	1.15 (1.03–1.30)	0.94 (0.83–1.06)	1.78 (1.54–2.05)	1.18 (1.02–1.37)	1.11 (0.95–1.30)
	≥70	Overall	0.93 (0.88–0.98)	0.74 (0.71–0.78)	0.74 (0.70–0.78)	1.19 (1.12–1.27)	0.78 (0.73–0.83)	0.75 (0.70–0.80)
		Men	0.99 (0.92–1.06)	0.81 (0.76–0.86)	0.79 (0.73–0.84)	1.20 (1.10–1.31)	0.82 (0.76–0.89)	0.76 (0.70–0.83)
		Women	0.85 (0.78–0.92)	0.66 (0.61–0.72)	0.69 (0.63–0.74)	1.20 (1.08–1.33)	0.74 (0.67–0.81)	0.74 (0.67–0.82)
Ischaemic stroke[4]	<70	Overall	2.18 (2.05–2.33)	1.23 (1.14–1.31)	0.90 (0.83–0.97)	1.46 (1.32–1.62)	1.00 (0.89–1.11)	0.80 (0.72–0.90)
		Men	3.05 (2.80–3.32)	1.38 (1.26–1.51)	0.95 (0.86–1.05)	2.40 (2.08–2.77)	1.26 (1.09–1.46)	0.91 (0.78–1.07)
		Women	1.76 (1.59–1.96)	1.05 (0.94–1.17)	0.85 (0.76–0.95)	1.18 (1.01–1.38)	0.77 (0.66–0.90)	0.73 (0.61–0.86)
	≥70	Overall	0.88 (0.84–0.93)	0.65 (0.63–0.68)	0.71 (0.68–0.75)	1.13 (1.06–1.19)	0.64 (0.60–0.67)	0.69 (0.66–0.73)
		Men	0.94 (0.88–1.00)	0.71 (0.66–0.75)	0.77 (0.72–0.82)	1.17 (1.07–1.27)	0.69 (0.64–0.75)	0.75 (0.69–0.81)
		Women	0.84 (0.78–0.89)	0.61 (0.58–0.65)	0.68 (0.63–0.72)	1.09 (1.01–1.19)	0.60 (0.56–0.65)	0.66 (0.61–0.71)
Other arterial	<70	Overall	1.72 (1.28–2.31)	0.95 (0.70–1.30)	0.69 (0.50–0.94)	1.63 (1.10–2.43)	1.13 (0.74–1.72)	0.85 (0.55–1.32)
		Men	2.31 (1.58–3.37)	0.99 (0.66–1.49)	0.71 (0.47–1.07)	3.22 (2.02–5.13)	1.56 (0.95–2.58)	1.16 (0.69–1.96)
		Women	1.46 (0.90–2.37)	0.88 (0.54–1.45)	0.69 (0.42–1.15)	0.93 (0.43–1.99)	0.65 (0.30–1.43)	0.59 (0.26–1.33)
	≥70	Overall	0.59 (0.46–0.76)	0.51 (0.40–0.65)	0.52 (0.41–0.67)	0.64 (0.47–0.85)	0.47 (0.35–0.62)	0.46 (0.34–0.62)
		Men	0.49 (0.35–0.69)	0.46 (0.32–0.65)	0.47 (0.33–0.66)	0.52 (0.37–0.74)	0.46 (0.32–0.66)	0.44 (0.31–0.64)
		Women	0.74 (0.51–1.08)	0.59 (0.41–0.84)	0.60 (0.42–0.85)	0.83 (0.50–1.37)	0.50 (0.31–0.79)	0.49 (0.30–0.78)
Haematological events								
Any thrombocytopenia	<70	Overall	2.10 (1.60–2.77)	1.48 (1.11–1.97)	1.00 (0.75–1.34)	2.03 (1.42–2.92)	1.54 (1.07–2.23)	0.97 (0.66–1.41)
		Men	3.31 (2.23–4.91)	2.07 (1.35–3.16)	1.19 (0.78–1.83)	2.95 (1.61–5.42)	2.02 (1.08–3.79)	0.93 (0.49–1.78)
		Women	1.50 (1.01–2.22)	1.17 (0.78–1.74)	0.97 (0.65–1.46)	1.58 (1.00–2.50)	1.32 (0.83–2.08)	1.14 (0.72–1.80)
	≥70	Overall	0.85 (0.64–1.13)	0.72 (0.54–0.95)	0.68 (0.51–0.90)	1.22 (0.87–1.71)	0.90 (0.64–1.25)	0.79 (0.56–1.12)
		Men	0.87 (0.61–1.26)	0.73 (0.51–1.05)	0.68 (0.47–0.97)	1.55 (1.03–2.32)	1.12 (0.75–1.67)	0.95 (0.62–1.46)
		Women	0.82 (0.53–1.29)	0.71 (0.46–1.09)	0.68 (0.44–1.07)	0.84 (0.48–1.48)	0.63 (0.36–1.09)	0.57 (0.32–1.02)
Other events								
Haemorrhagic stroke[5]	<70	Overall	1.39 (1.13–1.71)	0.83 (0.67–1.03)	0.77 (0.62–0.96)	1.31 (1.00–1.73)	0.90 (0.68–1.19)	0.86 (0.65–1.15)
		Men	1.65 (1.22–2.23)	0.86 (0.63–1.18)	0.77 (0.57–1.05)	1.43 (0.89–2.29)	0.86 (0.53–1.40)	0.77 (0.47–1.25)
		Women	1.28 (0.95–1.72)	0.80 (0.59–1.08)	0.78 (0.58–1.05)	1.34 (0.95–1.88)	0.89 (0.62–1.26)	0.92 (0.65–1.31)
	≥70	Overall	0.80 (0.69–0.91)	0.59 (0.52–0.68)	0.65 (0.57–0.74)	1.28 (1.09–1.50)	0.72 (0.62–0.84)	0.80 (0.68–0.94)
		Men	0.82 (0.67–1.01)	0.59 (0.48–0.71)	0.64 (0.52–0.78)	1.45 (1.14–1.83)	0.78 (0.62–0.98)	0.86 (0.68–1.09)
		Women	0.78 (0.65–0.94)	0.60 (0.50–0.72)	0.66 (0.55–0.79)	1.16 (0.93–1.44)	0.68 (0.56–0.84)	0.76 (0.61–0.93)
Mesenteric thrombosis	<70	Overall	1.61 (1.10–2.36)	0.84 (0.56–1.26)	0.65 (0.43–0.97)	1.77 (1.11–2.84)	1.08 (0.66–1.75)	0.83 (0.50–1.37)
		Men	2.91 (1.78–4.75)	1.29 (0.77–2.17)	0.97 (0.57–1.64)	2.21 (0.99–4.94)	1.15 (0.50–2.62)	0.82 (0.35–1.91)
		Women	0.91 (0.49–1.67)	0.55 (0.29–1.03)	0.45 (0.23–0.86)	1.61 (0.90–2.88)	1.09 (0.60–1.98)	0.94 (0.50–1.75)
	≥70	Overall	0.72 (0.57–0.90)	0.54 (0.43–0.67)	0.54 (0.44–0.68)	1.06 (0.81–1.37)	0.61 (0.48–0.78)	0.59 (0.45–0.76)
		Men	0.57 (0.37–0.86)	0.41 (0.27–0.61)	0.42 (0.28–0.62)	0.82 (0.53–1.29)	0.44 (0.29–0.66)	0.42 (0.27–0.65)
		Women	0.80 (0.61–1.04)	0.61 (0.47–0.80)	0.62 (0.48–0.81)	1.18 (0.86–1.63)	0.72 (0.53–0.98)	0.70 (0.51–0.96)
Lower limb fracture	<70	Overall	1.44 (1.31–1.59)	1.03 (0.93–1.14)	0.93 (0.84–1.02)	1.24 (1.08–1.42)	0.95 (0.83–1.09)	0.89 (0.78–1.03)
		Men	1.19 (0.99–1.42)	0.94 (0.78–1.12)	0.85 (0.71–1.02)	0.99 (0.74–1.31)	0.83 (0.62–1.11)	0.80 (0.60–1.07)
		Women	1.49 (1.33–1.67)	1.08 (0.96–1.21)	0.98 (0.87–1.10)	1.27 (1.09–1.48)	1.00 (0.85–1.17)	0.94 (0.80–1.11)
	≥70	Overall	0.99 (0.93–1.05)	0.65 (0.62–0.69)	0.69 (0.66–0.73)	1.48 (1.38–1.59)	0.67 (0.63–0.72)	0.71 (0.66–0.76)
		Men	0.95 (0.85–1.06)	0.58 (0.52–0.64)	0.63 (0.57–0.70)	1.60 (1.40–1.83)	0.65 (0.57–0.73)	0.69 (0.61–0.79)
		Women	1.00 (0.93–1.07)	0.68 (0.64–0.73)	0.72 (0.67–0.77)	1.42 (1.30–1.55)	0.68 (0.63–0.73)	0.71 (0.65–0.77)

Outcome	Age	Sex	1-28 days post-vaccination			>28 days post vaccination s		
			Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]	Unadjusted[1]	Age and sex adjusted[2]	Fully adjusted[3]
Death	<70	Overall	0.74 (0.69–0.79)	0.35 (0.32–0.37)	0.24 (0.22–0.26)	0.91 (0.84–1.00)	0.52 (0.48–0.57)	0.33 (0.30–0.37)
		Men	1.06 (0.97–1.17)	0.39 (0.36–0.43)	0.28 (0.25–0.31)	1.62 (1.44–1.82)	0.69 (0.61–0.79)	0.44 (0.39–0.51)
		Women	0.56 (0.50–0.63)	0.29 (0.26–0.33)	0.22 (0.19–0.25)	0.66 (0.58–0.75)	0.38 (0.34–0.44)	0.28 (0.25–0.33)
	≥70	Overall	0.22 (0.22–0.23)	0.16 (0.16–0.16)	0.19 (0.19–0.20)	0.30 (0.29–0.31)	0.16 (0.16–0.16)	0.19 (0.18–0.19)
		Men	0.23 (0.22–0.24)	0.16 (0.15–0.16)	0.19 (0.18–0.20)	0.33 (0.32–0.34)	0.16 (0.15–0.17)	0.19 (0.18–0.19)
		Women	0.21 (0.20–0.22)	0.16 (0.16–0.17)	0.20 (0.19–0.20)	0.27 (0.26–0.28)	0.16 (0.15–0.17)	0.19 (0.18–0.19)

[1]Stratifed by region
[2]Adjusted for age, age^2, sex and stratified by region
[3]Adjusted for age, age^2, sex, ethnicity, deprivation, smoking and co-morbidities, and stratified by region
[4]Ischaemic stroke: including ischaemic stroke, stroke of uncertain cause, and spinal stroke
[5]Hemorrhagic stroke: including intracranial hemorrhage and subarachnoid hemorrhage

Supplementary table 2 (a) Fully adjusted hazard ratios (HRs) for the association of ChAdOx1-S and BNT162b2 with venous events 1-28 and >28 days after vaccination, within pre-specified subgroups. P values for between subgroup differences were derived using Wald tests. OCP: oral contraceptive. HRT: hormone replacement therapy

Subgroup		ChAdOx1-S vaccine N (Events) 3001	Post vaccination ≤28 days HR (95% CI)[1]	P value	Post vaccination >28 days HR (95% CI)[1]	P value	BNT162b2 vaccine N (Events) 3133	Post vaccination ≤28 days HR (95% CI)[1]	P value	Post vaccination >28 days HR (95% CI)[1]	P value
Age groups	18-29	40	2.30 (1.63–3.25)		1.29 (0.58–2.90)		29	0.99 (0.60–1.62)		0.99 (0.57–1.71)	
	30-39	63	1.33 (1.01–1.75)		0.85 (0.46–1.59)		60	0.90 (0.65–1.25)		0.71 (0.47–1.07)	
	40-49	145	1.32 (1.09–1.61)		1.52 (1.09–2.13)		122	0.96 (0.77–1.21)		0.75 (0.56–1.01)	
	50-59	270	0.93 (0.81–1.07)	<0.0001	1.00 (0.77–1.30)	0.003	237	0.78 (0.66–0.92)	0.005	0.69 (0.55–0.86)	0.839
	60-69	618	0.87 (0.79–0.96)		0.89 (0.73–1.07)		359	0.78 (0.69–0.89)		0.67 (0.55–0.83)	
	70-79	1124	0.75 (0.69–0.82)		0.79 (0.71–0.88)		823	0.69 (0.62–0.77)		0.73 (0.65–0.82)	
	80-89	546	0.79 (0.69–0.89)		0.73 (0.63–0.84)		1255	0.63 (0.56–0.70)		0.67 (0.61–0.74)	
	>90	195	0.76 (0.62–0.94)		0.75 (0.59–0.95)		248	0.80 (0.65–0.99)		0.67 (0.55–0.81)	
Sex	Male	1338	0.81 (0.75–0.87)	0.029	0.74 (0.66–0.82)	0.005	1423	0.70 (0.64–0.76)	0.150	0.66 (0.60–0.72)	0.024
	Female	1663	0.90 (0.84–0.97)		0.89 (0.81–0.98)		1710	0.75 (0.70–0.81)		0.74 (0.69–0.81)	
Ethnicity	White	2856	0.86 (0.82–0.91)		0.82 (0.76–0.89)		2977	0.72 (0.68–0.77)		0.71 (0.66–0.76)	
	Asian or Asian British	38	0.51 (0.34–0.77)		0.66 (0.38–1.14)		56	0.69 (0.48–0.98)		0.57 (0.38–0.86)	
	Black or Black British	53	0.91 (0.66–1.25)		0.62 (0.35–1.10)		42	0.62 (0.41–0.94)		0.61 (0.39–0.97)	
	Mixed	10	0.47 (0.21–1.06)	0.018	0.76 (0.28–2.06)	0.429	12	0.74 (0.36–1.51)	0.002	0.43 (0.16–1.17)	0.691
	Other Ethnic Groups	17	0.73 (0.37–1.43)		1.63 (0.80–3.32)		23	1.30 (0.75–2.24)		0.94 (0.48–1.84)	
	Unknown	21	1.11 (0.65–1.89)		1.14 (0.50–2.58)		15	0.74 (0.36–1.51)		0.68 (0.32–1.47)	
	Missing	6	2.59 (1.05–6.42)		1.29 (0.18–9.32)		8	3.85 (1.67–8.87)		1.33 (0.33–5.43)	
Deprivation	1-2 (least)	564	0.76 (0.68–0.85)		0.76 (0.65–0.89)		489	0.65 (0.57–0.74)		0.54 (0.47–0.62)	
	3-4	560	0.86 (0.77–0.96)		0.80 (0.68–0.93)		576	0.76 (0.67–0.85)		0.70 (0.62–0.80)	
	5-6	625	0.90 (0.81–1.00)	0.138	0.82 (0.71–0.96)	0.43	606	0.70 (0.62–0.79)	0.095	0.67 (0.59–0.76)	<0.0001
	7-8	632	0.87 (0.78–0.97)		0.80 (0.69–0.93)		687	0.73 (0.65–0.82)		0.71 (0.63–0.80)	
	9-10 (most)	609	0.91 (0.81–1.01)		0.92 (0.79–1.06)		770	0.81 (0.72–0.91)		0.88 (0.79–0.98)	
Prior thrombocytopenia	No	3001	0.86 (0.82–0.91)	0.030	0.82 (0.76–0.89)	0.127	3132	0.73 (0.69–0.78)	0.014	0.71 (0.66–0.75)	0.372
	Yes	0	0.54 (0.35–0.82)		0.54 (0.31–0.93)		<5	0.37 (0.21–0.63)		0.58 (0.38–0.89)	
Prior thrombophilia	No	2986	0.86 (0.81–0.91)	0.093	0.82 (0.76–0.89)	0.130	3116	0.73 (0.69–0.77)	0.254	0.70 (0.66–0.75)	0.924
	Yes	15	0.50 (0.27–0.94)		0.38 (0.14–1.03)		17	0.48 (0.23–0.99)		0.68 (0.34–1.37)	
Prior venous event	No	3001	0.95 (0.90–1.01)	<0.0001	0.96 (0.88–1.05)	<0.0001	3133	0.80 (0.76–0.86)	<0.0001	0.78 (0.73–0.84)	<0.0001
	Yes	0	0.54 (0.48–0.61)		0.42 (0.35–0.50)		0	0.44 (0.38–0.51)		0.44 (0.38–0.51)	
Prior COVID infection	No	2839	0.88 (0.83–0.93)	<0.001	0.85 (0.79–0.93)	<0.0001	3012	0.74 (0.70–0.79)	0.001	0.72 (0.68–0.77)	<0.001
	Yes	162	0.60 (0.49–0.74)		0.49 (0.37–0.64)		121	0.47 (0.36–0.61)		0.45 (0.34–0.58)	
OCP	No	2996	1.15 (0.99–1.35)	0.382	1.03 (0.77–1.39)	0.786	3129	0.84 (0.70–1.00)	0.195	0.77 (0.61–0.97)	0.270
	Yes	5	0.69 (0.22–2.17)		1.26 (0.31–5.09)		<5	0.33 (0.08–1.34)		0.34 (0.08–1.41)	
HRT	No	2972	0.78 (0.73–0.83)	0.514	0.75 (0.69–0.82)	0.532	3112	0.69 (0.65–0.73)	0.554	0.66 (0.62–0.71)	0.949
	Yes	29	0.92 (0.56–1.50)		0.96 (0.45–2.05)		21	0.57 (0.30–1.08)		0.65 (0.34–1.23)	
Anticoagulant	No	2759	0.98 (0.92–1.03)	<0.0001	0.98 (0.90–1.07)	<0.0001	2875	0.84 (0.79–0.89)	<0.0001	0.83 (0.77–0.89)	<0.0001
	Yes	242	0.34 (0.29–0.40)		0.28 (0.22–0.35)		258	0.25 (0.21–0.31)		0.27 (0.22–0.32)	
Antiplatelet	No	2503	0.89 (0.84–0.95)	<0.001	0.84 (0.77–0.92)	0.081	2513	0.75 (0.71–0.80)	0.011	0.72 (0.67–0.77)	0.220
	Yes	498	0.69 (0.61–0.78)		0.73 (0.62–0.85)		620	0.62 (0.55–0.71)		0.66 (0.58–0.75)	
Prior MI or stroke	No	2603	0.88 (0.83–0.93)	0.004	0.87 (0.80–0.95)	<0.0001	2680	0.74 (0.70–0.79)	0.021	0.72 (0.67–0.77)	0.086
	Yes	398	0.71 (0.62–0.82)		0.58 (0.48–0.69)		453	0.62 (0.53–0.72)		0.63 (0.55–0.73)	
Prior diabetes	No	2363	0.88 (0.83–0.93)	0.053	0.85 (0.78–0.93)	0.034	2456	0.74 (0.70–0.79)	0.149	0.71 (0.66–0.76)	0.639
	Yes	638	0.78 (0.70–0.87)		0.72 (0.62–0.83)		677	0.67 (0.60–0.76)		0.69 (0.61–0.78)	

Supplementary table 2 (b) Fully adjusted hazard ratios (HRs) for the association of ChAdOx1-S and BNT162b2 with arterial events 1-28 and >28 days after vaccination, within pre-specified subgroups. P values for between subgroup differences were derived using Wald tests. OCP: oral contraceptive. HRT: hormone replacement therapy

Subgroup	All arterial	ChAdOx1-S vaccine N (Events)	Post vaccination ≤28 days HR (95% CI)[1]	P value	Post vaccination >28 days HR (95% CI)[1]	P value	BNT162b2 vaccine N (Events)	Post vaccination ≤28 days HR (95% CI)[1]	P value	Post vaccination >28 days HR (95% CI)[1]	P value
Age groups	18-29	12	1.72 (0.89–3.33)		1.84 (0.59–5.73)		21	2.01 (1.13–3.57)		1.85 (0.96–3.59)	
	30-39	54	1.51 (1.11–2.06)		1.54 (0.87–2.72)		67	1.36 (1.00–1.86)		1.09 (0.74–1.61)	
	40-49	277	1.37 (1.20–1.57)		1.29 (0.99–1.69)		284	1.17 (1.01–1.36)		0.86 (0.70–1.05)	
	50-59	963	0.98 (0.91–1.05)	<0.0001	0.89 (0.76–1.04)	0.031	1062	1.06 (0.98–1.14)	<0.0001	0.81 (0.72–0.91)	<0.0001
	60-69	2442	0.85 (0.81–0.90)		0.90 (0.82–0.99)		1705	0.89 (0.84–0.95)		0.83 (0.76–0.92)	
	70-79	5030	0.90 (0.86–0.94)		0.93 (0.88–0.99)		3993	0.85 (0.81–0.89)		0.91 (0.86–0.96)	
	80-89	2826	0.86 (0.81–0.91)		0.88 (0.82–0.94)		7733	0.74 (0.71–0.78)		0.79 (0.76–0.82)	
	>90	1286	0.85 (0.78–0.93)		0.86 (0.78–0.94)		1894	0.72 (0.66–0.79)		0.78 (0.73–0.84)	
Sex	Male	7116	0.78 (0.75–0.80)	<0.0001	0.75 (0.71–0.79)	<0.0001	9292	0.77 (0.74–0.79)	<0.0001	0.71 (0.68–0.73)	<0.0001
	Female	5774	0.89 (0.86–0.93)		0.87 (0.83–0.92)		7467	0.85 (0.82–0.88)		0.89 (0.85–0.92)	
Ethnicity	White	11729	0.82 (0.79–0.84)		0.80 (0.77–0.84)		15492	0.80 (0.78–0.82)		0.79 (0.76–0.81)	
	Asian or Asian British	647	0.84 (0.76–0.93)		0.74 (0.64–0.86)		728	0.75 (0.67–0.84)		0.63 (0.56–0.71)	
	Black or Black British	217	0.90 (0.75–1.07)		0.89 (0.70–1.15)		190	0.77 (0.62–0.94)		0.79 (0.64–0.99)	
	Mixed	72	1.19 (0.90–1.57)	<0.0001	0.56 (0.32–0.98)	0.180	85	1.10 (0.82–1.49)	<0.0001	0.81 (0.57–1.17)	<0.0001
	Other Ethnic Groups	113	0.93 (0.73–1.18)		0.90 (0.64–1.28)		121	0.89 (0.69–1.15)		0.76 (0.57–1.01)	
	Unknown	80	0.88 (0.67–1.16)		0.75 (0.49–1.16)		103	0.87 (0.64–1.18)		1.09 (0.83–1.44)	
	Missing	32	3.01 (1.98–4.57)		1.91 (0.89–4.08)		40	2.87 (1.81–4.56)		2.87 (1.80–4.58)	
Deprivation	1-2 (least)	2498	0.76 (0.71–0.80)		0.71 (0.66–0.77)	<0.0001	2788	0.75 (0.71–0.79)		0.69 (0.65–0.73)	
	3-4	2514	0.81 (0.76–0.85)		0.76 (0.71–0.82)		2980	0.76 (0.72–0.81)		0.74 (0.69–0.78)	
	5-6	2683	0.85 (0.81–0.90)	<0.001	0.79 (0.74–0.85)		3489	0.78 (0.74–0.83)	<0.0001	0.79 (0.75–0.84)	<0.0001
	7-8	2616	0.83 (0.78–0.87)		0.86 (0.80–0.92)		3693	0.82 (0.78–0.87)		0.83 (0.79–0.88)	
	9-10 (most)	2541	0.89 (0.85–0.95)		0.88 (0.82–0.95)		3772	0.90 (0.85–0.95)		0.85 (0.80–0.89)	
Prior thrombocytopenia	No	12889	0.82 (0.80–0.85)	0.334	0.80 (0.77–0.84)	0.499	16758	0.80 (0.78–0.82)	0.920	0.78 (0.76–0.81)	0.136
	Yes	<5	0.75 (0.61–0.91)		0.73 (0.57–0.95)		<5	0.81 (0.66–1.00)		0.67 (0.54–0.82)	
Prior thrombophilia	No	12864	0.82 (0.80–0.85)	0.059	0.80 (0.77–0.83)	0.620	16735	0.80 (0.78–0.82)	0.499	0.78 (0.76–0.81)	0.098
	Yes	26	0.48 (0.28–0.84)		0.66 (0.32–1.39)		24	0.66 (0.38–1.15)		0.43 (0.21–0.88)	
Prior venous event	No	12874	0.83 (0.81–0.85)	0.029	0.80 (0.77–0.84)	0.504	16751	0.81 (0.78–0.83)	0.022	0.79 (0.76–0.81)	0.002
	Yes	16	0.73 (0.66–0.82)		0.77 (0.67–0.88)		8	0.70 (0.63–0.79)		0.66 (0.59–0.74)	
Prior COVID infection	No	12227	0.84 (0.81–0.86)	<0.0001	0.83 (0.79–0.86)	<0.0001	16234	0.81 (0.79–0.84)	<0.0001	0.80 (0.77–0.82)	<0.0001
	Yes	663	0.63 (0.57–0.70)		0.51 (0.45–0.59)		525	0.57 (0.50–0.65)		0.51 (0.45–0.59)	
OCP	No	12889	0.91 (0.79–1.05)	0.983	0.89 (0.67–1.18)	<0.0001	16759	0.91 (0.79–1.06)	<0.0001	0.94 (0.75–1.16)	<0.0001
	Yes	<5	0.89 (0.12–6.51)		0.00 (0.00–0.00)		0	0.00 (0.00–0.00)		0.00 (0.00–0.00)	
HRT	No	12832	0.85 (0.82–0.87)	0.293	0.84 (0.80–0.88)	0.820	16671	0.81 (0.78–0.83)	0.061	0.79 (0.76–0.81)	0.070
	Yes	58	1.01 (0.73–1.40)		0.79 (0.45–1.37)		88	1.10 (0.79–1.51)		1.08 (0.77–1.52)	
Anticoagulant	No	10726	0.84 (0.82–0.87)	<0.001	0.82 (0.78–0.85)	0.015	13715	0.83 (0.81–0.86)	<0.0001	0.80 (0.77–0.83)	<0.001
	Yes	2164	0.74 (0.69–0.79)		0.74 (0.68–0.80)		3044	0.66 (0.62–0.70)		0.71 (0.67–0.76)	
Antiplatelet	No	7636	0.99 (0.96–1.03)	<0.0001	1.06 (1.01–1.11)	<0.0001	9770	0.96 (0.93–0.99)	<0.0001	1.05 (1.01–1.09)	<0.0001
	Yes	5254	0.65 (0.63–0.68)		0.59 (0.55–0.62)		6989	0.65 (0.63–0.67)		0.55 (0.53–0.57)	
Prior MI or stroke	No	6931	1.09 (1.05–1.13)	<0.0001	1.18 (1.12–1.24)	<0.0001	9428	1.08 (1.04–1.12)	<0.0001	1.22 (1.17–1.26)	<0.0001
	Yes	5959	0.64 (0.61–0.66)		0.60 (0.57–0.63)		7331	0.60 (0.57–0.62)		0.51 (0.49–0.53)	
Prior diabetes	No	8742	0.85 (0.83–0.88)	<0.0001	0.84 (0.80–0.88)	<0.001	11689	0.84 (0.82–0.87)	<0.0001	0.82 (0.79–0.85)	<0.0001
	Yes	4148	0.77 (0.73–0.80)		0.73 (0.69–0.78)		5070	0.72 (0.69–0.76)		0.70 (0.67–0.74)	

Supplementary table 3. Characteristics of patient who had an ICVT event before and after vaccination. Only percentage are presented, as disclosure control does not allow numbers <5 to be released

		ICVT before vaccination %	ICVT after ChAdOx1, %	ICVT after BNT162b, %	p-value[1]
N		<210	<40	<20	<0.0001
Sex	Male	40	33	21	0.24
	Female	60	67	79	
Age	18 - 29	19	17	0	
	30 - 49	42	22	26	
	50 - 69	30	36	16	<0.0001
	70 - 79	6	11	21	
	80+	3	14	37	
Ethnicity	Asian or Asian British	15	6	0	
	Black or Black British	5	3	0	
	Mixed	2	0	0	
	Other ethnic groups	1	0	0	0.38
	White	75	92	100	
	Unknown or missing	1	0	0	
Deprivation[2]	1 - 2	29	17	11	
	3 - 4	20	14	26	
	5 - 6	25	25	21	0.19
	7 - 8	13	25	16	
	9 - 10	13	19	26	
Smoking status	Current	19	28	11	
	Former	21	17	32	0.5
	Never	57	56	58	
Medical history	Stroke	5	6	11	<0.0001
	MI	3	6	0	0.01
	DVT or PE	8	6	5	<0.0001
	Thrombophilia	1	0	0	0.05
	Coronavirus infection[3]	6	6	0	<0.0001
	Diabetes	10	17	37	<0.0001
	Depression	26	36	16	<0.0001
	Obesity	37	31	42	<0.0001
	Cancer	17	28	37	<0.0001
	COPD	2	11	16	0.78
	Liver disease	<1	3	0	0.61
	CKD	5	11	32	0.25
	Major surgery[4]	18	8	11	<0.0001
	Dementia	2	3	0	0.03
Medications	Antiplatelet	5	6	16	0.01
	BP lowering	15	28	42	<0.0001
	Lipid lowering	12	22	26	<0.0001
	Anticoagulant	10	11	11	<0.0001
	Oral contraceptive	4	0	0	<0.0001
	HRT	1	0	0	0.05
Number of diagnoses[5]	0	71	50	58	0.04
	1 - 5	29	50	42	
Number of medications	0	31	11	2	
	1 - 5	62	83	5	0.02
	6+	7	6	2	

[1] Approximation if Chi squared test included one or more zero counts; [2] Index of Multiple Deprivation deciles where 1 indicates least deprived and 10 indicates most deprived; [3] After 31/12/2019 and prior to 08/12/2020; [4] In the last year; [5] The category '6+' contained no individuals and so was excluded.

Supplementary figure 1: Cumulative incidence of vaccines of different types during follow up

Supplementary figure 2: Adjusted hazard ratios for portal vein thrombosis, other venous events, and death after ChAdOx1-S or BNT162b2 vaccine