

1 **Full title:** Is the BioNTech-Pfizer COVID-19 vaccination effective in elderly populations?

2 Results from population data from Bavaria, Germany

3 **Short title:** Effectiveness of BNT162b2 vaccination in the elderly

4 Delphina Gomes^{1¶}, Andreas Beyerlein^{1¶}, Katharina Katz², Gabriele Hoelscher², Uta

5 Nennstiel², Bernhard Liebl², Klaus Überla³, Rüdiger von Kries^{1*}

6 ¹ Institute of Social Pediatrics and Adolescent Medicine, Division of Pediatric Epidemiology,

7 Ludwig-Maximilians-University, Munich, Germany

8 ² Bavarian Health and Food Safety Authority, Oberschleissheim, Germany

9 ³ Institute of Clinical and Molecular Virology, Universitätsklinikum Erlangen, Friedrich-

10 Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany

11 *ruediger.kries@med.uni-muenchen.de

12 [¶]These authors contributed equally to this work.

13 **Abbreviations:** CI, confidence interval; COVID-19, coronavirus disease; HR, hazard ratios;

14 KM, Kaplan-Meier; LGL, Bavarian Health and Food Safety Authority; PHM, Public Health

15 Microbiology; RKI, Robert Koch-Institute; RT-PCR, reverse transcription polymerase chain

16 reaction; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; STIKO, Standing

17 Committee on Vaccination; VE, vaccine efficacy.

18 **Abstract**

19 **Background**

20 The effect of the BioNTech-Pfizer BNT162b2 vaccination in the elderly (≥ 80 years)
21 could not be fully assessed in the BioNTech-Pfizer trial due to low numbers in this age group.
22 We aimed to evaluate the effectiveness of the BioNTech-Pfizer (BNT162b2) vaccine to
23 prevent SARS-CoV-2 infection and severe outcomes in octo- and novo-generians in a
24 German state setting.

25 **Methods and Findings**

26 A prospective observational study of 708,187 persons aged ≥ 80 years living in
27 Bavaria, Germany, was conducted between Jan 9 to Apr 11, 2021. We assessed the vaccine
28 efficacy (VE) for two doses of the BNT162b2 vaccine with respect to SARS-CoV-2 infection
29 and related hospitalisations and mortality. Additionally, differences in VE by age groups ≥ 80
30 to ≤ 89 years and ≥ 90 years were studied. Analyses were adjusted by sex.

31 By the end of follow-up, 63.8% of the Bavarian population ≥ 80 years had received one
32 dose, and 52.7% two doses, of the BNT162b2 vaccine. Two doses of the BNT162b2 vaccine
33 lowered the proportion of SARS-CoV-2 infections and related outcomes, resulting in VE
34 estimates of 68.3% (95% confidence interval (CI) 65.5%, 70.9%) for infection, 73.2% (95%
35 CI 65.3%, 79.3%) for hospitalisation, and 80.1% (95% CI 80.0%, 89.0%) for mortality. Sex
36 differences in the risk of COVID-19 outcomes observed among unvaccinated persons
37 disappeared after two BNT162b2 vaccine doses. Overall, the BNT162b2 vaccine was equally
38 efficacious in octo- and novo-generians.

39 **Conclusions**

40 Two doses of BioNTech-Pfizer's BNT162b2 vaccine is highly effective against COVID-19
41 outcomes in elderly persons.

42 **Introduction**

43 Global research on coronavirus disease (COVID-19) has shown that an increasing age is the
44 most relevant risk factor for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)
45 infections.¹ The high risk for poor outcomes of SARS-CoV-2 infection in the elderly is likely
46 to be due to chronic comorbidities such as hypertension and cardiovascular disease,²
47 diminishing physiological functions including that of the respiratory system,³ and impaired
48 immune response.⁴

49 The high SARS-CoV-2 infection case fatality rates in the elderly population⁵ has guided
50 prioritization of this age group in several countries, including Germany.^{6,7} The COVID-19
51 disease vaccination programme in Germany was launched on Dec 27, 2020 using the
52 BioNTech-Pfizer mRNA BNT162b2⁸ vaccine which requires two doses within at least 21
53 days for full protection.⁹ Other vaccines including Moderna (mRNA-1273), Oxford-
54 AstraZeneca (ChAdOx1 nCoV-19), and Johnson & Johnson (Ad26.COV2.S) were licensed
55 for use in Germany on Jan 6, 2021, Jan 29, 2021, and Mar 11, 2021, respectively.

56 In this study, we used vaccine uptake records of the large population of individuals aged ≥ 80
57 years in Bavaria ($n \geq 700,000$), Germany, to evaluate the vaccine efficacy (VE) of the
58 BNT162b2 vaccine with respect to SARS-CoV-2 infection or COVID-19-related
59 hospitalisation or mortality in this age group. We further assessed whether VE differed
60 between age groups ≥ 80 to ≤ 89 years and ≥ 90 years.

61 **Methods**

62 **Study design and population**

63 The federal state of Bavaria is located in the South-eastern part of Germany. Of its nearly 13
64 million inhabitants, 831,499 (6.3%) persons were aged 80 years and above (males 38.6%,
65 females 61.4%) by 31 Dec, 2019¹⁰ and were considered as reference population in the present
66 analysis.

67 **The German vaccination programme**

68 The German Ministry of Health outlined recommendations for COVID-19 vaccination in day-
69 to-day care practices and designated vaccination centres for implementation in respective
70 German states. All vaccines used until Apr 30, 2021 in Germany required two doses of the
71 same vaccine, according to the Standing Committee on Vaccination (STIKO).⁷

72 Data on the COVID-19 vaccination was collected by authorized staff of the vaccination
73 centers and vaccination teams and entered in the Bavarian Corona vaccination portal
74 “BayIMCO”. During the study period cumulative vaccination data was transferred to the
75 Robert Koch-Institut (RKI), because the transmission of records was not yet possible. The
76 Bavarian Health and Food Safety Authority (LGL) could receive anonymous records for
77 regional studies.

78 Aggregated level data on vaccination uptake in Bavaria per calendar week until Apr 11, 2021
79 were used for this study. Vaccination uptake data for persons aged ≥ 80 years included
80 information on vaccination type (BNT162b2, ChAdOx1 nCoV-19, and mRNA-1273), number
81 of vaccination doses received (1 or 2), sex, and age group (≥ 80 to < 85 years, ≥ 85 to < 90
82 years, ≥ 90 to < 95 years, and ≥ 95 years).

83 **Testing for SARS-CoV-2 infection**

84 Following the mandate given by the German Ministry of Health, notification of SARS-CoV-2
85 infections became obligatory since Jan 30, 2020.¹¹ A positive COVID-19 case was defined as
86 a person with a laboratory confirmation by detection of the SARS-CoV-2 virus by reverse
87 transcription polymerase chain reaction (RT-PCR).^{12,13} Samples were tested for SARS-CoV-2
88 infection at the Public Health Microbiology (PHM) laboratory of the LGL,¹⁴ private hospitals
89 and university laboratories, and in routine care.¹³ Further details on the Bavarian strategy of
90 SARS-CoV-2 testing can be found elsewhere.¹³ The data were collected at the public health
91 offices and pseudo-anonymously forwarded to the LGL.

92 Data on all COVID-19 cases aged ≥ 80 years in Bavaria between Jan 9, 2021 to Apr 11, 2021
93 was extracted from the LGL data bank. We used the data up until Apr 11, 2021 because as of
94 Mar 31, 2021 vaccination was also implemented by general practitioners, who document
95 vaccinations in a different platform and format. Thus, uniform vaccination uptake data were
96 no longer available.

97 Case data included information on age, sex, whether an RT-PCR test was performed, date of
98 infection, hospitalisation status, date and reasons of hospitalisation, mortality status, date and
99 reasons of mortality, vaccination dose (1 or 2), as well as date and type of the last COVID-19
100 vaccination.

101 **Outcomes**

102 We assessed VE for SARS-CoV-2 infection, COVID-19-related hospitalisation, and COVID-
103 19-related mortality. A case with a SARS-CoV-2 infection was defined as a person with a
104 positive RT-PCR test result. All SARS-CoV-2 infections included both symptomatic and
105 asymptomatic cases. Hospitalisation was defined as admission to hospital with a positive RT-

106 PCR test result. A COVID-19-related mortality was defined as a person with a positive RT-
107 PCR test result who died because of or in temporal relation to a SARS-CoV-2 infection.

108 **Exclusion criteria for analysis**

109 The analysis was limited to persons aged ≥ 80 years in Bavaria. We excluded observations
110 with any of the following conditions: a) the infection was not confirmed by a positive RT-
111 PCR test result, b) the infection was reported to have occurred before vaccination, c) the
112 infection was related to a hospitalisation or mortality which occurred before 9th Jan, 2021, d)
113 the vaccination was performed already in 2020, e) the person received another vaccine than
114 BNT162b2). The exclusion criteria d) and e) applied also for the vaccine uptake dataset, with
115 29,071 persons already vaccinated in 2020, and 40,049 persons and 54,192 persons
116 vaccinated with Moderna (mRNA-1273) or Oxford-AstraZeneca (ChAdOx1 nCoV-19),
117 respectively, leaving a final number of $n=708,187$ for our analysis. There were no
118 vaccinations with Johnson & Johnson (Ad26.COV2.S) recorded in the data because
119 procurement was still outstanding during the course of the study.

120 **Statistical analyses**

121 To assess VE for each outcome, we combined the information from the vaccination uptake
122 dataset with aggregate level data and from the SARS-CoV-2 infection dataset with individual
123 patient data. The uptake data were used to calculate the number of vaccinated and
124 unvaccinated persons per calendar week in 2021. Persons who were vaccinated with the
125 BNT162b2 vaccine between Jan 9, 2021 (calendar week 1) and Apr 11, 2021 (calendar week
126 14) contributed to the follow-up of both unvaccinated and vaccinated persons. For example, if
127 a person was vaccinated in calendar week 5, he/she was regarded as under follow-up for 4
128 weeks in the unvaccinated group and for the subsequent 10 weeks in the vaccinated group,
129 respectively (and as censored thereafter in both cases).

130 With the combined vaccination uptake and infection data, we calculated Kaplan-Meier (KM)
131 plots, p-values from log-rank tests, and hazard ratios (HRs) with corresponding 95%
132 confidence intervals (CIs) from Cox's proportional hazard models for time to SARS-CoV-2
133 infection by vaccination group. Time to any SARS-CoV-2 infection, time to SARS-CoV-2
134 infection with later hospitalisation, and time to SARS-CoV-2 infection leading to mortality
135 were investigated as separate outcome variables. In case the date of COVID-19 related
136 hospitalisation or mortality was earlier than the date of confirmed SARS-CoV-2 infection, the
137 date of SARS-CoV-2 infection was set to the date of hospitalisation or mortality, respectively.
138 All events were inversely weighted by the number of SARS-CoV-2 infections per calendar
139 week in the age group ≥ 80 years to account for temporal changes in infection risk. Having
140 two vaccination doses was used as the main exposure variable and compared to having no
141 vaccination, respectively. All analyses were done for the whole age group ≥ 80 years and the
142 HRs were additionally stratified by age ≥ 80 to ≤ 89 years and ≥ 90 years. Furthermore, the KM
143 plots were stratified by sex, and the HRs were calculated with and without adjustment for sex.
144 VE was derived as 1 minus HR.

145 All analyses were carried out using R 3.6.3. The analysis code is available at
146 <https://osf.io/e9qm6/>.

147 **Role of the funding source**


148 The funder had no role in the study design, data analysis, interpretation, or writing of the
149 report. All authors had full access to all the data in the study and had final responsibility for
150 the decision to submit for publication.

151 **Results**

152 **Vaccine uptake in Bavaria in elderly persons**

153 Until Apr 11, 2021, 63.8% elderly persons received one and 52.7% received the second
154 BNT162b2 vaccination dose in Bavaria (Figure 1).


155 **Fig 1. Uptake of BNT162b2 vaccine in persons ≥ 80 years in Bavaria.**



156 **Confirmed COVID-19 cases in Bavaria in elderly persons**

157 Within the follow-up period, there were 11,228 Bavarian persons aged ≥ 80 years with a
158 confirmed SARS-CoV-2 infection based on a positive RT-PCR test (Fig 2). After excluding
159 vaccinated cases with missing information on vaccine dose or type and those who were not
160 vaccinated with the BioNTech-Pfizer BNT162b2 vaccine (n=135), a total of 1,148 vaccinated
161 cases were considered in the vaccination group. Of these included vaccinated cases, 44.9%
162 were partially and 55.1% were fully vaccinated with the BNT162b2 vaccine.

163 **Fig 2. Flow chart of the study population.**



164 The mean age of the included cases was 86.1 years (SD 4.9 years) and 35.5% were males.
 165 There were 1,349 cases who were hospitalized and 1,731 cases who died due to SARS-CoV-2
 166 infection. The proportion of COVID-19-related hospitalisation did not differ according to
 167 vaccination status (from Chi-square test $p=0.135$). However, the proportion of COVID-19-
 168 related mortality was significantly lower in vaccinated persons compared to persons not
 169 vaccinated (from Chi-square test $p<0.0001$).

170 **Risk of SARS-CoV-2 infection and related outcomes**


171 In persons aged ≥ 80 years, having received two compared to no BNT162b2 vaccination doses
172 was associated with a substantially lower cumulative risk of getting a SARS-CoV-2 infection
173 (0.41% versus 1.25% after 70 days, Figure 3A). The estimate for VE after 2 doses of the
174 BNT162b2 vaccine to prevent SARS-CoV-2 infection was 68.3% (95% CI 65.5%, 70.9%)
175 (Figure 4A).

176 The risk of COVID-19-related hospitalisation (0.04% versus 0.19% after 70 days) and
177 COVID-19-related mortality (0.03% versus 0.20% after 70 days) were also lower after two
178 doses of the BNT162b2 vaccine (Figure 3B, Figure 3C). In the group of unvaccinated
179 persons, the risk of infection was higher in females than in males and the risk for
180 hospitalisation and mortality was higher in males than in females. After two doses of the
181 BNT162b2 vaccine, females showed a significantly higher cumulative infection risk than
182 males ($p < 0.0001$ from log-rank test), while there were no significant sex differences for VE
183 with respect to hospitalisation ($p = 0.91$) and mortality ($p = 0.77$), respectively.

184 By the end of the follow-up period, VE for COVID-19-related hospitalisation and COVID-19-
185 related mortality was 73.2% (95% CI 65.3%, 79.3%) and 85.1% (95% CI 80.0%, 89.0%),
186 respectively (Figure 4B, Figure 4C). Estimates of VE in octo- and novo-generians overlapped.


187 S1 Figure presents VE based on the HR for SARS-CoV-2 infection and related outcomes in
188 cases with one or more doses of the BNT162b2 vaccination. The trends in risk of SARS-CoV-
189 2 infections and related outcomes were similar after either ≥ 1 dose or both doses of the
190 BNT162b2 vaccine. However, VE for all outcomes were lower in persons with at least 1 than
191 in persons with both doses of the BNT162b2 vaccine.

192 **Fig 3. Risk of SARS-CoV-2 infection and related outcomes after two BNT162b2 vaccine**
 193 **doses in Bavarian persons aged 80 years and above.**



194

194 **Fig 4. Vaccine efficacy to prevent SARS-CoV-2 infection and related outcomes after two**
195 **BNT162b2 vaccine doses compared to none in Bavarian persons aged 80 years and**
196 **above.**



197 Vaccine efficacy was calculated as $1 - \text{HR}$, adjusted for sex. CI, confidence interval; HR,
198 hazards ratio; VE, vaccine efficacy.

199 **Discussion**

200 In this population-based analysis of all Bavarian persons aged ≥ 80 years with suitable follow-
201 up between Jan 9, 2021 and Apr 11, 2021, we found a high protection against SARS-CoV-2
202 infection and mortality after the second dose of the BNT162b2 vaccine. This association did
203 not differ by age above ≥ 80 to ≤ 89 years and ≥ 90 years. In the unvaccinated group, women
204 were more likely to be infected than men whereas men were more likely to be hospitalized or
205 die from COVID-19. Sex differences in the risk of hospitalisation and mortality in the absence
206 of vaccination disappeared widely after vaccination.

207 Population-based studies to evaluate the efficacy of COVID-19 vaccines have shown that a
208 single dose of the BNT162b2 vaccine had an efficacy between 42% and 75% against SARS-
209 CoV-2 infection, between 43% and 85% against COVID-19-related hospitalisation, and
210 between 51% and 72% against COVID-19-related mortality after at least 14 days.¹⁵ Overall,
211 the estimates of VE increased after the second dose.¹⁵ There are scarce studies addressing VE
212 after BNT162b2 vaccination in persons 80 years and above and none distinguishing VE in
213 octo- and novo-genarians.

214 In the efficacy study on the BioNTech-Pfizer trial, VE to reduce SARS-CoV-2 infection was
215 94.7% in persons ≥ 65 years and 100.0% in persons ≥ 75 years. However, the CI for the VE
216 estimates included null which could be possibly due to the low proportion of elderly persons
217 in their study. Our real life data of elderly Bavarian persons, we found an overall VE of
218 68.3% after two doses with small 95% confidence bands. The efficacy of 68.3% against any
219 infection after two BNT162b2 might explain outbreaks in homes for the elderly despite high
220 uptake of the vaccine reported in the press occasionally. Interestingly, VE did not differ
221 between octo- and novo-genarians. The BioNTech-Pfizer efficacy study could not assess VE
222 by outcome due to few hospitalized cases and mortality in the older age groups. Our data

223 clearly show that VE against hospitalisation after 2 doses is the same as for any SARS-CoV-2
224 infection and is substantially and significantly higher for COVID-19-related mortality.

225 Although there was a clear difference in the risk of SARS-CoV-2 infection and COVID-19-
226 related hospitalisation and mortality between men and women, no such difference was
227 observed after vaccination for hospitalisation or mortality, suggesting that BNT162b2 is
228 equally effective for severe outcomes in elderly men and women.

229 An important study on elderly persons has addressed the onset of effect after one and two
230 doses of the BNT162b2 or the Oxford-AstraZeneca vaccine.¹⁶ The study showed that the VE
231 increases up to 42 days after the first dose and up to after 14 days after the second dose. While
232 the risk for infection decreased only gradually within the first 42 days after the first dose, VE
233 after the second dose already reached 80% after 7 days. During vaccination programmes
234 requiring two doses, it is evident that some persons will only be vaccinated once up to the
235 second dose. In order to estimate the effect of the vaccination programme, it may also be
236 interesting to know the effect of one or more doses. In a population where most second doses
237 were given within 3 weeks after the first dose, VE was found only slightly lower after 2 doses
238 with overlapping 95% CIs.

239 A strength of our study is the large sample of elderly persons living in Bavaria. Data was
240 retrieved from government data bases and have been checked for implausibilities. Given the
241 structure of the data base for cases which only included date of the last vaccination and
242 number of vaccine doses, we were unfortunately unable to evaluate the VE after the first
243 BNT162b2 vaccine dose. Further, information of risk factors which could contribute to severe
244 COVID-19 disease outcomes in persons aged 80 years and above have not been ascertained
245 for the cases.

246 In conclusion, we found that two doses of BioNTech-Pfizer's BNT162b2 vaccine provided a
247 high protection against SARS-CoV-2 infection and severe outcomes in elderly persons. There

248 was no difference between octo- and novo-generians and efficacy appeared to be equal in
249 both sexes.

250 **Acknowledgements**

251 We are grateful to the public health offices in Bavaria, Germany who supplied data on cases
252 with SARS-CoV-2 infection and to teams of the Bavarian Corona vaccination portal
253 “BayIMCO” for providing the COVID-19 vaccination data.

254 **Disclosure of conflicts of interest**

255 All authors stated that they had no interests that might be perceived as posing a conflict or
256 bias. As well, they all disclose prior publication and submission of the manuscript.

257 **Author Contributions**

258 **Conceptualization:** Klaus Überla, Rüdiger von Kries

259 **Data Curation:** Delphina Gomes, Andreas Beyerlein, Katharina Katz, Gabriele Hölscher,
260 Uta Nennstiel, Bernhard Liebl

261 **Formal Analysis:** Andreas Beyerlein, Delphina Gomes

262 **Funding Acquisition:** Klaus Überla, Rüdiger von Kries

263 **Investigation:** Rüdiger von Kries

264 **Methodology:** Andreas Beyerlein, Rüdiger von Kries

265 **Project Administration:** Rüdiger von Kries

266 **Software:** Andreas Beyerlein, Delphina Gomes

267 **Supervision:** Rüdiger von Kries

268 **Validation:** Andreas Beyerlein, Delphina Gomes, Rüdiger von Kries

269 **Visualization:** Andreas Beyerlein, Delphina Gomes, Rüdiger von Kries

270 **Writing – Original Draft Preparation:** Delphina Gomes, Andreas Beyerlein, Rüdiger von
271 Kries

272 **Writing – Review & Editing:** Delphina Gomes, Andreas Beyerlein, Katharina Katz,
273 Gabriele Hölscher, Uta Nennstiel, Bernhard Liebl, Klaus Überla, Rüdiger von Kries


274 **References**

- 275 1. Singhal S, Kumar P, Singh S, Saha S, Dey AB. Clinical features and outcomes of
276 COVID-19 in older adults: a systematic review and meta-analysis. *BMC Geriatr* 2021;
277 **21**(1): 321.
- 278 2. Thakur B, Dubey P, Benitez J, et al. A systematic review and meta-analysis of
279 geographic differences in comorbidities and associated severity and mortality among
280 individuals with COVID-19. *Sci Rep* 2021; **11**(1): 8562.
- 281 3. Dhama K, Patel SK, Kumar R, et al. Geriatric Population During the COVID-19
282 Pandemic: Problems, Considerations, Exigencies, and Beyond. *Front Public Health*
283 2020; **8**: 574198.
- 284 4. Bajaj V, Gadi N, Spihlman AP, Wu SC, Choi CH, Moulton VR. Aging, Immunity, and
285 COVID-19: How Age Influences the Host Immune Response to Coronavirus Infections?
286 *Front Physiol* 2020; **11**: 571416.
- 287 5. Yanez ND, Weiss NS, Romand JA, Treggiari MM. COVID-19 mortality risk for older
288 men and women. *BMC Public Health* 2020; **20**(1): 1742.
- 289 6. European Centre for Disease Prevention and Control. Overview of the implementation of
290 COVID-19 vaccination strategies and deployment plans in the EU/EEA – 6 May
291 2021. ECDC: Stockholm; 2021.
- 292 7. Robert Koch-Institut. STIKO-Empfehlungen zur COVID-19-Impfung, 12.05.2021.
293 Germany, 2021.
- 294 8. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA
295 Covid-19 Vaccine. *N Engl J Med* 2020; **383**(27): 2603-15.
- 296 9. Clinical trial number NCT04368728 for "NCT04368728: Study to Describe the Safety,
297 Tolerability, Immunogenicity, and Efficacy of RNA Vaccine Candidates Against
298 COVID-19 in Healthy Individuals" at [ClinicalTrials.gov](https://clinicaltrials.gov).

- 299 10. Fortschreibung des Bevölkerungsstandes (EVAS-Nr. 12411) – Bevölkerung:
300 Bundesländer, Stichtag, Geschlecht, Altersjahre. Statistisches Bundesamt (Destatis).
301 2020.
- 302 11. Bundesministerium für Gesundheit. [Regulation on the extension of the notification
303 obligation according to § 6 para. 1 sentence 1 number 1 and § 7 para. 1 sentence 1 of the
304 German Protection against Infection Act with the novel coronavirus ('2019-nCoV')]. In:
305 German Ministry of Health, 2020.
- 306 12. Robert Koch Institute. Case definition: Coronavirus disease-2019 (COVID-19) (SARS-
307 CoV-2) [Faldefinition: Coronavirus-Krankheit-2019 (COVID-19) (SARS-CoV-2)]. In.
308 Berlin 2020.
- 309 13. Böhm S, Woudenberg T, Chen D, et al. Epidemiology and transmission characteristics of
310 early COVID-19 cases, 20 January-19 March 2020, in Bavaria, Germany. *Epidemiol*
311 *Infect* 2021; **149**: e65.
- 312 14. Böhmer MM, Buchholz U, Corman VM, et al. Investigation of a COVID-19 outbreak in
313 Germany resulting from a single travel-associated primary case: a case series. *The Lancet*
314 *Infectious Diseases* 2020; **20**(8): 920-8.
- 315 15. Henry D, Jones M, Stehlik P, Glasziou P. Effectiveness of COVID-19 vaccines: findings
316 from real world studies. *MJA*, 2021.
- 317 16. Lopez Bernal J, Andrews N, Gower C, et al. Effectiveness of the Pfizer-BioNTech and
318 Oxford-AstraZeneca vaccines on covid-19 related symptoms, hospital admissions, and
319 mortality in older adults in England: test negative case-control study. *BMJ* 2021; **373**:
320 n1088.
- 321

322 **Supporting information**

323 **S1 Figure. Vaccine efficacy to prevent SARS-CoV-2 infection and related outcomes after**
324 **at least one BNT162b2 vaccine dose compared to none in Bavarian persons aged 80**
325 **years and above.**



Vaccine efficacy was calculated as $1 - \text{HR}$ and are adjusted for sex. CI, confidence interval; HR, hazards ratio; VE, vaccine efficacy.