

# 1 Anticipating the hospital burden of future COVID-19 epidemic waves

2 Corentin Boennec · Samuel Alizon\* · Mircea T Sofonea\*

3

4 August 19, 2021

5 **Abstract** Forecasting SARS-CoV-2 epidemic trends with confidence more than a few weeks ahead is almost  
6 impossible as these entirely depend on political decisions. We address this problem by investigating the con-  
7 sequences for the health system of an epidemic wave of a given size. This approach yields semi-quantitative  
8 results that depend on the proportion of the population already infected and vaccinated. We introduce the  
9 COVimpact software, which allows users to visualise estimated numbers of ICU admissions, deaths, and in-  
10 fections stratified by age class at the French departmental, regional, or national level caused by the wave. We  
11 illustrate the usefulness of our approach by showing that for France, even with a 95% vaccination coverage,  
12 the current vaccine efficiency against the delta variant would make a large epidemic wave infecting 25% of  
13 the population difficult to sustain for the current hospital bed occupancy capacity. Overall, using the final  
14 epidemic wave size and ignoring detailed epidemiological dynamics yields valuable and practical insights to  
15 optimise public health response to epidemics.

16 **Keywords** epidemic wave · mathematical modelling · intensive care units · final epidemic size

---

MIVEGEC, Uni Montpellier, CNRS, IRD, Montpellier, France

E-mail: [corentin.boennec@ird.fr](mailto:corentin.boennec@ird.fr)

E-mail: [samuel.alizon@cnrs.fr](mailto:samuel.alizon@cnrs.fr)

E-mail: [mircea.sofonea@umontpellier.fr](mailto:mircea.sofonea@umontpellier.fr)

\* These authors contributed equally to this work.

**NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.**

## 17 **1 Introduction**

18 In France, as in other countries, detailed mechanistic models have been developed that can capture COVID-19  
19 epidemic dynamics, especially hospital admission and bed occupancy data [8, 10]. The problem is that these  
20 models are extremely unreliable to make forecasting more than 3 to 4 weeks ahead because epidemic dynamics  
21 rely on political decisions. For example, let us consider a country with an intermediate SARS-CoV-2 infection  
22 prevalence that decides to switch public health strategy the next week. It is obvious that the number of hos-  
23 pital admissions a month later will be strikingly different if the strategy consists in alleviating control over  
24 transmission chains (e.g. as some regions in Brazil in 2020), or in maintaining incidence to a high plateau (as  
25 France in spring 2021), or in applying 'zero COVID' strategy (as in countries such as New-Zealand, Australia,  
26 or Vietnam). In a way, COVID-19 epidemiological forecasting more than a month ahead largely consists in an-  
27 ticipating political decisions. On top of this, additional uncertainty originates from social (e.g. school holidays),  
28 environmental (e.g. humidity), and biological (e.g. virus evolution) factors.

29 We address this issue by ignoring detailed temporal epidemiological dynamics and summarising these  
30 into a single metric namely the final epidemic size, i.e. the proportion of the population that will be infected  
31 during the epidemic wave. This allows us to estimate the resulting number of casualties and of intensive care  
32 unit (ICU) admissions, one of the most crucial indicators for health care systems, caused by a future epidemic  
33 wave.

34 Several hurdles need to be overcome that are further discussed in the Methods and in Supplementary  
35 Text S1. First, we need to stratify the total number of cases in the epidemic wave by age the SARS-CoV-2  
36 infection fatality ratio is strongly age-dependent [8]. Second, and also for each age, we need to estimate which  
37 proportion of the population that has already been infected by the virus and has developed immunity. Third,  
38 we need to set the proportion of the population has been vaccinated, and by which type of vaccine. Again, this  
39 variable should be stratified by age. To help public health field workers, we also introduce a dedicated software  
40 COVimpact, which performs these estimations at the French department, regional, and national level. In spite  
41 of the lack of temporal dynamics (we do not know whether the total number of admissions will be spread over  
42 one or 4 months), the semi-quantitative results we obtain provide valuable insights on the long term (more  
43 than a month ahead). As we illustrate in the case of France, where our results indicate that vaccination alone  
44 is unlikely to be sufficient in the near future, this approach can be instrumental in planning and optimisation  
45 local hospital response to future COVID-19 epidemic waves.

## 46 2 Methods

47 Our goal is to estimate the total number of ICU admissions and deaths caused by an epidemic wave of a given  
48 size, i.e. associated with the total number of people infected. A first difficulty is that this variable, as most of  
49 the ones used in our approach, needs to be stratified by the age of the individuals ( $a$ ) and their geographical  
50 location (department, region, or country). Two other layers of stratification are natural immunity (i.e. having  
51 recovered from a previous infection) and vaccine immunity. If  $N$  vaccines are being implemented in the pop-  
52 ulation, and assuming an individual is only vaccinated by a single type of vaccine, this leads to  $2N + 2$  host  
53 categories for each age.

54 The number of hosts of age  $a$  that have been infected in the past is calculated by combining a previous  
55 mathematical model from the team [10] and detailed data analysed by Hozé et al. [4]. Note that some hosts can  
56 be reinfected (approximately 16% [2]), which means that the number of hosts with natural immunity is lower  
57 than the number of past infections.

58 The proportion of hosts of age  $a$  vaccinated by vaccine  $i$  is set by the user. The vaccine efficacy, i.e. the  
59 protection against severe infection forms, is also set by the user. Note that our model formalism does not  
60 require any assumption regarding vaccine protection against reinfection or against transmission. We assume  
61 that vaccination is not targeted towards hosts without natural immunity. For hosts with both natural and  
62 vaccine immunity, we assume that the level of protection conferred against severe infection forms is the highest  
63 of the two protections.

64 Finally, we assume that all hosts are exposed in the same way to the infection, i.e. there is no effect of age,  
65 or of natural and vaccine immunity.

66 Regarding the link between the epidemic size, estimating the proportion of hosts infected by an uncon-  
67 trolled epidemic wave ( $p^*$ ) is an old problem in epidemiology and it is known to be tightly linked with the  
68 basic reproduction number ( $R_0$ ), the expected number of secondary infections caused by an infected host in an  
69 otherwise fully susceptible population. Mathematically, for a simple Susceptible-Infected-Recovered model,  
70 these variables are linked through a simple formula which in this work [5]:

$$R_0 p^* + \log(1 - p^*) = 0 \quad (1)$$

71 Additional details about the methods are in Supplementary Text S1 and the scripts used can be accessed at  
72 [https://gitlab.in2p3.fr/ete/residual\\_epidemic](https://gitlab.in2p3.fr/ete/residual_epidemic).




Fig. 1: COVimpact user window.

### 73 3 Results

74 We first present the dedicated software for the French epidemic, which can be accessed at <https://cloudapps.france-bioinformatique.fr/covimpact/>, before focusing on a scenario for the French epidemic.

#### 76 3.1 COVimpact interface

77 The software contains a setting panel, which is divided into two sub-panels. The first subpanel allows the user  
78 to choose the French geographical scale (department, region, country) to use for the inputs and outputs. The  
79 user can also design the level of protection conferred by natural immunity against severe cases. The motivation  
80 for this is that some variants, especially  $\beta$  and  $\gamma$  could be evading natural immunity [3].

81 The second subpanel revolves around the vaccination settings. The percentage of the population vaccinated  
82 is set separately for three age groups: children from 12 to 17 years old, adults from 18 to 65, and adults above

65. These classes were chosen to reflect differences in infection fatality ratio (IFR, [8]) but also implementation  
of vaccination strategies. The vaccine efficacy, i.e. the percentage of decrease in severe infections amongst  
vaccinated people, is set manually and is assumed not to vary with age.

COVimpact allows the user to implement several vaccines in the population. Each vaccine has its own  
efficacy and its own coverage in the three age classes. Note that individuals are assumed to be vaccinated with  
only one kind of vaccine, i.e. the sum of the percentage of vaccinated individuals in an age class cannot be  
greater than 100%.

COVimpact displays 5 types of outputs following the settings provided by the user and an estimation of  
the proportion of the population already infected, i.e. with natural immunity (see the Methods). The interface  
is shown in Figure 2. The number of ICU admissions and deaths associated with the epidemic wave (shown  
in the top panel as a function of the epidemic wave size) are stratified by age in the middle panels. The bottom  
panel shows the age-stratification of the inferred prevalence of natural immunity. In all the figures, the confi-  
dence intervals shown are solely derived from the confidence intervals associated with the estimation of the  
percentage of the population already infected.

### 3.2 France as a test case

To illustrate the implications of the insights provided by COVimpact, we investigate the number of ICU ad-  
missions that an epidemic wave affecting 25% of the population would cause in France. The associated repro-  
duction number is 1.15, which is consistent with observed dynamics in France [9] and the fact that vaccination  
only partly limits reinfection and transmission from reinfected hosts [7].

Even with 95% of the adults have a full vaccination scheme leading to an 85% protection against severe  
symptoms of both vaccine and natural immunity [7], such a wave would cause approximately 20,000 ICU ad-  
missions. Since the median duration of a stay in ICU is 14 days, this corresponds to 280,000 days of ICU bed  
occupancy. Even if the epidemic wave lasted 2 months, this would mean more than 4,600 beds dedicated to  
COVID-19 patients, i.e. more than half of the ICU capacity, with a peak after a month. Unfortunately, vaccina-  
tion coverage is currently much lower than 95% in France.

## 4 Discussion

Forecasting epidemic trends more than 4 weeks in advance is essential, e.g. to hire or train qualified personnel.  
Unfortunately, variations in public health policies make mathematical models unreliable to anticipate long-




Fig. 2: Total number of ICU admissions (in blue) and death (in black) as a function of epidemic wave size in France in 2021. For this figure, we assume a 95% vaccine coverage in adults with an 85% efficiency against severe forms. On the top x-axis, using equation 1, we show the reproduction number associated with the amplitude of the epidemic wave. Confidence intervals on the figure originate from the uncertainty in the estimation of the number of hosts with natural immunity.

111 term trends [9]. We show that valuable insights can be gained in absence of a detailed description of the  
112 epidemic dynamics. More precisely, assuming that the size of the epidemic wave and that the vaccine coverage  
113 and efficiency are known, it is possible to estimate the total number of ICU admissions and deaths caused by  
114 the wave.

115 The semi-quantitative results we obtain are particularly useful for public health agencies. Indeed, they  
116 allow one to explore extreme scenarios to make informed decisions. For instance, we show that even with  
117 extremely high vaccination coverage, and although vaccination strongly decreases the pressure on the health  
118 care system, a country like France is likely to experience major strain in ICU occupancy in the event of a large  
119 epidemic wave (25% of the population infected in 2 months). To maximise the popularisation of this approach,  
120 we also introduce COVIDici, a software allowing one to visualise the effects of an epidemic wave at the French  
121 departmental, regional, and national level.

122 Importantly, our approach makes a series of simplifying assumptions that need to be carefully weighed.  
123 First, individuals are assumed to be vaccinated independently of their natural immunisation status. Further-  
124 more, the infection fatality ratio is kept constant as in [8] but this is known to depend on hospital capacity  
125 strain [6] or the infecting variant [1]. Second, there were also some missing data at the departmental level,  
126 e.g. the age-stratified proportion of infected individuals. These were obtained by adjusting the regional data

127 to the departmental population. Third, regarding the type of immunity, we assume that the protection, quanti-  
128 fied as the reduction of severe disease probability, in a vaccinated individual with natural immunity was equal  
129 to the rate of the most effective protection. Finally, infected individuals are assumed to be randomly selected  
130 in the population, regardless of their level of protection. In other words, the protections (conferred by a past  
131 infection, a vaccine or both) only offer a reduction of severe forms but do not affect the risk of being exposed.

132 There are several ways in which this work could be extended. We focused here on France but these in-  
133 sights could be equally valuable for other countries. The main difficulty, which we solved by leaning on earlier  
134 work by Hozé et al. [4], is to estimate the proportion of individuals already infected per age class before the  
135 epidemic wave. However, the importance of this assumption decreases as the protection conferred by natural  
136 immunity decreases. Another extension, more specific to our software, would be to improve the calculations  
137 to implement the method by Hozé et al. [4] to re-estimate these proportions but, for France, this requires access  
138 to restricted data.

139 Overall, analysing the consequence of epidemic waves of given sizes as a function of vaccination coverage  
140 can greatly help shape public health policies, e.g. to optimise the number of beds in ICU in a region or to  
141 estimate the need to hire or train qualified personnel.

142 **Acknowledgements** We thank the ETE team for discussion and the Institut Français de Bioinformatique (IFB) for hosting the software.  
143 This work was supported by the PhyEpi grant from the Région Occitanie and the ANR.

#### 144 **Conflict of interest**

145 The authors declare that they have no conflict of interest.

#### 146 **References**

- 147 1. Challen R, Brooks-Pollock E, Read JM, Dyson L, Tsaneva-Atanasova K, Danon L (2021) Risk of mortality  
148 in patients infected with SARS-CoV-2 variant of concern 202012/1: matched cohort study. *BMJ* 372:n579,  
149 DOI 10.1136/bmj.n579
- 150 2. Hall VJ, Foulkes S, Charlett A, Atti A, Monk EJM, Simmons R, et al (2021) SARS-CoV-2 infection rates of  
151 antibody-positive compared with antibody-negative health-care workers in England: a large, multicentre,  
152 prospective cohort study (SIREN). *The Lancet* 397(10283):1459–1469, DOI 10.1016/S0140-6736(21)00675-9
- 153 3. Hoffmann M, Arora P, Groß R, Seidel A, Hörnich BF, Hahn AS, Krüger N, Graichen L, Hofmann-Winkler  
154 H, Kempf A, Winkler MS, Schulz S, Jäck HM, Jahrsdörfer B, Schrezenmeier H, Müller M, Kleger A, Münch

- 155 J, Pöhlmann S (2021) SARS-CoV-2 variants B.1.351 and P.1 escape from neutralizing antibodies. Cell DOI  
156 10.1016/j.cell.2021.03.036
- 157 4. Hozé N, Paireau J, Lapidus N, Kiem CT, Salje H, Severi G, Touvier M, Zins M, Lamballerie Xd, Lévy-Bruhl  
158 D, Carrat F, Cauchemez S (2021) Monitoring the proportion of the population infected by SARS-CoV-  
159 2 using age-stratified hospitalisation and serological data: a modelling study. *The Lancet Public Health*  
160 6(6):e408–e415, DOI 10.1016/S2468-2667(21)00064-5
- 161 5. Kermack WO, McKendrick AG (1927) A contribution to the mathematical theory of epidemics. *Proc R Soc*  
162 *Lond A* 115:700–721, DOI 10.1098/rspa.1927.0118
- 163 6. Lefrancq N, Paireau J, Hozé N, Courtejoie N, Yazdanpanah Y, Bouadma L, Boëlle PY, Chereau F, Salje  
164 H, Cauchemez S (2021) Evolution of outcomes for patients hospitalised during the first 9 months of the  
165 SARS-CoV-2 pandemic in France: A retrospective national surveillance data analysis. *The Lancet Regional*  
166 *Health - Europe* 5:100087, DOI 10.1016/j.lanep.2021.100087
- 167 7. Puranik A, Lenehan PJ, Silvert E, Niesen MJM, Corchado-Garcia J, O'Horo JC, Virk A, Swift MD, Halamka  
168 J, Badley AD, Venkatakrisnan AJ, Soundararajan V (2021) Comparison of two highly-effective mRNA vac-  
169 cines for COVID-19 during periods of Alpha and Delta variant prevalence. medRxiv p 2021.08.06.21261707,  
170 DOI 10.1101/2021.08.06.21261707
- 171 8. Salje H, Kiem CT, Lefrancq N, Courtejoie N, Bosetti P, Paireau J, Andronico A, Hozé N, Richet J, Dubost  
172 CL, Strat YL, Lessler J, Levy-Bruhl D, Fontanet A, Opatowski L, Boelle PY, Cauchemez S (2020) Estimating  
173 the burden of SARS-CoV-2 in France. *Science* 369(Issue 6500):208–211, DOI 10.1126/science.abc3517
- 174 9. Sofonea MT, Alizon S (2021) Anticipating COVID-19 intensive care unit capacity strain: a look back at  
175 epidemiological projections in France. *Anaesthesia Critical Care & Pain Medicine* in press
- 176 10. Sofonea MT, Reyné B, Elie B, Djidjou-Demasse R, Selinger C, Michalakis Y, Alizon S (2021) Memory is  
177 key in capturing COVID-19 epidemiological dynamics. *Epidemics* p 100459, DOI 10.1016/j.epidem.2021.  
178 100459