EFFECT OF EARLY TREATMENT WITH FLUVOXAMINE ON RISK OF EMERGENCY CARE AND HOSPITALIZATION AMONG PATIENTS WITH COVID-19: THE TOGETHER RANDOMIZED PLATFORM CLINICAL TRIAL

Authors: Gilmar Reis, MD[1,2]; Eduardo Augusto dos Santos Moreira Silva, MD[1,2]; Daniela Carla Medeiros Silva, MD[1,2]; Professor Lehana Thabane, PhD[3]; Aline Cruz Milagres, RN[4,5]; Thiago Santiago Ferreira, MD[1]; Castilho Vitor Quirino dos Santos[1,2]; Adhemar Dias de Figueiredo Neto, MD[6]; Eduardo Diniz Callegari, MD[7]; Leonardo Cançado Monteiro Savassi, MD[2]; Vitoria Helena de Souza Campos,[1,2] Maria Izabel Campos Simplicio, BScPharm[1]; Luciene Barra Ribeiro, RN[1]; Rosemary Oliveira[1]; Ofir Harari, PhD[4]; Jamie I Forrest, MPH[4]; Hinda Ruton, MSc[4]; Sheila Sprague, PhD[3]; Paula McKay, MSc[3]; Alla V Glushchenko, MD,[3] PhD, Craig R. Rayner, PharmD, FRCP *Edin*[10,11]; Professor Eric J. Lenze, MD;[12] Angela M. Reiersen, MD[12]; Professor Gordon H. Guyatt, MD[3]; Professor Edward J. Mills, PhD, FRCP[3]; for the TOGETHER Investigators*

*TOGETHER Investigators are listed in the supplementary materials

Affiliations:

1. Research Division, Cardresearch - Cardiologia Assistencial e de Pesquisa, Brazil

2. Department of Medicine, Pontifícia Universidade Católica de Minas Gerais, Brazil

3. Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada

4. Cytel Inc, Vancouver, British Columbia, Canada

5. Department of Public Health, Montes Claros State University, Brazil

6. Public Health Fellowship Program, Governador Valadares Public Health Authority, Brazil

7. Public Health, Mental and Family Medicine Department, Ouro Preto Federal University, Brazil

8. Public Health Care Division, City of Ibirité, Brazil

9. Drug Research and Development Center, Federal University of Minas Gerais, Belo Horizonte, Brazil

10. Certara Inc. Princeton, New Jersey, USA

11. Monash Institute of Pharmaceutical Sciences, Monash University, Melbourne, Australia

12. Department of Psychiatry, Washington University School of Medicine, St. Louis, MO, USA

1	**Corresponding Authors**:	**Gilmar Reis MD, PhD**
2		Director - Research Division
3		CARDRESEARCH – Cardiologia Assistencial e de
4		Pesquisa
5		Associate Professor of Medicine
6		Pontificia Universidade Católica de Minas Gerais
7	Address:	Rua Domingos Vieira 300, Sala 606 – Santa Efigenia
8		Belo Horizonte – Minas Gerais
9		Brazil – ZIP: 30.150-242
10	Email:	greis@cardresearch.org
11		
12		**Edward Mills PhD, FRCP**
13		Professor
14		Health Research Methods, Evidence & Impact
15		McMaster University
16		Hamilton, Ontario, Canada
17	Email:	millsej@mcmaster.ca
18	Phone:	778 317 8530

ABSTRACT

Background

Recent evidence indicates a potential therapeutic role of fluvoxamine for COVID-19. In the

TOGETHER randomized platform clinical trial for acutely symptomatic patients with COVID-19,

we assessed the efficacy of fluvoxamine vs. placebo in preventing either extended emergency

room observation or hospitalization due to COVID-19. Herein, we report the preliminary findings.

Methods

This placebo-controlled, randomized, adaptive, platform trial conducted among symptomatic

Brazilian adults confirmed positive for SARS-CoV-2 included eligible patients with a known risk

factor for progression to severe disease. Patients were randomly assigned to either fluvoxamine

(100 mg twice daily for 10 days) or placebo. The primary endpoint was a composite outcome of

emergency room observation for >6 hours or hospitalization from COVID-19 up to 28 days post

randomization. Secondary outcomes included viral clearance at day 7, time to hospitalization,

mortality, and adverse drug reactions. We used a Bayesian analytic framework to determine effects

along with probability of success of intervention compared to placebo. The trial is registered at

clinicaltrials.gov (NCT04727424) and is ongoing.

Results

The study team screened 9020 potential participants for this trial. The trial was initiated on June

2, 2020, with the current protocol reporting randomization from January 15, 2021 to August 6[th]

2021, when the trial arms were stopped for superiority. A total of 3238 patients were allocated to

fluvoxamine (n=739), placebo (n=733) and other treatments (n=1766). Herein, we report the

effectiveness of fluvoxamine vs. a concurrent placebo control. The average age of participants was

50 years (range 18-102 years); 57% were female. The proportion of patients observed in an emergency room for >6 hours or admitted to hospital due to COVID-19 was lower for the fluvoxamine group compared to placebo (77/739 vs 108/733; Relative Risk [RR]: 0.71; 95% Bayesian Credible Interval [95% BCI]: 0.54 - 0.93), with a probability of superiority of 99.4% surpassing the prespecified superiority threshold of 97.6% (risk difference 4.3%). Of the composite primary outcome events, 88% were hospitalizations. We found no significant relative effects between the fluvoxamine and placebo groups on viral clearance at day 7 (Odds ratio [OR]: 0.75; 95% Confidence Intervals [95% CI]: 0.53 - 1.07), mortality (OR: 0.70; 95% CI: 0.36 - 1.30), time to death (Hazard ratio [HR]: 0.79; 95% CI: 0.58 - 1.08), days hospitalized (Mean Difference (MD) 1.22 days; 95% CI: 0.98 - 1.53), number of days ventilated (MD 1.10; 95% CI: 0.70 - 1.73) or other secondary outcomes. Data capturing all 28 days of follow-up will be reported after August 26[th], 2021.

Conclusions

Treatment with fluvoxamine (100 mg twice daily for 10 days) among high-risk outpatients with early diagnosed COVID-19, reduced the need for extended emergency room observation or hospitalization.

Funding The trial was supported by FastGrants and The Rainwater Foundation.

BACKGROUND

Although safe and effective vaccines for COVID-19 have been developed and distributed, there

remain, particularly in low resource settings, major challenges regarding their production,

allocation, and affordability.[1] Identifying inexpensive, widely available and effective therapies

against COVID-19 is, therefore, of great importance. In particular, repurposing existing medicines

that are widely available and with well understood safety profiles, has particular appeal.[2]

Fluvoxamine is a selective serotonin reuptake inhibitor (SSRI) and a Sigma-1 receptor

(S1R) agonist.[3] There are several potential mechanisms for fluvoxamine in treatment of COVID-

19 illness, including anti-inflammatory and possible antiviral effects.[4] A small placebo-controlled

randomized trial has raised the possibility that fluvoxamine may reduce the risk of clinical

deterioration in outpatients with COVID-19, suggesting the need for larger randomized, placebo-

controlled studies.[5,6]

To evaluate the efficacy of fluvoxamine to prevent progression of COVID-19 and

hospitalization among outpatients with laboratory-documented SARS-CoV-2, we conducted a

randomized, placebo-controlled adaptive platform trial in Minas Gerais, Brazil. This flexible

platform trial design allows for additional agents to be added and tested with standardized

operating procedures outlined in a single overarching master protocol.[7,8] Among eight different

interventions evaluated in this platform trial, we report here on the clinical evaluation of

fluvoxamine using a concurrent placebo control group.

METHODS

Study design and oversight

The TOGETHER Trial is a randomized adaptive platform trial to investigate the efficacy of

repurposed treatments for COVID-19 disease among high-risk adult outpatients.[9] The trial was

designed and conducted in partnership with local public health authorities from eleven

participating cities in Brazil to simultaneously test potential treatments for early disease using a

master protocol. A master protocol defines prospective decision criteria for discontinuing

interventions for futility, stopping due to superiority against placebo, or adding new interventions.

Interventions evaluated in the TOGETHER trial, thus far, include, hydroxychloroquine (protocol

1), lopinavir/ritonavir (protocol 1),[10] metformin, ivermectin, fluvoxamine, doxasozin and

pegylated interferon lambda versus matching placebos (protocol 2).

The trial began on June 2, 2020 and enrollment into the fluvoxamine arm began on January

15th 2021. The trial, which complied with the International Conference of Harmonization – Good

Clinical Practices, as well as local regulatory requirements, was approved for research ethics by

local and national ethics boards in Brazil (CONEP CAAE: 41174620.0.1001.5120, approval letter

5.501.284) and the Hamilton Integrated Research Ethics Board (HiREB, approval letter 13390) in

Canada. The full protocol, statistical analysis plan, and additional details are appended in the web-

appendix. The adaptive designs CONSORT extension (ACE) statement guided this trial report.[16,17]

The steering committee made all protocol-related decisions and sponsors had no role in trial

conduct, data analysis or decision to submit manuscript for publication. An independent Data

Safety Monitoring Committee (DSMC) provided trial oversight.

Setting

The appended web-appendix lists the cities and investigators of the eleven clinical sites in Brazil that participated in the trial. Local investigators, in partnership with local public health authorities, recruited participants at community health facilities (emergency settings, flu-symptom referral centers, primary care community centers). We used several community outreach strategies including physical and social media as per local public health authorities, in order to create awareness of the trial.

Participant Screening

Upon presentation to one of the trial outpatient care clinics, local investigators screened potential participants to identify those who met the eligibility criteria.

The key inclusion criteria were: 1) patients over the age of 18, 2) presenting to an outpatient care setting with an acute clinical condition consistent with COVID-19 and symptoms beginning within 7 days of the screening date, 3) positive rapid test for SARS-CoV-2 antigen performed at the time of screening or patient with positive SARS-CoV-2 diagnostic test within 7 days of symptom onset, 4) at least one additional criterion for high-risk: diabetes mellitus; systemic arterial hypertension requiring at least one oral medication for treatment; known cardiovascular diseases (heart failure, congenital heart disease, valve disease, coronary artery disease, cardiomyopathies being treated, clinically manifested heart disease and with clinical repercussion); symptomatic lung disease and / or being treated (emphysema, fibrosing diseases); symptomatic asthma patients requiring chronic use of agents to control symptoms; smoking; obesity, defined as BMI>30 kg / m² (weight and height information provided by the patient); transplant patients; patient with stage IV chronic kidney disease or on dialysis; immunosuppressed patients / using corticosteroid therapy (equivalent to at least 10 mg of prednisone per day) and / or

immunosuppressive therapy; patients with a history of cancer in the last 0.5 years or undergoing

current cancer treatment, or age $\geq$ 50 years; and unvaccinated status.

Patients who met any of the following criteria were excluded from the trial: 1) Diagnostic

examination for SARS-CoV2 negative associated with acute flu-like symptoms (patient with

negative test taken early and becoming positive a few days later were eligible, if he/she was <7

days after the onset of flu-like symptoms); 2) Acute respiratory condition compatible with COVID-

19 treated in the primary care and previously requiring hospitalization; 3) Acute respiratory

condition due to other causes; 4) Received vaccination for SARS-CoV2; 5) Dyspnea secondary to

other acute and chronic respiratory causes or infections (e.g., decompensated COPD, acute

bronchitis, pneumonia, primary pulmonary arterial hypertension); 6) Current use of selective

serotonin reuptake inhibitors; uncontrolled psychiatric disorders; or suicidal ideation; 7) Inability

or unwillingness to follow research guidelines and procedures. A full list of exclusion criteria are

provided in the trial protocol.

If a patient met the above eligibility criteria, study personnel obtained written informed

consent. After obtaining informed consent a rapid antigen test for COVID-19 (Panbio ®, Abbott

laboratories) and a pregnancy test for women of childbearing age were performed. If the COVID-

19 test was negative or if the pregnancy test was positive, the participant was not included in the

trial. After informed consent, study personnel collected the following data prior to randomization:

demographics, medical history concomitant medications, co-morbidities, exposure to Index Case

information, WHO clinical worsening scale, and the PROMIS Global Health Scale.

Randomization and Trial Interventions

Participants were randomized using a centralized core randomization process handled by an

independent unblinded pharmacist who was not aware of any protocol-related procedures and

contracted specifically for this process. Sites requested randomization by text message to the

pharmacist at the coordinating center. This maintained concealment of allocation. Patients were

randomly assigned using a block randomization procedure for each participating site, stratified by

age (< 50 years /≥ 50 years). The trial team, site staff and patients were blinded to treatment

allocation. The active drugs and the placebo pills were packaged in identically shaped bottles and

labeled with alphabet letters corresponding to the active arm or placebo arm. Only the third-party

pharmacist responsible for releasing the randomization was aware of which letter was associated

with which drug and/or placebo. As this is a multi-arm trial and all active interventions have a

matching inert placebo, the matching placebo represents the proportion of the control group for

the number of arms in the trial at any given time.

Data Collection and Participant Follow-Up

Our primary outcome is a composite that includes emergency room visits due to the clinical

worsening of COVID-19 (defined as participant remaining under observation for > 6 hours) or

hospitalization due to the progression of COVID-19 (worsening of clinical status) and/or suspected

COVID-19 complications within 28 days of randomization. Key secondary outcomes include: 1)

viral clearance, 2) time to clinical improvement, 3) number of days with respiratory symptoms, 4)

time to hospitalization for any cause or due to COVID-19 progression, 5) all-cause mortality and

time to death from any causes, 7) WHO clinical worsening scale score, 8) days in hospital and on

ventilator and 9) adverse events, adverse reactions to the study medications and the proportion of

participants who are non-adherent with the study drugs. All secondary outcomes are assessed up

to 28 days following randomization.

Study personnel collected, via in person, telephone contact and social media applications

using video-teleconferencing, outcome data on days 1, 2, 3, 4, 5, 7, 10, 14, and 28. We collected

outcome data irrespective of whether participants took study medication. In case of adverse events,

unscheduled visits (during the treatment period) outside of clinical care could occur at any time.

Considering the transmissible characteristic of SARS-CoV-2 and the isolation

recommendations of positive individuals, we collected limited vital sign data. Cardiac safety was

assessed using a 6-lead ECG (Kardiamobile, Mountain View, CA) at the baseline visit. The digital

recordings were de-identified and transferred to a central facility (Cardresearch, Belo Horizonte,

Brazil) for reading. Oxygen status was assessed using a pulse oximeter for non-invasive arterial

oxygen saturation (SpO2) and pulse (Jumper Medical Equipment, Shenzhen, China), and

temperature using a standard digital oral thermometer administered by research personnel. Mid-

turbinate nasal swab kits and sterile recipient storage were provided for collection of

nasopharyngeal swab or sputum/saliva. They were performed by the first quarter of participants

enrolled in the trial at days 3 and 7. PCR of viral clearance was assessed to determine if active

drugs demonstrate any anti-viral effects.

All serious and non-serious adverse events were reported to study personnel as per local

regulatory requirements. Reportable adverse events included serious adverse events, adverse

events resulting in study medication discontinuation, and adverse events assessed as possibly

related to study medication.

Trial interventions

All participants received usual standard care for COVID-19 provided by healthcare professionals

workers at public health facilities. Patients were randomized to fluvoxamine (Luvox ®, Abbott) at

a dose of 100 mg twice a day for 10 days or corresponding placebo starting right after

randomization (day 1). Research personnel provided participants a welcome video with

information on trial, study drug, adverse events and follow-up procedures. Clinicians providing

usual care in public health facilities typically focus on the management of symptoms and with

antipyretics, and recommend antibiotics only if they suspect bacterial pneumonia.

Statistical Analyses

The Adaptive Design Protocol and the Master Statistical Analysis Plan provide details of sample

size calculation and statistical analysis (appended). This trial is adaptive and applies sample size

re-estimation approaches. To plan for each arm, we assumed a minimum clinical utility of 37.5%

(relative risk reduction) to achieve 80% power with 0.05 two-sided Type 1 error for a pairwise

comparison against the placebo (talc) assuming a control event rate (CER) of 15%. This resulted

in an initial plan to recruit 681 participants per arm. The statistical team conducted planned interim

analyses. Stopping thresholds for futility were established if the posterior probability of superiority

was less than 40% at interim analysis. An arm can be stopped for superiority if the posterior

probability of superiority meets the threshold of 97.6%

Baseline characteristics are reported as count (percent) or median and interquartile range (IQR)for

continuous variables. We applied a Bayesian framework for our primary outcome analysis and a

frequentist approach for all sensitivity analyses and secondary outcomes. Posterior efficacy of

fluvoxamine for the primary outcome is calculated using the beta-binomial model for event rates,

assuming informed priors based on the observational data for both placebo and fluvoxamine, for

both Intention-to-Treat (ITT), modified ITT (defined as receiving the study drug for at least 24

hours before an event) and Per-Protocol (PP) analyses (defined as taking >80% of possible doses).

Modified ITT was defined as receiving treatment for at least 24 hours before a primary outcome.

We accounted for any temporal changes in events rates by using only the concurrent randomised

population. We assessed subgroup effects according to the preplanned statistical analysis plan. We

calculated the number needed to treat.

Secondary outcomes were assessed using a pre-specified frequentist approach. For viral

clearance we fitted a longitudinal, mixed-effect logistic regression model with a treatment and time

interaction term for binary patient outcomes (Covid-19 positive/negative) reported on day 3 and 7

from randomization, with subject random effect. We assessed time-to-event outcomes using Cox

proportional hazard models and binary outcomes using logistic regression. Per protocol analyses

were considered sensitivity analyses to assess the robustness of the results. All analyses were

performed using R version 4.0.3. Full details of the Statistical Analysis Plan (SAP) are appended.

Data and Safety Monitoring Committee

A Data and Safety Monitoring Committee provided independent oversight for this trial. We

planned a 4th interim analysis of the fluvoxamine arm with data collected up to August 2nd, 2021.

Role of the funding source

The funders had no role in the study design, data collection, analysis, interpretation or writing, or

decision to submit for publication. The executive committee take responsibility for the integrity of

the data and the accuracy of the data analysis. The trial executive committee oversaw all aspects

of trial conduct, completeness, data accuracy and adherence of trial conduct to the protocol and

the committee vouch for the accuracy and completeness of the data and for fidelity to the protocol.

RESULTS

We screened 9020 potential participants for inclusion in this trial to date. The trial enrolled its first

participant on June 2, 2020 and enrolment into the fluvoxamine arm began on January 20, 2021.

By August 6, 2021, 1,472 recruited participants were randomized to fluvoxamine (n= 739) or the

concurrent placebo (n=733), and 1,766 were randomized to other treatment arms (Figure 1). The

median age was 50 years (range 18-102) and 846 (57.5%) were women (Table 1). Most

participants self-identified as mixed-race 1,403 (95.3%), 11 (0.7%) as white, 10 (0.7%) as black

or African American, the rest self-identified as unknown 48 (3.3%). As the trial is ongoing, herein

we provide descriptive summaries of only those randomized to fluvoxamine and its concurrent

control. With respect to covariates of age, BMI, and co-morbidities, the groups were generally

well balanced (Table 1). The mean number of days with symptoms prior to randomization was 4

days (Standard Deviation 1.76). All patients were identified at the end of the study for inclusion

in the analysis.

Primary Outcomes

The Data Safety Monitoring Committee met four times since the protocol initiation and last met

on August 5, 2021, recommending that the TOGETHER Trial stop randomizing patients to the

fluvoxamine arm, as this comparison had met the pre-specified superiority criterion for the primary

endpoint (prespecified superiority threshold 97.6%). Based on the Bayesian beta-binomial model,

there was evidence of a benefit of fluvoxamine reducing hospitalization or observation in an

emergency room for greater than six hours due to COVID-19 (Relative Risk [RR]: 0.71; 95%

Bayesian Credible Interval [BCI]: 0.54 - 0.93) in the Intention-to-Treat (ITT) population (figure

2A) and (RR: 0.68; 95% BCI: 0.50 - 0.91) in a modified ITT population (figure 2B). The

probability that the event rate was lower in the fluvoxamine group compared to placebo was 99.4%

for the ITT population, and 99.6% for the modified ITT population (Figure 2A/B). These posterior

efficacy numbers were higher than the pre-specified 97.6% threshold set for the fourth interim

analysis. In the fluvoxamine group 77 (10.4%) participants experienced a primary outcome event

compared to 108 (14.7%) in the placebo group (Table 2). Most events (88%) were hospitalizations.

The number needed to treat is 24. Per protocol analysis demonstrated a larger treatment effect

(0.34, 95% BCI, 0.20-0.54).

Secondary Outcomes

Table 3 presents findings from secondary outcome analyses. There were no significant differences

between fluvoxamine and placebo for viral clearance at Day 7 (p=0.18) and eFigure3 in web-

appendix, hospitalizations due to COVID (p=0.17), all-cause hospitalizations (p=0.12), time to

hospitalization (p=0.14, number of days in hospital (p=0.07), mortality (p=0.26), time to death

(p=0.26), number of days on mechanical ventilation (p=0.67), time to recovery (p=0.86) or the

PROMIS Global Health Scale (p=0.44). With respect to adverse events, there were significantly

greater number of Grade 1 (mild) treatment emergent adverse events (TEAE) among patients in

the fluvoxamine arm (p<0.01). However, no differences between fluvoxamine and placebo were

observed for TEAEs of Grades 2, 3, 4, or 5.

Sub-group Analyses

In the prespecified subgroup analysis, we found no evidence of moderation of treatment effect for

fluvoxamine compared to placebo, for sub-groups of age, sex, days since symptom onset, or co-

morbidities (Figure 3 and web-appendix eTable 1).

DISCUSSION

This is the first large randomized controlled trial to test the efficacy of fluvoxamine for acute

treatment of COVID-19. We found a clinically important absolute risk reduction of 4.3%, and

29% relative risk reduction, on the primary outcome of retention in an emergency setting for

COVID-19 disease observation or hospitalization, consequent on the administration of

fluvoxamine for 10 days. This study is only the 2nd study to demonstrate an important treatment

benefit for a repurposed drug in the early treatment population.[11] Our findings represent the latest

interim analysis of the trial resulting in the DSMC recommending stopping the active fluvoxamine

arm. The final analysis from the trial, wherein all patients have contributed 28 days of follow-up

data, will be made available 28 days after the last randomized patient has completed this period

(August 25, 2021). Given fluvoxamine's safety, tolerability, ease of use, low cost, and widespread

availability, these findings may have influence on national and international guidelines on the

clinical management of COVID-19.

Comparison with prior evidence

Our results are consistent with an earlier smaller trial conducted in the United States (led by EJL

and AMR).[6] That study used a higher dose of fluvoxamine (100mg tid for 15 days) and included

a lower risk group for the primary outcome but found no clinical deterioration among 80 patients

receiving fluvoxamine vs. 6 of 72 patients receiving placebo. A large observational study from

France involved a different population, 7230 hospitalized COVID-19 patients, and reported a

reduction in use of intubation or death with use of SSRIs.[5]

The underlying mechanism of fluvoxamine for COVID-19 disease remains uncertain. Although

hypotheses include several potential mechanisms,[4] the main reason for the initial study of

fluvoxamine as a treatment of COVID-19 was its anti-inflammatory action through activation of

the σ-1 receptor (S1R).[12] S1R is an endoplasmic reticulum (ER) chaperone membrane protein

involved in many cellular functions,[13] including regulation of ER stress response/unfolded protein

response and regulation of cytokine production in response to inflammatory triggers.[14] In the

presence of fluvoxamine, S1R may prevent the ER stress sensor Inositol-Requiring Enzyme 1α

(IRE1) from splicing and activating the mRNA of X-Box Protein 1 (XBP1), a key regulator of

cytokine production including IL-6, IL-8, IL-1β and IL-12. In a 2019 study by Rosen and

colleagues, fluvoxamine showed benefit in preclinical models of inflammation and sepsis through

this mechanism.[14]

Another mechanism may be fluvoxamine's anti-platelet activity.[15] SSRIs can prevent loading of

serotonin into platelets and inhibit platelet activation, that may reduce the risk of thrombosis, and

these antiplatelet effects can be cardioprotective. In vitro and animal studies are needed to help

clarify the most likely mechanism(s). Biomarker studies included as part of future RCTs may also

help to clarify mechanisms.

Strengths and limitations

Since the start of the COVID-19 pandemic, there have been more than 2800 RCTs registered on

clinicaltrials.gov. However, less than 300 have been reported and the vast majority of clinical trials

have been small and underpowered, with sample sizes less than 100. In many cases, these trials

have been unsuccessful at recruiting as the local epidemics occur in waves and sustainable

infrastructure to maintain staff or local interest for recruitment is lacking. The trials that provide

the clearest medical understanding tend to be the larger platform trials, such as SOLIDARITY,[16]

RECOVERY,[17] PRINCIPLE,[11] and REMAP-CAP.[18] As a result, we actively collaborate with other

investigators running trials with overlapping interventions so that they can be aware of our study

decisions and determine whether they should influence their respective trials.

Major strengths include the rapid recruitment and enrolment of high-risk patients for the

development of severe COVID-19. Our recruitment strategy involves the engagement with the

local public health system, thus allowing recruitment that frequently exceeds twenty patients per

day. We enrolled only participants with diagnosed COVID-19 and less than 7 days of symptom

onset using a commercially available COVID-19 AG rapid test (Panbio ®). The concordance of

COVID-19 positive tests with RT-PCR was evaluated on the group of participants with PCR

evaluations and found a concordance rate of > 99% on both tests collected at baseline.

Our understanding of the epidemiology of COVID-19 as well as its disease progression and

outcomes have evolved since beginning this platform trial in June 2020. Early studies assessed the

effects of interventions of viral load and clearance, while later studies also evaluate more clinical

outcomes. We made adjustments to the trial according to prespecified rules and in communication

with the appropriate ethics review committees that allowed us to respond to the epidemic waves

while maintaining high rates of recruitment. Unlike many outpatient clinical trials, our study

involves direct patient contact through the use of medical students, nurses and physicians who do

at-home visits as well as follow-up via telecommunications. Given the rapid recruitment of patients

in combination with the high event rate of emergency room visits and hospitalizations, we were

able to evaluate the effects of interventions when portions of the planned population had been

recruited. The period of time between first recruitment of a patient on fluvoxamine and the data

cut for our trial was 197 days. Our trial assessed a primary outcome as a binary event having

occurred by 28 days post-randomization.

One of the major limitations of our fluvoxamine trial is primarily related to the challenges of

conducting a trial in a disease that is not well characterized. Currently, there is no standard of care

that exists for early treatment of COVID-19 and various advocacy groups promote different

interventions, including some of those evaluated in this and our previous trials. Furthermore, there

is little understanding of who is at greatest risk of disease progression from this disease as some

patients with numerous risk factors do recover quickly while some others with less established risk

factors may not. Our population had a higher rate of hospitalization events than observed in most

clinical trials, thus permitting inferences on treatment effects in this higher-risk population.

Implications

Our trial has found that fluvoxamine, an inexpensive existing drug, reduces the need for advanced

disease care in this high-risk population. A ten-day course of fluvoxamine costs approximately $4

even in well-resourced settings.[19] Our study compares favorably and exceeds the treatment effects

of more expensive treatments including monoclonal antibodies for outpatient treatment.[20,21] The

absolute number of serious adverse events associated with fluvoxamine was lower than our

placebo group and this might reflect the modulatory effect of fluvoxamine on systemic

inflammation in these participants. Lower respiratory tract infections were reported less frequently

in patients in the fluvoxamine group than those in the placebo group. This is concordant with the

reduction of hospital admissions in patients with confirmed COVID-19 treated with fluvoxamine,

and the numerically lower number of patients requiring mechanical ventilation.

Fluvoxamine is widely available but is not on the WHO Essential Medicines List,[22] whereas a

closely related SSRI fluoxetine is on the list. It is now of critical importance to determine whether

a class-effect exists and whether these drugs can be used interchangeably for COVID-19. The

recent important findings that inhaled budesonide increased time to recovery among a similar

population as our trial and had a trend towards decreased hospitalizations suggests this as an

alternative or additional intervention for outpatient care that should be evaluated. The PRINCIPLE

trial evaluated time to recovery using self-reported recovery up to 28 days after randomization to

budesonide.[11] Our trial differed as we evaluated improvement in the WHO categorization of

disease disability up to days 14 and then 28 (web-appendix eFigure 2). Finally, our study was

among unvaccinated patients. Further evidence of treatment benefits are needed to determine the

effect of fluvoxamine among vaccinated populations.

Use of interventions, including fluvoxamine, to prevent progression of illness and hospitalization

is critically dependent on identifying higher risk individuals. Unselected populations will have a

lower risk. What absolute reduction in risk of clinical deterioration would motivate patients to

choose treatment (probably the >4% we observed, but perhaps not much lower) remains uncertain.

These considerations raise the importance of the development of a validated prediction rule for

deterioration in patients in the early stages of COVID-19 infection.

Conclusion

Administration of fluvoxamine reduced the rate of prolonged observation in an emergency care

setting or hospitalization due to COVID-19 in people with a high risk of serious disease.

ACKNOWLEGEMENTS

The trial was supported by FastGrants and the Rainwater Foundation. GR and EJM had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Our research network consists of partnerships between academics and clinicians at McMaster University in Ontario, Canada, and Pontificia Universidade Catolica de Minas Gerais, Claros State University, and University of Ouro Preto in Minas Gerais, Brazil. Other partners include Cytel, Platform Life Sciences, MMS Holdings, WHO Therapeutic Guidelines Committee, and the Society for Clinical Trials. Trial documents are found on the Open Science Framework (https://doi.org/10.17605/OSF.IO/EG37X). Trial data are shared with the International COVID-19 Data Alliance (ICODA). The work has been presented and reviewed by the WHO Platform Trials Group.

Data Safety and Monitoring Committee (DSMC) Members: William Cameron, University of Ottawa (Canada), James Orbinski, York University (Canada), Sonal Singh, University of Massachusetts (USA), Kristian Thorlund, McMaster University (Canada), Jonas Haggstrom of Cytel Inc. (Sweden).

REFERENCES

1.   Torres I, Artaza O, Profeta B, Alonso C, Kang J. COVID-19 vaccination: returning to WHO's Health For All. *Lancet Glob Health* 2020; **8**(11): e1355-e6.

2.   Rayner CR, Dron L, Park JJH, et al. Accelerating Clinical Evaluation of Repurposed Combination Therapies for COVID-19. *Am J Trop Med Hyg* 2020; **103**(4): 1364-6.

3.   Omi T, Tanimukai H, Kanayama D, et al. Fluvoxamine alleviates ER stress via induction of Sigma-1 receptor. *Cell Death Dis* 2014; **5**(7): e1332.

4.   Sukhatme VP, Reiersen AM, Vayttaden SJ, Sukhatme VV. Fluvoxamine: A Review of Its Mechanism of Action and Its Role in COVID-19. Front Pharmacol.2021;12:652688. doi: 10.3389/fphar.2021.652688. eCollection 2021

5.   Hoertel N, Sánchez-Rico M, Vernet R, et al. Association between antidepressant use and reduced risk of intubation or death in hospitalized patients with COVID-19: results from an observational study. *Mol Psychiatry* 2021.

6.   Lenze EJ, Mattar C, Zorumski CF, et al. Fluvoxamine vs Placebo and Clinical Deterioration in Outpatients With Symptomatic COVID-19: A Randomized Clinical Trial. *Jama* 2020; **324**(22): 2292-300.

7.   Park JJH, Siden E, Zoratti MJ, et al. Systematic review of basket trials, umbrella trials, and platform trials: a landscape analysis of master protocols. *Trials* 2019; **20**(1): 572.

8.   Woodcock J, LaVange LM. Master Protocols to Study Multiple Therapies, Multiple Diseases, or Both. *N Engl J Med* 2017; **377**(1): 62-70.

9.   Reis G, Silva EAdSM, Silva DCM, et al. A multi-center, adaptive, randomized, platform trial to evaluate the effect of repurposed medicines in outpatients with early coronavirus disease 2019 (COVID-19) and high-risk for complications: the TOGETHER master trial protocol. *Gates Open Research* 2021; **5**(117): 117.

10.   Reis G, Moreira Silva E, Medeiros Silva DC, et al. Effect of Early Treatment With Hydroxychloroquine or Lopinavir and Ritonavir on Risk of Hospitalization Among Patients With COVID-19: The TOGETHER Randomized Clinical Trial. *JAMA Netw Open* 2021; **4**(4): e216468.

11.   Yu L-M, Bafadhel M, Dorward J, et al. Inhaled budesonide for COVID-19 in people at high risk of complications in the community in the UK (PRINCIPLE): a randomised, controlled, open-label, adaptive platform trial. *The Lancet*.

12.   Nicol GE, Karp JF, Reiersen AM, Zorumski CF, Lenze EJ. "What Were You Before the War?" Repurposing Psychiatry During the COVID-19 Pandemic. *J Clin Psychiatry* 2020; **81**(3).

13.   Ishima T, Fujita Y, Hashimoto K. Interaction of new antidepressants with sigma-1 receptor chaperones and their potentiation of neurite outgrowth in PC12 cells. *Eur J Pharmacol* 2014; **727**: 167-73.

14.   Rosen DA, Seki SM, Fernández-Castañeda A, et al. Modulation of the sigma-1 receptor-
IRE1 pathway is beneficial in preclinical models of inflammation and sepsis. *Sci Transl Med* 2019;
11(478).

15.   Schlienger RG, Meier CR. Effect of selective serotonin reuptake inhibitors on platelet
activation: can they prevent acute myocardial infarction? *Am J Cardiovasc Drugs* 2003; **3**(3): 149-
62.

16.   Pan H, Peto R, Henao-Restrepo AM, et al. Repurposed Antiviral Drugs for Covid-19 -
Interim WHO Solidarity Trial Results. *N Engl J Med* 2021; **384**(6): 497-511.

17.   Horby PW, Pessoa-Amorim G, Peto L, et al. Tocilizumab in patients admitted to hospital
with COVID-19 (RECOVERY): preliminary results of a randomised, controlled, open-label,
platform trial. *medRxiv* 2021: 2021.02.11.21249258.

18.   Angus DC, Derde L, Al-Beidh F, et al. Effect of Hydrocortisone on Mortality and Organ
Support in Patients With Severe COVID-19: The REMAP-CAP COVID-19 Corticosteroid
Domain Randomized Clinical Trial. *Jama* 2020; **324**(13): 1317-29.

19.   Wang J, Levi J, Ellis L, Hill A. Minimum manufacturing costs, national prices and
estimated global availability of new repurposed therapies for COVID-19. *medRxiv* 2021.

20.   Chen P, Nirula A, Heller B, et al. SARS-CoV-2 Neutralizing Antibody LY-CoV555 in
Outpatients with Covid-19. *N Engl J Med* 2021; **384**(3): 229-37.

21.   Weinreich DM, Sivapalasingam S, Norton T, et al. REGN-COV2, a Neutralizing Antibody
Cocktail, in Outpatients with Covid-19. *N Engl J Med* 2021; **384**(3): 238-51.

22.   Organization WH. World Health Organization model list of essential medicines: 21st list
2019: World Health Organization, 2019.

TABLES & FIGURES

Figure 1: Participant Flowchart

Figure 2: Probability of efficacy and Bayesian relative risk of extended emergency room observation or hospitalization for fluvoxamine vs. placebo (Panel A: ITT population; Panel B: Modified ITT population)

A) Intention-to-treat analysis

B) Modified intention-to-treat analysis

494   **Figure 3:** Sub-group analyses of fluvoxamine vs. placebo in the TOGETHER Trial

Subgroup	N Placebo (N events)	N Treatment (N events)	HR [95% CI]
Age (years)			
<=50	367 (32)	367 (21)	0.65 [0.38; 1.13]
>50	318 (70)	324 (50)	0.67 [0.47; 0.97]
Sex			
Female	438 (53)	407 (27)	0.53 [0.33; 0.84]
Male	295 (55)	330 (50)	0.81 [0.55; 1.18]
BMI (kg/m2)			
<30	360 (49)	354 (34)	0.69 [0.44; 1.07]
>=30	359 (58)	371 (42)	0.69 [0.46; 1.03]
Time from onset of symptoms			
0–3 days	296 (35)	317 (28)	0.73 [0.45; 1.21]
4–7 days	249 (38)	241 (31)	0.83 [0.52; 1.34]
Diabetes mellitus			
N	365 (27)	354 (22)	0.84 [0.48; 1.47]
Y	367 (81)	382 (55)	0.63 [0.44; 0.88]
Cardiovascular disease			
N	444 (54)	442 (39)	0.72 [0.48; 1.08]
Y	289 (54)	295 (38)	0.67 [0.44; 1.01]
Lung disease			
N	711 (102)	716 (73)	0.70 [0.52; 0.94]
Y	21 (6)	20 (4)	0.70 [0.20; 2.47]
Use of corticoid therapy			
N	724 (106)	729 (77)	0.71 [0.53; 0.95]
Y	7 (1)	6 (0)	

0.35   0.50   0.75   1.0   1.5
<------ Fluvoxamine better --   -- Placebo better ---

497 **Table 1.** Patient characteristics by treatment allocation in the TOGETHER Trial

	Fluvoxamine (n=739)	Placebo (n=733)	Total (n=1472)
Sex			
Female	408(55.2)	438(59.8)	846(57.5)
Male	331(44.8)	295(40.2)	626(42.5)
Race			
Mixed Race[+]	708(95.8)	695(94.8)	1403(95.3)
White	5(0.7)	6(0.8)	11(0.7)
Black or African American	5(0.7)	5(0.7)	10(0.7)
Unknown	21(2.8)	27(3.7)	48(3.3)
Age, years			
>= 50 years	325(44.0)	318(43.4)	643(43.7)
Age Descriptive Statistics			
Median	50	49	50
IQR	17	18	18
Body Mass Index (BMI)			
<30 kg/m^2	354(47.9)	360(49.1)	714(48.5)
>=30 kg/m^2	372(50.3)	359(49.0)	731(49.7)
Unspecified	13(1.8)	14(1.9)	27(1.8)
Time since onset of symptoms			
0-3 days	318(43.0)	296(40.4)	614(41.7)
4-7 days	241(32.6)	249(34.0)	490(33.3)
Unspecified	180(24.3)	188(25.7)	368(25.0)
Risk factors			

Chronic cardiac disease	8(1.1)	8(1.1)	16(1.1)
Hypertension	106(14.4)	88(12.0)	194(13.2)
Chronic pulmonary disease	6(0.8)	3(0.4)	9(0.6)
Asthma	11(1.5)	16(2.2)	27(1.8)
Chronic kidney disease	2(0.3)	2(0.3)	4(0.3)
Rheumatologic disorder	1(0.1)	0(0.0)	1(0.1)
Chronic neurological disorder	8(1.1)	6(0.8)	14(1.0)
Diabetes mellitus: Type 1	25(3.4)	22(3.0)	47(3.2)
Diabetes mellitus: Type 2	103(14.0)	93(12.7)	196(13.3)
Obesity	2(0.3)	1(0.1)	3(0.2)
Any other risk factor(s) or co-morbidities	24(3.3)	19(2.6)	43(2.9)

498 **LEGEND:** [+]Self-identified as someone with mixed-race ancestry.

499   **Table 2:** Proportion of primary outcome events and relative risk of extended emergency room

500   observation or hospitalization of fluvoxamine vs. placebo

	Intention-to-treat analysis			Modified intention-to-treat analysis		
	N	n (%)	Relative risk (95% CrI)	N	n (%)	Relative risk (95% CrI)
Fluvoxamine	739	77 (10.4 %)	0.71 [0.54; 0.93]	728	66(9.1 %)	0.68 [0.50; 0.91]
Placebo	733	108 (14.7 %)	1.00 (ref)	722	97(13.4 %)	1.00 (ref)
All	1472	185 (12.6%)	--	1450	163(11.2 %)	--

501   **LEGEND:** 95% CrI – Credible intervals

503   **Table 3:** Secondary outcomes of fluvoxamine vs placebo in the TOGETHER Trial

	Fluvoxamine	Placebo	Estimated treatment effect (95% CI)	p-value
Viral clearance (Day 7)	45/218 (21%)	57/217 (26%)	0.73 (0.47, 1.14) *	0.17
Hospitalized for COVID	74/739 (10.0%)	90/733 (12.3%)	0.79 (0.57, 1.10) *	0.17
All-cause hospitalization	74/739 (10.0%)	92/733 (12.6%)	0.77 (0.56, 1.07) *	0.12
Time to hospitalization	5 days [3 to 7]	5 days [3 to 7]	0.79 (0.58, 1.08) †	0.14
Number of days of hospitalization	7 days [5 to 12.5]	6 days [3 to 10.75]	1.22 (0.98, 1.53) ‡	0.07
Emergency room visit for at least 6 hours	6/739 (0.8%)	30/733 (4.1%)	0.19 (0.07, 0.43) *	<0.01
Time to the emergency visit for at least 6 hours	5.5 days [4.25 to 6.75]	6 days [3.25 to 8.75]	0.19 (0.08, 0.47) †	<0.01
Death	17/739 (2.3%)	24/733 (3.3%)	0.70 (0.36 to 1.30) *	0.26
Time to death	17 days [9 to 21]	14 days [8 to 20]	0.79 (0.58 to 1.08)	0.26
Number of days on mechanical ventilator	7 days [3 to 12]	6.5 days [3 to 12]	1.10 (0.70 to 1.73) ‡	0.67
Adherence	541/739 (73.2%)	609/733 (83.1%)	0.48 (0.36, 0.63) *	<0.01
Grade 1 TEAE	17/739 (2.3%)	5/733 (0.7%)	3.43 (1.35 to 10.47) *	0.02
Grade 2 TEAE	62/739 (8.4%)	74/733 (9.4%)	0.82 (0.57 to 1.16) *	0.26
Grade 3 TEAE	34/739 (4.2%)	43/733 (4.6%)	0.77 (0.48 to 1.23) *	0.28
Grade 4 TEAE	18/739 (2.3%)	20/733 (2.2%)	0.89 (0.46 to 1.70) *	0.72
Grade 5 TEAE	18/739 (2.2%)	25/733 (2.3%)	0.71(0.38 to 1.30) *	0.27

504   **LEGEND:** TEAE: Treatment Emergent Adverse Event

505   Summary statistics are presented as n/N or median (IQR) unless otherwise stated.

506   * Unadjusted odd ratio

507   + Unadjusted hazard ratio.

508   ‡Exponentiated unadjusted estimates from a log-transformed linear regression.