Predictors of bovine *Schistosoma japonicum* infection in rural Sichuan, China

Elise Grover[1], Sara Paull[1], Katerina Kechris[2], Andrea Buchwald[1,3], Katherine James[1], Yang Liu[4], Elizabeth

J. Carlton[1*]

[1] Department of Environmental and Occupational Health, Colorado School of Public Health, University of

Colorado Anschutz Medical Campus, Aurora, USA

[2] Department of Biostatistics and Informatics, Colorado School of Public Health, University of Colorado Anschutz

Medical Campus, Aurora, USA

[3] Center for Vaccine Development and Global Health, University of Maryland School of Medicine, Baltimore, USA

[4] Institute of Parasitic Diseases, Sichuan Center for Disease Control and Prevention, Chengdu, China

* Corresponding author

Email: elizabeth.carlton@cuanschutz.edu

Note: Supplementary data is associated with this article.

Abstract

In China, bovines are believed to be the most common animal source of human schistosomiasis infections,

though little is known about what factors promote bovine infections. The current body of literature features

inconsistent, and sometimes contradictory results, and to date, few studies have looked beyond physical

characteristics to identify the broader environmental conditions that predict bovine schistosomiasis. Because

schistosomiasis is a sanitation-related, water-borne disease transmitted by many animals, we hypothesized that

several environmental factors – such as the lack of improved sanitation systems, or participation in agricultural

production that is water-intensive – could promote schistosomiasis infection in bovines. Using data collected as a

part of a repeat cross-sectional study conducted in rural villages in Sichuan, China from 2007 to 2016, we used a

Random Forests, machine learning approach to identify the best physical and environmental predictors of bovine

S. japonicum infection. Candidate predictors included: 1) physical/biological characteristics of bovines, 2) human

sources of environmental schistosomes, 3) socio-economic indicators, 4) animal reservoirs, and 5) agricultural

practices. The density of bovines in a village and agricultural practices such as the area of rice and dry summer

crops planted and the use of night soil as an agricultural fertilizer, were among the top predictors of bovine *S.*

japonicum infection in all collection years. Additionally, human infection prevalence, pig ownership and bovine age

were found to be strong predictors of bovine infection in at least one year. Our findings highlight that

presumptively treating bovines in villages with high bovine density or human infection prevalence may help to

interrupt transmission. Furthermore, village-level predictors were stronger predictors of bovine infection than

household-level predictors, suggesting future investigations may need to apply a broad ecological lens to identify

potential underlying sources of persistent transmission.

Keywords: schistosomiasis, buffalo, cattle, machine learning, prevention and control, China

1. Introduction

Bovines have long-been recognized as a key contributor to human schistosomiasis in Asia (Gray et al., 2007; Guo et al., 2006), and emerging evidence suggests they may be contributing to human schistosomiasis in Africa (Colley and Loker, 2018).Schistosomiasis is among the most burdensome helminth infections worldwide, with transmission being reported in a total of 78 countries in 2018 and approximately 230 million people in need of preventative treatment (World Health Organization, 2020). Although great strides have been made in the last few decades in global schistosomiasis control (World Health Organization, 2020), pockets of reemergent or persistent transmission highlight the need for careful consideration of possible local drivers of transmission (Kittur et al., 2019; Song et al., 2016). A poignant example is found in *Schistosoma japonicum* transmission in China, where despite well-established control programs and considerable progress towards elimination since the mid-1950s (Xu et al., 2016), a national report highlighted that there remained 450 endemic counties in 2020, a quarter of which had not achieved elimination criteria by year's end (Zhang et al., 2021). *S. japonicum* is transmitted by at least 40 species of wild and domestic mammals (Gray et al., 2009b), and animal activities near likely transmission sites may be important sources of reemergence and persistence. Although there are several domesticated and wild animals that can carry and transmit *S. japonicum*, (Li et al., 2015; Van Dorssen et al., 2017), estimates from Eastern China suggest that bovines may be responsible for as much as 75% of human transmission (Guo et al., 2006). This substantial contribution is thought to be related to the high degree of environmental overlap between humans and bovines during agricultural production, as well as the large amount of fecal output of bovines, which is estimated to be as high as 100 times that of human fecal production each day (Gray et al., 2009b; He et al., 2001; Ross et al., 2001). Additionally, the high frequency of livestock movement via the livestock trade within mountainous regions of China may facilitate the spread of *S. japonicum* infections across the region (Zheng et al., 2000; Zhou et al., 2012).

Bovine parasitic flatworm infections also have important veterinary impacts. The consequences of *Schistosoma bovis* and other parasitic flatworms can be quite severe on the animal, including: mortality, extreme morbidity and organ damage, emaciation, reduced milk yields and other production issues, and greater susceptibility to other pathogens (Al-Gaabary et al., 2009; Alemneh et al., 2015; McCauley et al., 1983).

Despite strong evidence that bovines are an important driver of human *S. japonicum* infection (Colley and Loker, 2018; Gray et al., 2007; Guo et al., 2006; Rudge et al., 2013) and of human schistosomiasis in Africa, there

are major gaps in our understanding of the factors that contribute to bovine *Schistosoma* infection. Most studies

assessing bovine infection risks have focused primarily on veterinary schistosome species, the results of which

have been inconsistent. For example, several recent assessments of *Schistosoma bovis* infection risk in Eastern

Africa assessed bovine sex, age, breed and body condition as predictors of bovine infection status with

contradictory results (Chanie et al., 2012; Defersha and Belete, 2018; Gebremeskel et al., 2017; Kebede et al.,

2018; Lulie and Guadu, 2014; Tsega and Derso, 2015; Yihunie et al., 2019). Whereas assessments from 2012

and 2017 in Northwest Ethiopia found that neither age, sex nor the bovine's body condition was associated with

bovine schistosomiasis infection (Chanie et al., 2012; Gebremeskel et al., 2017), a 2018 assessment found no

significant effect of sex or breed, but suggested that bovines with poor body condition and bovines aged 2-5 had

the highest infection risk (Defersha and Belete, 2018). By contrast, a 2019 study from northwestern Ethiopia

recently highlighted that female sex, breed and poor body condition were all associated with bovine infection,

while age was not found to be associated (Yihunie et al., 2019). Reasons for such discrepancies have not been

fully elucidated, though Defersha & Belete (2018) hypothesize that it may be related to variations in management

practices for different bovine groups (e.g. separation of sexes or of age groups) and different grazing ranges and

patterns allowed on different farms (e.g. smaller grazing area of very young and very old bovines) (Defersha and

Belete, 2018).

Outside of eastern Africa, few studies have characterized predictors of bovine schistosomiasis infection.

One study from Malaysia found that older age, low weight and male sex, were all risk factors for *Schistosoma*

spindale infection in a range of different cattle species (Tan et al., 2015). By comparison, a study conducted

primarily among water buffaloes (96.2% water buffalos, 3.8% cattle) in Southern China found that *S. japonicum*

infection intensity was highest in young bovines (<2 years) (Gray et al., 2007). These seemingly contradictory

results may potentially be explained by isolation and limited grazing for calves in Malaysia (Tan et al., 2015), as

well as potential genus-related differences in acquired immunity and self-cure rates (Li et al., 2014), as studies

assessing worm establishment success in the two genera have found that cattle are more susceptible to infection

than water buffalo (He et al., 1992; Xu et al., 1993). Nevertheless, water buffaloes may still act as important hosts

in marshland areas of China, as they are more likely to spend time in water, and therefore more likely to be

involved in the *S. japonicum* transmission cycle (He et al., 2001).

Not only is there a great deal of disagreement in the current body of literature over the key risk factors for

bovine schistosomiasis infection, but studies to date have almost exclusively focused on physical characteristics

rather than broader environmental conditions. There is a considerable literature that documents the role of social

and environmental conditions in human schistosomiasis. For example, water, sanitation and hygiene (WASH)

infrastructure, agriculture and fishing production, irrigation and night soil use – that is, the collection of either

treated or untreated human and/or animal waste for use as fertilizer – have all been implicated as significant risk

factors for human *Schistosoma* infection (Carlton et al., 2015; Grimes et al., 2015; Southgate, 1997; Zhou et al.,

2018). While it is plausible that many of these factors pose similar risks to bovines, this has not yet been tested.

Given that bovine *parasitic flatworm* infections can yield significant morbidity and economic losses, and contribute

substantially to human *S. japonicum* infection risk, studies aimed at assessing potential predictors of *S. japonicum*

infection in bovines are of paramount importance.

To address this gap, we set out to assess a range of potential physical and environmental predictors of

bovine *S. japonicum* infection in 2007, 2010 and 2016 at the individual, household and village-levels in a region

where schistosomiasis persistence has been demonstrated to exist in both humans and bovine populations. We

draw from the literature on human and bovine schistosomiasis to define a broad set of candidate predictors that

describe individual bovine characteristics. In so doing, our study helps to fill the gaps in knowledge about the

conditions that predict *S. japonicum* infection in bovines, as well as the relevance of two additional assessment

scales (household and village-level) that have rarely been considered in the studies conducted to date. The

results of this study will ultimately serve as a critical stepping stone toward developing appropriately-scaled and

targeted schistosomiasis intervention and control activities.

2. Materials and Methods

2.1 Village selection

This study was conducted in two rural counties in Sichuan, China where schistosomiasis reemerged and

persisted despite aggressive control efforts. Surveys of environmental and social risk factors, as well as human

and bovine infection were conducted in 2007, 2010 and 2016. In 2007, three of eight Sichuan counties where

schistosomiasis had reemerged (Liang et al., 2006), were selected for inclusion in our study based on

surveillance record availability and the local control stations' willingness and capacity to collaborate, and a set of

17 to 19 villages was selected in each county, as described previously (Carlton et al., 2011). However, in May

2008, a 7.9 magnitude earthquake severely impacted one of the study counties (Carlton et al., 2015), such that, in

2010, follow-up surveys were conducted in 2 counties (36 villages). In an effort to hone in on the highest risk

locations within the two study counties in a region where infections were declining, surveillance records were

reexamined in 2016, from which, a total of ten villages were selected for inclusion in the 2016 data collection. Of

these, seven were villages that were also included in 2007 and 2010, while three were new. We restrict this

analysis to the two counties surveyed in all three years, noting that our villages are not a representative sample of

Sichuan, but rather a survey of high-risk locations where schistosomiasis has persisted despite control efforts. All

villages are located in the hilly regions of rural Sichuan and ranged from approximately 20 to 150 households and

50 to 200 residents. Finally, the analysis presented here was restricted to villages where bovines were both

present and tested for *S. japonicum* infection: 35 villages in 2007, 30 villages in 2010 and 8 villages in 2016.

Details about the total number of included villages, households and bovines are provided in Supplementary Table

S1. *2.2 Census, household questionnaires and infection surveys*

A village census was conducted in each collection year and all residents over the age of five were invited

to participate in surveys for *S. japonicum* infection. In addition, attempts were made to survey all bovines in the

village for *S. japonicum* infection. In the summers of 2007, 2010 and 2016, the head of each household was

asked to complete a household survey that contained closed-ended questions related to socioeconomic status,

domestic and farm animal ownership, sanitation and water access and agricultural practices. Bovine age, type

and sex were collected at the time of the bovine infection surveys in 2007 and 2010 (these data were not

collected in 2016). Trained staff from the Sichuan Center for Disease Control and Prevention and the county

Schistosomiasis Control Stations piloted and conducted all surveys in the local Sichuan dialect. Only bovines that

had both household survey data and *S. japonicum* infection data were included in this assessment

(Supplementary Table S1).

S. japonicum infection surveys were conducted by attempting to test three stool samples on three

consecutive days from eligible humans and all bovines in the village. Infection surveys were conducted in

November and December of 2007 and 2010, and July 2016. Bovines were isolated in a pen or tied up until a stool

was produced on three separate days (consecutive, when possible). All stool samples were transported to the

central laboratory soon after collection to be examined using the miracidium hatching test, following standard

protocols (Department of Disease Control, 2000). To account for the short survival and rapid hatching of bovine

miracidia, the bovine samples were examined for miracidia at one, three and five hours after preparation for at

least two minutes each time, whereas human samples were assessed at two, five and eight hours after

preparation. One sample from each human was also examined using the Kato Katz thick smear procedure in

2007 and 2010 (Katz et al., 1972). A bovine was considered positive for *S. japonicum* if any miracidium hatching

test was positive. A human was considered positive for *S. japonicum* if any miracidium hatching test or the Kato-

Katz test was positive. Each person who tested positive was notified and referred to the local anti-schistosomiasis

control station for treatment. When bovines tested positive, owners were notified and the animal was referred to

the county veterinary station for treatment.

2.3 Ethics statement

This study was approved by the Sichuan Institutional Review Board, the University of California, Berkeley,

Committee for the Protection of Human Subjects, and the Colorado Multiple Institutional Review Board. All

participants provided written, informed consent. The collection of bovine samples was determined to be exempt

from review by the Animal Care and Use Committee at the University of California, Berkeley and the Institutional

Animal Care and Use Committee at the University of Colorado Anschutz.

2.4 Predictor selection and definitions

The primary outcome of interest in this analysis was bovine *S. japonicum* infection in 2007, 2010 and

2016. All candidate predictors were defined using either the household surveys or the human infection surveys

and were divided into five categories: 1) biological/physical characteristics; 2) potential human sources of

environmental schistosomes; 3) socio-economic indicators; 4) potential animal reservoirs/sources of infection; 5)

agricultural risk factors (Table 1). We included agriculture as its own category because bovines are frequently

employed in agricultural work in China (Zheng et al., 2000), and because different crop types and agricultural

practices have their own inherent exposure risks (e.g. planting wet crops like rice may increase the likelihood of

contact with snail habitat and exposure to cercariae (Gordon et al., 2019)). Variables with hypothesized similar

mechanisms of transmission risk were aggregated where possible. Three crop type categories were created:

winter crops (primarily rapeseed and wheat), summer dry crops (primarily corn, peanuts and vegetables) and

summer wet crops (rice). Night soil use was also included as an agricultural risk factor and divided into three

categories: night soil use on winter crops, dry summer crops and wet summer crops.

There were minor variations in the household survey content and question formulation across the study

period (e.g., pig ownership was not assessed in 2007). Where possible, continuous/discrete predictors were

included over binary measures. We excluded binary variables from the analysis if they represented very rare

(<10%) or very common conditions (>90%). Continuous variables were excluded when >90% of the observations

took a single value. For example, household dog and pig ownership were both excluded in 2016 because >90%

of the households owned dogs while >90% did not own any pigs. A composite household asset score (0-9) was

developed for use in this assessment, which included eight household assets assessed in all three collection

years (washing machine, television, air conditioner, refrigerator, computer, car, motorcycle), as well as a binary

measure indicating that the home was constructed of concrete, wood or bricks (vs. adobe).

Because prior work has demonstrated that group-level measures can serve as important predictors of

schistosomiasis infection in humans (Carlton et al., 2015), we also generated village-level predictors from the

household survey data. Village-level variables represent all households that participated in the household survey

from a given village, even if they didn't own bovines. Village-level variables were either the village-average value

of continuous household measures, or for binary variables, the proportion of the village population reporting the

condition. Notably, the village-level variables excluded all observations from the bovine's own household, and

instead used only the data from the other households in the village that participated in the household survey. This

allowed for an assessment of how the surrounding village environment impacts individual bovine infection risk,

independent of the home environment, whereas the household-level variables aim to unpack the influence of the

unique household environment on bovine infection status. The aforementioned criteria led to a total of 31

predictors.

2.5 Analysis

Among the bovines with *S. japonicum* infection data (i.e. the outcome variable), those that lacked all

household survey data (i.e. the predictor variables) were excluded from the analysis (30/503 bovines in 2007;

36/233 bovines in 2010; and 1/72 in 2016). Infection prevalence was similar among the excluded bovines (11/67,

16.4%) as compared to those included in this analysis (111/741, 15.0%). Among included bovines, missingness

was generally low: the variable with the most missing data was bovine sex in 2007 (21.6% missing). Missing

values were imputed separately for each collection year using the rfImpute function from the "randomForest"

package in R (Breiman, 2018; RColorBrewer and Liaw).

Two preliminary descriptive analyses were carried out. First, spatial patterns of bovine infection

prevalence by village were visualized using ESRI's ArcGIS ArcMap software release 10.5.1 (Environmental

Systems Research Institute (ESRI), 2017). Second, categorical versions of the candidate predictors were

generated in order to evaluate changes in the distribution of predictors over time as well as to allow a qualitative

examination of the distribution of predictors by bovine infection status.

To determine which of our candidate predictors serve as the best predictors of bovine schistosomiasis

infection in 2007, 2010 and 2016, a random forests (RF) machine learning approach was used. For each year,

25% of the data was reserved for validation, while the remaining 75% was used for model construction. To

address class imbalance in our outcome of interest (13.3%, 17.3%, and 19.7% of bovines were *S. japonicum*

positive in 2007, 2010 and 2016, respectively), over sampling of the minority class was conducted. For model

tuning, 10-fold cross validation was performed using the Caret package in R to help select the optimal maximum

node size and the number of variables to try at each branch. Once the optimal value of each of these parameters

was determined, a final model was run using 5000 trees per forest (Breiman and Cutler, 2011).

For each collection year, we conducted a total of ten rebalancing and model tuning iterations to assess

the degree of stability in our variable importance rankings. The mean decrease in accuracy (MDA) value was

used to rank the top ten predictors from each model on a scale of ten to one – from most to least important – such

that the predictor with the highest MDA was assigned a score of 10, and predictors not in the top ten were

assigned a score of zero. These variable rankings were then summed across the 10 rebalancing iterations to give

a 10-model summary score of variable importance, ranging from 100-1. The ten highest scoring variables from

the ten-model summary score were then reassigned a final ranking of first to tenth. Next, we created a "lean"

ranking, using only those predictors ranked first through tenth within each collection year. To do this, we

performed an additional ten iterations of the aforementioned balancing and tuning process with the top ten

predictors, thereby reducing excess noise in the variable ranking assessment caused by including a large number

of candidate predictors. Because we hypothesized that the inclusion of human infection as a predictor of bovine

infection would strongly influence the predictive capacity of our RF models due to a presumed association

between bovine and human infection, we also conducted ten iterations of a sensitivity analysis for each collection

year that excluded the human infection variables from the algorithm. The ability of the full, lean and sensitivity RF

models to predict infection status was assessed using accuracy, sensitivity and the receiver operator curve (ROC)

area under the curve (AUC), respectively. In the case of disagreement or a tie when comparing the chosen

performance metrics, kappa and specificity were subsequently compared to select the top performing model for

each year.

Simple logistic regression analyses were performed to assess the direction of association between the

top predictors and bovine infection, dividing continuous variables into tertiles to assess for potential non-linear

relationships. The direction of association was recorded for the top predictors within each collection year, using a

p-value of <0.2 to indicate weak evidence of a between-group difference. In the case that no difference was

indicated between tertile groups at the p<0.2 level, the predictors were further divided into quartiles and re-

assessed. If still no evidence for a between group difference was identified using quartiles, this point was noted in

the results. Density plots by infection status were also examined for a subset of predictors that were found to have

a change in the direction of association across the collection years. Stata 15 and R Studio 4.0 were used for all

analyses (RStudio Team, 2020; StataCorp, 2015).

3. Results

3.1 Bovine infection prevalence and population characteristics

Villages with high bovine infection prevalence (>30%) were present in all collection years. While some

villages were found to have repeatedly high infection prevalence in bovines across all years despite treatment

programs and aggressive control measures, other villages emerged as high prevalence villages in 2016 after

years of low infection prevalence in bovines. Figure 1 shows a map of bovine infection distribution by village

across two counties.

The bovine study population was comprised of fewer water buffaloes than cattle (18.5% of all bovines

were water buffalo in 2007; 15.5% in 2010), was predominantly female (86.3% in 2007; 87.2% in 2010), and

ranged widely in age from less than a year to 26 years old.

Table 2 compares the distributions of each predictor variable by bovine infection status, for 2007, 2010

and 2016.

There were several notable shifts in the distribution of predictors over time in our study region. Access to

improved sanitation increased across our study period, rising from 22.8% of households reporting improved

sanitation in 2007 to 52.1% with improved sanitation by 2016. Notably, the percent of bovines infected was

roughly equal across the two sanitation groups in 2007, but by 2016, a greater proportion of bovines were infected

if they belonged to households with unimproved sanitation (23.5%) compared to the households with improved

sanitation (16.2%).Household assets also increased over time, as approximately half of all villages had a mean

asset score of less than two in 2007, whereas in 2010 and 2016, none of the included villages had a mean asset

score that low. Across the study period, there was a steady increase in the percent of households reporting

planting rice (69.1% in 2007; 71.5% in 2010; 81.7% in 2016), and other summer crops (77.2% in 2007; 98% in

2010; 100% in 2016), as well as the cultivated area of rice and dry summer crops. Night soil use on rice and

winter crops decreased over time: the proportion of households reporting any night soil use on rice crops dropped

from 35.6% to 11.7% between 2007 and 2016, and for winter crops it dropped from 58.3% to 12.6%. By contrast,

the proportion of night soil users for dry summer crops remained relatively constant across the study period

(52.4% in 2007; 53.5% in 2010; 50.7% in 2016). Notably, the average amount of night soil being applied to crops

dropped across the study period for all crop types.

3.2 Predictors of bovine infection

The predictor rankings were relatively stable across the full models, lean models and sensitivity models

within each year, though there was more variability across collection years (Figure 2). For example, animal

reservoirs were frequently among the top ten predictors in 2016, but only a household's pig ownership (2010) and

bovine density in the surrounding village (both 2007 and 2010) were among the top ten in earlier years.

The density of bovine in a village was one of the most consistent predictors: it was scored in the top ten in

at least one RF analysis from each year, all of which indicate that an increase in bovine density in the surrounding

village corresponds with an increase in bovine infection risk. Pig ownership was associated with a decrease in

bovine infection risk in both years that it was assessed.

Agricultural practices was the most frequently ranked predictor category across all years. Specifically, the

household area of winter crops planted, the mean area of rice planted in the surrounding village, and the mean

amount of night soil applied to dry summer crops in the surrounding village were all among the top ten predictors

in all models. Additionally, the total household area of summer crops planted and the village mean area of winter

crops were also all among the top ten predictors in at least one of the three model types for 2007, 2010 and 2016.

However, despite the inter-year agreement indicating the importance of agricultural predictors, the direction of

association between the top agricultural predictors and bovine infection was not always consistent across years.

In 2007, increases in all key agricultural predictors were associated with an increase in bovine infection risk, apart

from night soil use on winter crops. By contrast, in 2010 and 2016 our models indicate a mixture of positive and

negative associations across the key agricultural predictors, and in one instance (household winter crop area in

2010), no evidence of a relationship was found.

Figure 3 depicts changes in the distribution of different agricultural practices by bovine infection status

between 2007 and 2016. Most prominent among these is dry summer crop farming, which saw a notable increase

in the total and mean area of crop being planted by households and villages between 2007 and 2016 (Figure 3,

panel B). In 2007 and 2016 the area of dry summer crops cultivated at the household and village-level was

positively associated with bovine infection, but in 2010 it was not. There was a general decrease in the amount of

night soil being applied to rice, winter crops and, to a lesser extent, dry summer crops, between 2007 and 2016

(Figure 3, panels D-F). Greater night soil use on dry summer crops at the village level was associated with

increased bovine infection risk in 2007 and 2016, but not in 2010. Less night soil was used on rice, compared to

other crops, and night soil use on rice was not a key predictor of bovine infection in any year.

There are several other key predictors that stand out in one or more collection years. Human infection

prevalence in the surrounding village was among the top five predictors of bovine infection in 2007 and 2010, and

the number of infected humans within the household was among the top ten predictors in 2010. In all cases, an

increase in human infections was associated with an increase in bovine infections. When the human infection

predictors were removed for the sensitivity analysis, the rankings of the remaining predictors did not shift

substantially in any collection year. Bovine age was also among the top predictors in 2007, and the probability of

infection was found to increase with age. In all of the 2010 analyses, the number of hatch tests was an important

predictor of bovine infection, a feature not shared by the 2007 and 2016 analyses. This may be related to the

lower proportion of bovines with three hatch test results in 2010 (64.5%), as compared to 2007 (70.6%) and 2016

(73.2%).

Of the three different analyses performed (full, lean and sensitivity) for each collection year, the full

models (i.e. those that included the full list of predictors available in a given year) tended to perform the best

(Table 3, Figure 4). Overall, the models had high accuracy values, with the top performing full models producing a

maximum accuracy of 0.864 (95% CI: 0.79 – 0.92) in 2007, 0.816 (95% CI: 0.68 – 0.91) in 2010, and 1.0 (0.81 –

1.0) in 2016. Furthermore, despite some variation in model performance across the ten iterations of RF models

for each analysis year, ultimately the models were relatively stable. For example, across each of the ten iterations

of full model analyses, the AUC range was relatively narrow: 0.724 – 0.75 in 2007, 0.816 – 0.863 in 2010, and

0.982 – 1.0 in 2016. However, due to class imbalance in the reserved test datasets (see the no information rate

(NIR) in Table 3), the Kappa value is also a useful performance metric for our models, as this takes class

imbalance into account. According to the benchmarks laid out by Landis and Koch (1977), the Kappa statistics

from our 2007 analyses suggest a "Fair" level of agreement (0.21 – 0.40) between our best RF models and the

true known values in 2007. For 2010, the highest Kappa statistic came from the full predictor analysis, with a

Kappa of 0.463, indicating a "Moderate" level of agreement (0.41 – 0.60) between the prediction model and the

reserved test dataset (Landis and Koch, 1977). In 2016, both the full and sensitivity models achieved perfect

prediction (Kappa = 1) for the test dataset in at least one of the ten model iterations, whereas the Kappa statistic

for the top performing lean model was 0.853, or "Almost Perfect", according to Landis & Koch (Landis and Koch,

1977).

Because the full list of predictors changed slightly across the collection years, a supplemental analysis was

conducted in which only predictors that were available in all three collection years were included in the RF

models. This analysis demonstrated that 1) intra-year rankings and extra-year patterns did not change

substantially, and 2) agricultural variables remained the most prominent predictor category when comparing

across the entire study period. See Supplementary Figure S1 for details of the supplemental analysis. **4.**

Discussion

Agricultural risk factors, bovine density, human infection, pig ownership and bovine age were all

found to be key predictors of bovine infection status in one or more of the years assessed in this analysis. Of the

agricultural risk factors, night soil use on dry summer crops, the village-level area of rice crops, and both the

household and village-level areas of dry summer and winter crops were each ranked among the top five

predictors for one or more collection years in our RF models. Interestingly, for 2007, all of the ranked agricultural

variables except night soil use on winter crops were associated with an increase in bovine infection risk in our

logistic regression assessments, whereas in 2010 and 2016, these agricultural factors were found to be variably

positively and negatively associated with infection. For example, the area of rice cultivated in a village was

positively associated with bovine infection in 2007 and 2010, but negatively associated in 2016. This finding may

be related to changing norms and interventions that have taken hold in recent years as a result of increasing

awareness of the potential risks posed by both bovines as a reservoir of schistosomiasis, and specific agricultural

practices. For example, across our study period, we saw a steady increase in the percent of households planting

rice (69.1% in 2007; 71.5% in 2010; 81.7% in 2016), and a simultaneous decrease in the percent of households

applying any night soil to their rice crops (35.6% in 2007, 14.7% in 2010; 11.7% in 2016). These shifting norms in

rice production and night soil use likely resulted in a decrease in the overall concentration of night soil on rice

crops within our study villages, which in turn, may help to explain why the village-level rice crop area shifts from

having a positive association with bovine infection in 2007 and 2010, to a negative association by 2016.

Assessments conducted in China early in the new millennium repeatedly highlighted bovines as a key

source of environmental contamination and as the main animal reservoir of *S. japonicum* in the country (Gray et

al., 2009a; Gray et al., 2007; Guo et al., 2006). Beginning in 2004, a new government-led approach to eliminating

schistosomiasis transmission in China was adopted, which – in conjunction with infrastructure improvements in

rural areas and several new schistosomiasis elimination interventions – featured replacing bovines with

machinery in agricultural production (Liu et al., 2017). Thus, the negative associations that were found

intermittently between bovine infection and some of our agricultural variables in 2010 and 2016 may be linked to

the added precautions that were being adopted when bovines were being used for agriculture, or because

bovines were being reallocated for other purposes (e.g. beef production) as machinery became the norm for large

crop areas or those deemed high risk (e.g. wet rice crops).

Dry summer crops stand out as a key agricultural risk factor for bovine infection. The area of dry summer

crops cultivated and the quantity of night soil applied to dry summer crops were key predictors of bovine infection

in all years, and positively associated with bovine infection in 2007 and 2016. It is unclear why this is: it is possible

that the cultivation of such crops provides snail habitat and/or the use of night soil on dry summer crops facilitates

infection of snails living near the field (in our study region, more night soil is applied to dry summer crops than

rice), leading to bovine infection. A downward trend in night soil use (total and mean number of buckets) on crops

can be observed in Figure 3 (parts D-F), though notably, we do not see any substantial shift in the overall

proportion of households that reported any night soil use on summer crops over the years (52.4% in 2007; 53.5%

in 2010; 50.7% in 2016) (Table 2). Increasing recognition of the potential risks posed by night soil use during our

study period (Carlton et al., 2011) may have also contributed to some decreases in night soil use and/or the more

careful treatment of night soil prior to field applications. However, the continued practice of applying night soil to

dry summer crops, paired with the steady increase in the total area of summer crops being planted by villagers

over the study period (see Figure 3, part B) may help to explain why night soil use on summer crops is positively

association with bovine infection status in 2016.

Bovine density in the surrounding village was in the top ten predictors of our RF models, and was

positively associated with bovine infection in our regression models in all collection years. These findings align

well with the existing literature that points to bovines as the most important reservoir of *S. japonicum* infection in

China (Gray et al., 2009a), and suggests that being in close proximity to higher densities of bovine hosts may

correspond with increasing infection risk, as has been found for other bovine pathogens (Guo et al., 2006;

Meadows et al., 2018; Spencer et al., 2015). However, it is worth noting that household-level bovine ownership

was not among the top predictors in any of our RF models, highlighting that the larger-scale lens (i.e. village-level

analysis scale) may be particularly important to future investigations and control strategies. Likewise, recent

informal interviews with residents in our study sites have revealed that bovines are infrequently kept near the

home, as allowing bovines to graze (and defecate) freely is an economical and efficient way of raising bovines,

further illustrating that the household scale may not always be broad enough to capture larger scale trends.

Instead, villagers opt to bring their bovines to the mountains to graze during the day, which subsequently presents

more opportunities for contact between bovines from different households, and may ultimately result in more

widespread environmental contamination (e.g. bovine feces washed into nearby irrigation ditches after

precipitation).

In the developmental stages of this analysis, we hypothesized that human infection prevalence and the

number of infected people in the household would be among the top predictors of bovine infection status, given

the known link between human schistosomiasis and bovine reservoirs (e.g. Gray et al., 2009a). It was therefore

somewhat surprising to find that household-level human infection was only highly ranked in 2007 within our RF

models, while village human infection prevalence was ranked in the top ten in 2007 and 2010, and neither

household-level human infection nor village-level human infection prevalence were important predictors in 2016.

One potential explanation for the changing importance of human infection as a predictor of bovine infection could

be related to the aforementioned bovine-removal phenomenon, in which bovines are increasingly being removed

from the village area and brought to alternative mountain locations for grazing. As this act would likely result ~~in~~

~~an~~in an overall drop in time humans spent in bovine grazing areas, one might expect to see a separation of the

bovine and human infection cycles if snail habitats are common in bovine grazing sites. In fact, the drop in the

important rankings of human infection status in 2016 coincides with a jump in the variable importance rankings for

village-level bovine ownership (6th – 8th in 2007 and 2010; 1st -2nd in 2016), providing further support of the theory

that bovines may be becoming increasing important reservoirs of schistosomiasis infection. On the other hand, an

altogether different explanation for the differences in the 2016 rankings compared to 2007 and 2010 is that the

2016 data collection simply didn't have a large enough sample size to allow for the detection of a true relationship

between relatively rare events.

As such, one limitation of this assessment was the relatively small sample sizes, particularly in 2016

(N=71), though to a lesser extent, 2010 (N=197) and 2007 (N=473), given the correspondingly large number of

predictors that were included in the full predictor models (N=29, N=31, N=26, in 2007, 2010 and 2016

respectively). While RF models are generally acknowledged as being able to handle assessments of high

dimensional data even with relatively small sample sizes (Biau and Scornet, 2016), it remains that small samples

sizes can still give rise to the aforementioned issue of non-detection of rare events. Another limitation to this

assessment is that RF models tends to favor continuous predictors over categorical measures, as they allow for a

wider range of potential split points for classifying observations. For this reason, it is not particularly surprising that

age was the only predictor from the individual/physical characteristics predictor group that was ranked among the

top ten predictors, as the remaining individual characteristics were binary measures. Another notable limitation of

the variable importance rankings used in RF models is that they become less reliable when predictors are highly

correlated with one another (Strobl et al., 2008). This may be particularly important to the rankings ascribed to the

agricultural variables, as correlation between the area of the different crop types planted and the amount of night

soil used on each crop tended to be high across all collection years, with the highest predictor correlations found

in the 2016 collection year (See Supplementary Figures S2 – S4 for correlation matrices). This is notable, as a

higher degree of instability in the variable importance rankings was also found for 2016 as compared to 2010 or

2007, suggesting predictor correlation may be responsible. We therefore recommend that the variable rankings

presented in this analysis be interpreted more holistically (e.g. agricultural variables are strong predictors of

bovine infection), and advise caution when comparing unique variable ranking values against one another (e.g.

rice crop area is less important than winter crop area).

Our main interests in this assessment were to 1) identify the best physical and environmental predictors

of bovine *S. japonicum* infection within rural farming communities in Sichuan China, and 2) to ascertain whether

there are broader trends in bovine infection distribution across individual, household or village-levels scales or

over time. Our RF assessments have highlighted some key patterns that were repeated across multiple collection

years and multiple iterations of three different models. Agricultural factors and animal reservoirs – specifically,

high bovine density in the surrounding village – were repeatedly found to be among the top predictors of bovine *S.*

japonicum infection across all assessment years. Furthermore, human infection, pig ownership and bovine age

were also found to be strong predictors of bovine infection in at least one year. Taken together, these findings

highlight the potential utility of presumptively treating bovines residing in villages and households that engage in

high-risk agricultural practices, those of older age, or those belonging to villages with particularly high human

infection or bovine density. Additionally, village-level predictors tended to be better predictors of bovine infection

than household-level predictors, suggesting that future investigations and interventions may need to apply a

broad ecological lens in order to successfully extricate and address environmental sources of ongoing

transmission. In so doing, not only can unnecessary bovine morbidity and the associated economic losses be

reduced, but such measures will also help to minimize the potential for bovines to serve as another pathway to

persistent human schistosomiasis transmission.

Acknowledgments: We are grateful for the support and efforts of the field research team members from the

Institute of Parasitic Diseases and the county anti-schistosomiasis control stations for their efforts in collecting the

data presented here.

Author Contributions: Conceptualization: EC, YL, SP, AB; Data Curation: DL, EG; Formal Analysis: EG, SP,

KK, AB, EC; Funding Acquisition: EC, YL, SP; Investigation: YL, DL, EC; Methodology: EG, SP, EC, AB, KK;

Project Administration: YL, DL; Resources: EC, YL, DL; Validation: EG, SP, EC, KK; Writing – Original Draft

Preparation: EG, SP, EC, KJ; Writing – Review & Editing: EG, SP, KK, YL, DL, EC, AB, KJ.

Data Availability: Due to the inclusion of potentially identifying information (e.g. socio-economic indicators and

human infection status), access to study data must be requested through the Carlton Lab group to ensure the

protection of research subjects and compliance with all ethical guidelines. Please contact Elizabeth Carlton at

elizabeth.carlton@cuanschutz.edu for more information.

Declaration of interests: The authors have declared that no competing interests exist.

Funding: This research was supported by grants from the National Institute of Allergy and Infectious Diseases:

R01AI134673 (EJC, PI), R21AI115288 (EJC, PI) and R01AI068854 (Robert Spear, PI). The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National Institutes of

Health. The funders had no role in study design, data collection and analysis, decision to publish, or preparation

of the manuscript.

References

References

Al-Gaabary, M., Osman, S., El-Tonoby, A., 2009. STUDIES ON PARAMPHISTOMIASIS IN RUMINANTS.
Kafrelsheikh Veterinary Medical Journal 7, 116-136.

Alemneh, T., Engdaw, Alemseged, G., Abuhay, 2015. OVERVIEW ON: SCHISTOSOMA INFECTION IN
CATTLE. 1-11.

Biau, G., Scornet, E., 2016. A random forest guided tour. TEST 25, 197-227.

Breiman, L., Cutler, A., 2011. Manual–setting up, using, and understanding random forests V4. 0. 2003. URL
https://www. stat. berkeley. edu/~ breiman/Using_random_forests_v4. 0. pdf.

Breiman, L.C., A.; Liaw, A.; Wiener, M. , 2018. Breiman and Cutler's Random Forests for Classification and

Regression, https://cran.r-project.org/web/packages/randomForest/randomForest.pdf.

Carlton, E.J., Bates, M.N., Zhong, B., Seto, E.Y.W., Spear, R.C., 2011. Evaluation of Mammalian and
Intermediate Host Surveillance Methods for Detecting Schistosomiasis Reemergence in Southwest China. PLoS
Negl Trop Dis 5.

Carlton, E.J., Liu, Y., Zhong, B., Hubbard, A., Spear, R.C., 2015. Associations between Schistosomiasis and the
Use of Human Waste as an Agricultural Fertilizer in China. PLoS Negl Trop Dis 9.

Chanie, M., Dejen, B., Fentahun, T., 2012. Prevalence of cattle schistosomiasis and associated risk factors in
Fogera cattle, south Gondar zone, Amhara national regional state, Ethiopia. Journal of Advanced Veterinary
Research 2, 153-156.

Colley, D.G., Loker, E.S., 2018. New Tools for Old Questions: How Strictly Human Are "Human Schistosomes"-
And Does It Matter? The Journal of infectious diseases 218, 344-346.

Defersha, T., Belete, B., 2018. The Neglected Infectious Disease, Bovine Schistosomiasis: Prevalence and
Associated Risk Factors for its Occurrence among Cattle in the North Gulf of Lake Tana, Northwest Ethiopia. J
Vet Med Health 2, 112.

Department of Disease Control, 2000. Textbook for Schistosomiasis Control, Shanghai Shanghai Publishing
House for Science and Technology, pp. 72-76.

Environmental Systems Research Institute (ESRI), 2017 ArcGIS Desktop Release, 10.5.1 ed, Redlands, CA.

Gebremeskel, A.K., Simeneh, S.T., Mekuria, S.A., 2017. Prevalence and Associated Risk Factors of Bovine
Schistosomiasis in Northwestern Ethiopia. World 7, 01-04.

Gordon, C.A., Kurscheid, J., Williams, G.M., Clements, A.C.A., Li, Y., Zhou, X.N., Utzinger, J., McManus, D.P.,
Gray, D.J., 2019. Asian Schistosomiasis: Current Status and Prospects for Control Leading to Elimination. Trop
Med Infect Dis 4.

Gray, D.J., Williams, G.M., Li, Y., Chen, H., Forsyth, S.J., Li, R.S., Barnett, A.G., Guo, J., Ross, A.G., Feng, Z.,
2009a. A cluster-randomised intervention trial against Schistosoma japonicum in the Peoples' Republic of China:
bovine and human transmission. PloS one 4, e5900.

Gray, D.J., Williams, G.M., Li, Y., Chen, H., Li, R.S., Forsyth, S.J., Barnett, A.G., Guo, J., Feng, Z., McManus,
D.P., 2007. A cluster-randomized bovine intervention trial against Schistosoma japonicum in the People's
Republic of China: design and baseline results. The American journal of tropical medicine and hygiene 77, 866-
874.

Gray, D.J., Williams, G.M., Li, Y., McManus, D.P., 2009b. Transmission Dynamics of Schistosoma japonicum in
the Lakes and Marshlands of China. PloS one 3, e4058.

Grimes, J.E.T., Croll, D., Harrison, W.E., Utzinger, J., Freeman, M.C., Templeton, M.R., 2015. The roles of water,
sanitation and hygiene in reducing schistosomiasis: a review. Parasites & vectors 8, 156.

Guo, J., Li, Y., Gray, D., Ning, A., Hu, G., Chen, H., Davis, G.M., Sleigh, A.C., Feng, Z., McManus, D.P., Williams,
G.M., 2006. A drug-based intervention study on the importance of buffaloes for human Schistosoma japonicum
infection around Poyang Lake, People's Republic of China. The American journal of tropical medicine and
hygiene 74, 335-341.

He, Y., Xu, S., Shi, F., Shen, W., Hsü, S., Hsü, H., 1992. Comparative studies on the infection and maturation of
schistosoma japonicum in cattle and buffaloes. Current Zoology 38, 266-271.

He, Y.X., Salafsky, B., Ramaswamy, K., 2001. Host--parasite relationships of Schistosoma japonicum in
mammalian hosts. Trends Parasitol 17, 320-324.

Katz, N., Chaves, A., Pellegrino, J., 1972. A simple device for quantitative stool thick-smear technique in
Schistosomiasis mansoni. Rev Inst Med Trop Sao Paulo 14, 397-400.

Kebede, A., Dugassa, J., Haile, G., Wakjira, B.M., 2018. Prevalence of bovine of schistosomosis in and around
Nekemte, East Wollega zone, Western Ethiopia. Journal of Veterinary Medicine and Animal Health 10, 123-127.

Kittur, N., King, C.H., Campbell, C.H., Kinung'hi, S., Mwinzi, P.N.M., Karanja, D.M.S., N'Goran, E.K., Phillips,
A.E., Gazzinelli-Guimaraes, P.H., Olsen, A., Magnussen, P., Secor, W.E., Montgomery, S.P., Utzinger, J.,
Walker, J.W., Binder, S., Colley, D.G., 2019. Persistent Hotspots in Schistosomiasis Consortium for Operational
Research and Evaluation Studies for Gaining and Sustaining Control of Schistosomiasis after Four Years of Mass
Drug Administration of Praziquantel. The American journal of tropical medicine and hygiene 101, 617-627.

Landis, J.R., Koch, G.G., 1977. The Measurement of Observer Agreement for Categorical Data. Biometrics 33,
159-174.

Li, H., Dong, G.D., Liu, J.M., Gao, J.X., Shi, Y.J., Zhang, Y.G., Jin, Y.M., Lu, K., Cheng, G.F., Lin, J.J., 2015.
Elimination of schistosomiasis japonica from formerly endemic areas in mountainous regions of southern China
using a praziquantel regimen. Veterinary parasitology 208, 254-258.

Li, Y.S., McManus, D.P., Lin, D.D., Williams, G.M., Harn, D.A., Ross, A.G., Feng, Z., Gray, D.J., 2014. The
Schistosoma japonicum self-cure phenomenon in water buffaloes: potential impact on the control and elimination
of schistosomiasis in China. International journal for parasitology 44, 167-171.

Liang, S., Yang, C., Zhong, B., Qiu, D., 2006. Re-emerging schistosomiasis in hilly and mountainous areas of
Sichuan, China.

Liu, Y., Zhong, B., Wu, Z.-S., Liang, S., Qiu, D.-C., Ma, X., 2017. Interruption of schistosomiasis transmission in
mountainous and hilly regions with an integrated strategy: a longitudinal case study in Sichuan, China. Infectious
Diseases of Poverty 6, 79.

Lulie, B., Guadu, T., 2014. Bovine schistosomiasis: A threat in public health perspective in Bahir Dar town,
northwest Ethiopia. Acta Parasitologica Globalis 5, 1-6.

McCauley, E.H., Majid, A.A., Tayeb, A., Bushara, H.O., 1983. Clinical diagnosis of schistosomiasis in sudanese
cattle. Tropical Animal Health and Production 15, 129-136.

Meadows, A.J., Mundt, C.C., Keeling, M.J., Tildesley, M.J., 2018. Disentangling the influence of livestock vs. farm
density on livestock disease epidemics. Ecosphere 9, e02294.

RColorBrewer, S., Liaw, M.A., Package 'randomForest'.

Ross, A.G., Sleigh, A.C., Li, Y., Davis, G.M., Williams, G.M., Jiang, Z., Feng, Z., McManus, D.P., 2001.
Schistosomiasis in the People's Republic of China: prospects and challenges for the 21st century. Clin. Microbiol.
Rev. 14, 270-295.

RStudio Team, 2020. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA, 2020.

Rudge, J.W., Webster, J.P., Lu, D.B., Wang, T.P., Fang, G.R., Basáñez, M.G., 2013. Identifying host species
driving transmission of schistosomiasis japonica, a multihost parasite system, in China. Proceedings of the
National Academy of Sciences of the United States of America 110, 11457-11462.

Song, L.G., Wu, X.Y., Sacko, M., Wu, Z.D., 2016. History of schistosomiasis epidemiology, current status, and
challenges in China: on the road to schistosomiasis elimination. Parasitology research 115, 4071-4081.

Southgate, V.R., 1997. Schistosomiasis in the Senegal River Basin: before and after the construction of the dams
at Diama, Senegal and Manantali, Mali and future prospects. J Helminthol 71, 125-132.

Spencer, S.E., Besser, T.E., Cobbold, R.N., French, N.P., 2015. 'Super' or just 'above average'? Supershedders
and the transmission of Escherichia coli O157:H7 among feedlot cattle. J R Soc Interface 12, 0446.

StataCorp, 2015. Stata Statistical Software: Release 15. StataCorp LP, College Station, TX.

Strobl, C., Boulesteix, A.-L., Kneib, T., Augustin, T., Zeileis, A., 2008. Conditional variable importance for random
forests. BMC Bioinformatics 9, 307.

Tan, T.K., Low, V.L., Lee, S.C., Panchadcharam, C., Kho, K.L., Koh, F.X., Sharma, R.S.K., Jaafar, T., Lim, Y.A.L.,
2015. Detection of Schistosoma spindale ova and associated risk factors among Malaysian cattle through
coprological survey. Japanese Journal of Veterinary Research 63, 63-71.

Tsega, M., Derso, S., 2015. Prevalence of bovine schistosomiasis and its associated risk factor in and around
Debre Tabor town, north west of Ethiopia. Europ J Biol Sci 7, 108-113.

Van Dorssen, C.F., Gordon, C.A., Li, Y., Williams, G.M., Wang, Y., Luo, Z., Gobert, G.N., You, H., McManus,
D.P., Gray, D.J., 2017. Rodents, goats and dogs – their potential roles in the transmission of schistosomiasis in
China. Parasitology 144, 1633-1642.

World Health Organization, 2020. Schistosomiasis: Key Facts, Fact sheets.

Xu, J., Steinman, P., Maybe, D., Zhou, X.N., Lv, S., Li, S.Z., Peeling, R., 2016. Evolution of the National
Schistosomiasis Control Programmes in The People's Republic of China. Adv. Parasitol. 92, 1-38.

Xu, S., Shi, F., Shen, W., Lin, J., Wang, Y., Lin, B., Qian, C., Ye, P., Fu, L., Shi, Y., 1993. Vaccination of bovines
against schistosomiasis japonica with cryopreserved-irradiated and freeze-thaw schistosomula. Veterinary
parasitology 47, 37-50.

Yihunie, A., Urga, B., Alebie, G., 2019. Prevalence and risk factors of bovine schistosomiasis in Northwestern
Ethiopia. BMC veterinary research 15, 12.

Zhang, L.J., Xu, Z.M., Yang, F., Dang, H., Li, Y.L., Lü, S., Cao, C.L., Xu, J., Li, S.Z., Zhou, X.N., 2021. [Endemic
status of schistosomiasis in People's Republic of China in 2020]. Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi
33, 225-233.

Zheng, J., Guo, J.G., Wang, X.F., Zhu, H.Q., 2000. Relationship of the livestock trade to schistosomiasis
transmission in mountainous area. Zhongguo ji sheng chong xue yu ji sheng chong bing za zhi = Chinese journal
of parasitology & parasitic diseases 18, 146-148.

Zhou, G., Shan, L., Shi-Zhu, L., Hui, D., Li-Juan, Z., Jing, X., 2018. Analysis on the situation of schistosome
infections in floating population in national schistosomiasis surveillance sites of China. Zhongguo xue xi chong
bing fang zhi za zhi= Chinese journal of schistosomiasis control 30, 124-130.

Zhou, Y.B., Liang, S., Jiang, Q.W., 2012. Factors impacting on progress towards elimination of transmission of
schistosomiasis japonica in China. Parasit Vectors 5, 275.

Figure Legends

Figure 1. Village-level prevalence of schistosomiasis in bovines in 2007, 2010 and 2016. The prevalence of

schistosomiasis infection in bovines is indicated for each village included in 2007 (left), 2010 (center) and 2016

(right). The darker the shade, the higher the prevalence. While villages in the lower left corner of each map were

found to have repeatedly high infection prevalence, other villages achieved 0% infection in bovines in later years

(upper left and center right of maps), while still others emerged as high prevalence villages in 2010 (center of

maps) and 2016 (upper left of maps) after years of low infection prevalence in bovines. Service Layer Credits:

World Terrain Base Sources: Esri, USGS, NOAA. OpenStreetMap Data Extracts for China, Asia: Data/Maps

Copyright 2018 Geofabrik GmbH and OpenStreetMap Contributors.

Note: Figure 1 has a color version (online) and a grayscale version (print).

Figure 2. V**ariable importance rankings and direction of association for candidate predictors of bovine** *S.*

japonicum **infection in 2007, 2010 and 2016.** Variable importance rankings are based on a composite of mean

decrease in accuracy scores for 10 random forest (RF) models for each model type (full, lean and sensitivity) and

collection year. The direction of association was determined through logistic regression, using tertile categories

for continuous variables to assess evidence for non-linearity. A p-value of <0.2 was used to indicate evidence of a

between-group difference, and, when a between group difference was found, the direction of association is

indicated. See Supplementary Table S2 for detailed logistic regression results.

Note: Figure 2 will be in color for both the online and print versions.

Figure 3. Changes in agricultural practices and the relationship between bovine infection and agricultural

predictors over time. For each of the agricultural predictors included in this analysis, boxplots are used to

represent the distribution of uninfected (white) and infected bovines, for household-level (left) and village-level

variables (right) in 2007, 2010 and 2016.

Note: Figure 3 has a color version (online) and a grayscale version (print).

Figure 4. ROC for the best performing models. The ROC curves and corresponding AUC are shown for the top

performing full, lean and sensitivity models for A) 2007 in the top panel, B) 2010 in the middle panel, and C) 2016

in the bottom panel.

Note: Figure 4 has a color version (online) and a grayscale version (print).

Table 1. Summary of predictor variables included in the analysis.

Predictor list	Predictor category [a]	Scale of analysis	Years available	Variable type
Bovine type (water buffalo vs. cow)	Physical/biological	Individual	2007, 2010	Binary
Bovine sex	Physical/biological	Individual	2007, 2010	Binary
Bovine age	Physical/biological	Individual	2007, 2010	Continuous
Number of hatch tests [b]	Physical/biological	Individual	All	Discrete
County of residence	Physical/biological	Household	All	Binary
Prevalence of human infection in village	Potential human sources	Village	All	Continuous
Number of infected human household members	Potential human sources	Household	All	Discrete
Percent of households with improved sanitation	Socio-economic indicators	Village	All	Continuous
Household has improved sanitation (y/n) [c]	Socio-economic indicators	Household	All	Binary
Village mean asset score (0-9)	Socio-economic indicators	Village	All	Continuous
Household asset score (0-9)	Socio-economic indicators	Household	All	Discrete
Percent of households that own cats	Animal reservoirs/sources	Village	All	Continuous
Household cat ownership	Animal reservoirs/sources	Household	All	Binary
Percent of households that own dogs	Animal reservoirs/sources	Village	All	Continuous
Household dog ownership	Animal reservoirs/sources	Household	2007, 2010	Binary
Mean number of pigs per household	Animal reservoirs/sources	Village	2010, 2016	Continuous
Household pig ownership	Animal reservoirs/sources	Household	2010	Discrete
Mean number of bovines per household	Animal reservoirs/sources	Village	All	Continuous
Household owns other bovines	Animal reservoirs/sources	Household	All	Discrete
Mean area of rice planted per household	Agricultural risk factors	Village	All	Continuous
Area of rice planted by household	Agricultural risk factors	Household	All	Continuous
Mean area of dry summer crop planted per household	Agricultural risk factors	Village	All	Continuous
Area of dry summer crop planted by household	Agricultural risk factors	Household	All	Continuous
Mean area of winter crop planted per household	Agricultural risk factors	Village	All	Continuous
Area of winter crop planted by household	Agricultural risk factors	Household	All	Continuous
Mean number of buckets of night soil used on rice crops	Agricultural risk factors	Village	All	Continuous
Number of buckets of night soil used on rice crops	Agricultural risk factors	Household	All	Discrete
Mean number of buckets of night soil used on dry summer crops	Agricultural risk factors	Village	All	Continuous
Number of buckets of night soil used on dry summer crops	Agricultural risk factors	Household	All	Discrete
Mean number of buckets of night soil used on winter crops	Agricultural risk factors	Village	All	Continuous
Number of buckets of night soil used on winter crops	Agricultural risk factors	Household	All	Discrete

[a] Predictors were grouped into five categories relevant to bovine *S. japonicum* infection risk probability: 1) physical/biological characteristics (e.g. old-age); 2) potential human sources of environmental schistosomes (e.g. human *S. japonicum* infection prevalence in the bovine's household); 3) socio-economic indicators (e.g. prevalence of improved sanitation systems in the surround village); 4) potential animal reservoirs/sources of infections (e.g. prevalence of dog ownership in the surrounding village); 5) agricultural risk factors (e.g. a household's total rice crop area).

[b] Because not all bovines produced three stool samples, and examination of a greater number of stool samples can increase the probability of detecting infection, the number of hatch tests used on a given bovine was also included as a predictor.

[c] Improved sanitation was defined as access to an improved toilet in the household, including a biogas digester or a three-compartment toilet.

Table 2. Tabulation of all predictors by bovine infection status.

	2007			2010			2016		
	N Positive	Total tested	% Positive	N Positive	Total tested	% Positive	N Positive	Total tested	% Positive
Total	63	473	13.32%	34	197	17.26%	14	71	19.72%
Bovine type									
Water buffalo	7	86	8.14%	5	29	17.24%	ND [a]	ND [a]	
Cattle	54	380	14.21%	28	158	17.72%	ND [a]	ND [a]	
Missing	2	7	28.57%	1	10	10.00%			
Bovine sex									
Female	50	320	15.63%	29	163	17.79%	ND [a]	ND [a]	
Male	9	51	17.65%	4	24	16.67%	ND [a]	ND [a]	
Missing	4	102	3.92%	1	10	10.00%			
Bovine age [b]									
<2	3	43	6.98%	6	35	17.14%	ND [a]	ND [a]	
2 to 4	26	220	11.82%	9	59	15.25%	ND [a]	ND [a]	
5+	28	137	20.44%	18	93	19.35%	ND [a]	ND [a]	
Missing	6	73	8.22%	1	10	10.00%			
Hatch tests (N)									
1	3	59	5.08%	0	29	0.00%	1	10	10.00%
2	8	80	10.00%	6	41	14.63%	2	9	22.22%
3	52	334	15.57%	28	127	22.05%	11	52	21.15%
County									
1	15	191	7.85%	17	71	23.94%	3	21	14.29%
2	48	282	17.02%	17	126	13.49%	11	50	22.00%
Prevalence of human infection in village									
0-2.53%	14	147	9.52%	0	76	0.00%	4	27	14.81%
2.54- 11.11%	13	158	8.23%	11	57	19.30%	8	29	27.59%
≥11.12%	36	168	21.43%	23	64	35.94%	2	15	13.33%
Number of infected human household members									
0	38	366	10.38%	19	154	12.34%	12	61	19.67%
1+	19	84	22.62%	13	29	44.83%	2	10	20.00%
Missing	6	23	26.09%	2	14	14.29%	0	0	--
Percent of households with improved sanitation									
<10%	14	180	7.78%	3	24	12.50%	0	0	0.00%
10-50%	38	215	17.67%	29	133	21.80%	6	24	25.00%
≥50%	11	78	14.10%	2	40	5.00%	8	47	17.02%
Household has improved sanitation									
No	47	365	12.88%	27	129	20.93%	8	34	23.53%
Yes	16	108	14.81%	7	68	10.29%	6	37	16.22%
Village mean asset score (0-9)									
<2	39	225	17.33%	0	0	0.00%	0	0	0.00%
2-3	24	248	9.68	26	143	18.18%	2	23	8.70%
≥4	0	0	0.00%	8	54	14.81%	12	48	25.00%
Household asset score (0-9)									
0 - 2	22	184	11.96%	4	24	16.67%	2	10	20.00%
2 - 3	31	211	14.69%	15	89	16.85%	3	14	21.43%
4+	10	78	12.82%	15	84	17.86%	9	47	19.15%
Percent of households that owns dogs									
<70%	17	98	17.35%	2	51	3.92%	7	50	14.00%
70-85%	34	217	15.67%	18	94	19.15%	7	21	33.33%
≥85%	12	158	7.59%	14	52	26.92%	0	0	0.00%
Household dog ownership [c]									
No	6	67	8.96%	3	29	10.34%	2	7	28.57%
Yes	57	404	14.11%	31	168	18.45%	12	64 [c]	18.75%
Missing	0	2	0.00%	0	0	--	0	0	--

Percent of households that owns cats									
<45%	12	109	11.01%	12	71	16.90%	4	36	11.11%
45-60%	21	124	16.94%	14	73	19.18%	5	13	38.46%
≥60%	30	240	12.50%	8	53	15.09%	5	22	22.73%
Household cat ownership									
No	17	155	10.97%	12	69	17.39%	5	23	21.74%
Yes	46	312	14.74%	22	128	17.19%	9	48	18.75%
Missing	0	6	0.00%	0	0	--	0	0	--
Mean number of pigs per household									
0	ND [a]	ND [a]		1	21	4.76%	11	20	55.00%
0.01 - 1	ND [a]	ND [a]		26	119	21.85%	1	45	2.22%
≥ 1	ND [a]	ND [a]		7	57	12.28%	2	6	33.33%
Household pig ownership [c]									
0	ND [a]	ND [a]		27	116	23.28%	14	64 [c]	21.88%
1+	ND [a]	ND [a]		7	81	8.64%	0	7	0.00%
Mean number of bovines per household									
< 0.5	5	116	4.31%	4	59	6.78%	10	57	17.54%
0.5 – 1	40	300	13.33%	19	99	19.19%	4	14	28.57%
≥ 1	18	57	31.58%	11	39	28.21%	0	0	0.00%
Household bovine ownership									
0	49	334	14.67%	20	126	15.87%	8	38	21.05%
1+	14	139	10.07%	14	71	19.72%	6	33	18.18%
Mean area of rice planted per household (mu)									
<0.75	14	143	9.79%	9	90	10.00%	7	18	38.89%
0.75-1.5	19	183	10.38%	12	48	25.00%	5	41	12.20%
≥1.5	30	147	20.41%	13	59	22.03%	2	12	16.67%
Area of rice planted by household (mu)									
0	16	146	10.96%	5	56	8.93%	5	13	38.46%
<2	24	195	12.31%	16	84	19.05%	4	31	12.90%
2+	23	132	17.42%	13	57	22.81%	5	27	18.52%
Mean area of dry summer crops planted per household (mu)									
<1	11	184	5.98%	1	14	7.14%	0	3	0.00%
1 -2.5	44	195	22.56%	23	110	20.91%	2	26	7.69%
≥2.5	8	94	8.51%	10	73	13.70%	12	42	28.57%
Area of dry summer crops planted by household (mu)									
0	8	108	7.41%	0	4	0.00%	0	0	--
<3	39	242	16.12%	18	99	18.18%	2	26	7.69%
3+	16	123	13.01%	16	94	17.02%	12	45	26.67%
Mean area of winter crops plant planted per household (mu)									
<2	21	149	14.09%	17	77	22.08%	12	53	22.64%
2-2.75	18	178	10.11%	11	66	16.67%	0	2	0.00%
≥2.75	24	146	16.44%	6	54	11.11%	2	16	12.50%
Area of winter crops planted by household (mu)									
<2	14	142	9.86%	9	52	17.31%	7	29	24.14%
2-3	33	223	14.80%	18	97	18.56%	7	29	24.14%
4+	16	108	14.81%	7	48	14.58%	0	13	0.00%
Mean number of buckets of night soil used on rice crops per household									
<1	3	58	5.17%	29	146	19.86%	8	41	19.51%
1-9.9	21	153	13.73%	4	38	10.53%	6	27	22.22%
≥10	39	262	14.89%	1	13	7.69%	0	3	0.00%
Number of buckets of night soil used on rice crops by household									
0	40	304	13.16%	29	168	17.26%	8	53	15.09%

1+	22	168	13.10%	5	29	17.24%	1	7	14.29%
Missing	1	1	100.0%	0	0	--	5	11	45.45%
Mean buckets of night soil used on dry summer crops per household									
<10	13	190	6.84%	24	99	24.24%	1	15	6.67%
10-30	18	109	16.51%	7	47	14.89%	13	47	27.66%
≥30	32	174	18.39%	3	51	5.88%	0	9	0.00%
Number of buckets of night soil used on dry summer crops by household									
0	31	248	12.50%	24	105	22.86%	6	36	16.67%
1-25	6	53	11.32%	5	32	15.63%	5	22	22.73%
>25	26	172	15.12%	5	60	8.33%	3	13	23.08%
Mean number of buckets of night soil used on winter crops per household									
<10	1	26	3.85%	24	121	19.83%	14	71	100.0%
10-30	19	218	8.72%	7	41	17.07%	0	0	--
≥30	43	229	18.78%	3	35	8.57%	0	0	--
Number of buckets of night soil used on winter crops by household									
0	27	197	13.71%	31	137	22.63%	9	62	14.52%
1-26	5	79	6.33%	0	16	0.00%	1	3	33.33%
>26	31	197	15.74%	3	44	6.82%	4	6	66.67%

[a] No data (ND). Data was not collected on bovine sex, age or type in 2016, or on pig ownership in 2007.

[b] Although predictors are broken into categories to facilitate a comparison of the distributions across the study period, most were assessed as continuous variable in our RF models. See Table 1 for details on variable formulation for the RF models.

[c] Variables with >90% of observations taking on a single value were excluded from the RF assessment. This exclusion criteria applied twice in 2016: >90% of the included households reported owning at least one dog, and >90% of households owned zero pigs

Table 3. Comparison of model performance metrics for the top performing model [a] from the full, lean and sensitivity analyses in 2007, 2010 and 2016.

Performance metrics	2007 Models			2010 Models			2016 Models		
	Full	**Lean**	**Sens.**	**Full**	**Lean**	**Sens.**	**Full**	**Lean**	**Sens.**
Accuracy	0.864	0.864	0.856	0.816	0.816	0.816	1	0.944	1
95% CI	0.79 - 0.92	0.79 - 0.92	0.78 - 0.91	0.68 - 0.91	0.68 - 0.91	0.68 - 0.91	0.81 - 1	0.73 - 1	0.81 - 1
NIR [b]	0.864	0.864	0.864	0.837	0.837	0.837	0.778	0.778	0.778
Kappa [c]	0.313	0.313	0.246	0.463	0.360	0.416	1	0.853	1
Sensitivity	0.313	0.313	0.25	0.75	0.5	0.505	1	1	1
Specificity	0.951	0.951	0.951	0.829	0.878	0.625	1	0.929	1

[a] The top performing model was defined as the one with the highest accuracy for each analysis type (full, lean & sensitivity) and collection year (2007, 2010, 2016). In the case of a tie for the highest accuracy value, the sensitivity, ROC AUC, kappa and specificity were subsequently compared to select the top performing model for each analysis type and year.

[b] No Information Rate

[c] Due to the high degree of imbalance between the outcome classes across the study period, the Kappa values is a useful metric for our models, as it helps to correct bias that results when rewarding the prediction of the majority class. The benchmark values outlined by Landis & Koch (1977) are useful here for determining the relative strength of the predictive models: <0.00 = Poor; 0.00 – 0.20= Slight; 0.21 – 0.40 = Fair; 0.41 – 0.60 = Moderate; 0.61 – 0.81 = Substantial; 0.81 – 1.0 = Almost Perfect.

Figure1_color

Predictor categories	2007			2010			2016		
	Full	Lean	Sens.	Full	Lean	Sens.	Full	Lean	Sens.
Physical/biological characteristics									
Bovine type							ND	ND	ND
Bovine sex							ND	ND	ND
Bovine age	3 ↑	1 ↑	2 ↑				ND	ND	ND
Number of hatch tests				2 ↑	2 ↑	1 ↑			
County of residence									
Potential human sources of schistosomes									
Human infection prevalence in the village	5 ↑	3 ↑	Excl.	1 ↑	1↑	Excl.			Excl.
Number of infected human household members			Excl.	6 ↑	7 ↑	Excl.			Excl.
Socio-economic indicators									
Village prevalence of improved sanitation						9 ∩			
Household has improved sanitation (y/n)									
Village mean asset score (0-9)			10 ↓				8 ↑	5 ↑	8 ↑
Household asset score (0-9)						10 ∅			
Potential animal reservoirs/sources of infection									
Village prevalence of cat ownership							9* ↑	10↑	10 ↑
Household cat ownership									
Village prevalence of dog ownership							9* ↑	9 ↑	
Household dog ownership							Excl.	Excl.	Excl.
Village mean number of pigs owned	ND	ND	ND				3 ↓	7 ↓	4 ↓
Household pig ownership	ND	ND	ND	7 ↓	8 ↓	6 ↓	Excl.	Excl.	Excl.
Village mean number of bovines owned	8 ↑	8 ↑	6 ↑			8 ↑	2 ↑	1 ↑	1 ↑
Household bovine ownership									
Agricultural risk factors									
Village mean rice area	6 ↑	5 ↑	3 ↑	5 ↑	5 ↑	3* ↑	5 ↓	8 ↓	5 ↓
Household rice area				8 ↑	10↑	7 ↑			
Village mean dry summer crop area	2 ↑	7 ↑	4 ↑				6 ↑	6 ↑	7 ↑
Household dry summer crop area	4 ↑	6 ↑	5 ↑	4 ↓	4 ↓	3* ↓			9 ↑
Village mean winter crop area	1 ↑	2 ↑	1 ↑	10↓	6 ↓		1 ↓	2 ↓	2 ↓
Household winter crop area	7 ↑	4 ↑	7 ↑	9 ∅	9 ∅	5 ∅	7 ↓	4 ↓	6 ↓
Village night soil rice: mean # buckets									
Household night soil rice: # buckets									
Village night soil summer dry crop: mean # buckets	9 ↑	10↑	8 ↑	3 ↓	3 ↓	2 ↓	4 ↑	3 ↑	3 ↑
HH night soil summer dry crop: # buckets									
Village night soil winter crop: mean # buckets	10* U	9 U	9 U						
Household night soil winter crop: # buckets	10* U								

Symbol key:

- ↑ Positive association
- ↓ Negative association
- ∩ Non-linear association (Rise-Fall)
- U Non-linear association (Fall-Rise)
- ND Data not collected
- Excl. Variable excluded from model
- * Tied for importance rank
- ∅ No evidence for a significant between-group difference found

Color key: Variable importance rankings (1st-10th) by scale

Individual	1st – 2nd	3rd – 4th	5th – 6th	7th – 8th	9th – 10th
Household	1st – 2nd	3rd – 4th	5th – 6th	7th – 8th	9th – 10th
Village	1st – 2nd	3rd – 4th	5th – 6th	7th – 8th	9th – 10th

  = Not ranked 1-10 for the given model/year

Key

☐ *S. japonicum* **negative bovines**

▨ *S. japonicum* **negative bovines**