Does Blood-Brain Barrier Disruption Define the Glioma Extracellular

Metabolome?

Running title: Live human glioma extracellular metabolome

Authors: Cecile Riviere-cazaux[1], Lucas P. Carlstrom[1], Karishma Rajani[1], Amanda

Munoz-casabella[1], Masum Rahman[1], Ali Gharibi-Loron[1], Desmond A. Brown[2], Kai J

Miller[1], Jaclyn J. White[3], Benjamin T. Himes[4], Ignacio Jusue-Torres[1], Samar Ikram[1],

Seth Ransom[1], Renee Hirte[1], Ju-Hee Oh[5], William F. Elmquist[5], Jann N. Sarkaria[6],

Rachael A. Vaubel[7], Moses Rodriguez[8], Arthur Warrington[1,8], Sani H. Kizilbash[9], Terry

C. Burns[1]

Affiliations:

Departments of [1]Neurological Surgery, [6]Radiation Oncology, [7]Laboratory Medicine and

Pathology, [8]Neurology, and [9]Oncology, Mayo Clinic, Rochester MN, USA

[2]Neurosurgical Oncology Unit, Surgical Neurology Branch, National Institutes of

Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, USA

[3]Department of Neurological Surgery, Wake Forest Baptist Health, Winston-Salem, NC,

USA

[4]Department of Neurological Surgery, Albert Einstein College of Medicine, Montefiore

Medical Center, Bronx, NY, USA

[5]Brain Barriers Research Center, Department of Pharmaceutics, College of Pharmacy,

University of Minnesota, Minneapolis, MN, USA

Corresponding Author   Terry Burns, M.D., Ph.D.

22   Department of Neurosurgery – Mayo Clinic

23   200 First St SW

Rochester, MN, USA 55905

burns.terry@mayo.edu

Telephone: 507-284-2871 Fax: 507-284-5206

Ethics approval and consent to participate: This study was approved by the Mayo Clinic Institutional Review Board and all participants provided their consent to participate in this study.

Consent for publication: All participants have provided consent for publication.

Data availability statement: All data from this study are provided as a supplementary data file.

Funding: National Institute of Health: T32GM065841 to CRC, TCB was supported by NINDS NRCDP K12, NINDS R61 NS122096, Mayo Clinic Center for Individualized Medicine and CCaTS award UL1TR002377, the American Brain Tumor Association, Brains Together for the Cure, Humor to fight the Tumor, and Lucius & Terrie McKelvey. Support was also provided through NCI U54 CA210180 (JS).

Conflict of Interest: None.

Authorship:

- Study design and conception: TCB, CRC, LPC, KR, WFE

- Performed experiments: TCB, CRC, LPC, KR, KJM, JJW, DAB

- Analyzed data: TCB, CRC, KR

- Critically reviewed manuscript: all authors

Acknowledgements: We would like to thank each of our patients who selflessly participated in this study and without whom this work and any advancement in the field of neuro-oncology would not be possible. We also thank Elizabeth Oishi for regulatory support and the Mayo Clinic Neurosurgery Clinical Research team, and the Mayo neurosurgery operative staff for their invaluable support.

Abstract

Background: The extracellular microenvironment modulates cancer behavior. Although radiographic contrast enhancement is an ominous finding in gliomas, it remains unclear if the associated blood-brain barrier disruption merely reflects or functionally supports tumor aggressiveness.

Methods: We utilized intra-operative microdialysis to sample the extracellular metabolome of radiographically diverse regions during fifteen neurosurgical resections. The global extracellular metabolome of recovered microdialysate was evaluated via ultra-performance liquid chromatography tandem mass spectrometry (UPLC-MS/MS) and assessed using enrichment and correlation analyses.

Results: Among 162 named metabolites identified via ultra-performance liquid chromatography tandem mass spectrometry, guanidinoacetate (GAA), was 126.32x higher in enhancing tumor than in adjacent brain. 48 additional metabolites were 2.05-10.18x more abundant in enhancing tumor than brain. With exception of GAA, and 2-HG in IDH-mutant gliomas, differences between non-enhancing tumor and brain microdialysate were comparatively modest and less consistent. The enhancing but not the non-enhancing glioma metabolome was significantly enriched for plasma-associated metabolites largely comprising amino acids and carnitines.

Conclusions: Our findings suggest that metabolite diffusion through a disrupted blood-brain barrier may largely define the enhancing extracellular glioma metabolome. Future studies are needed to determine how the altered extracellular metabolome impacts glioma behavior.

Keywords: Glioma, microdialysis, metabolomics, isocitrate dehydrogenase

Introduction

Astrocytomas and oligodendrogliomas comprise most adult primary malignant brain tumors and remain incurable, regardless of the best available therapies[1]. Isocitrate dehydrogenase (IDH) mutations occur in all oligodendrogliomas and a subset of astrocytomas, affording improved prognosis due to typically slower growth and more favorable response to standard-of-care radiation and alkylating chemotherapy[2,3]. Glioma heterogeneity hampers therapeutic generalizations across diverse patient cohorts. Moreover, diverse genetic phenotypes accumulate within individual patients' tumor ecosystems, hampering therapeutic efforts to target glioma-associated mutations[4,5]. Conversely, molecularly diverse gliomas may resort to similar metabolic pathways in the setting of nutrient deprivation, hypoxia, and genotoxic stress[6,7]. Although glioma metabolism is increasingly scrutinized for therapeutic targets, few strategies currently exist to interrogate the metabolic microenvironment of human gliomas *in situ*.

Microdialysis has been used to quantify human extracellular biomarkers of traumatic and hypoxic brain injury in neurocritical care units. Microdialysis is also a well-established method to quantify central nervous system (CNS) drug delivery in early phase clinical trials[8-10] and has been used longitudinally to study metabolites present in human gliomas when compared to adjacent brain[11,12]. However, while some studies have utilized intraoperative low molecular weight microdialysis (8-20 kDA) to quantify select extracellular metabolites[13-15], no study thus far has deployed intra-operative microdialysis to characterize the global extracellular glioma metabolome, nor to compare the metabolome of contrast-enhancing versus non-enhancing tumor regions. We used variable flow-rate pumps set to 2 μL/min and high molecular weight catheters (100 kDa) to maximize the volume of microdialysate recovered during 15 surgeries in 14 patients. We report that contrast-enhancing (radiographically blood-brain barrier (BBB)

disrupted) regions of high-grade gliomas (HGGs) exhibited a conserved extracellular

metabolome across both IDH-mutant and IDH-WT astrocytomas, similar to findings from other

groups utilizing intra-or-post-operative microdialysis[14,16,17]. However, comparison to the

radiographically BBB intact (non-enhancing) portions of these tumors revealed that this

enhancing glioma signature was significantly enriched for plasma-associated metabolites,

suggesting that BBB disruption may contribute to the glioma extracellular metabolome.

MATERIALS AND METHODS

Patient Cohort, Study Design, and Intraoperative Microdialysis

All study procedures were approved by the Mayo Clinic Institutional Review Board (IRB).

Patients provided written informed consent to participate in NCT04047264--an ongoing study

utilizing intraoperative high molecular weight (100 kDa) microdialysis under an investigational

device exemption. Study eligibility included adults (>18yo) undergoing a clinically indicated

surgery for known or suspected glioma. Each patient (n=14; 15 surgeries; **Table 1**) underwent

intraoperative microdialysis using up to three 100 kDA catheters and variable rate microdialysis

pumps (M Dialysis 71 High Cut-Off Brain Microdialysis Catheters and 107 Microdialysis Pump,

respectively) across radiographically diverse regions (**Fig. 1A-B**). The target and trajectory of

each microdialysis catheter was planned prior to surgery using the computerized

neuronavigation system. When feasible, the microdialysis membrane (typically 10 mm) of each

catheter was planned to be located fully within (i) enhancing tumor (typically "catheter X"), (ii)

non-enhancing tumor (typically "catheter Y"), or (iii) adjacent brain planned for inclusion within

the resection volume (typically "catheter Z"). Once the dura was opened and the cortical surface

exposed, pia at each planned entry point was coagulated and opened with microscissors. The

pre-flushed catheter was then advanced to the intended trajectory. Microdialysis was performed

at 2 μL/min based on a prior intra-operative study demonstrating that this flowrate could enable

sufficient collection of analytes for mass spectrometry[15], with collection vials exchanged every

20 minutes. See supplemental methods for details of microdialysis procedure, pathology, and

molecular tumor analyses. Each patient is referred to via a unique de-identified label reflecting

their histology, IDH-status, and WHO grade according to 2021 WHO classification[18] (Table 1).

Targeted Analysis of D/L-2-HG

Targeted metabolomic analysis of microdialysate was performed by the Mayo Clinic

metabolomic core facility. L and D isomers of 2-hydroxyglutaratic acid were derivatized and

quantified by liquid chromatography mass spectrometry (LC/MS) using slight modifications to

previously described methods[19-21]. See supplemental methods for details.

Untargeted Metabolomic Analysis

Untargeted metabolomic analysis was performed by Metabolon, Inc. Ultra-performance liquid

chromatography tandem mass spectrometry (UPLC-MS/MS), which combines physical

separation of liquid chromatography with the mass analysis capabilities of mass spectrometry.

See supplemental methods for details.

Ranked Metabolite lists

Ranked metabolite lists of tumor (Catheter X or Y) versus brain (Catheter Z) were generated

from fifteen cases based on calculated fold changes of the peak area (e.g. X/Z or Y/Z) for each

metabolite. A tumor versus brain ranked metabolite list could not be generated for one patient

(Oligo[2]) given all catheters were located within tumor. In patients with both enhancing and non-

enhancing tumor catheters (n=7 patients), a ranked metabolite lists of enhancing vs. non-

enhancing tumor was generated (Catheter X/Y). Initial analysis was performed using samples

generated during the first 5 surgeries. Three of these patients had contrast-enhancing tumors.

To determine an overall enhancing tumor (E)-to-brain (B) ranking for each metabolite, each

metabolite present in both catheters was ranked based on fold change (E/B) and then averaged

across all three patients for that metabolite. In the remaining cases, the individual tumor-to-brain

rankings were compared against the first cohort's average enhancing tumor-to-brain ranking of

metabolites. Similar analyses were performed for enhancing versus non-enhancing tumor

(E/NE) and non-enhancing tumor versus brain (NE/B). To understand the potential impact of

extracellular blood-derived metabolites on microdialysate metabolic signatures, we additionally

generated a ranked list of bloodiness-associated metabolites using a pair of cranial CSF

samples obtained from a single patient immediately after incising dura, prior to resection of a 4th

ventricular ependymoma. Both samples were obtained within approximately 1 minute of each

other—one with and one without blood contamination from the surgical wound. Samples were

centrifuged at 400 Gy for 10 minutes to remove red blood cells and other debris, and aliquots of

supernatant submitted to metabolon for global metabolomic analysis. The ranked metabolite list

was based on metabolite fold-change (bloody / non-bloody) between samples, after filtering for

metabolites present in both CSF and >90% of microdialysate samples.

Enrichment Analysis

Gene Set Enrichment Analysis (GSEA) is the most widely utilized analytical method for rank-

based analysis of data sets in biomedicine. GSEA is frequently performed using curated gene

sets from the molecular signatures database (MSigDB), though custom analyte sets can be

used enabling enrichment analysis of metabolites rather than genes. In GSEA, custom analyte

sets are queried against a ranked list of analytes to determine where those analytes fall on the

ranked list (**Supplementary Fig. 1**). If the analytes are most frequently found at the top of the

ranked list, this is known as "positive enrichment;" if they fall at the end of the ranked list, this is

known as "negative enrichment." The software uses this information to calculate a normalized

enrichment score and a p value. FDR values are also provided to correct for multiple hypothesis

testing when multiple analyte sets are evaluated simultaneously for enrichment in a particular

data set. Custom metabolite libraries were created using the top and bottom 35 metabolites

from each ranked metabolite list described above and can be found in **Supplementary Data**

(.gmx file). Using these metabolite libraries (.gmx file), the full ranked list for each patient

(Catheter X vs. Z, Catheter Y vs. Z, or Catheter X vs Y .rnk files) was then used for enrichment

analysis to determine each patient's relative positive and negative enrichment for other patients'

tumor or brain extracellular metabolomes based on the metabolites sets in the .gmx file (**Supplemental Data**). Additionally, results from Björkblom et al. were included as a metabolite set to provide an external reference[17]. The bloody and non-bloody CSF metabolite libraries were included to identify the relative enrichment of tumor or brain microdialysate for plasma-derived metabolites.

MetaboAnalyst

Metabolomic analyses were performed using MetaboAnalyst (5.0), a publicly available web-based tool to analyze and visualize metabolomic data[22]. The normalized peak area data for the metabolites present across at least 40/44 catheters were entered into MetaboAnalyst without further filtering. Assuming a monotonical rather than linear relation between metabolites and samples, Spearman Correlation maps were generated in MetaboAnalyst based on features (metabolites present across all samples) and samples.

Statistical Analysis

Data are presented as normalized peak areas (median = 1 for each metabolite within each analyzed batch. Raw and normalized peak area data are provided in Supplemental File 1. MetaboAnalyst 5.0 was used for Spearman correlations. Enrichment Analysis was performed using GSEA 4.1.0 (Gene Set Enrichment Analysis), repurposed for metabolite set analysis using custom metabolite sets. The normal distribution of metabolites in Figure 4 and Table 2 was tested via D'Agostino-Pearson test; a Wilcoxon signed rank test for non-parametric distributions was then performed on paired enhancing tumor and brain metabolites using GraphPad PRISM 9.1. Volcano plot cut-offs were set at $FC \geq 2$ and $p \leq 0.05$. Graphs were generated using GraphPad PRISM 9.1. $FDR \leq 0.05$ was considered statistically significant for enrichment analysis.

RESULTS

Elevated D-2-Hydroxyglutarate (D-2-HG) in IDH-mutant tumor microdialysate

Microdialysis is a method that can be used to sample "dialyzable" extracellular analytes,

such as metabolites and certain drugs, from the tumor or brain interstitial space. This is most

commonly performed in the post-operative setting with low-molecular weight catheters, including

an FDA-approved system consisting of 20 kDA catheters perfused at 0.3 µL/min using the M-

dialysis 106 pump. While a limited subset of patients may be willing to undergo post-operative

microdialysis, it is usually hoped that there will be minimal tumor left to sample following

surgery, often preventing sampling from multiple regions of the glioma. Intra-operative

microdialysis can enable sampling of diverse regions within the live extracellular glioma

microenvironment during standard-of-care glioma resections. However, low molecular weight

catheters preclude collection of most extracellular analytes; slow flowrates combined with short

intra-operative sampling times prevent acquisition of enough microdialysate volumes to enable

multi-omic analyses. We hypothesized that we could safely and feasibly deploy intra-operative

high molecular weight (HMW, 100kDA) microdialysis using an elevated perfusion rate of 2

µL/min[15] to recover a usable volume of microdialysate containing diverse analytes, including

metabolites, during a standard-of-care surgery. To that end, we initiated our intra-operative

microdialysis trial in an initial cohort of five patients. To evaluate the feasibility of the intra-

operative microdialysis approach, we aimed to measure microdialysate levels of the well-

characterized and dialyzable oncometabolite, D-2-hydroxyglutarate (D-2-HG), known to be

produced by IDH-mutant gliomas[23]. As such, patients were enrolled whose tumors could

plausibly be IDH-mutant. Subsequent pathology revealed that three of these initial five patients

indeed had IDH-mutant gliomas (Oligo[2], Oligo[3]1, and Astro[4-mut]1); two patients were found to

have IDH-wild type (WT) glioblastomas (GBM[WT]1, GBM[WT]2) (**Table 1**). To determine if D-2-HG

levels in intra-operatively acquired microdialysate reflected IDH status, we utilized 10 µL of a 40

µL microdialysate sample from each catheter to quantify D-2-HG and L-2-HG via targeted

metabolomic analysis. D-2-HG was elevated in microdialysate from IDH-mutant as compared to

IDH-wild-Type (WT) tumors (median: 27.29 versus 0.63 μM) (**Supplementary Fig. S2A**), levels of L-2-HG were not different between groups (**Supplementary Fig. S2B**). Patient Astro[4-mut]1 yielded a >1000x difference in D-2-HG between non-enhancing tumor (FLAIR) and brain adjacent to tumor (831.80 versus 0.68 μM). These data demonstrated the feasibility of intraoperative HMW microdialysis to sample a dialyzable oncometabolite during standard-of-care surgery for tumor resection.

Since IDH status has been reported to impact the metabolism of IDH-mutant gliomas, we further analyzed these samples via untargeted metabolomic analyses--an increasingly leveraged strategy to gain more comprehensive insights regarding the global metabolome. We utilized 20 μL of microdialysate from each catheter for untargeted metabolomic analysis via the Metabolon platform[24]. Among hundreds of other metabolites, untargeted analysis reported total 2-HG. When plotted against the combined stereoisomer (D-2-HG + L-2-HG) levels from targeted analyses, a robust correlation (R^2=0.9989) between the two platforms was observed across a 2,500-fold concentration range (**Supplementary Fig. S2C).** Although limited sample volume precluded repeating this approach for other metabolites, these data supported the potential for untargeted metabolomic analyses to generate at least relatively quantifiable data.

Analysis of 2-HG peak area in a second cohort comprising 5 IDH-mutant and 5 IDH-WT cases, including one patient's repeat surgery, confirmed increased levels in IDH-mutant tumors when compared to IDH-WT tumors (21.41x average normalized peak area in IDH-mutant tumor catheters compared to IDH-WT tumor catheters) (**Fig. 1C**). However, one patient with an unusual right frontal H3K27M-mutant infiltrative grade 4 astrocytoma also demonstrated moderately elevated 2-HG levels despite Next Generation Sequencing-confirmed IDH-WT status. This one apparent "outlier" illustrates that elevated extracellular 2-HG is not infallibly specific for IDH-mutant gliomas and is consistent with prior reports of occasionally elevated 2-HG in IDH-WT gliomas[25].

Additionally, in two patients with grade 4 IDH-mutant astrocytomas, the extracellular

abundance of 2-HG was several-fold less in enhancing than non-enhancing tumor (**Fig.1D,** FC

(E/NE): 0.017x and 0.20x for patients Astro[4-mut]1 and Astro[4-mut]2, respectively). This observation

raised the possibility that 2-HG could be lost down its concentration gradient into systemic

circulation. Indeed, elevated 2-HG was previously reported in venous outflow of IDH-mutant

high-grade gliomas[26]. An alternate explanation could be that enhancing tumor regions from IDH-

mutant gliomas generate less 2-HG due to changes in the IDH metabolism of the tumor core.

Arguing against this alternate explanation, a third patient with an enhancing grade 4 IDH-mutant

astrocytoma showed enhancement throughout the lesion, precluding placement of a catheter in

non-enhancing tumor. Although both tumor catheters were placed within the margins of

contrast-enhancing tumor, the second tumor catheter targeted a region of less intense

enhancement, suggestive of poorer central perfusion or relative necrosis (E†). Despite its

placement in presumably poorly perfused tumor core, this catheter recovered 12.1-fold more 2-

HG than the catheter in the most avidly contrast-enhancing region, suggesting that IDH-mutant

metabolism was still present in the tumor core. Collectively, data from these three patients

suggested that 2-HG may be lost from the extracellular compartment in regions of contrast-

enhancing tumor due to increased diffusion across a disrupted blood-brain barrier (**Fig. 1D)**.

Clustering of catheters: correlation analysis of the extracellular global metabolome

Having established that intra-operative HMW microdialysis could detect the relative abundance

of an established oncometabolite of IDH-mutant gliomas, we next sought to evaluate the global

extracellular metabolome across our patients' 44 catheters. To ensure comparison and

reproducibility of metabolite levels across patients, we analyzed 162 named metabolites that

were present in at least 90% of the catheters ($\geq$40/44 catheters). We first asked how patient's

catheters correlated to each other to identify possible patterns in the global metabolome based

on patient identity, radiographic location, IDH mutation, and primary vs. recurrent status.

Spearman correlation of all 44 catheters revealed that all samples from each patient with an

oligodendroglioma (n=3 patients, 9/9 catheters) clustered based on patient identity. Conversely,

9/10 catheters from enhancing regions of WHO grade 4 astrocytomas clustered together (**Fig.**

2A). Clustering of non-enhancing tumor regions was inconsistent, with some catheters

clustering near the patient's own enhancing catheters (patients Astro[4-mut]1, Astro[4-mut]2, and

Astro[4-mut]3), while others clustered with brain catheters (**Fig. 2A),** particularly those from a

completely non-enhancing tumor (MolecGBM[WT]1). Although Patient GemAstro[3-mut]'s tumor did

have enhancing components, these areas were extremely fibrous, hampering catheter

placement. As such, both catheters were placed into non-enhancing regions. Interestingly,

however, this patient's catheters clustered with the enhancing tumor catheters, potentially

suggesting flow of metabolites from nearby enhancing tumor regions (**Fig. 2A)**. Contrary to our

expectations based on recently published tissue metabolomic studies, no obvious clustering

was observed based on IDH status[27], nor based on recurrence. Indeed, one patient's enhancing

catheter from their primary resection (GBM[WT]2: Catheter X) was most strongly correlated with

the enhancing catheter from their repeat resection (GBM[WT]2B: Catheter X). In sum, results of

correlation analyses suggest that the extracellular metabolome of molecularly diverse contrast-

enhancing astrocytomas clustered by catheter location, whereas that of oligodendrogliomas

clustered by patient. Perhaps unsurprisingly, the extracellular metabolome of non-enhancing

tumor regions of WHO grade 4 astrocytomas appeared to exist on a spectrum, appearing in

some cases more similar to the enhancing tumor and in other patients, closer to that of the

"brain" catheter. However, visual inspection of relative metabolite abundance as a function of

catheter demonstrated substantial heterogeneity across patients within each region (**Fig. 2B**).

A reproducible metabolome of enhancing glioma versus brain

Given the substantial metabolic heterogeneity between patients, we asked if each patient could

serve as their own control to identify the extracellular tumor metabolome. As the Spearman

correlation had suggested clustering of enhancing tumor catheters (**Fig. 2A**), we began by comparing catheters in enhancing tumor as compared to adjacent brain. For each patient, metabolites were ranked from highest to lowest fold change based on metabolite peak area in enhancing tumor (E) versus brain adjacent to tumor (B). Although the magnitude of these fold changes varied between patients (**Supplementary Fig. S3)**, the rank of enhancing tumor versus brain-associated metabolites appeared similar in 3/3 patients with enhancing tumors from our initial cohort (**Supplementary Fig. S4A)**. We used the average ranks from these 3 patients to then evaluate the reproducibility of findings in the second cohort of 5 new patients and one patient's recurrent (GBM[WT]2B) disease 10 months later. The ranked order of metabolites in the enhancing versus brain comparison appeared visually similar for all patients in both cohorts when ranked based on the average of the 9 cases **(Fig. 3Ai)**. Additionally, paired enhancing versus non-enhancing tumor catheters were available from 7 patients. Similar rank-based analyses were performed for enhancing versus non-enhancing tumor which revealed a surprisingly conserved rank of enhancing-to-non-enhancing metabolites across patients (**Fig. 3Bii; Supplementary Fig. S4B**). Conversely, the distribution of non-enhancing catheter-associated metabolites appeared less consistent (**Fig. 3Ci; Supplementary Fig. S4C**). Despite this diversity, when this non-enhancing-to-tumor metabolite distribution was queried in five patients from whom only non-enhancing tumor and brains catheters were available, a similar ranked metabolome was observed in two patients, Oligo[3]1 and GemAstro[3-mut] (**Supplementary Fig. S5)**. These data demonstrated that metabolic diversity exists among patients with oligodendrogliomas and suggests that there could be recurrent metabolic patterns in some but not all non-enhancing tumors.

To more rigorously and quantitatively compare metabolic phenotypes between catheters and between patients, we utilized enrichment analysis (EA) to statistically identify similarities between and across patients' catheters. Although more commonly applied to gene-based analyses, we repurposed this for metabolomic analysis. Using EA, we selected the top and

bottom 35 metabolites based on fold change from each patient's tumor vs. brain or enhancing vs. non-enhancing tumor ranked metabolite list ("metabolite sets") (**Supplementary Fig. S1**). We then determined where these metabolites fell in the ranked metabolite list of tumor versus brain or enhancing vs. non-enhancing tumor for the remaining patients. "Positive enrichment" meant that the metabolites fell at the tumor-end of the ranked list, while "negative enrichment" indicated that they fell at the brain-end of the ranked list**.** In patients with enhancing astrocytomas, each patient's individual enhancing tumor signature could be robustly and positively identified in the enhancing tumor of the other patients, suggesting a convergent glioma metabolome (**Fig. 3Aii).** Conversely, each patient's brain signature could be found at the "brain" end of every other patients' ranked metabolite list **(Fig. 3Aii).** Leveraging these individual patient-level data, these analyses proved highly robust across patients with a False Discovery Rate (FDR) of 0.000 for most interpatient comparisons of enhancing tumor versus brain metabolites (**Fig. 3Aii, Supplementary Table 2)**. For independent reference, the previously published post-operative tumor versus adjacent brain microdialysate metabolome from Björkblom et al[17] was evaluated and found to be present in the enhancing catheters of astrocytomas (FDR of 0.002 or less, Supplementary Table 2). Indeed, among 24 reported brain- or tumor-associated metabolites, all but one metabolite revealed a reproducible pattern in our data set (**Supplementary Fig. S6**). These data confirmed the robust nature of the extracellular enhancing glioma metabolome, being readily and comparably discernible in both the intra-operative and post-operative settings.

Enrichment analyses of enhancing versus non-enhancing catheters again demonstrated that enhancing catheters were metabolically separate from their non-enhancing catheter counterparts and that this could be consistently found across most patients (most FDRs=0, except GBM^WT1, **Fig. 3Bii;** Supplementary Table 3**)**. As expected from Figure 3Ci, non-enhancing catheters were rarely enriched for each other when compared to brain (**Fig. 3Cii,**

Supplementary Table 4). Overall, these data suggest that the extracellular metabolome of

enhancing glioma is distinct from, and more consistent than that of non-enhancing glioma.

Metabolites of the enhancing and non-enhancing glioma extracellular microenvironments

After demonstrating that enhancing astrocytomas exhibit a conserved extracellular

metabolome via rank-based and formal enrichment analyses (**Fig. 4A)**, we next asked which

metabolites define this extracellular microenvironment across the full cohort of all 9 surgeries.

The Wilcoxon rank sum test for non-parametric distributions, paired enhancing vs. brain

catheters revealed 48 significantly elevated metabolites in the enhancing tumor catheter and 22

significantly elevated metabolites in the brain catheter using cutoff criteria of p-value$\leq$0.05 and

FC$\geq$2 (**Fig. 4A; Table 2)**. The top enhancing astrocytoma metabolite was guanidinoacetate

(GAA) (E/B = 126.32x), which, to our knowledge, has not previously been reported in the

context of glioma. The enhancing glioma signature was also defined by greater relative

abundance of proline (FC=10.18x), carnitine-family metabolites (deoxycarnitine, carnitine, and

acetylcarnitine at FC= 8.71x, 7.65x, and 7.47x, respectively), and N1-methyl-2-pyridone-5-

carboxamide (2-PY) (FC=7.18x) (all at p<0.005). Metabolites defining the brain metabolome

included arabitol (FC= 0.04) and arabonate (FC=0.08), in addition to canonical brain-associated

metabolites, such as N-acetylaspartate (NAA)[28] (all p<0.005).

We then asked if these metabolites were unique to enhancing tumor when compared to

non-enhancing tumor for each patient (**Fig. 4B)**. Again using a Wilcoxon rank sum test for non-

parametric distributions, twenty-eight metabolites were significantly higher in the enhancing

glioma metabolome, while twenty metabolites were greater in the non-enhancing glioma

metabolome (**Table 2**). Interestingly, 27/28 enhancing glioma metabolites, had also been found

in the enhancing glioma-vs-brain comparison, again demonstrating a robust extracellular

metabolome of enhancing glioma (relative abundance of representative metabolites shown in

Fig 4C). Of the twenty non-enhancing versus enhancing glioma metabolites, 15 overlapped with

brain (versus enhancing tumor). These data suggest the continued presence of brain-

associated metabolites within non-enhancing portions of glioma. Indeed, when non-enhancing

glioma catheters were compared to brain **(Fig. 4D)**, no metabolites were unique to the brain

metabolome, while fifteen metabolites were higher in non-enhancing tumor, including GAA (FC

NE/B=13.63x). Overall, these data demonstrate that a specific set of metabolites define the

extracellular microenvironment of enhancing HGGs when compared to non-enhancing glioma

and brain adjacent to tumor.

Plasma in the glioma extracellular metabolome

By definition, contrast-enhancing glioma regions have a more disrupted blood-brain barrier than

non-enhancing regions[29], facilitating solute diffusion between the intravascular compartment

and tumor microenvironment. Our data demonstrated a similar metabolic signature of

enhancing, but not non-enhancing portions, of gliomas when compared to brain (**Fig. 3A, C**),

and metabolic differences between enhancing vs. non-enhancing portions of gliomas (**Fig. 3B,**

4B). As such, we hypothesized that enhancing glioma metabolites may originate from plasma

preferentially entering regions of greater BBB disruption where there is decreased resistance to

diffusion. To test this hypothesis, we utilized a ranked metabolite list of "bloodiness" from

identically paired clean versus bloody CSF samples (see methods). In the 7 patients for whom

paired enhancing and non-enhancing microdialysates were obtained, we first compared the

relative enrichment of enhancing versus non-enhancing brain catheters for bloody versus clean

CSF using each patient's ranked enhancing versus non-enhancing list (Cath. X/Y). Enrichment

analyses demonstrated significant enrichment in 7/7 patients for at least one of (i) enhancing

tumor enriched in bloody CSF, or (ii) non-enhancing tumor enriched in clean CSF; 5/7 patients

showed significant enrichment in both analyses (FDR<0.05) (**Fig. 5A**). Evaluation of specific

enhancing versus non-enhancing tumor-associated metabolites from Fig 4B revealed higher

abundance of bloody CSF-associated metabolites in enhancing tumor (**Fig. 5B**). As expected,

performing enrichment analyses on enhancing glioma versus brain ranked list again demonstrated that enhancing glioma catheters were enriched for bloody CSF in 9/9 patients **(Supplementary Fig. S7).** Results in non-enhancing tumor suggested a modest enrichment of bloody CSF-associated metabolites in some catheters, perhaps suggestive of either some sub-radiographic BBB disruption[29], or metabolite diffusion from adjacent enhancing regions. Collectively, these data suggest that plasma-derived metabolites from a disrupted BBB significantly contribute to the extracellular metabolome of enhancing gliomas **(Fig. 5C)**.

One recent intraoperative study compared metabolites within the glioma's arterial versus venous supply to determine which were consumed versus produced by the gliomas[26]. We analyzed their raw data maintaining thresholds we had used for our own data (FC$\geq$2; p$\leq$0.05), and initially found no significantly differentially abundant metabolites. By removing the FC criterion, 4 significant metabolites were identified. Taurine was more abundant in venous than arterial glioma blood, suggesting production by the tumor. In agreement with this result, we found taurine to be significantly more abundant in both enhancing and non-enhancing tumor as compared to brain (FC E/B= 2.99x, NE/B = 2.08x). Three metabolites were more abundant in glioma arterial than venous blood, suggesting utilization by the tumor **(Sup. Fig. S8).** In agreement with our data, one of these, alanine, was part of our enhancing glioma metabolome (**Table 2**) and enriched for bloody CSF **(Fig. 5B)**, consistent with the hypothesis of BBB disruption improving the glioma's access to needed metabolites. One limitation of the arterial-venous sampling study was that no parallel analyses were performed to compare metabolite utilization and production in adjacent brain. For example, although glucose was identified as utilized by "glioma" in their data, glucose readily crosses easily crosses the BBB and is highly utilized by brain, making fluorodeoxyglucose (FDG)-positron emission tomography (PET) unhelpful to discriminate tumor from brain[30]. Indeed, extracellular glucose was not differentially abundant in our tumor versus brain microdialysate. Nevertheless, these limited available data further support our hypothesis that the enhancing glioma metabolome is significantly enriched

for metabolites from peripheral blood, at least some of which directly support glioma

metabolism. Conversely, extracellular metabolites that are differentially abundant in both

enhancing and non-enhancing tumor when compared to brain are most likely generated by

tumor cells, including GAA and taurine.

DISCUSSION

We utilized an intraoperative window-of opportunity to perform microdialysis, sampling

the extracellular metabolome of radiographically distinct regions of human gliomas *in situ*.

Findings revealed that 1) 2-HG is robustly elevated in IDH-mutant gliomas, with potentially

highest levels in non-enhancing tumor regions. 2) GAA is a previously unreported extracellular

glioma metabolite that highly differentially abundant in enhancing (126.32x) and non-enhancing

(13.63x) glioma as compared to brain; 3) The extracellular metabolome of enhancing glioma is

conserved across patients, comprises largely amino acids and carnitines, and is enriched for

plasma-derived metabolites.

Acute intraoperative microdialysis yielded relative levels of hundreds of metabolites for

untargeted metabolomics analyses, including 2-HG. IDH-mutant gliomas epitomize the

oncogenic relevance of altered metabolism[31,32]. We observed elevated 2-HG in the

microdialysate of IDH-mutant tumors via both targeted (D vs L-2-HG) and untargeted (total D+L-

2HG) analysis (**Fig. 1C; Supplementary Fig. 2)**. Total 2-HG was previously used as a

pharmacodynamic biomarker for IDH inhibitors in tissue[33] and via magnetic resonance

spectroscopy[34-36]. As microdialysis can be utilized for pharmacokinetic studies of dialyzable

drugs[9], the ability to quantify extracellular 2-HG *in vivo* across a >1000x concentration range

(Fig. 1C) suggests that extracellular 2-HG within microdialysate could provide an

pharmacodynamic readout for relevant early phase clinical trials[37]. Interestingly, 2-HG was 5-

12x lower in the most avidly enhancing tumor region (n=3) (**Fig. 1D)**, suggesting that some

tumor metabolites may be lost down a disrupted blood brain barrier—a consideration of

potential importance for longitudinal pharmacodynamic monitoring of tumor interstitial fluid.

Importantly, although IDH-mutations are thought to drive early tumorigenic metabolic and epigenetic changes[32,38], the global extracellular metabolomes of enhancing IDH-mutant versus IDH-WT grade 4 astrocytomas were otherwise indistinguishable (**Fig. 2A, 3A**). Recently published work from over 200 glioma tissues demonstrated less separation between IDH-mutant grade 4 astrocytomas from GBMs[27], than lower grade IDH-mutant gliomas, potentially suggesting metabolic convergence of high-grade gliomas. Greatest similarities were seen across patients in the extracellular metabolome of contrast-enhancing tumors, which could be expected if improved availability of plasma-associated metabolites impacts tumor metabolism.

Interestingly, in each of four patients with non-enhancing oligodendrogliomas the 3 catheters clustered together based on patient identity (**Fig. 2A**). Such close and consistent clustering of catheters within each of these patients could reflect the frequently more diffuse nature of oligodendrogliomas, with each catheter sampling an admixture of metabolically individualized tumor and brain.

Cancer-associated metabolic vulnerabilities are of increasing interest as therapeutic targets. In contrast to gliomas' notorious genomic heterogeneity[39,40], a more focused inventory of metabolic strategies may be utilized to combat bioenergetic demands[41] oxidative stress[42], hypoxia[43], and nutrient deprivation[44,45]. Our rank-based analyses revealed an enhancing glioma extracellular metabolome that was consistent across primary and recurrent IDH-WT and mutant lesions, as well as an H3K27M-mutant tumor (**Fig. 3A(i-ii)**). This signature appeared comparable to that previously obtained in the context of post-operative microdialysis[16,17], consistent with reproducibility of the enhancing glioma metabolome across clinical and technical variables (**Supplementary Table 1**), research teams and contexts spanning intra-operative versus post-operative sampling. However, our data demonstrate for the first time that most of the metabolites in the enhancing glioma extracellular metabolome are absent from the non-enhancing glioma metabolome. These results suggest that the extracellular metabolome of enhancing gliomas may be largely defined by metabolites from systemic circulation that more

easily access the tumor microenvironment through a disrupted BBB. Although catheter insertion may hypothetically result in some degree of BBB disruption, intra-patient catheter comparisons revealed significant enrichment for plasma-associated metabolites, suggesting that the extent BBB disruption induced from catheter insertion was negligible compared to the tumor-induced disruption in enhancing versus non-enhancing tumor. Furthermore, our findings are in alignment with the observation that aggressive portions of gliomas avidly uptake amino acids from the systemic circulation, serving as the basis for highly sensitive and specific glioma PET tracers, including tyrosine[46] and methionine,[47] both of which were more abundant in enhancing glioma than non-enhancing tumor or brain **(Fig. 4; Table 2).**

Given the established importance of multiple identified plasma- and enhancing tumor-associated metabolites, including amino acids and carnitines to cancer biology[48-50], BBB disruption itself may underlie the metabolic similarities across molecularly diverse HGGs by providing a continuous source of protumorigenic metabolites. Anecdotally, our patient with a completely non-enhancing molecular GBM had evidence of this lesion on a CT obtained 5 years prior to diagnosis—consistent with a lower rate of growth than patients with enhancing lesions. Indeed, the adverse prognostic impact of a contrast-enhancement in gliomas, including histologically low-grade tumors, is well recognized[51,52], as is the positive impact on survival of gross total resection of enhancing tumor across all molecularly defined GBM subgroups and ages[53]. As such, our findings suggest that blood brain barrier disruption itself may transform the extracellular glioma metabolome into a wellspring of nutrients for accelerated growth, particularly in enhancing tumor regions.

At this point, we cannot definitively answer the "chicken" or "egg" question of whether BBB disruption occurs for the purpose of accessing systemically-derived analytes or if it is simply a pre-existing process that a glioma can utilize to its advantage. Many of the plasma-associated metabolites enriched in enhancing tumor have been reproducibly associated with cancer biology[54], including the amino acids proline[48], glycine[55], and amino acid derivatives, such

as N6-methyllysine. Similarly, carnitines (**Fig. 4A-B, D(iii-iv)**) have been linked to energy

production via fatty acid oxidation in glioma[56,57]. It is possible that the induction of blood-brain

barrier disruption by proliferating gliomas[58] may represent an adaptive response to nutrient

deprivation observed in gliomas[59-61], catalyzing a feedforward cycle of nutrient utilization and

recruitment. Mechanistic studies selectively manipulating BBB permeability or extracellular

plasma abundance will be required to dissect these interactions. If our hypothesis of disrupted

BBB-mediated tumor aggressiveness is correct, maintaining and restoring BBB integrity may

prove an independently important therapeutic goal, distinct from angiogenesis-blocking agents

such as bevacizumab[62].

Although we used UPLC-MS/MS to maximize our yield of potentially novel glioma

metabolites, we had not expected to find a new extracellular glioma metabolite that could

outperform all other enhancing glioma metabolites by over 10-fold. Although GAA was present

at a low abundance in bloody CSF and not detected in clean CSF, its disproportionately high

fold change in both enhancing and non-enhancing tumor versus brain (126.32x, and 13.63x,

respectively; Figure 4A, D(i), **Table 2**) suggest it may be produced by the tumor. No CNS efflux

transporter exists for GAA at the BBB[63]. As such, if generated as a result of glioma metabolism,

it is liable to accumulate. GAA also accumulates in patients with GAMT deficiency, leading to

impaired creatine production, seizures, and cognitive impairment[64]. GAA is a precursor for

creatine, levels of which were slightly elevated in non-enhancing (2.25x; p=0.03), but not

enhancing tumor (**Table 2**). Interestingly, GAA is co-produced with ornithine, which was slightly

higher in enhancing tumor when compared to non-enhancing tumor (2.36x, p=0.03) and brain

(3.07x, p=0.004) (**Table 2**). Ornithine is the substrate for polyamine synthesis via ornithine

decarboxylase (ODC) which is upregulated in cancers, including glioblastoma, to support

catabolic demand, regulate intracellular pH in the acidic tumor core, and protect against immune

surveillance[65]. It is therefore tempting to speculate that increased GAA could be a byproduct of

increased glioma polyamine synthesis and utilization. If true and based on the substantial fold-

change, extracellular GAA could serve as a biomarker of not only local glioma abundance but also pharmacodynamic response to ODC blockade. ODC blockade with Difluoromethylornithine (DFMO) remains in clinical trials for glioma (NCT02796261); though upregulation of polyamine transporters may mediate metabolic compensation leading to improved survival with dual blockade of ODC and polyamine transporters (via AMXT-1501), in preclinical models of diffuse midline gliomas and other cancers[66,67]. To determine if GAA can be used as a dynamic biomarker of ODC production *in situ*, we are opening a post-operative microdialysis study under an IND, monitoring microdialysate levels of GAA in unresectable regions of residual tumor with or without DFMO +/- AMXT1501. Further dynamic insights may be achievable with isotope tracing and are under development.

Conclusions

With the help of our patients, and intraoperative microdialysis we have shown that GAA is surprisingly abundant in the glioma microenvironment, and that plasma-associated metabolites largely define the enhancing glioma microenvironment, potentially positioning them to support glioma aggressiveness. Our patients understand the formidable odds of their disease. We present their data as a first installment toward what we intend to be a growing collaborative, open-source repository of individualized data from live human gliomas. We (both patients and investigators) hope that the data and insights shared can help inspire novel translational paradigms. By leveraging neurosurgical access to live human disease, we hope to accelerate tangible progress within the lifetimes of individual patients.

List of Abbreviations

2-HG: 2-hydroxyglutarate

2-PY: N1-methyl-2-pyridone-5-carboxamide

B: brain

BBB: Blood-brain barrier

CNS: central nervous system

CSF: cerebrospinal fluid

CT: computed tomography

DFMO: difluoromethylornithine

E: enhancing tumor

FC: fold-change

FDR: false-discovery rate

GAA: guanidinoacetate

GBM: glioblastoma

GSEA: gene set enrichment analysis

HGG: high-grade glioma

HMW: high molecular weight

IDH: isocitrate dehydrogenase

IND: investigational new drug

LC/MS: liquid chromatography-mass spectrometry

NAA: N-acetylaspartate

NE: non-enhancing tumor

ODC: ornithine decarboxylase

WT: wild type

UPLC-MS/MS: Ultra-performance liquid chromatography tandem mass spectrometry

References

1   Stupp, R. *et al.* Radiotherapy plus concomitant and adjuvant temozolomide for
glioblastoma. *The New England journal of medicine* **352**, 987-996,
doi:10.1056/NEJMoa043330 (2005).
2   Yan, H. *et al.* IDH1 and IDH2 mutations in gliomas. *The New England journal of medicine*
360, 765-773, doi:10.1056/NEJMoa0808710 (2009).

SongTao, Q. *et al.* IDH mutations predict longer survival and response to temozolomide in secondary glioblastoma. *Cancer Sci* **103**, 269-273, doi:10.1111/j.1349-7006.2011.02134.x (2012).

Shibao, S. *et al.* Metabolic heterogeneity and plasticity of glioma stem cells in a mouse glioblastoma model. *Neuro Oncol* **20**, 343-354, doi:10.1093/neuonc/nox170 (2018).

Kondo, Y., Katsushima, K., Ohka, F., Natsume, A. & Shinjo, K. Epigenetic dysregulation in glioma. *Cancer Sci* **105**, 363-369, doi:10.1111/cas.12379 (2014).

Vlashi, E. *et al.* Metabolic state of glioma stem cells and nontumorigenic cells. *Proc Natl Acad Sci U S A* **108**, 16062-16067, doi:10.1073/pnas.1106704108 (2011).

Strickland, M. & Stoll, E. A. Metabolic Reprogramming in Glioma. *Front Cell Dev Biol* **5**, 43, doi:10.3389/fcell.2017.00043 (2017).

Lynes, J. *et al.* Cytokine Microdialysis for Real-Time Immune Monitoring in Glioblastoma Patients Undergoing Checkpoint Blockade. *Neurosurgery* **84**, 945-953, doi:10.1093/neuros/nyy392 (2019).

Portnow, J. *et al.* A pilot microdialysis study in brain tumor patients to assess changes in intracerebral cytokine levels after craniotomy and in response to treatment with a targeted anti-cancer agent. *J Neurooncol* **118**, 169-177, doi:10.1007/s11060-014-1415-4 (2014).

Blakeley, J. O. *et al.* Effect of blood brain barrier permeability in recurrent high grade gliomas on the intratumoral pharmacokinetics of methotrexate: a microdialysis study. *J Neurooncol* **91**, 51-58, doi:10.1007/s11060-008-9678-2 (2009).

Marcus, H. J., Carpenter, K. L. H., Price, S. J. & Hutchinson, P. J. In vivo assessment of high-grade glioma biochemistry using microdialysis: a study of energy-related molecules, growth factors and cytokines. *Journal of Neuro-Oncology* **97**, 11-23, doi:10.1007/s11060-009-9990-5 (2010).

Roslin, M., Henriksson, R., Bergstrom, P., Ungerstedt, U. & Bergenheim, A. T. Baseline levels of glucose metabolites, glutamate and glycerol in malignant glioma assessed by stereotactic microdialysis. *J Neurooncol* **61**, 151-160 (2003).

Melani, A. *et al.* Adenosine extracellular levels in human brain gliomas: an intraoperative microdialysis study. *Neuroscience letters* **346**, 93-96 (2003).

Bianchi, L. *et al.* Extracellular levels of amino acids and choline in human high grade gliomas: an intraoperative microdialysis study. *Neurochemical research* **29**, 325-334 (2004).

Homapour, B. *et al.* Intra-operative, real-time, three-dimensional ultrasound assisted positioning of catheters in the microdialysis of glial tumours. *Journal of Clinical Neuroscience* **17**, 506-510, doi:https://doi.org/10.1016/j.jocn.2009.06.022 (2010).

Wibom, C. *et al.* Metabolomic patterns in glioblastoma and changes during radiotherapy: a clinical microdialysis study. *J Proteome Res* **9**, 2909-2919, doi:10.1021/pr901088r (2010).

Bjorkblom, B. *et al.* Metabolic response patterns in brain microdialysis fluids and serum during interstitial cisplatin treatment of high-grade glioma. *Br J Cancer* **122**, 221-232, doi:10.1038/s41416-019-0652-x (2020).

Louis, D. N. *et al.* The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro-Oncology* **23**, 1231-1251, doi:10.1093/neuonc/noab106 (2021).

Oldham, W. M. & Loscalzo, J. Quantification of 2-Hydroxyglutarate Enantiomers by Liquid Chromatography-mass Spectrometry. *Bio Protoc* **6**, doi:10.21769/BioProtoc.1908 (2016).

Struys, E. A., Jansen, E. E., Verhoeven, N. M. & Jakobs, C. Measurement of urinary D- and L-2-hydroxyglutarate enantiomers by stable-isotope-dilution liquid chromatography-tandem mass spectrometry after derivatization with diacetyl-L-tartaric anhydride. *Clin Chem* **50**, 1391-1395, doi:10.1373/clinchem.2004.033399 (2004).

Cheng, Q. Y. *et al.* Sensitive Determination of Onco-metabolites of D- and L-2-hydroxyglutarate Enantiomers by Chiral Derivatization Combined with Liquid Chromatography/Mass Spectrometry Analysis. *Sci Rep* **5**, 15217, doi:10.1038/srep15217 (2015).

Pang, Z. *et al.* MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional insights. *Nucleic Acids Res* **49**, W388-w396, doi:10.1093/nar/gkab382 (2021).

Waitkus, M. S., Diplas, B. H. & Yan, H. Isocitrate dehydrogenase mutations in gliomas. *Neuro-Oncology* **18**, 16-26, doi:10.1093/neuonc/nov136 (2015).

Ryals, J., Lawton, K., Stevens, D. & Milburn, M. Metabolon, Inc. (2007).

Pope, W. B. *et al.* Non-invasive detection of 2-hydroxyglutarate and other metabolites in IDH1 mutant glioma patients using magnetic resonance spectroscopy. *J Neurooncol* **107**, 197-205, doi:10.1007/s11060-011-0737-8 (2012).

Xiong, N. *et al.* Using arterial-venous analysis to characterize cancer metabolic consumption in patients. *Nat Commun* **11**, 3169, doi:10.1038/s41467-020-16810-8 (2020).

Björkblom, B. *et al.* Distinct metabolic hallmarks of WHO classified adult glioma subtypes. *Neuro-Oncology* **24**, 1454-1468, doi:10.1093/neuonc/noac042 (2022).

Moffett, J. R., Ross, B., Arun, P., Madhavarao, C. N. & Namboodiri, A. M. N-Acetylaspartate in the CNS: from neurodiagnostics to neurobiology. *Prog Neurobiol* **81**, 89-131, doi:10.1016/j.pneurobio.2006.12.003 (2007).

Sarkaria, J. N. *et al.* Is the blood-brain barrier really disrupted in all glioblastomas? A critical assessment of existing clinical data. *Neuro Oncol* **20**, 184-191, doi:10.1093/neuonc/nox175 (2018).

Heiss, W.-D., Raab, P. & Lanfermann, H. Multimodality assessment of brain tumors and tumor recurrence. *Journal of Nuclear Medicine* **52**, 1585-1600 (2011).

Dang, L. *et al.* Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature* **462**, 739-744, doi:10.1038/nature08617 (2009).

McBrayer, S. K. *et al.* Transaminase Inhibition by 2-Hydroxyglutarate Impairs Glutamate Biosynthesis and Redox Homeostasis in Glioma. *Cell* **175**, 101-116.e125, doi:10.1016/j.cell.2018.08.038 (2018).

Konteatis, Z. *et al.* Vorasidenib (AG-881): A First-in-Class, Brain-Penetrant Dual Inhibitor of Mutant IDH1 and 2 for Treatment of Glioma. *ACS Med Chem Lett* **11**, 101-107, doi:10.1021/acsmedchemlett.9b00509 (2020).

Molloy, A. R. *et al.* MR-detectable metabolic biomarkers of response to mutant IDH inhibition in low-grade glioma. *Theranostics* **10**, 8757-8770, doi:10.7150/thno.47317 (2020).

Andronesi, O. C. *et al.* Pharmacodynamics of mutant-IDH1 inhibitors in glioma patients probed by in vivo 3D MRS imaging of 2-hydroxyglutarate. *Nat Commun* **9**, 1474, doi:10.1038/s41467-018-03905-6 (2018).

Wenger, K. J. *et al.* Non-Invasive Measurement of Drug and 2-HG Signals Using (19)F and (1)H MR Spectroscopy in Brain Tumors Treated with the Mutant IDH1 Inhibitor BAY1436032. *Cancers (Basel)* **12**, doi:10.3390/cancers12113175 (2020).

Mellinghoff, I. K. *et al.* Ivosidenib in Isocitrate Dehydrogenase 1–Mutated Advanced Glioma. *Journal of Clinical Oncology* **38**, 3398-3406, doi:10.1200/jco.19.03327 (2020).

Turcan, S. *et al.* IDH1 mutation is sufficient to establish the glioma hypermethylator phenotype. *Nature* **483**, 479-483, doi:10.1038/nature10866 (2012).

Zhang, Y. *et al.* Elevated signature of a gene module coexpressed with CDC20 marks genomic instability in glioma. *Proceedings of the National Academy of Sciences* **116**, 6975-6984, doi:10.1073/pnas.1814060116 (2019).

Milinkovic, V. *et al.* Genomic instability and p53 alterations in patients with malignant glioma. *Exp Mol Pathol* **93**, 200-206, doi:10.1016/j.yexmp.2012.05.010 (2012).

Bi, J. *et al.* Altered cellular metabolism in gliomas — an emerging landscape of actionable co-dependency targets. *Nature Reviews Cancer* **20**, 57-70, doi:10.1038/s41568-019-0226-5 (2020).

Huang, H. *et al.* Suppression of mitochondrial ROS by prohibitin drives glioblastoma progression and therapeutic resistance. *Nature Communications* **12**, 3720, doi:10.1038/s41467-021-24108-6 (2021).

Steinbach, J. P., Wolburg, H., Klumpp, A., Probst, H. & Weller, M. Hypoxia-induced cell death in human malignant glioma cells: energy deprivation promotes decoupling of mitochondrial cytochrome c release from caspase processing and necrotic cell death. *Cell Death & Differentiation* **10**, 823-832, doi:10.1038/sj.cdd.4401252 (2003).

Tanaka, K. *et al.* Glioma cells require one-carbon metabolism to survive glutamine starvation. *Acta Neuropathologica Communications* **9**, 16, doi:10.1186/s40478-020-01114-1 (2021).

Mondal, S., Bhattacharya, K. & Mandal, C. Nutritional stress reprograms dedifferention in glioblastoma multiforme driven by PTEN/Wnt/Hedgehog axis: a stochastic model of cancer stem cells. *Cell Death Discovery* **4**, 110, doi:10.1038/s41420-018-0126-6 (2018).

Dunet, V., Rossier, C., Buck, A., Stupp, R. & Prior, J. O. Performance of 18F-fluoro-ethyl-tyrosine (18F-FET) PET for the differential diagnosis of primary brain tumor: a systematic review and Metaanalysis. *J Nucl Med* **53**, 207-214, doi:10.2967/jnumed.111.096859 (2012).

Van Laere, K. *et al.* Direct comparison of 18F-FDG and 11C-methionine PET in suspected recurrence of glioma: sensitivity, inter-observer variability and prognostic value. *European Journal of Nuclear Medicine and Molecular Imaging* **32**, 39-51, doi:10.1007/s00259-004-1564-3 (2005).

Burke, L. *et al.* The Janus-like role of proline metabolism in cancer. *Cell Death Discovery* **6**, 104, doi:10.1038/s41420-020-00341-8 (2020).

Sanderson, S. M., Gao, X., Dai, Z. & Locasale, J. W. Methionine metabolism in health and cancer: a nexus of diet and precision medicine. *Nature Reviews Cancer* **19**, 625-637, doi:10.1038/s41568-019-0187-8 (2019).

Melone, M. A. B. *et al.* The carnitine system and cancer metabolic plasticity. *Cell Death & Disease* **9**, 228, doi:10.1038/s41419-018-0313-7 (2018).

Pallud, J. *et al.* Prognostic significance of imaging contrast enhancement for WHO grade II gliomas. *Neuro Oncol* **11**, 176-182, doi:10.1215/15228517-2008-066 (2009).

Wang, Y. Y. *et al.* Patterns of Tumor Contrast Enhancement Predict the Prognosis of Anaplastic Gliomas with IDH1 Mutation. *AJNR Am J Neuroradiol* **36**, 2023-2029, doi:10.3174/ajnr.A4407 (2015).

Molinaro, A. M. *et al.* Association of Maximal Extent of Resection of Contrast-Enhanced and Non-Contrast-Enhanced Tumor With Survival Within Molecular Subgroups of Patients With Newly Diagnosed Glioblastoma. *JAMA Oncol* **6**, 495-503, doi:10.1001/jamaoncol.2019.6143 (2020).

Lieu, E. L., Nguyen, T., Rhyne, S. & Kim, J. Amino acids in cancer. *Experimental & Molecular Medicine* **52**, 15-30, doi:10.1038/s12276-020-0375-3 (2020).

Tiwari, V. *et al.* Glycine by MR spectroscopy is an imaging biomarker of glioma aggressiveness. *Neuro Oncol* **22**, 1018-1029, doi:10.1093/neuonc/noaa034 (2020).

Fink, M. A. *et al.* L-Carnitine-Mediated Tumor Cell Protection and Poor Patient Survival Associated with OCTN2 Overexpression in Glioblastoma Multiforme. *Clin Cancer Res* **25**, 2874-2886, doi:10.1158/1078-0432.Ccr-18-2380 (2019).

Juraszek, B., Czarnecka-Herok, J. & Nałęcz, K. A. Glioma cells survival depends both on fatty acid oxidation and on functional carnitine transport by SLC22A5. *Journal of Neurochemistry* **156**, 642-657, doi:https://doi.org/10.1111/jnc.15124 (2021).

Arvanitis, C. D., Ferraro, G. B. & Jain, R. K. The blood-brain barrier and blood-tumour barrier in brain tumours and metastases. *Nat Rev Cancer* **20**, 26-41, doi:10.1038/s41568-019-0205-x (2020).

Panosyan, E. H. *et al.* Asparagine depletion potentiates the cytotoxic effect of chemotherapy against brain tumors. *Mol Cancer Res* **12**, 694-702, doi:10.1158/1541-7786.Mcr-13-0576 (2014).

Yamashita, D. *et al.* Targeting glioma-initiating cells via the tyrosine metabolic pathway. *J Neurosurg* **134**, 721-732, doi:10.3171/2019.11.Jns192028 (2020).

Friedrich, M. *et al.* Tryptophan metabolism drives dynamic immunosuppressive myeloid states in IDH-mutant gliomas. *Nature Cancer* **2**, 723-740, doi:10.1038/s43018-021-00201-z (2021).

Gilbert, M. R. *et al.* A Randomized Trial of Bevacizumab for Newly Diagnosed Glioblastoma. *New England Journal of Medicine* **370**, 699-708, doi:10.1056/NEJMoa1308573 (2014).

Tachikawa, M. & Hosoya, K.-i. Transport characteristics of guanidino compounds at the blood-brain barrier and blood-cerebrospinal fluid barrier: relevance to neural disorders. *Fluids and Barriers of the CNS* **8**, 13, doi:10.1186/2045-8118-8-13 (2011).

Gordon, N. Guanidinoacetate methyltransferase deficiency (GAMT). *Brain Dev* **32**, 79-81, doi:10.1016/j.braindev.2009.01.008 (2010).

Miska, J. *et al.* Polyamines drive myeloid cell survival by buffering intracellular pH to promote immunosuppression in glioblastoma. *Sci Adv* **7**, doi:10.1126/sciadv.abc8929 (2021).

Levin, V. A., Ictech, S. E. & Hess, K. R. Clinical importance of eflornithine (alpha-difluoromethylornithine) for the treatment of malignant gliomas. *CNS Oncol* **7**, CNS16, doi:10.2217/cns-2017-0031 (2018).

Khan, A. *et al.* Dual targeting of polyamine synthesis and uptake in diffuse intrinsic pontine gliomas. *Nat Commun* **12**, 971, doi:10.1038/s41467-021-20896-z (2021).

Tables

Table 1. Intraoperative microdialysis patient characteristics.

The patient IDs and final pathologic diagnoses, including IDH- status and other available key molecular features, are summarized together with the catheter locations. Additional specifics regarding each patient, tumor, and catheter are available in Supplemental Table 1. *Diagnosis of IDH-WT GBM was reached for patient GBM^WT2/2B without need for further molecular studies.

1 †: Region of minimal enhancement, suggestive of tumor necrosis.

Patient ID	Pathology	IDH Status	Other molecular/genetic features	Catheter X	Catheter Y	Catheter Z
Oligo[2]	Oligodendroglioma WHO 2	Mutant	1p/19q co-deletion; TERT-mutant	Non-enhancing (NE)	Non-enhancing (NE)	Non-enhancing (NE)
Oligo[3]1	Oligodendroglioma WHO 3	Mutant	1p/19q co-deletion; TERT-mutant	Non-enhancing (NE)	Non-enhancing (NE)	Brain adjacent to tumor (B)
Oligo[3]2	Oligodendroglioma WHO 3	Mutant	1p/19q co-deletion; TERT-mutant	Non-enhancing (NE)	Non-enhancing (NE)	Brain adjacent to tumor (B)
Oligo[3]3	Oligodendroglioma WHO 3	Mutant	1p/19q co-deletion; TERT-mutant	Non-enhancing (NE)	Non-enhancing (NE)	Brain adjacent to tumor (B)
GemAstro[3-mut]	Gemistocytic Astrocytoma WHO 3	Mutant	None remarkable	Non-enhancing (NE)	Non-enhancing (NE)	Brain adjacent to tumor (B)
Astro[4-mut]1	Astrocytoma WHO 4 (Recurrent)	Mutant	MGMT non-methylated; loss of CDKN2A/B; complex karyotype	Enhancing (E)	Non-enhancing (NE)	Brain adjacent to tumor (B)
Astro[4-mut]2	Astrocytoma WHO 4	Mutant	MGMT non-methylated; loss of CDKN2A/B	Enhancing (E)	Non-enhancing (NE)	Brain adjacent to tumor (B)
Astro[4-mut]3	Astrocytoma WHO 4 (Recurrent)	Mutant	Indeterminate MGMT methylation	Enhancing (E)	Enhancing† (E†)	Brain adjacent to tumor (B)
H3K27M-Astro[4-WT]	Infiltrating Astrocytoma WHO 4	Wild-type	H3K27M, TERT-mutant; MGMT non-methylated	Enhancing (E)	Non-enhancing (NE)	Brain adjacent to tumor (B)
GBM[WT]1	Glioblastoma	Wild-type	MGMT non-methylated; loss of CDKN2A/B; TERT-mutant	Enhancing (E)	Non-enhancing (NE)	Brain adjacent to tumor (B)
GBM[WT]2	Glioblastoma (Primary)	Wild-type	Indeterminate MGMT methylation*	Enhancing (E)	Non-enhancing (NE)	Brain adjacent to tumor (B)
GBM[WT]2B	Glioblastoma (Recurrent)	Wild-type	Indeterminate MGMT methylation*	Enhancing (E)		Brain adjacent to tumor (B)
GBM[WT]3	Glioblastoma	Wild-type	MGMT non-methylated	Enhancing (E)	Non-enhancing (NE)	Brain adjacent to tumor (B)
GBM[WT]4	Glioblastoma (Recurrent)	Wild-type	MGMT non-methylated; EGFR amplification; loss of CDKN2A/B	Enhancing (E)	Non-enhancing (NE)	Brain adjacent to tumor (B)
MolecGBM[WT]	Molecular Glioblastoma	Wild-type	Gain of ch. 7, loss of ch. 10; TERT-mutant; MGMT non-methylated	Non-enhancing (NE)	Non-enhancing (NE)	Brain adjacent to tumor (B)

3 **Table 2.** Significantly altered metabolites across enhancing, non-enhancing, and brain

4 microenvironments in paired patient specimens (n=9 for E/B and n=7 for E/NE or NE/B). *bold =

5 $p < 0.05$.

Metabolite	FC (E/B)	P-value	FC (E/NE)	P-value	FC (NE/B)	P-value
Guanidinoacetate (GAA)	**126.32**	0.0039	9.76	0.2188	**13.63**	0.0469
proline	**10.18**	0.0039	**4.46**	0.0313	2.09	0.1094

deoxycarnitine	**8.71**	0.0039	**4.55**	0.0156	1.94	0.3750
carnitine	**7.65**	0.0039	**3.32**	0.0313	2.18	0.0781
acetylcarnitine (C2)	**7.47**	0.0039	2.76	0.1094	**2.97**	0.0469
N1-methyl-2-pyridone-5-carboxamide (2-PY)	**7.18**	0.0391	**3.20**	0.0156	1.28	0.6875
N6-methyllysine	**7.17**	0.0039	**4.67**	0.0313	1.16	0.2969
methionine	**7.09**	0.0039	**2.94**	0.0156	2.41	0.0781
dimethylglycine	**6.90**	0.0039	**3.75**	0.0156	1.51	0.4688
asparagine	**6.84**	0.0039	**3.64**	0.0156	1.55	0.3750
propionylcarnitine (C3)	**6.78**	0.0078	**3.77**	0.0469	1.84	0.1094
alanine	**6.54**	0.0039	**3.11**	0.0156	1.96	0.0781
valine	**6.01**	0.0039	**3.24**	0.0156	1.70	0.1563
tyrosine	**5.66**	0.0039	**3.14**	0.0313	1.62	0.2969
glycine	**5.66**	0.0078	2.43	0.2188	2.18	0.1094
imidazole lactate	**5.57**	0.0039	**3.78**	0.0156	1.09	0.5781
urate	**5.05**	0.0039	**2.63**	0.0156	1.60	0.0781
lysine	**4.54**	0.0039	**2.68**	0.0156	1.41	0.2969
allantoin	**4.52**	0.0039	**3.96**	0.0313	0.90	0.5781
histidine	**4.49**	0.0039	**2.30**	0.0313	**1.73**	0.0156
phenylalanine	**4.48**	0.0039	**2.59**	0.0156	1.52	0.0781
trigonelline (N'-methylnicotinate)	**4.40**	0.0078	1.75	0.1563	2.57	0.2969
3-hydroxybutyrate (BHBA)	**4.40**	0.0039	**2.22**	0.0313	**1.77**	0.0313
3-methyl-2-oxobutyrate	**4.34**	0.0039	3.86	0.0781	0.80	0.5781
hydroxyproline	**4.22**	0.0039	**2.11**	0.0313	1.48	0.2188
threonine	**4.20**	0.0039	**2.27**	0.0313	1.58	0.1563
leucine	**4.12**	0.0039	**2.43**	0.0156	1.54	0.0781
tryptophan	**3.96**	0.0039	**2.33**	0.0313	1.40	0.2969
N1-methyl-4-pyridone-3-carboxamide (4-PY)	3.88	0.0742	**2.16**	0.0156	1.27	0.5781
N,N,N-trimethyl-5-aminovalerate	**3.88**	0.0039	2.17	0.0781	1.74	0.0781
isoleucine	**3.76**	0.0039	**2.29**	0.0156	1.52	0.0781
guanine	**3.73**	0.0078	1.24	0.5781	**2.62**	0.0469
N,N,N-trimethyl-alanylproline betaine (TMAP)	**3.50**	0.0039	1.59	0.1094	1.58	0.2969
p-cresol sulfate	**3.36**	0.0039	2.03	0.2969	1.44	0.2188
Glutamine degradant	**3.24**	0.0078	**1.88**	0.0156	1.61	0.0781

arginine	**3.16**	0.0039	**2.17**	0.0156	1.25	0.6875
Lactate*	**3.12**	0.0313	1.34	0.0625	2.32	0.0625
ornithine	**3.07**	0.0039	**2.36**	0.0313	1.13	1.0000
taurine	**2.99**	0.0117	1.76	0.1094	**2.08**	0.0156
thioproline	**2.97**	0.0078	1.31	0.0781	2.50	0.0781
gamma-glutamylglutamine	**2.93**	0.0117	**1.45**	0.0469	**2.03**	0.0313
serine	**2.80**	0.0039	**2.12**	0.0469	1.12	0.9375
N6,N6,N6-trimethyllysine	**2.62**	0.0195	1.80	0.0781	1.27	0.2188
glycerophosphorylcholine (GPC)	2.60	0.0742	0.87	1.0000	**4.32**	0.0156
phenol sulfate	**2.59**	0.0078	2.08	0.0781	1.09	0.2188
N-acetylputrescine	**2.44**	0.0273	1.30	0.8125	1.75	0.4688
ergothioneine	**2.41**	0.0391	0.60	0.3750	**3.26**	0.0156
3-hydroxyisobutyrate	**2.27**	0.0391	1.51	0.3750	1.15	0.3750
stachydrine	**2.24**	0.0078	1.45	0.2188	1.39	0.0781
1-methyl-5-imidazolelactate	2.13	0.1073	0.56	0.8125	**4.27**	0.0156
7-methylguanine	**2.10**	0.0195	1.53	0.3750	1.40	0.2945
N-acetylglycine	**2.05**	0.0039	1.71	0.1563	1.10	0.2188
glycerophosphoglycerol	2.04	0.2340	0.45	0.4688	**5.21**	0.0156
malate	1.87	0.1641	0.70	0.9375	**2.99**	0.0156
phosphoethanolamine (PE)	1.56	0.6523	0.49	0.5781	**4.83**	0.0156
creatine	1.05	0.5703	0.55	0.2188	**2.25**	0.0313
glycerol-3-phosphate	1.02	0.5703	0.40	0.3750	**3.25**	0.0313
phosphate	0.87	1.0000	**0.33**	0.0313	**2.91**	0.0469
glycerophosphoethanolamine	0.82	0.8203	0.43	0.1563	**2.83**	0.0156
inosine	0.64	0.4258	**0.48**	0.0156	1.60	0.1563
myo-inositol	0.51	0.4258	**0.46**	0.0469	1.13	0.5781
orotate	0.47	0.1641	**0.39**	0.0313	0.98	0.8125
N-formylmethionine	0.42	0.1289	**0.36**	0.0469	1.13	0.6875
aspartate	**0.40**	0.0195	0.43	0.2188	1.06	0.6875
lyxonate	**0.39**	0.0078	0.47	0.0781	0.86	0.9375
N-acetylasparagine	**0.37**	0.0039	**0.37**	0.0469	0.86	0.6875
3-hydroxy-3-methylglutarate	**0.37**	0.0391	0.29	0.0781	1.55	0.1563
N-acetylglutamine	**0.34**	0.0078	**0.34**	0.0156	0.99	0.6875
N-acetylthreonine	**0.34**	0.0078	**0.34**	0.0469	0.93	1.0000
N-acetyl-beta-alanine	**0.30**	0.0391	0.45	0.1422	0.73	0.6875

N-acetylalanine	**0.30**	0.0391	**0.25**	0.0156	0.96	0.9375
2-methylcitrate/homocitrate	**0.29**	0.0039	**0.27**	0.0156	0.94	0.6875
N-acetylneuraminate	**0.25**	0.0039	0.32	0.0781	0.75	0.9375
fructose	**0.25**	0.0039	**0.40**	0.0469	0.56	0.4688
N1-methylinosine	**0.23**	0.0078	**0.20**	0.0313	1.09	0.2969
N-acetylserine	**0.21**	0.0039	**0.23**	0.0313	0.87	0.6875
Erythronate	**0.18**	0.0039	**0.21**	0.0156	0.83	1.0000
N-acetylaspartate (NAA)	**0.17**	0.0039	**0.16**	0.0156	1.18	0.5781
N-acetyl-aspartyl-glutamate (NAAG)	**0.16**	0.0078	0.19	0.1563	0.73	0.6875
4-acetamidobutanoate	**0.15**	0.0273	0.28	0.1563	0.96	0.9375
2,3-dihydroxy-5-methylthio-4-pentenoate (DMTPA)	**0.15**	0.0039	**0.24**	0.0156	0.55	0.0781
2-O-methylascorbic acid (2-O-MA)	**0.14**	0.0039	**0.15**	0.0156	0.83	0.8125
ribonate	**0.11**	0.0039	**0.13**	0.0156	0.82	0.9375
arabonate/xylonate	**0.08**	0.0039	**0.14**	0.0156	0.70	0.4688
arabitol/xylitol	**0.04**	0.0039	**0.08**	0.0156	0.57	0.3750

1   *Fold-change and p-value for lactate are based on n=7 cases, rather than n=9, as the perfusate

2   solution contained lactate in 2 cases.

10   **Figure Legends**

11   **Figure 1. Intraoperative microdialysis set-up and D-2-hydroxyglutarate (D-2-HG)**

12   **detection.**

(A) A representative intraoperative photograph (Oligo[2]) demonstrating the experimental setup,

including placement of three 100 kDA microdialysis catheters, as well as pumps (X, Y, Z) and

collection vials within the surgical field.

(B) Illustrative intraoperative trajectory views captured from the Neuronavigation system (Astro[4-

[mut]1) demonstrate the planned trajectory (blue line) toward the intended target location for each

10 mm microdialysis membrane (indicated by yellow box) in enhancing (E, Catheter X) and non-

enhancing tumor (NE, Catheter Y), in addition to relatively normal brain adjacent to tumor (B,

Catheter Z).

(**C**) 2-HG peak areas from UPLC-MS/MS were measured in the microdialysate from each of the

44 catheters. IDH status is indicated as mutant (blue) or wild type (red). Symbol shape indicates

catheter placement location (see legend).

(D) The 2-HG peak areas from three patients with enhancing grade 4 IDH-mutant astrocytomas

are shown in closer detail.

Figure 2. Microdialysate samples cluster based on patient identity and catheter location.

(A) Spearman correlation map of the 44 catheters from all 15 microdialysis cases (14 patients)**.**

Dot colors indicate catheter location, IDH status, and primary or recurrent status. (Minimal

correlation: -0.7, maximal correlation: 1).

(B) Hierarchical clustering heat map (Ward method, Euclidean distance measure) of the 44

catheters based on 162 metabolites present in at least 90% (40/44) catheters. (Normalized

abundance by MetaboAnalyst: minimum, -6 to maximum, 6).

Figure 3. Metabolic signatures of enhancing versus non-enhancing tumors versus brain.

(**A**) (i) 162 metabolites present in at least 40/44 catheters were ranked according to the

enhancing tumor-versus-brain fold change in each patient. The rank order of each metabolite in

each 2-catheter tumor/brain comparison (e.g., catheter X versus Z) is conveyed as a heat map

from 1 (orange, enhancing tumor) to 162 (green, brain). Metabolites are listed based on the

average of ranks of enhancing vs. brain across nine cases. (ii) Normalized enrichment scores

(NES) are shown based upon enrichment analysis for the top 35 enhancing tumor (E) and brain

(B)-associated metabolites in the ranked metabolite of enhancing tumor vs. brain (Cath. X/Z)

(bold: significant at FDR<0.05; Complete FDR details available in Supplemental Table 2). See

Supplemental Figure 1 for graphical depiction of how enrichment analyses are performed by

repurposing the Gene Set Enrichment Analysis software.

(**B**) Using the same method as in 3A(i), the rank order of each metabolite in enhancing-versus-

non-enhancing catheter metabolite lists (Catheter X and Y) is depicted as a heat map in the 7

patients for whom both enhancing and non-enhancing catheters were present. Metabolites are

listed based on the average of ranks of enhancing versus non-enhancing tumor across seven

patients. (ii) As in 3A(ii), using enhancing versus non-enhancing tumor metabolite ranked lists

and metabolite sets. NES are shown for the enrichment of each E vs. NE ranked list for other E

vs. NE metabolite sets (bold: FDR<0.05; Complete FDR details available in Supplemental table

3).

(**C**) (i) As in 3A-B, the ranked order of metabolites in non-enhancing versus brain metabolite lists

is shown as a heat map. Metabolites are ranked based on the average rank of NE tumor to

brain across 7 patients. (ii) As in prior enrichment analyses, with non-enhancing versus brain

ranked lists and metabolite sets and their respective NESs across patients (bold: FDR<0.05;

Complete FDR details available in Supplemental table 4).

Figure 4. Differentially abundant extracellular metabolites in the enhancing glioma, non-

enhancing glioma, and adjacent brain comparisons.

Wilcoxon signed-rank tests and fold-changes were calculated between (A) enhancing tumor vs.

brain (n=9 paired catheters), (B) enhancing versus non-enhancing tumor (n=7 paired catheters),

and (D) non-enhancing tumor versus brain (n=7 paired catheters) were utilized to construct a

volcano plot (cut-offs for significance: p-value≤0.05; FC≥2). The normalized peak areas for the

top four most differentially abundant extracellular metabolites in enhancing tumor vs. brain are

shown in (**C)** for the 9 cases, including the peak area in non-enhancing tumor for the seven

patients for whom this catheter was available; colors correlate with patient identity from Table 1.

2-O-MA= 2-O-methylascorbic acid; DMTPA = 2,3-dihydroxy-5-methylthio-4-pentenoic acid; G-3-

P=glycerol-3-phosphate; GAA = guanidinoacetate; GPC = glycerophosphocholine; GPG =

glycerophosphorylglycerol; NAA = n-acetylaspartate; NAAG = n-acetylaspartylglutamate; NANA

= n-acetylneuraminate.

Figure 5. Enhancing glioma microdialysate is enriched for plasma-derived metabolites.

(A) Enrichment analysis was utilized to determine the enrichment of each patient's enhancing

vs. non-enhancing tumor catheters (ranked list: Cath. X vs. Y) for high-plasma versus low-

plasma CSF (metabolite sets). Positive normalized enrichment scores (NES) indicate metabolic

similarities to high-plasma CSF; negative NES indicate metabolic similarities to low-plasma CSF

(* = FDR≤0.05). (**B)** The rank of each metabolite from the significant enhancing versus non-

enhancing glioma metabolite in Figure 4B is shown for each patient (E: purple, rank 1, to NE:

orange, rank 162), along with the metabolite's rank in a bloody vs. clean CSF ranked fold-

change list (bloody: red, 1 – clean: blue, 157). (**C)** Proposed model depicting the impact of

blood-brain barrier disruption on the enhancing versus non-enhancing glioma extracellular

metabolomes (created using BioRender).

A

B Astro$^{4\text{-mut}}$1

Catheter X: Enhancing (E) Catheter Y: Non-enhancing (NE) Catheter Z: Brain adjacent to tumor (B)

C

Enhancing ■
Non-enhancing ●
Brain adjacent △
to tumor
IDH1-R132H (blue)
IDH-WT (red)

Normalized Peak Area

Oligo^2, Oligo^{3}1, Oligo^{3}2, Oligo^{3}3, GemAstro$^{3\text{-mut}}$, Astro$^{4\text{-mut}}$1, Astro$^{4\text{-mut}}$2, Astro$^{4\text{-mut}}$3, H3K27M-Astro$^{4\text{-WT}}$, GBM^{WT}1, GBM^{WT}2, GBM^{WT}2B, GBM^{WT}3, GBM^{WT}4, MolecGBM^{WT}

D 2-HG
IDH-mutant Astrocytomas, WHO Grade 4

E NE B   E NE B   E E† B
Astro$^{4\text{-mut}}$1   Astro$^{4\text{-mut}}$2   Astro$^{4\text{-mut}}$3

A. Enhancing Tumor vs. Brain Catheter X/Z

B. Enhancing vs. Non-Enhancing Tumor Catheter X/Y

C. Non-Enhancing Tumor vs. Brain Catheter Y/Z