Environmental Exposure to Volatile Organic Compounds is Associated with Endothelial Injury

*Daniel W. Riggs,[1-5] *Marina V. Malovichko,[1-4] Hong Gao,[1-4] Katlyn E. McGraw,[1-4] Breandon S. Taylor,[1,2] Tatiana Krivokhizhina,[1-4] Shesh N. Rai,[1,2,6] Rachel J. Keith,[1,2,4] Aruni Bhatnagar,[1-4] and Sanjay Srivastava[1-4]

[1]Superfund Research Center, University of Louisville; [2]Diabetes and Obesity Center, University of Louisville; [3] Christina Lee Brown Envirome Institute, University of Louisville, [4]Division of Environmental Medicine, Department of Medicine, University of Louisville; [5]Department of Epidemiology and Population Health, University of Louisville; [6]Department of Bioinformatics and Biostatistics, University of Louisville

*These people contributed equally to the manuscript

Address correspondence to: Sanjay Srivastava, PhD
                                    Superfund Research Center, Cardiovascular
                                    Innovation Institute
                                    302 E. Muhammad Ali Blvd
                                    Louisville, KY 40202
                                    Phone: 502-852-5724; Fax: 502-852-5834
                                    E-mail: sanjay.srivastava@louisville.edu

The authors declare they have no actual or potential competing financial interests.

ABSTRACT

Objective: Volatile organic compounds (VOCs) are airborne toxicants abundant in outdoor and indoor air. High levels of VOCs are also present at various Superfund and other hazardous waste sites; however, little is known about the cardiovascular effects of VOCs. We hypothesized that ambient exposure to VOCs exacerbate cardiovascular disease (CVD) risk by depleting circulating angiogenic cells (CACs).

Approach and Results: In this cross-sectional study, we recruited 375 non-smokers with low-to-high CVD risk and measured 15 subpopulations of CACs by flow cytometry and 16 urinary metabolites of 12 VOCs by LC/MS/MS. Associations between CAC and VOC metabolite levels were examined using generalized linear models. Our data show that VOC exposure varied by sex, age, and race. In single pollutant models, metabolites of xylene, benzene, and 1,3-butadiene were negatively associated with CAC levels. The metabolite of acrylonitrile was negatively associated with $CD45^{dim}/CD146^+/CD34^+/AC133^+$ cells and $CD45^+/CD146^+/AC133^+$, and the toluene metabolite with $AC133^+$ cells. Multipollutant models showed a negative association with metabolites of ethylbenzene/styrene, benzene, and xylene with $CD45^{dim}/CD146^+/CD34^+$ cells, independent of other VOC metabolite levels. Cumulative VOC risk score showed a strong negative association with $CD45^{dim}/CD146^+/CD34^+$ cells, suggesting that total VOC exposure has a cumulative effect on pro-angiogenic cells. Dose-response relationship showed an increase in CAC levels at low, but depletion at higher levels of VOC exposure. Sex and race, hypertension, and diabetes significantly modified VOC associated CAC depletion.

Conclusion: Low-level ambient exposure to VOCs is associated with CAC depletion, which could compromise endothelial repair and angiogenesis, and exacerbate CVD risk.

62 **Keywords:** Circulating angiogenic cells, VOCs, endothelial injury, benzene,

63 ethylbenzene, xylene

64 The study was approved by the University of Louisville Institutional Review Board (IRB

65 15.1260).

1. INTRODUCTION

Exposure to ambient air pollution has been widely recognized to be an important contributor to premature mortality and morbidity. The Global Burden of Disease (GBD) study estimates that in 2015, 4.2 million deaths could be attributed to ambient air pollutants, with an additional 2.8 million deaths due to household air pollution (Cohen et al., 2017). Exposure to fine particulate matter in the ambient air, defined as particle size < 2.5 µm in diameter ($PM_{2.5}$), is associated with an increased risk for hypertension (Yang et al., 2018), myocardial infarction (Mustafić et al., 2012), type 2 diabetes (Brook et al., 2013b; Rajagopalan and Brook, 2012), stroke (Shah et al., 2015), and cardiovascular mortality (Rajagopalan et al., 2018). The emphasis on $PM_{2.5}$ and cardiovascular disease (CVD) has been motivated largely by the availability of historical monitoring data, however in an urban setting, over 90% of pollutant mass consists of gases or vaporous compounds, such as volatile organic compounds (VOCs) (Brook et al., 2010). In comparison with $PM_{2.5}$, data on VOCs tend to exhibit higher temporal and spatial variability.

Most studies on VOC have focused on tobacco smoke (De Jesús et al., 2020; Keith et al., 2020; Lorkiewicz et al., 2019) which represents a major source of human exposures. However, non-tobacco derived VOCs are ubiquitous in urban environments. High concentrations of VOCs such as alkanes, alkenes, aromatic hydrocarbons, and aldehydes, are emitted into the atmosphere from both anthropogenic and biogenic sources, and although biogenic VOCs are more abundant worldwide, in urban areas, anthropogenic VOCs are often higher (Atkinson and Arey, 2003). Outdoor VOC exposures can originate from vehicle emissions, industrial facilities, landfills and

hazardous waste sites, while indoor exposure can originate from diverse sources, such

as paints, cleaning agents, personal care products, and solvents (Li et al., 2021). Our

analysis of the secular trends of VOC exposures in the US suggests that even though

there was a decline in the ambient air VOC levels, individual-level exposure increased

from 2005-2014 (Konkle et al., 2020). Hence, there is urgent need to assess the health

impacts of VOC exposures.

VOC exposure have been associated with a wide range of acute and chronic

health effects, including allergies, respiratory diseases, liver dysfunction, and cancer

(Shuai et al., 2018; Villeneuve et al., 2013). Moreover, exposures to ambient levels of

acrolein, benzene, and xylene have been associated with elevated CVD risk (Abplanalp

et al., 2017; DeJarnett et al., 2014). In an intra-urban analysis of VOC gradients, higher

levels of VOCs were found to be associated with increased cardiovascular and cancer

mortality (Villeneuve et al., 2013). Moreover, longitudinal time-series studies have shown

that ambient VOCs, such as alkenes and alkynes, are associated with increased heart

failure emergency hospitalizations (Ran et al., 2018b), as well as overall cardiovascular

hospitalizations (Ye et al., 2017), and that occupational exposure to VOC such as styrene

increases the relative risk of death from ischemic heart disease (Matanoski and Tao,

2003).

The biological plausibility that VOCs can induce cardiovascular injury and

exacerbate CVD is extensively supported by a range of animal studies showing that

exposures to benzene, acrolein, and formaldehyde, can lead to changes in blood

pressure regulation (Awe et al., 2006; Jin et al., 2019; Lynch et al., 2020), depletion of

circulating stem cells (Conklin et al., 2017), insulin resistance (Abplanalp et al., 2019),

and platelet activation (Sithu et al., 2010). Exposure to these VOCs also induces systemic

dyslipidemia, lipoprotein modification (Conklin et al., 2010), dilated cardiomyopathy

(Ismahil et al., 2011); and lead to the acceleration of atherogenesis (Srivastava et al.,

2011) and exacerbation of myocardial ischemic injury (Wang et al., 2008). Nonetheless,

despite this evidence, little is known about the effects of VOCs on the mechanisms and

processes that contribute to cardiovascular disease in humans. Previous work has shown

that in humans, high levels of VOC exposures from sources such as cigarette smoke

(Heitzer et al., 1996; Pittilo, 2000) and diesel exhaust (Törnqvist et al., 2007) are

associated with endothelial dysfunction. We have previously found that exposure to

acrolein and 1,3-butadiene are associated with lower reactive hyperemia index, a

measure of endothelial function (Li et al., 2021; McGraw et al., 2021). The endothelium

is a first line of defense of cardiovascular tissues against environmental toxins, and

endothelial function is a sensitive reflection of vascular and systemic health. Therefore,

to identify and understand the effects of VOC exposures and cardiovascular health, we

measured circulating angiogenic cells (CAC), also referred to as endothelial progenitor

cells (EPC). It is believed that these cell aid in the repair of the endothelium after injury,

and are good biomarkers of vascular function (Hill et al., 2003; Zafar et al., 2018). CAC

levels are associated with several CVD risk factors (Werner and Nickenig, 2006), and are

predictive of both cardiovascular events and cardiovascular mortality (Werner et al.,

2005). Thus, using CACs as surrogate biomarkers of endothelial health, we examined

their association with urinary levels of VOC metabolite levels in non-smokers with low to

severe CVD risk.

2. METHODS

2.1 Study Population and Design

In a cross-sectional study, adult study participants, aged 25-70 years, were recruited from the Louisville, KY area. Recruitment occurred in the summer months (May to September) in the years 2018 and 2019, to minimize seasonal outdoor VOC heterogeneity. Persons unwilling or unable to provide informed consent or with significant and/or severe comorbidities were excluded. These included subjects that had hepatitis, HIV/AIDS, active treatment for cancer, and active bleeding including wounds. Other exclusions included subjects with body weight less than 100 pounds, pregnant women, prisoners, and other vulnerable populations, those with conditions known to effect peripheral blood cell counts and bone marrow function.

At the time of enrollment non-fasting blood and spot urine were collected from patients who met the enrollment criteria and gave written consent. Demographic information, including age, sex, ethnicity, residential address, smoking status history and secondhand smoke exposure (verified by measurement of urinary cotinine concentrations), and BMI (calculated from self-reported height and weight) was obtained by using a questionnaire. Information on medication usage, blood pressure, and CVD history, including incidence of heart attack, heart failure, angina, hypertension, hypercholesterolemia, diabetes, obesity, stroke, and revascularization were also obtained through the questionnaire. For the current study, hypertension was defined as a previous diagnosis of hypertension or systolic blood pressure greater than 140 mmHg. Diabetes was defined as previous diagnosis of diabetes or HbA1C greater than 6.5. The study was approved by the University of Louisville Institutional Review Board (IRB 15.1260). A total of 735 individuals met the inclusion criteria (Scheme 1).

Of those enrolled in the study, 115 individuals did not provide blood samples, or did not

have circulating angiogenic cell measurements, 9 individuals did not provide urine

samples, and 8 were missing other covariates. Of the 603 participants with both urinary

VOC metabolites and CAC measurements, 228 were excluded because they were

current tobacco users. Tobacco use was confirmed using urinary cotinine values, with a

value of >40 µg/g creatinine indicating current user. The final sample size consisted of

375 participants.

2.2 Urinary Volatile Organic Compound Metabolites

Urinary metabolites of 12 VOCs (16 metabolites), and free forms of the nicotine

metabolite cotinine were quantified by ultra-performance liquid chromatography-mass

spectrometry (UPLC-MS/MS) as described before (Keith et al., 2020; Lorkiewicz et al.,

2019; Yeager et al., 2020). All VOC metabolites that were below the limit of detection

(LOD) were imputed with the LOD value divided by the square root of 2. All urinary

analytes were normalized to urinary creatinine levels, measured on a COBAS MIRA-plus

analyzer (Roche, New Jersey) with Infinity Creatinine Reagent (Thermo Fisher Scientific,

Massachusetts). A complete list of metabolites and parent compounds is provided in

supplementary material (Table S1).

2.3 Circulating Angiogenic Cells

Fifteen different subtypes of circulating angiogenic cells were measured in the

peripheral blood by flow cytometry - on fresh blood specimens as previously described

(DeJarnett et al., 2014; O'Toole et al., 2010). CACs were defined as CD14 negative cells

and differentiated into subtypes based on expression of CD45, CD34, CD146 and AC133

surface markers.

2.4 Air Pollution and Meteorological Data

Data on ambient levels of $PM_{2.5}$ (particulate matter with an aerodynamic diameter of ≤2.5 μm; μg/m^3) and ozone (ppm) were obtained from three regional EPA-validated monitoring stations in the Louisville, KY region. We used the average of the three monitors for the day the study participants provided blood and urine samples.

Data on average daily temperature (°F) and relative humidity (%) were obtained from the National Oceanic and Atmospheric Administration's (NOAA) National Centers for Environmental Information. Data was download for the Louisville International Airport monitoring station, which is in close proximity to the study participants' residence, for the day the study participants provided blood and urine sample. This data is publicly available (https://www.ncdc.noaa.gov/cdo-web/datatools/selectlocation).

2.5 Statistical Analysis

Participant characteristics are expressed as mean ± standard deviation (SD) for continuous variables, and frequency (%) for categorical variables. Descriptive statistics were conducted for all VOC metabolites. Pearson correlations were performed across VOC metabolites and daily $PM_{2.5}$, ozone, temperature, and relative humidity (Supplemental Figure 1). Because CACs were positive, and heavily right-skewed, generalized linear models with the gamma distribution and log-link were used to examine the association between CACs and VOC metabolites (Figure 1). Models used natural log-transformed VOC metabolites, and were adjusted based on prior literature showing an association with VOC exposure or CAC levels (Fadini et al., 2008; Fadini et al., 2007; Giannotti et al., 2010; O'Toole et al., 2010; Samman Tahhan et al., 2018; Tobler et al.,

2010; Xia et al., 2012). Model adjustments included: age, sex, race, BMI, diabetes status,

hypertension status, and daily $PM_{2.5}$ levels. There are a total of 240 unique comparisons

between CACs and VOC metabolites. To account for multiple testing, we applied the

Benjamini-Hochberg procedure to control the false-discovery rate (FDR). We examined

potential non-linear relationships between VOC metabolites and CACs using restricted

cubic splines, with tests for non-linearity using the likelihood ratio test. An environmental

risk score (ERS) representing cumulative VOC exposure risk was calculated for each

CAC using a modified version of the method developed by Park *et al* (2014). Briefly, (1)

VOC metabolites were log transformed, (2) each log-transformed VOC metabolite was

regressed against each CAC separately, while adjusting for covariates, (3) the

environmental risk score for each participant was calculated as the sum of log-

transformed VOC metabolite x (regression coefficient/standard error). If multiple VOC

metabolites came from the same parent compound, only one was used in the creation of

the ERS. The final set of VOC metabolites used to calculate the ERS consist of: 3HPMA,

AAMA, CYMA, MU, MHBMA3, HPMMA, AMCC, PGA, 2HPMA, MA, BMA, and

3MHA+4MHA. Multi-pollutant models including all VOC metabolites that were

independently associated with CACs were further developed to test whether single VOC

effects were confounded by correlated VOCs (Pearson correlations provided in

Supplemental Table 2). These models used log-transformed metabolite values, and the

same adjustments as single pollutant models. Interactions between covariates and log

VOC metabolites were tested in single pollutant models. We stratified analyses by

participant characteristics to examine whether effect estimates differed by subgroups.

In addition to single-VOC models, we created models for multiple VOC exposures.

To assess multiple VOC exposures as a mixture, we applied a Bayesian Kernel Machine

Regression (BKMR) approach to flexibly model the relationship between the mixture of

VOC metabolites and each of the 15 CACs (Bobb et al., 2015). The BKMR is a semi-

parametric technique that models the exposure response relationship as a non-

parametric kernel function of the mixture components, adjusting for covariates

parametrically. Implementation of BKMR relied on a Gaussian kernel distribution, with flat

priors. The Gaussian kernel flexibly captures underlying functional forms without

specifying the shape of the exposure-response curve. This method can assess

independent effects, estimate the overall mixture, and assess for interactions among

mixture components. We used R software (version 4.0.2) and the bkmr package (Bobb

et al., 2018) to create posterior inclusion probabilities (PIPs) that quantify the relative

importance of each exposure in the model, assess univariate and bivariate associations

between mixture components and CACs, assess interactions among VOC metabolites,

and estimate the effect of the overall mixture on individual CACs.

As a sensitivity analysis, we conducted generalized linear models additionally

adjusting one at a time for temperature or relative humidity, while adjusting for the

covariates as above. We then repeated the models adjusting for ozone in place of $PM_{2.5}$,

while adjusting one at time for temperature, then relative humidity. All statistical analyses

were performed using SAS, version 9.4 (SAS Institute, Inc., Cary, North Carolina) and

GraphPad Prism, version 8 (GraphPad Software, La Jolla, California).

3. RESULTS

The demographic characteristics of study participants are summarized in Table 1.

Within the 375 participants of the study, the average age was 50.6 years. A majority were

female (64%) and White (83%); 42% were hypertensive, and 17% diabetic. The average

daily PM$_{2.5}$ level (±SD) on the day of the study visit was 8.6 ± 3.4 µg/m^3. The urinary

metabolites of VOCs measured in study participants are listed in Table 2, along with their

parent VOCs.

The mean and median concentrations of VOCs show a high degree of variability.

Although significant levels of most VOC metabolites could be detected, in many (>20%)

samples the level of acrylonitrile (CYMA), styrene/ethylbenzene (PHEMA) metabolites

were below the limit of detection (LOD). There was minimal correlation between VOC

metabolites and average daily ambient PM$_{2.5}$, ozone, temperature, and relative humidity

(Supplemental Figure 1). The levels of urinary VOC metabolites showed a strong

dependence on demographics. A comparison of urinary metabolites showed that females

had higher levels of the metabolites of N,N'-DMF (AMCC), ethylbenzene/styrene

(phenylglyoxylic acid, PGA and mandelic acid, MA) and toluene (BMA) than males.

Likewise, in comparison with Blacks, Whites had higher levels of metabolites of benzene

(*trans, trans*-muconic acid, MU), N,N'-DMF (N-Acetyl-S-(N-methylcarbamoyl)-L-cysteine,

AMCC) and styrene/ethylbenzene (MA). The metabolites of several VOCs – 1,3,

butadiene (N-Acetyl-S-(3,4-dihydroxybutyl)-L-cysteine, DHBMA), crotonaldehyde (N-Acetyl-

S-(3-hydroxypropyl-1-methyl)-L-cysteine, HPMMA), N,N'-DMF (AMCC),

styrene/ethylbenzene (MA), toluene (BMA), and xylene (3MHA+4MHA) were also higher

in those >55 years of age. Urinary levels of the metabolite of butadiene (MHBMA3) and

toluene (N-Acetyl-S-(benzyl)-L-cysteine, BMA and, 2-Methylhippuric acid, 2MHA) levels were

higher in those with a BMI of <30 (Supplemental Table 2), reflecting different levels of

exposure.

In single-VOC analysis, the levels of several urinary VOC metabolites were

associated with the levels of CACs measured in peripheral blood (Figure 1). Some of the

most pronounced effects were observed with PGA – a metabolite that could be derived

from either ethylbenzene or styrene - which showed a significant inverse association with

CACs-1,2,3,4,5,6, and 7, as well as CAC-13 and CAC-15. Several other VOC metabolites

showed similar, but variable associations. The percent difference ranged from -10.7%

(95% CI: -19.6%, -0.8%) to -42.8% (95% CI: -53.1%, -30.1%). Many VOC metabolites

were inversely associated with CAC levels, including metabolites of xylene (3-MHA+4-

MHA), acrylonitrile (CYMA), and 1,3-butadiene (MHBMA3). For benzene (MU), we

observed a significant inverse association with CAC-1 (-11.9%; 95% CI: -20.0%, -3.0%),

and its subpopulation CAC-7 (-10.7%; 95% CI: -18.3%, -2.3%), along with a positive

association with CAC-10. Positive associations were also observed with toluene (BMA)

and CAC-5, 11, 13, and 14, which share AC133$^+$ marker. After correcting for multiple

testing, PGA showed statistically significant associations with CAC-1, 3, 5, 6, and 15

(Figure 1C). Further, MU remained significantly associated with CAC-10 and BMA

remained significantly associated with CAC-5. In our sensitivity analysis, additionally

adjusting temperature, relative humidity, and ambient ozone did not substantially alter our

associations (Supplemental Figure 2). Overall, these observations suggest that exposure

to several VOCs is generally associated with lower levels of several CAC subpopulations

in peripheral blood, particularly the early progenitor CACs.

To determine the extent to which total VOC exposure can affect CAC levels, and

which cell populations contribute to the directionality of the associations (found in Figure

1) we created an Environmental Risk Score (ERS) for each cell population (Figure 2). We

found a strong negative risk score for CAC-1 (-62.9), CAC-3 (-54.6), and CAC-7 (-62.1),

which are a related group of cells with common markers CD45^{dim}/CD146^{+}/CD34^{+}. To a

lesser extent, we also found negative risk scores for CAC-6 (-47.6), and CAC-9 (-37.7),

which are CD34^{+} cells. In comparison, we found positive risk scores for CD45^{+} cells (CAC-

12), along with subgroups of CD45^{+} cells: CAC-2, CAC-8, and CAC-14. In addition, we

also found a positive score for CAC-10, which is CD146^{+}. This indicates that the levels of

CD45^{+} cells were positively associated with the urinary metabolites of VOC metabolites,

whereas CD45^{dim} and CD34^{+} cells were negatively associated with the metabolite levels.

We used restricted cubic splines to determine whether the relationship between

VOC metabolites and CACs was non-linear (Figure 3).   We observed non-linear

relationships between the circulating levels of several cell populations and the metabolite

of benzene (MU).   In general, low levels of MU was associated with higher levels of

circulating cells, whereas higher MU levels were associated with lower cell levels. These

non-linear relationships are presented in Figure 3A, which shows a near significant

(p<0.1) non-linear relationships with CD146+ cell populations (CAC-2, 4, 5, and 6), and

with AC133+ cell population – CAC-4 and CAC-5.

To determine whether AC133^{+} cells contribute to the non-linear relationships, we

examined the association of subgroups of CD146^{+} cells, CAC-4 (CD45^{+}/CD146^{+}/AC133^{+})

and CAC-8 (CD45^{+}/CD146^{+}/AC133^{-}) with VOC metabolites. We found near significant

(p<0.1) non-linear relationships between CAC-4 and metabolites of acrolein, 1,3-

butadiene, N, N-dimethylformamide, propylene oxide, ethylbenzene/styrene, and xylene. These non-linear relationships were not observed with CAC-8, with the exception of xylene (3-MHA+4-MHA), suggesting that early progenitor endothelial cells ($CD146^+/AC133^+$) are recruited at low exposure levels, but are depleted at high VOC exposures.

Given that co-exposure to VOCs is common, we used multi-VOC models to assess whether the significant VOC effects found in single-VOC models were confounded by co-exposures to other VOCs (Figure 4). For this analysis, we focused on rare cell populations with negative ERS scores CAC-3, CAC-5, CAC-7, and CAC-13 (Figure 2). We observed that the metabolite of ethylbenzene/styrene (PGA) remained negatively associated with CAC-3 (-37.0% per 2-fold increase; 95% CI: -48.7%, -22.6%), CAC-5 (-29.8%; 95% CI: -40.6%, -17.0%), CAC-7 (-16.8%; 95% CI: -29.1%, -2.4%), and CAC-13 (-13.1%; 95% CI: -22.2%, -2.8%), suggesting that the associations between PGA and CACs are independent of other VOC metabolite levels. We also found that the metabolites of xylene (2MHA) remained negatively associated with CAC-3, crotonaldehyde (HPMMA) remained positively associated with CAC-5, benzene (MU) remained negatively associated with CAC-7, and toluene (BMA) remained positively associated with CAC-13.

In our Bayesian kernel machine regression analysis, important VOC metabolites contributing to changes in CACs are indicated by high posterior inclusion probabilities (Table 3). The directionalities of single VOC associations in our mixture models were generally consistent with the observed directionalities in our single VOC generalized linear models. For example, in mixture models, individuals with greater PGA levels had lower numbers of CACs, while individuals with greater BMA and HPMMA levels had

higher numbers of CACs. As evident in Figure 5, we also found that the mixture of VOCs

had a nonlinear relationship with CAC-4, similar to the nonlinear associations observed

in our single VOC models. We also found that greater exposure to the VOC mixture was

associated with higher CAC-10 levels (Figure 5). The effect of the VOC mixture on CAC-

10 was largely driven by benzene (MU), with a posterior inclusion probability of 0.55,

where all other VOC metabolites were less than 0.15. We found minimal evidence of

interactions between pairs of VOC metabolites in our mixture analysis (data not shown).

Because the metabolites of benzene, 1,3-butadiene, N,N-dimethylformamide, and

ethylbenzene/styrene consistently showed strong associations with CACs, we

investigated whether these associations were modified by the demographic

characteristics of the participants or their co-morbidity profile. In our subgroup analysis,

we found a common theme in many of the models, in which participants who were male,

White, normotensive, and recruited on low $PM_{2.5}$ days showed stronger inverse

relationships between VOCs and CAC levels (Figure 6).

For example, for CAC-13, we found a negative association between

ethylbenzene/styrene metabolite (PGA) in White participants of -17.2% per 2-fold

increase in CAC levels (95% CI: -26.4%, -7.0%), whereas in Black participants, we found

a positive association of 79.1% (95% CI: 23.7%, 159.3%). We observed a similar

relationship for the metabolite of benzene (MU), when stratified by race; however, this

may be due to lower levels of MU in urine samples from Black participants (62.2 µg/g

creatinine) than White participants (146.0 µg/g creatinine), as presented in *Supplemental*

Table 2.

We also found some evidence that age and BMI modify the relationships between

VOCs and CAC levels, although the direction was not consistent in all models.

Interestingly, in several of our models, we found stronger inverse associations between

VOC metabolite and CACs in normotensives than hypertensives, and to a lesser extent,

non-diabetics than diabetics, - conditions that are known to affect cell migration. For

example, for CAC-5, we found a negative association with ethylbenzene/styrene

metabolite (PGA) in non-diabetics of -28.2% per 2-fold increase, and with N,N-

dimethylformamide (AMCC) in non-diabetics of -24.0% per 2-fold increase, while the

relationships in diabetics were not significant. To determine whether these results were

due to differential levels of VOC exposure between subgroups, we calculated average

metabolite levels for each subgroup *Supplemental Table 2*. We found that levels of MU

(benzene) and PGA (ethylbenzene and styrene) were higher in diabetics than non-

diabetes and were higher in hypertensive than normotensive individuals, although

normotensive had higher levels of 2HPMA (propylene oxide).

4. DISCUSSION

The major findings of this study are that the urinary metabolites of several common

VOCs, such as ethylbenzene/styrene, benzene, xylene, and 1,3-butadiene were

negatively associated with circulating angiogenic cells. For some VOCs, such as

benzene, we observed that low levels of their metabolites were positively associated with

an increase in CAC levels, whereas at high levels of exposure were associated with a

significant decrease in CAC levels. Our environmental risk score identified that CAC-1,

CAC-3, and CAC-7, with common cell markers $CD45^{dim}/CD146^{+}/CD34^{+}$, were the most

sensitive to total VOC exposure. Our mixture analysis revealed that the total VOC

metabolite mixture was positively associated with CD146[+] cells. We also found sex, race,

age, and hypertension to be effect modifiers of the some of the relationships between the

levels of VOC metabolites and CACs. Taken together, these findings suggest that

exposure to VOCs could lead to deficits in endothelial repair capacity in a dose-dependent

manner, which vary with sex, race, and comorbidities. Given that low CAC levels are

predictive of future cardiovascular events and death (Werner et al., 2005), these results

support the notion that VOC exposure is associated with significant vascular injury and

therefore it may be an important environmental risk factor for CVD.

391   The potential cardiovascular effects of VOCs derived from tobacco smoke have

been extensively studied (De Jesús et al., 2020; Keith et al., 2020; Lorkiewicz et al.,

2019); however, in contrast, the cardiovascular effects of VOCs not derived from tobacco

smoke have received scant attention. Nonetheless, in addition to tobacco smoke,

automobile emissions, industrial facilities, hazardous waste sites, and household

products are major sources of VOC emissions, and therefore, VOC exposures are

common and frequent (Li et al., 2021). In our study, we found significant levels of the

VOC metabolites in the urine of non-smokers, indicating particularly high levels of

exposure to acrolein, crotonaldehyde, and 1,3-butadiene, and acrylamide, as well as the

BTEXS (benzene, toluene, ethylbenzene and styrene) group of VOCs, suggesting a

contribution of both indoor and outdoor exposure. Specifically, we found that the CAC

levels were particularly sensitive to metabolites of BTEXS compounds. We observed

significant associations between CACs and urinary metabolite of benzene (3 CACs),

toluene (4 CACs), ethylbenzene/styrene (9 CACs), and xylene (4 CACs).

Although exposure to BTEXS is common, little is known about their cardiovascular

effects. A study from Hong Kong found that short term elevations in atmospheric levels

of benzene and TEX (toluene, ethylbenzene, and xylene) was associated with increases

in circulatory disease mortality by 5.8% and 3.5% (Ran et al., 2018a), although in a study

from Toronto, no intra-urban variations in VOC levels were found be associated with

cardiovascular disease mortality (Villeneuve et al., 2013). Other BTEXS such as toluene,

styrene, ethylbenzene, and xylenes have been also linked to a higher prevalence of CVD

(Xu et al., 2009), and exposure to VOCs such as benzene, ethylbenzene, and styrene

have also been found to be associated with higher odds of developing metabolic

syndrome (Shim et al., 2019). Along with these data, the results of our study, support the

notion that exposure to VOC could have adverse cardiovascular consequences,

potentially due to low-level vascular injury reflected by changes in CAC levels.

We found inverse associations between metabolites of acrylonitrile and 1, 3-

butadiene and CACs. Although the cardiovascular effects of acrylonitrile are less well

studied, exposure to 1,3-butadiene has been found to accelerate the formation of

atherosclerotic lesions in animal models, and polymorphisms in phase II enzymes that

metabolize 1,3-butadiene are associated with the development of atherosclerosis in

humans (Penn and Snyder, 2007). Hence, our observation that urinary metabolites of 1,3-

butadiene are associated with CAC levels suggest that the cardiovascular risk of

exposure to this VOC may be in part due to depletion of early, progenitor cells, which aid

in vascular repair and thereby can prevent the progression of atherosclerotic lesions.

Therefore, our observations linking exposure to VOCs such as 1,3-butadiene to

concurrent changes in CACs, not only provides a new biological basis for the

cardiovascular effects of VOCs, independent of concurrent exposure to $PM_{2.5}$, but also

provides new insights into their mechanism of toxicity.

In real world scenario, the urban air contains many different VOCs and therefore,

we conducted multi-VOC models controlling for all significantly associated metabolites

from the single VOC models. In particular, we found that the urinary metabolite PGA

remained significantly associated with the levels of $CD45^{dim}/CD146^+/CD34^+$ cells (CAC-

3, CAC-7), $AC133^+/CD146^+$ cells (CAC-5), and $AC133^+/CD34^+$ cells (CAC-13). PGA is

oxidized from mandelic acid (MA) by alcohol dehydrogenase, and both are considered

the major urinary metabolites of styrene and ethylbenzene exposure (Capella et al.,

2019). Styrene is mainly used in the manufacturing of plastics, synthetic rubbers, and

resins, and is the top released carcinogen in the United States (Mori et al., 2000), while

ethylbenzene is found naturally in coal tar and petroleum, and used in the manufacturing

of styrene (Capella et al., 2019). The primary route of exposure to styrene and

ethylbenzene is through inhalation, contributing up to 99% of total exposure, with 1-2%

from food consumption (Tang et al., 2000). We found that the ethylbenzene/styrene

metabolite - PGA, but not MA was negatively associated with CACs. This could be due

to the chemical structure of PGA, which is a ketone, and is therefore, more reactive than

the acid metabolite MA. This could also be representative of higher ethylbenzene/styrene

exposure, which would affect the ratio of PGA to MA through an increase in alcohol

dehydrogenase activity. The results of our multi-VOC analysis indicate that CACs may

be particularly sensitive to ethylbenzene/styrene exposures.

We found that the negative association between the N,N-DMF metabolite (AMCC)

and CACs was significantly stronger in normotensive participants than those with

hypertension. Previous work has shown that hypertension is associated with lower CAC

levels and that hypertension is a major independent predictor of impaired endothelial

progenitor cell migration (Vasa et al., 2001). It has also been demonstrated that

hypertension accelerates EPC senescence in both clinical and experimental settings

(Imanishi et al., 2005). These findings suggest that because of the inherent susceptibility

of CACs to hypertension, CACs in hypertensive participants may be more vulnerable to

exposure to VOCs such as N,N-DMF. N,N-DMF is a commonly used solvent with wide

industrial applications, and occupational exposure to high levels of N,N-DMF is

associated with liver injury (Li and Zeng, 2019); however, our observation that even

ambient exposure to N,N-DMF is associated with vascular injury, suggests that there may

be a high risk of hypertension and cardiovascular disease even with low-level exposures.

The observed association between VOC metabolites and diminished circulating

angiogenic cells suggest that individuals exposed to VOCs may have a lower capacity to

repair the endothelium. These effects of VOC exposure appear to be similar to those of

other environmental pollutants, which are also associated with suppression of CAC levels.

Previous work has shown that exposure to $PM_{2.5}$ over a period of 24h to several weeks

suppresses EPC levels (Haberzettl et al., 2012; O'Toole et al., 2010), and exposure to

pollutants such as particulate nickel (Liberda et al., 2010), acrolein (DeJarnett et al.,

2014) and benzene (Abplanalp et al., 2017) have also been shown to reduce CAC levels.

These observations suggest that pollutant exposure either depletes CAC levels in

peripheral blood or interferes with the release from the bone marrow and the spleen.

However, in contrast to such suppression, CAC levels are slightly elevated in individuals

breathing coarse PM (2013a) or those living close to a major roadway (DeJarnett et al.,

2015). This biphasic response is consistent with the dose-response relationships

observed in the current study, which are indicative of increased CAC mobilization (to

promote repair) at low levels of exposure, and CAC suppression at higher exposures.

Recently, Yeager $et\ al.$, have reported that living in areas of high vegetation is

associated with increased levels of $CD34^+/CD45^+/AC133^+$ CACs, but decreased

$CD31^+/CD34^+/CD45^{dim}/AC133^+$ CACs (2018), suggesting greenness may be protective

against air pollution related health effects. Thus, it seems that low levels of toxicity could

lead to increased recruitment of CAC into the peripheral blood, whereas, high levels of

exposure could suppress CAC recruitment and/or depleting CAC by increasing utilization.

The directionality of the effects of exposure on CAC levels may also depend upon locus

of injury, some exposures may prevent recruitment, while others may induce injury that

stimulates recruitment. Regardless, our results with VOCs suggest that exposure to

these chemicals suppresses CAC levels in peripheral blood, but further studies are

required to determine whether this is due to decreased recruitment or increased

utilization.

We found that several VOC metabolite levels were negatively associated with

angiogenic cells ($CD45^{dim}/CD146^+/CD34^+$), but positively associated with hematopoietic

endothelial cells ($CD45^+/CD146^+$), as evident from our environmental risk score results

(Schematic 2).

In comparison with $CD45^{dim}$ cells, $CD45^+$ CACs are more primitive, with the potential to

self-renew and differentiate into specialized cells that can control biological activities such

as homeostasis and response to inflammation (Chotinantakul and Leeanansaksiri, 2012).

Additionally, the angiogenic cells are rare, and can be easily depleted when used in the

repair of endothelial injury. Our exposure response curves are consistent with this notion

that at low levels of VOC exposure, $CD45^+$ cells have the ability to regenerate to repair

endothelial injury, however at high VOC exposure, the injury surpasses the cells capacity

to regenerate or the mobilization of these cells from depots such as the spleen or bone

marrow is prevented. Consistent with this notion, results from our Bayesian kernel

machine regression also found a nonlinear relationship with the mixture of VOCs and

CAC-4 ($CD45^+$/$CD146^+$/$AC133^+$). Further, we found that the mixture of VOC metabolites

was positively associated with $CD146^+$ cells. These cells are thought to be mature cells

which detach from the intimal monolayer after endothelial injury. Studies have shown an

increased number of $CD146^+$ cells in CVD, such as unstable angina, acute myocardial

infarction, stroke, and diabetes (Boos et al., 2006). Positive associations could also be

due to the specific VOC exposure. We found primarily positive relationships between the

toluene metabolite (BMA) and CACs. This could be reflective of the low level of toxicity of

toluene in comparison to other VOCs.

The major strength of this study is the use of individual level data to assess

exposure. Although, measurements of the urinary metabolites do not permit source

assignment, they do provide an integrated measure to total exposure of an individual

whether via inhalation, ingestion, or endogenous production. Hence, in linking exposure

to outcome, such measurements provide a unique advantage as exposure-response

could be concurrently evaluated in the same individual. Using this approach, we were

able to assess association between exposure and a sensitive and robust measure of

preclinical cardiovascular injury (CACs) which is predictive of CVD events and mortality.

In our analysis, we also adjusted for exposure to $PM_{2.5}$ and were able to apply a

composite VOC risk score that could be used in future studies.

Our study also has some limitations. Because of the cross-sectional nature of the

study, the associations found in this study cannot address causality. Longitudinal studies

are needed to determine whether VOC exposures are associated with adverse clinical

outcomes, and whether these outcomes are related to changes in CAC levels. Our study

also has the potential for type I errors due to testing of multiple CACs and VOC

metabolites. However, due to the correlation between VOC metabolites, and limited prior

research on VOC exposure and CVD, we did not adjust for multiple comparisons, to

preserve our type II error, and identify potential VOC candidates for future studies.

Moreover, we used spot urine collection from each individual to estimate VOC exposure.

Although it has been previously shown that spot urine biomarkers of VOC exposure in

cigarette smokers correlate well with 24 h urine excretion(Sarkar et al., 2013), our study

excluded smokers. Following exposure, many VOCs rapidly undergo biotransformation

to generate metabolites that can conjugate with glutathione. Other enzymatic reactions

remove glutamic acid and glycine to produce a cysteine conjugate, which is N-acetylated

and excreted in the urine as mercapturic acid (Li et al., 2021). It has also been shown that

smoking one cigarette, which generates high levels of VOCs, is sufficient to immediately

affect the levels of CACs (Mobarrez et al., 2014). Therefore, because VOC metabolites

and CAC levels were concurrently measured, it is likely that the observed association

represent consequences of recent exposure. Additionally, our urinary VOC metabolites

were normalized to creatinine to account for dilution differences, which could introduce

error that relates to the dependence of creatinine excretions due to muscle mass, physical

activity, and kidney disease. Another limitation of our study is that most participants were,

by study design, of older age and at increased risk of cardiovascular disease. Thus, our

sample could represent a group of susceptible individuals, which may limit generalizability

of our findings.

In conclusion, we showed that metabolites of several VOCs were negatively

associated with circulating levels of $CD45^{dim}$/$CD146^{+}$/$CD34^{+}$ cells and the mixture of VOC

metabolites was positively associated with $CD146^{+}$ cells in the peripheral blood. These

findings suggest that exposure to low ambient levels of VOCs could damage the

endothelium, and deplete rare CACs or inhibit their recruitment, and thereby compromise

endothelial repair. Given the global burden of CVD, and the ubiquitous distribution of

VOCs, it is imperative to further assess their contribution to CVD risk. Because several

commonly used products, such as paints, self-care products, and glue, are important

source of VOCs, and that there are several sources of indoor emissions (compressed

wood, fire-retardants), this study raises the possibility that environmental exposure to

VOCs may be an important determinant of CVD risk. Our work also underscores the need

for greater awareness of the potential dangers of VOC exposure and heightened vigilance

against such exposures. Future longitudinal and mechanistic studies are needed to

confirm current findings. If substantiated, an association between VOC exposure and

CVD risk could be included in estimates of CVD risk and appropriate limits and

recommendations could be developed to reduce both ambient and indoor VOC

exposures.

FIGURE LEGENDS:

Scheme 1. Recruitment flow chart.

Figure 1. Association between the urinary metabolites of VOCs and circulating levels of CACs(A) and heat map of p-values for pair-wise associations between the urinary metabolites of VOCs and circulating levels of CACs (B and C). (A) The X axis represents % difference (± 95% confidence interval) per 2-fold increase in VOC metabolite for each CAC sub-population (CAC 1-15). Significant associations are shown in red. (B) Raw p-values. (C) False Discovery rate adjusted p-values. Black grids represent positive associations and blue grids represent negative associations. The intensity of color represents p-values, with darker grids indicating smaller p-values. Generalized linear models (n=375) were adjusted for age, sex, race, BMI, diabetes, hypertension, and daily $PM_{2.5}$ levels.

Abbreviations: CEMA, N-Acetyl-S-(2-carboxyethyl)-L-cysteine; 3-HPMA, N-Acetyl-S-(3-hydroxypropyl)-L-cysteine; AAMA, N-Acetyl-S-(2-carbamoylethyl)-L-cysteine; CYMA, N-Acetyl-S-(2-cyanoethyl)-L-cysteine; MU, *trans, trans*-muconic acid; DHBMA, N-Acetyl-S-(3,4-dihydroxybutyl)-L-cysteine; MHBMA3, N-Acetyl-S-(4-hydroxy-2-buten-1-yl)-L-cysteine; HPMMA, N-Acetyl-S-(3-hydroxypropyl-1-methyl)-L-cysteine; AMCC, N-Acetyl-S-(N-methylcarbamoyl)-L-cysteine; PGA, Phenylglyoxylic acid ; 2HPMA, N-Acetyl-S-(2-hydroxypropyl)-L-cysteine; PHEMA, N-Acetyl-S-(1-phenyl-2-hydroxyethyl)-L-cysteine +; MA, Mandelic Acid; BMA, N-Acetyl-S-(benzyl)-L-cysteine; 2MHA, 2-Methylhippuric acid; 3MHA+4MHA, 3-Methylhippuric acid + 4-Methylhippuric acid.

Figure 2. Association between Environmental Risk Scores and Circulating Angiogenic Cells. Environmental Risk score was calculated for each cell population as follows: 1. The urinary levels of VOC metabolites were log transformed. 2. Each log transformed VOC metabolite level was regressed against the levels of the cells separately, while adjusting for covariates. 3. ERS was calculated as the sum of the log transformed VOC metabolite levels x (regression coefficient/standard error). Mean and standard deviation of the ERS was calculated for each CAC. Values are mean ± SD of environmental risk score for each cell population.

Figure 3. Exposure-response relationships between selected CACs and VOCs. Non-linear relationships between MU metabolites and CACs (**A**); VOCs and CAC-4 (**B**); and VOCs and CAC-8 (**C**) were examined using restricted cubic splines. Tests for non-linearity used the likelihood ratio test. Models are adjusted for age, sex, race, BMI, diabetes, hypertension, beta-blocker use, and daily $PM_{2.5}$ levels. P-value represents likelihood ratio test for non-linearity.

Figure 4. Association between circulating angiogenic cells and multiple VOCs in the same model. Values represent % difference per 2-fold increase in VOC metabolite. VOC metabolites are ordered by strength of their regression coefficients. Only significant metabolites from one-pollutant models were entered in the model. Models adjusted for age, sex, race, BMI, diabetes, hypertension, and daily $PM_{2.5}$ levels.

590 **Figure 5**. Overall association of the VOC metabolite mixture (and 95% credible intervals) with
591 selected CACs, when all exposures are set to a given percentile relative to median. Models
592 (n=375) were adjusted for age, sex, race, BMI, diabetes, hypertension, and daily PM$_{2.5}$ levels.
593

Figure 6. Subgroup analysis of the relationship between different CAC population with the urinary
metabolites of benzene (MU), 1,3-butadiene (MHBMA3), ethylbenzene/styrene (PGA), and N,N-
Dimethylformamide (AMCC). Generalized linear models were adjusted for covariates - age, sex,
race, BMI, diabetes, hypertension, and daily PM$_{2.5}$ levels, unless the covariates were used for
stratification. Significant interaction between subgroups in the full model are indicated by an
asterisk.
600

601 **Scheme 2**. Gating Strategy for the Human Circulating Angiogenic Cells. Blue indicates negative
602 environmental risk score; orange indicates positive environmental risk score.
603

604 **Source of Funding:** This research was supported in part by NIH grants ES023716,

605 ES029846, ES019217, HL149351, HL120163 and GM127607; and Jewish Heritage

606 Foundation of Excellence grant GN190574L.

608 **Acknowledgements:** The authors would like to thank all the partners, members, and

609 volunteers involved in the HEAL study.

References

Abplanalp, W., et al., 2017. Benzene exposure is associated with cardiovascular disease risk.
PLoS One. 12, e0183602.

Abplanalp, W. T., et al., 2019. Benzene Exposure Induces Insulin Resistance in Mice. Toxicol Sci.
167, 426-437.

Atkinson, R., Arey, J., 2003. Atmospheric degradation of volatile organic compounds. Chemical
reviews. 103, 4605-4638.

Awe, S. O., et al., 2006. Acrolein induces vasodilatation of rodent mesenteric bed via an EDHF-
dependent mechanism. Toxicol Appl Pharmacol. 217, 266-76.

Bobb, J. F., et al., 2018. Statistical software for analyzing the health effects of multiple concurrent
exposures via Bayesian kernel machine regression. Environmental Health. 17, 1-10.

Bobb, J. F., et al., 2015. Bayesian kernel machine regression for estimating the health effects of
multi-pollutant mixtures. Biostatistics. 16, 493-508.

Boos, C. J., et al., 2006. Circulating Endothelial Cells in Cardiovascular Disease. Journal of the
American College of Cardiology. 48, 1538-1547.

Brook, R. D., et al., 2013a. The effect of acute exposure to coarse particulate matter air pollution
in a rural location on circulating endothelial progenitor cells: results from a randomized
controlled study. Inhalation toxicology. 25, 587-592.

Brook, R. D., et al., 2013b. Long-term fine particulate matter exposure and mortality from diabetes
mellitus in Canada. Diabetes care. DC_122189.

Brook, R. D., et al., 2010. Particulate matter air pollution and cardiovascular disease: an update
to the scientific statement from the American Heart Association. Circulation. 121, 2331-
2378.

Capella, K. M., et al., 2019. Ethylbenzene and styrene exposure in the United States based on
urinary mandelic acid and phenylglyoxylic acid: NHANES 2005–2006 and 2011–2012.
Environmental Research. 171, 101-110.

Chotinantakul, K., Leeanansaksiri, W., 2012. Hematopoietic Stem Cell Development, Niches, and
Signaling Pathways. Bone Marrow Research. 2012, 16.

Cohen, A. J., et al., 2017. Estimates and 25-year trends of the global burden of disease
attributable to ambient air pollution: an analysis of data from the Global Burden of
Diseases Study 2015. The Lancet. 389, 1907-1918.

Conklin, D. J., et al., 2010. Acrolein consumption induces systemic dyslipidemia and lipoprotein
modification. Toxicol Appl Pharmacol. 243, 1-12.

Conklin, D. J., et al., 2017. Biomarkers of Chronic Acrolein Inhalation Exposure in Mice:
Implications for Tobacco Product-Induced Toxicity. Toxicol Sci. 158, 263-274.

De Jesús, V. R., et al., 2020. Urinary Biomarkers of Exposure to Volatile Organic Compounds
from the Population Assessment of Tobacco and Health Study Wave 1 (2013-2014). Int J
Environ Res Public Health. 17.

DeJarnett, N., et al., 2014. Acrolein exposure is associated with increased cardiovascular disease
risk. J Am Heart Assoc. 3.

DeJarnett, N., et al., 2015. Residential Proximity to Major Roadways Is Associated With Increased
Levels of AC133+ Circulating Angiogenic Cells. Arterioscler Thromb Vasc Biol.

Fadini, G. P., et al., 2008. Gender differences in endothelial progenitor cells and cardiovascular
risk profile: the role of female estrogens. Arteriosclerosis, thrombosis, and vascular
biology. 28, 997-1004.

Fadini, G. P., et al., 2007. Significance of Endothelial Progenitor Cells in Subjects With Diabetes.
Diabetes Care. 30, 1305-1313.

Giannotti, G., et al., 2010. Impaired Endothelial Repair Capacity of Early Endothelial Progenitor
Cells in Prehypertension. Hypertension. 55, 1389-1397.

Haberzettl, P., et al., 2012. Exposure to ambient air fine particulate matter prevents VEGF-
induced mobilization of endothelial progenitor cells from the bone marrow. Environmental
health perspectives. 120, 848.

Heitzer, T., et al., 1996. Cigarette smoking potentiates endothelial dysfunction of forearm
resistance vessels in patients with hypercholesterolemia: role of oxidized LDL. Circulation.
93, 1346-1353.

Hill, J. M., et al., 2003. Circulating endothelial progenitor cells, vascular function, and
cardiovascular risk. New England Journal of Medicine. 348, 593-600.

Imanishi, T., et al., 2005. Endothelial progenitor cell senescence is accelerated in both
experimental hypertensive rats and patients with essential hypertension. Journal of
Hypertension. 23, 1831-1837.

Ismahil, M. A., et al., 2011. Chronic oral exposure to the aldehyde pollutant acrolein induces
dilated cardiomyopathy. Am J Physiol Heart Circ Physiol. 301, H2050-60.

Jin, L., et al., 2019. Formaldehyde Induces Mesenteric Artery Relaxation via a Sensitive Transient
Receptor Potential Ankyrin-1 (TRPA1) and Endothelium-Dependent Mechanism: Potential
Role in Postprandial Hyperemia. Front Physiol. 10, 277.

Keith, R. J., et al., 2020. Characterization of Volatile Organic Compound Metabolites in Cigarette
Smokers, Electronic Nicotine Device Users, Dual Users, and Nonusers of Tobacco.
Nicotine Tob Res. 22, 264-272.

Konkle, S. L., et al., 2020. National secular trends in ambient air volatile organic compound levels
and biomarkers of exposure in the United States. Environ Res. 182, 108991.

Li, A. J., et al., 2021. A review of environmental occurrence, toxicity, biotransformation and
biomonitoring of volatile organic compounds. Environmental Chemistry and
Ecotoxicology.

Li, M. J., Zeng, T., 2019. The deleterious effects of N,N-dimethylformamide on liver: A mini-review.
Chem Biol Interact. 298, 129-136.

Liberda, E. N., et al., 2010. Exposure to inhaled nickel nanoparticles causes a reduction in number
and function of bone marrow endothelial progenitor cells. Inhalation toxicology. 22, 95-99.

Lorkiewicz, P., et al., 2019. Comparison of Urinary Biomarkers of Exposure in Humans Using
Electronic Cigarettes, Combustible Cigarettes, and Smokeless Tobacco. Nicotine Tob
Res. 21, 1228-1238.

Lynch, J., et al., 2020. Acute and chronic vascular effects of inhaled crotonaldehyde in mice: Role
of TRPA1. Toxicol Appl Pharmacol. 402, 115120.

Matanoski, G. M., Tao, X. G., 2003. Styrene exposure and ischemic heart disease: a case-cohort
study. Am J Epidemiol. 158, 988-95.

McGraw, K. E., et al., 2021. Exposure to volatile organic compounds – acrolein, 1,3-butadiene,
and crotonaldehyde – is associated with vascular dysfunction. Environmental Research.
196, 110903.

Mobarrez, F., et al., 2014. The effects of smoking on levels of endothelial progenitor cells and
microparticles in the blood of healthy volunteers. PloS one. 9, e90314-e90314.

Mori, T. A., et al., 2000. Purified eicosapentaenoic and docosahexaenoic acids have differential
effects on serum lipids and lipoproteins, LDL particle size, glucose, and insulin in mildly
hyperlipidemic men. The American Journal of Clinical Nutrition. 71, 1085-1094.

Mustafić, H., et al., 2012. Main air pollutants and myocardial infarction: A systematic review and
meta-analysis. JAMA. 307, 713-721.

O'Toole, T. E., et al., 2010. Episodic exposure to fine particulate air pollution decreases circulating
levels of endothelial progenitor cells. Circ Res. 107, 200-3.

Park, S. K., et al., 2014. Environmental risk score as a new tool to examine multi-pollutants in
epidemiologic research: an example from the NHANES study using serum lipid levels.
PloS one. 9, e98632.

Penn, A., Snyder, C. A., 2007. 1,3-Butadiene exposure and cardiovascular disease. Mutat Res.
621, 42-9.

Pittilo, M., 2000. Cigarette smoking, endothelial injury and cardiovascular disease. International
Journal of Experimental Pathology. 81, 219-230.

Rajagopalan, S., et al., 2018. Air Pollution and Cardiovascular Disease: JACC State-of-the-Art
Review. Journal of the American College of Cardiology. 72, 2054-2070.

Rajagopalan, S., Brook, R. D., 2012. Air pollution and type 2 diabetes: mechanistic insights.
Diabetes. 61, 3037-3045.

Ran, J., et al., 2018a. Short-term effects of ambient benzene and TEX (toluene, ethylbenzene,
and xylene combined) on cardiorespiratory mortality in Hong Kong. Environ Int. 117, 91-
98.

Ran, J., et al., 2018b. Are ambient volatile organic compounds environmental stressors for heart
failure? Environmental Pollution. 242, 1810-1816.

Samman Tahhan, A., et al., 2018. Circulating Progenitor Cells and Racial Differences. Circulation
research. 123, 467-476.

Sarkar, M., et al., 2013. Evaluation of spot urine as an alternative to 24 h urine collection for
determination of biomarkers of exposure to cigarette smoke in adult smokers.
Environmental toxicology and pharmacology. 36, 108-114.

Shah, A. S. V., et al., 2015. Short term exposure to air pollution and stroke: systematic review
and meta-analysis. BMJ. 350.

Shim, Y. H., et al., 2019. Association between Heavy Metals, Bisphenol A, Volatile Organic
Compounds and Phthalates and Metabolic Syndrome. International Journal of
Environmental Research and Public Health. 16, 671.

Shuai, J., et al., 2018. Health risk assessment of volatile organic compounds exposure near
Daegu dyeing industrial complex in South Korea. BMC Public Health. 18, 528.

Sithu, S. D., et al., 2010. Exposure to acrolein by inhalation causes platelet activation. Toxicol
Appl Pharmacol. 248, 100-10.

Srivastava, S., et al., 2011. Oral exposure to acrolein exacerbates atherosclerosis in apoE-null
mice. Atherosclerosis. 215, 301-8.

Tang, W., et al., 2000. Estimation of human exposure to styrene and ethylbenzene. Toxicology.
144, 39-50.

Tobler, K., et al., 2010. Reduction of both number and proliferative activity of human endothelial
progenitor cells in obesity. International Journal of Obesity. 34, 687-700.

Törnqvist, H., et al., 2007. Persistent Endothelial Dysfunction in Humans after Diesel Exhaust
Inhalation. American Journal of Respiratory and Critical Care Medicine. 176, 395-400.

Vasa, M., et al., 2001. Number and migratory activity of circulating endothelial progenitor cells
inversely correlate with risk factors for coronary artery disease. Circulation research. 89,
e1-e7.

Villeneuve, P. J., et al., 2013. A cohort study of intra-urban variations in volatile organic
compounds and mortality, Toronto, Canada. Environ Pollut. 183, 30-9.

Wang, G. W., et al., 2008. Acrolein consumption exacerbates myocardial ischemic injury and
blocks nitric oxide-induced PKCepsilon signaling and cardioprotection. J Mol Cell Cardiol.
44, 1016-22.

Werner, N., et al., 2005. Circulating endothelial progenitor cells and cardiovascular outcomes.
New England Journal of Medicine. 353, 999-1007.

Werner, N., Nickenig, G., 2006. Influence of cardiovascular risk factors on endothelial progenitor
cells: limitations for therapy? Arteriosclerosis, thrombosis, and vascular biology. 26, 257-
266.

Xia, W. H., et al., 2012. Age-Related Decline in Reendothelialization Capacity of Human
Endothelial Progenitor Cells Is Restored by Shear Stress. Hypertension. 59, 1225-1231.

Xu, X., et al., 2009. Association between exposure to alkylbenzenes and cardiovascular disease
among National Health and Nutrition Examination Survey (NHANES) participants. Int J
Occup Environ Health. 15, 385-91.

Yang, B.-Y., et al., 2018. Global association between ambient air pollution and blood pressure: A
systematic review and meta-analysis. Environmental Pollution. 235, 576-588.

Ye, D., et al., 2017. Estimating Acute Cardiorespiratory Effects of Ambient Volatile Organic
Compounds. Epidemiology. 28, 197-206.

Yeager, R., et al., 2020. Association between residential greenness and exposure to volatile
organic compounds. Sci Total Environ. 707, 135435.

Yeager, R., et al., 2018. Association Between Residential Greenness and Cardiovascular Disease
Risk. J Am Heart Assoc. 7, e009117.

Zafar, N., et al., 2018. Circulating angiogenic stem cells in type 2 diabetes are associated with
glycemic control and endothelial dysfunction. PLoS One. 13, e0205851.

784   Table 1. Participant Characteristics

Variable	Overall (n=375)
Sex, Male	135 (36%)
Race	
White	311 (83%)
Black	43 (11%)
Other	21 (6%)
Age (years)	50.6 ± 13.2
BMI (kg/m^2)	30.7 ± 7.0
SBP (mmHg)	128.5 ± 18.1
DBP (mmHg)	78.1 ± 11.2
Hypertension	156 (42%)
Diabetes	65 (17%)
MI	24 (7%)
Medication Use	
ASA	46 (13%)
Statin	60 (17%)
ACE or ARB	63 (18%)
Beta Blockers	35 (10%)

786   Values represent mean ± S.D. for continuous variables, and frequency (%) for categorical
787   variables. Abbreviations: BMI indicates body mass index; SBP, systolic blood pressure; DBP,
788   diastolic blood pressure; MI, myocardial infarction; ASA, aspirin; ACE, angiotensin converting
789   enzyme inhibitor; ARB, angiotensin receptor blocker.

801 Table 2. Urinary levels of the major metabolites of volatile organic compounds (VOCs)
802 in study participants.

Parent Structure	VOC	Metabolite	Mean ± SD	Median±IQR	Min-Max	n (%) LOD
	Acrolein	CEMA	115.8 ± 117	89.6 ± 87.5	3.8 - 1213.7	42 (11.2%)
		3HPMA	341.5 ± 344.6	231.9 ± 222.5	46.3 - 2908.4	0 (0%)
	Acrylamide	AAMA	72.4 ± 61.6	56.2 ± 54	8.3 - 677.7	1 (0.3%)
	Acrylonitrile	CYMA	5.7 ± 17.2	1.3 ± 2.4	0.2 - 207.2	231 (61.6%)
	Benzene	MU	133.4 ± 214	56.5 ± 117.1	6 - 1827	6 (1.6%)
	1,3-Butadiene	DHBMA	353.6 ± 175.9	318.8 ± 192.4	50.2 - 1384.6	0 (0%)
		MHBMA3	9.1 ± 8.2	7.2 ± 5.3	0.8 - 75.4	18 (4.8%)
	Crotonaldehyde	HPMMA	308.7 ± 306.2	226.8 ± 172.6	13.5 - 3589.6	1 (0.3%)
	N,N-Dimethylformamide	AMCC	160.4 ± 117.3	134.8 ± 123.3	2.6 - 864.4	5 (1.3%)
	Ethylbenzene, styrene	PGA	225.9 ± 145.5	204.7 ± 146.9	9.8 - 1437.3	11 (2.9%)
	Propylene oxide	2HPMA	58.4 ± 122.9	27.2 ± 30.5	3.4 - 1044.7	4 (1.1%)
	Styrene, ethylbenzene	PHEMA	2.2 ± 4.1	1.4 ± 1.6	0.2 - 52.6	174 (46.4%)
		MA	232.8 ± 165.5	199.6 ± 132.2	20.9 - 1593.3	1 (0.3%)
	Toluene	BMA	11.5 ± 12.1	7.7 ± 8.1	0.1 - 106.9	4 (1.1%)
	Xylene	2MHA	32.8 ± 75.4	13 ± 27.3	0.2 - 1058.9	53 (14.1%)
		3MHA+4MHA	209.1 ± 409	95.8 ± 142.6	3.4 - 5499.1	30 (8%)

804 VOC metabolites were measured in 375 non-smoking participants, 25-70 years of age with low
805 to severe CVD risk. IQR, interquartile range; LOD, limit of detection. The complete chemical
806 names of each metabolite are listed in Supplemental Table 1.

    Table 3. Posterior inclusion probabilities of individual VOC metabolites in the BKMR
    model for 15 CACs.

VOC	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
CEMA	0.53	0.09	0.64	0.74	0.00	0.03	0.49	0.03	0.36	0.00	0.00	0.00	0.11	0.42	0.41
3HPMA	0.43	0.26	0.62	0.68	0.00	0.02	0.54	0.16	0.50	0.13	0.30	0.00	0.15	0.45	0.35
AAMA	0.43	0.10	0.61	0.65	0.00	0.00	0.48	0.06	0.35	0.00	0.00	0.00	0.04	0.42	0.39
CYMA	0.47	0.05	0.68	0.68	0.00	0.01	0.52	0.05	0.38	0.00	0.07	0.00	0.08	0.42	0.34
MU	0.48	0.13	0.60	0.77	0.00	0.01	0.44	0.06	0.38	0.55	0.00	0.00	0.18	0.38	0.39
DHBMA	0.57	0.12	0.61	0.78	0.00	0.02	0.58	0.01	0.33	0.00	0.02	0.00	0.02	0.45	0.47
MHBMA3	0.47	0.10	0.61	0.78	0.01	0.00	0.50	0.01	0.51	0.00	0.00	0.00	0.33	0.35	0.33
HPMMA	0.46	0.14	0.59	0.74	0.01	0.06	0.50	0.01	0.46	0.03	0.01	0.00	0.03	0.36	0.35
AMCC	0.50	0.10	0.57	0.70	0.03	0.01	0.58	0.03	0.35	0.00	0.03	0.00	0.08	0.46	0.39
PGA	0.54	0.19	0.64	0.62	0.03	0.02	0.80	0.03	0.26	0.00	0.02	0.00	0.04	0.44	0.44
2HPMA	0.43	0.17	0.61	0.70	0.00	0.01	0.53	0.04	0.30	0.02	0.05	0.00	0.04	0.37	0.35
PHEMA	0.47	0.07	0.65	0.61	0.01	0.15	0.49	0.05	0.40	0.03	0.00	0.00	0.06	0.40	0.34
MA	0.55	0.12	0.72	0.73	0.02	0.02	0.53	0.04	0.45	0.00	0.00	0.00	0.07	0.49	0.35
BMA	0.42	0.10	0.61	0.71	0.00	0.00	0.46	0.02	0.37	0.01	0.02	0.00	0.91	0.57	0.35
2MHA	0.53	0.19	0.76	0.76	0.02	0.00	0.74	0.02	0.38	0.00	0.06	0.00	0.09	0.34	0.41
3MHA+4MHA	0.46	0.10	0.62	0.89	0.12	0.03	0.71	0.01	0.30	0.00	0.01	0.00	0.05	0.44	0.37

Models (n=375) were adjusted for age, sex, race, BMI, diabetes, hypertension, and daily $PM_{2.5}$ levels.

Scheme 1

A

A

B

C

Scheme 2

Supplemental Results

Supplemental Table 1. Urinary VOC metabolites and Parent Compound

VOC Structure	VOC	VOC metabolite	Common abbr.	LOD (ng/ml)
	Acrolein	N-Acetyl-S-(2-carboxyethyl)-L-cysteine	CEMA	20.35
		N-Acetyl-S-(3-hydroxypropyl)-L-cysteine	3HPMA	9.12
	Acrylamide	N-Acetyl-S-(2-carbamoylethyl)-L-cysteine	AAMA	3.03
	Acrylonitrile	N-Acetyl-S-(2-cyanoethyl)-L-cysteine	CYMA	1.25
	Benzene	*trans, trans*-muconic acid	MU	6.41
	1,3-Butadiene	N-Acetyl-S-(3,4-dihydroxybutyl)-L-cysteine	DHBMA	6.76
		N-Acetyl-S-(4-hydroxy-2-buten-1-yl)-L-cysteine	MHBMA3	1.44
	Crotonaldehyde	N-Acetyl-S-(3-hydroxypropyl-1-methyl)-L-cysteine	HPMMA	1.11
	N,N-Dimethylformamide	N-Acetyl-S-(N-methylcarbamoyl)-L-cysteine	AMCC	9.87
	Ethylbenzene, styrene	Phenylglyoxylic acid	PGA	6.86
	Propylene oxide	N-Acetyl-S-(2-hydroxypropyl)-L-cysteine	2HPMA	3.97
	Styrene, ethylbenzene	N-Acetyl-S-(1-phenyl-2-hydroxyethyl)-L-cysteine + N-Acetyl-S-(2-phenyl-2-hydroxyethyl)-L-cysteine	PHEMA	1.21
		Mandelic Acid	MA	12.93
	Toluene	N-Acetyl-S-(benzyl)-L-cysteine	BMA	0.45
	Xylene	2-Methylhippuric acid	2MHA	1.24
		3-Methylhippuric acid + 4-Methylhippuric acid	3MHA+4MHA	6.85

Supplemental Table 2. Summary Statistics of urinary VOC metabolites for specific subgroups.

VOC metabolite	Male	Female	White	Black	Age <55	Age ≥55	BMI <30	BMI ≥30
n	135	240	311	43	202	173	199	176
CEMA	104.1 ± 87.6	122.4 ± 130.4	111.2 ± 102.5	136.1 ± 110.7	102.1 ± 78.4	131.8 ± 148.7	113.1 ± 132.6	118.9 ± 96.8
3HPMA	326.2 ± 269.2	350.2 ± 380.8	333.6 ± 313.3	330.3 ± 395.3	337.3 ± 325.8	346.5 ± 366.3	346.3 ± 310.9	336.1 ± 380.0
AAMA	73.4 ± 56.9	71.8 ± 64.2	71.2 ± 63.5	83.2 ± 53.5	72.2 ± 56.8	72.7 ± 66.9	77.9 ± 72.1	66.2 ± 46.3
CYMA	4.8 ± 12.3	6.1 ± 19.4	**4.7 ± 15.5**	**12 ± 23.3**	6.4 ± 17.0	4.8 ± 17.4	6.8 ± 20.6	4.4 ± 12.2
MU	125.8 ± 222.5	137.7 ± 209.4	**146 ± 228.7**	**62.2 ± 52.1**	126.0 ± 208.3	142.0 ± 220.8	134.8 ± 229.1	131.7 ±196.2
DHBMA	348.7 ± 182.2	356.4 ± 172.6	363.6 ± 181.3	311.8 ± 148.3	**328.0 ± 157.2**	**383.5 ± 191.6**	355.0 ± 166.6	352.1 ± 186.3
MHBMA3	8.4 ± 8.6	9.5 ± 8	9.3 ± 8.6	8.2 ± 6.1	9.2 ± 7.9	9.0 ± 8.6	**10.2 ± 9.6**	**7.9 ± 6.1**
HPMMA	286.1 ± 271.3	321.5 ± 324.1	305.1 ± 265.4	347.3 ± 529.9	**275.4 ± 241.8**	**328.6 ± 267.7**	315.1 ± 292.7	282.8 ± 203.8
AMCC	**140 ± 99.4**	**171.9 ± 125.1**	**169.3 ± 121.1**	**118.5 ± 74.2**	**143.2 ± 104.3**	**180.5 ± 128.3**	165.7 ± 119.9	154.5 ± 114.4
PGA	**202.2 ± 117.2**	**239.1 ± 157.9**	232.2 ± 151.8	196.3 ± 105	215.5 ±149.0	238.0 ± 140.9	226.8 ± 154.1	224.9 ± 135.7
2HPMA	67.7 ± 160.1	53.1 ± 95.8	60.1 ± 131.4	40.8 ± 31.8	58.1 ± 116.8	58.6 ± 130.0	67.1 ± 147.8	48.4 ± 86.0
PHEMA	**2.3 ± 5.4**	**2.2 ± 3**	2.3 ± 4.4	1.7 ± 1.2	2.1 ± 3.9	2.3 ± 4.2	**2.2 ± 3.9**	**2.2 ± 4.2**
MA	**206.3 ± 124.1**	**247.7 ± 183.3**	**243.7 ± 175.6**	**177.8 ± 91.8**	**210.7 ± 139.5**	**258.7 ± 188.7**	231.4 ± 156.0	234.5 ± 176.0
BMA	**10.7 ± 12.4**	**12 ± 11.9**	11.4 ± 12.1	11.6 ± 11.8	**10.5 ± 11.9**	**12.7 ± 12.2**	**12.2 ± 12.8**	**10.7 ± 11.1**
2MHA	26.3 ± 37.1	36.5 ± 89.8	33.8 ± 78.7	25.5 ± 32.6	27.6 ± 50.4	32.9 ± 56.7	**37.9 ± 63.7**	**21.2 ± 36.8**
3MHA+4MHA	189 ± 313.1	220.5 ± 454.3	213 ± 436	193.8 ± 205.9	**179.8 ± 310.6**	**212.5 ± 293.9**	227.4 ± 367.2	158.4 ± 204.2

Values represent Mean ± SD. Units: µg/g creatinine. Bold values represent significant difference between specific subgroups, based on T-test of log-transformed VOC metabolites.

Supplemental Table 2 continued. Summary Statistics of urinary VOC metabolites for specific subgroups.

VOC metabolite	Non-Diabetic	Diabetic	Normotensive	Hypertensive	Low PM$_{2.5}$	High PM$_{2.5}$
n	310	65	219	156	189	186
CEMA	116.8 ± 123.3	111.2 ± 81	104.7 ± 85.1	131.4 ± 149.8	117.0 ± 92.4	114.6 ± 137.9
3HPMA	338.7 ± 341.3	355 ± 362.4	327.9 ± 315.8	360.6 ± 381.7	371.4 ± 411.7	311.2 ± 257.0
AAMA	74.6 ± 64.3	62.2 ± 45.5	70.7 ± 55.6	74.9 ± 69.3	65.0 ± 41.6	80.0 ± 76.1
CYMA	6.1 ± 18.5	3.6 ± 8.7	5.7 ± 15.9	5.6 ± 18.9	4.4 ± 12.3	7.0 ± 21.0
MU	**128.1 ± 213.7**	**158.4 ± 215.3**	**127.5 ± 227.6**	**141.6 ± 193.7**	142.7 ± 257.3	123.9 ± 158.5
DHBMA	354.3 ± 172.8	350.4 ± 191.2	347.4 ± 164.4	362.4 ± 191.2	**368.6 ± 182.0**	**338.5 ± 168.6**
MHBMA3	9.3 ± 8.6	8.3 ± 6.2	8.9 ± 7.7	9.5 ± 8.9	9.1 ± 8.4	9.2 ± 8.1
HPMMA	300.9 ± 260.8	346 ± 466.9	277.8 ± 228	352.1 ± 387.2	308.9 ± 278.3	290.9 ± 229.5
AMCC	157.3 ± 108.6	175.1 ± 152.4	154.8 ± 109.2	168.3 ± 127.8	153.5 ± 102.8	167.4 ±130.4
PGA	**219.3 ± 143.1**	**257.2 ± 154**	222.7 ± 156.8	230.3 ± 128.6	**208.3 ± 133.0**	**243.7 ± 155.6**
2HPMA	62 ± 132.5	40.9 ± 55.8	**66.1 ± 137**	**47.4 ± 99.3**	64.8 ± 144.2	51.8 ± 96.5
PHEMA	2.3 ± 4.4	1.8 ± 1.9	2.3 ± 4.6	2.1 ± 3.2	2.1 ± 3.6	2.3 ± 4.5
MA	228.5 ± 155.5	253.4 ± 206.9	223.9 ± 154.4	245.3 ± 179.6	217.1 ± 146.2	248.9 ± 182.1
BMA	12 ± 12.7	9.4 ± 8.1	11.7 ± 12.7	11.3 ± 11.1	10.8 ± 11.4	12.3 ± 12.7
2MHA	35 ± 80	22.5 ± 46.6	37.4 ± 91.6	26.5 ± 43.2	32.8 ± 64.7	27.3 ± 38.8
3MHA+4MHA	223.7 ± 445.8	140.2 ± 115.9	226.4 ± 480	185.2 ± 282.4	211.2 ±368.4	178.4 ± 218.4

Values represent Mean ± SD. Units: μg/g creatinine. Bold values represent significant difference between specific subgroups, based on T-test of log-transformed VOC metabolites. Low PM$_{2.5}$ ranges from 1.2-7.8 μg/m^3. High PM$_{2.5}$ ranges from 8.0-17.7 μg/m^3.

Supplemental Figure 1. Pearson correlation coefficients between log-transformed urinary VOC metabolites and ambient pollutant and meteorological data (n=375).

Supplemental Figure 2. Heat map of pair-wise associations between the urinary metabolites of VOCs and circulating levels of CACs. False Discovery rate adjusted p-values. Generalized linear models (n=375) were adjusted for age, sex, race, BMI, diabetes, hypertension and: daily PM$_{2.5}$ + Temperature (A); daily PM$_{2.5}$ + Humidity (B); Ozone + Temperature (C); Ozone + Humidity (D). Black grids represent positive associations and blue grids represent negative associations. The intensity of colors represent p-values, with darker grids indicating smaller p-values.