Child and caregiver mental health during 12 months of the COVID-19 pandemic in Australia

Authors:
*Anna M. H. Price[1,2,3], Mary-Anne Measey[2, 4], Monsurul Hoq[5], Anthea Rhodes[2,3,6, ¶], Sharon Goldfeld[1,2,3,¶]

Affiliations^:
1. Centre for Community Child Health, The Royal Children's Hospital, Parkville, Vic
2. Population Health, Murdoch Children's Research Institute, Parkville, Vic
3. Department of Paediatrics, The University of Melbourne, Parkville, Vic
4. Health Services Research Unit, The Royal Children's Hospital, Parkville, Vic
5. Clinical Epidemiology and Biostatistics Unit, Murdoch children's Research Institute, Parkville, Vic
6. Department of General Medicine, The Royal Children's Hospital, Parkville, Vic
^All Australia,
*Corresponding author, anna.price@mcri.edu.au
¶ AR and SG are joint senior authors.

Word Count: 2500 words

ABSTRACT

Background: There are urgent calls for research into the mental health consequences of living through the COVID-19 pandemic. Australia's effective suppression of COVID-19 offers insights into the pandemic mental health impacts in the relative absence of the disease. We aimed to describe the mental health experiences of caregivers and children over 12 months, reporting differences related to demographic, socioeconomic and lockdown characteristics.

Methods: Data were drawn from Australia's only nationally representative, repeated cross-sectional survey of caregivers with children (0-17 years). N=2020 caregivers participated in June 2020, N=1434 in September 2020, and N=2508 in July 2021. Caregivers reported their mental health (poor versus not, Kessler-6), and mental health impacts of the pandemic (negative versus none/positive) for themselves and their children. Data were weighted to approximate population distributions of caregiver age, gender, sole-caregiving, number and ages of children, state/territory and neighbourhood-level disadvantage.

Results: Negative mental health impacts were more common for female (vs male) caregivers, older (vs younger) children. Poor caregiver mental health (K6) was more common for families experiencing socioeconomic adversity (especially financial), while perceived negative impacts were more common for more socially advantaged groups. Caregivers who experienced the least total lockdown reported similar K6 over time. Otherwise, poor mental health and negative mental health impacts increased over time with cumulative lockdown.

Conclusion: Despite Australia's low infection rates, the mental health impacts of the pandemic and related public health restrictions are real and concerning. Addressing poor mental health must be central to ongoing pandemic recovery efforts for families and children.

What is known about this topic?

- The global evidence shows that, for general adult populations, psychological distress peaked in the first months of the COVID-19 pandemic before appearing to improve.
- Less is known about mental health over time of living through the pandemic, especially for caregivers and children. There are urgent calls for research.
- Due to low infection rates, Australia's experience can provide insight into the mental health impacts of lockdown with minimal compounding harms of the virus.

What this study adds:

- From June 2020 to July 2021, Australia's lockdowns were detrimental for caregiver and child mental health.
- There was an unequal distribution of mental health impacts based on caregiver gender, child age and family socioeconomic characteristics.
- Pandemic response and recovery planning must consider both family mental health and socioeconomic security.

INTRODUCTION

During the first eighteen months of the COVID-19 pandemic, balancing the direct harms of the virus with the indirect harms of public health restrictions was a global challenge. Australia enacted some of the strictest public health measures in the world to effectively suppress infection.[1] By 31 July 2021, the measures successfully contained infection to an overall incidence rate of 137 cases and 3.7 deaths per 100,000 people.[2] With just 15% of the population of 25 million fully vaccinated by the end of July 2021, stay-at-home orders ('lockdown') were the country's main measure of disease control.[3] Families caring for children and young people (referred to as children hereafter) have been substantially affected by lockdown,[4 5] required to juggle working from home, responsibility for their children's remote learning and care during school and childcare closures, with minimal social opportunities due to closure of sports and play centres, and reduced access to healthcare. Despite mental health having "greater impact on human activity than any other non-communicable illness",[4] it appeared to be given little consideration in the majority of Australia's lockdown policies. With future variants and pandemics likely, Australia's experience offers insight into the mental health impacts of lockdown, effectively without the compounding direct harms of the virus.

The evidence for the indirect and longer-term impacts of the pandemic is quickly evolving. In a review of the global mental health evidence from the first year (to April 2021), Aknin et al., reported a peak in adults' psychological distress in the early months.[4] While many studies reported a decline to pre-pandemic levels by mid-2020,[4] the authors found that mental health inequities were sustained or exacerbated for adults who were younger, female, child-rearing, or with fewer socioeconomic resources.[4 5] However, published data on the specific experience of parents (termed caregivers throughout) are scarce. Evidence reviews of children's experiences also suggest that their mental health is declining during the pandemic.[6 7] In Racine and colleagues' (2021) meta-analysis of 29 studies published in the first year of the pandemic, prevalence estimates for depression and anxiety in

children doubled, and were higher over time, and for older adolescents and females.[8] However, the available systematic reviews are limited by over-representation of data from the early months of the pandemic,[6-8] and previous pandemics show that negative mental health effects can persist.[9] As such, there are urgent calls for research into the "long-term mental health consequences of living through the pandemic",[4] which prioritises children.[5]

In response to these calls, we aimed to describe the impact of the pandemic on caregiver and child mental health using Australia's only nationally representative, repeated cross-sectional survey of families during the pandemic. Unlike many high-income countries, Australia's low incidence of COVID-19 makes it possible to examine the mental health experiences in the relative absence of the disease. Based on the gender, age and socioeconomic disparities identified in the published evidence reviews,[4 8] we hypothesised that poor mental health and negative impacts of the pandemic on mental health would be more common: (i) for caregivers who identified as female compared with male, and older compared with younger children, (ii) for families experiencing greater socioeconomic adversity compared with more socioeconomic advantage and (iii) with cumulative (increasing total length) of lockdown.

METHODS

Design and procedure: The Royal Children's Hospital (RCH) National Child Health Poll comprises periodic cross-sectional surveys of approximately 2000 Australian caregivers of children aged 0-17 years. Data collection is contracted to the Online Research Unit who obtain written informed consent and draw a nationally representative sample of caregivers using stratified random sampling from their panel of over 350,000 adults aged 18 years or older, who live in Australia and have internet access. Surveys are administered in sixth grade-equivalent (end of elementary school) English, anonymous, and respondents are remunerated with points exchangeable for department store gift vouchers. Questions about mental health – the focus of the current study – were introduced after COVID-19 began and were collected in three surveys. The dates of data collection were: (1) 15-23 June 2020, after a first national lockdown (March-May 2020) eased; (2) 15-29 September 2020, when only metropolitan residents of Victoria were in a second, stricter lockdown (July-November 2020); and (3) 20-29 July 2021, when multiple states/territories were in and out of lockdown (June 2021 onwards)). Each survey undergoes pilot testing with a group of health professionals and families before going to field. The RCH Human Research Ethics Committee approved this research (February 2020, #35254).

Measures: Each survey collected the demographic, socioeconomic and pandemic-related financial characteristics described in Table . Table 1 also describes how measures of current and cumulative lockdown were defined. Families were assigned the Australian Bureau of Statistics' (ABS) Socio-Economic Indexes for Areas (SEIFA) Index of Relative Disadvantage, a national area level index derived from census data for all individuals living in a postcode, with higher scores indicating greater advantage. Caregivers self-reported their mental health (Kessler-6), and the impact of the pandemic on their own and their children's mental health (details in Table 1).

Analysis: To reduce effects of non-response and non-coverage and therefore approximate the population distributions of mental health experiences, the mental health measures were weighted using ABS distributions of caregiver age, gender, family structure (sole-caregiving, number of children and any under 5 years), state/territory and SEIFA. Mental health measures were described with weighted proportions for each survey, overall and by demographic, socioeconomic and pandemic-related financial and lockdown characteristics. For each of the three mental health measures, the association between mental health measure and time of the survey, according to state/territory grouping (as a proxy for cumulative lockdown, see Table 1), were calculated using generalised linear models. The binomial models investigated whether cumulative lockdown worsened the mental health measures over time, after adjusting for caregiver gender, sole caregiver status, education, SEIFA, and cultural or linguistic diversity (measures described in Table 1). The child mental health model was additionally adjusted for poor caregiver mental health (Kessler-6, see Table 1) and clustering at the level of family. The estimated models were then used to predict probabilities of each mental health outcome, by state/territory grouping, at the three survey timepoints, adjusted for the covariates in the model.

Table 1. Demographic, socioeconomic, lockdown and mental health measures

Measure	Description
Demographic	
Age	Collected for caregivers and children, reported in years. Child ages were categorised into pre-school (0-4 years), primary/elementary school (5-11 years) and high school (12-17 years).
Gender	Collected for caregivers and children: response options "male", "female", "other". Two caregivers and no children identified as "other". Before the 2020 Australian census, "other" gender data were not collected for the Australian population, so it was not possible to calculate weights for these caregivers. Hence, they were dropped from the analysis, and the gender variable was dichotomised into "female" compared with "male".
Sole caregiver	Question "Are you the sole (single) parent or carer of a child 17 years of age or younger?", binary response options "yes" (one-caregiver household) compared with "no" (multi-caregiver household).
Caregiver education	Question "What is the highest level of schooling / education you have completed?" Trichotomised into "Year 12 or less" (response options: less than year 10, Year 10 or equivalent (e.g. school certificate), Year 12 or equivalent); "vocational training certificate" (response options: trade/apprenticeship (e.g. carpenter), certificate/diploma (e.g. Cert IV Childcare)); or "university degree" (response options: undergraduate university degree, postgraduate university degree (e.g. Masters, Doctorate, PhD).
Caregiver country of birth	Question "Were you born in Australia", binary response options "yes" (born in Australia) compared with "no" (outside Australia).
Home language	Question "Do you speak a language other than English at home?", binary response options "yes" (other than English) compared with "no" (English).
Cultural or linguistic diversity	A composite of the two above variables, caregiver country of birth and home language, to represent respondents who answered "yes" to either or both items, compared with "no" (answering no to both items).
Socioeconomic and lockdown	
Job/income loss during COVID-19	Three items drawn from the CoRonavIruS Health Impact Survey (CRISIS) caregiver version:[10] "What changes in employment or income have occurred in your household due to coronavirus/COVID-19?" (binary response options "yes" compared with "no") including: "job loss by one caregiver"; "job loss by two caregivers" and "reduced total household income". A binary variable describing any job loss (by one or two caregivers) or reduction in income due to COVID-19 (compared with not) was created.
Low income (<$1000)	A binary variable based on current total household income before tax, categorized into 10 options ranging from "less than $500 p/week" to "more than $3,000 p/week", plus "prefer not to say". The binary variable summarized low income ("less than AU$1,000 p/week" compared with more) based on Australian thresholds for income-poverty.[11] n=767 caregivers preferred not to report income, so this variable should be interpreted with caution.
Couldn't afford essential items	Eight items adapted from the Household, Income and Labour Dynamics in Australia (HILDA) Survey Wave 18 Household Questionnaire Material Deprivation Module [12] asking "In the last month, because of money pressure did you miss or put off" (binary response options: "yes" compared with "no"): mortgage or rent repayments; electricity, gas, water bills; food; healthcare; prescription medicines; home or car insurance; mobile phone bills; and internet. A binary summary variable was created to denote "any material deprivation" inability to pay for one or more essential items compared with "none".
Current lockdown	All Australian states and territories experienced a first national, 10-week lockdown from 23 March to 1 June 2020. Victoria experienced a second extended and more severe, 20-week lockdown from 8 July-23 November 2020. There were several short (less than a week) lockdowns in the intervening months across states, before New South Wales experienced its second extended period of lockdown, which began incrementally on 26 June 2021. Victoria experienced its fifth (in total) lockdown from 16-27 July 2021, before an extended lockdown subsequently began a week later on 5 August 2021. Current lockdown was therefore categorised as follows. • The national lockdown had eased for the collection of the June 2020 Poll, so no (denoted 'N' in Tables 2-3) Australians were in lockdown. Thus, the

	• number/proportion not in lockdown represents the whole cohort (and is equal to the 'overall' numbers/proportions at top of Tables 2-3). • For the September 2020 Poll, only residents of metropolitan Melbourne in the state of Victoria were in lockdown (denoted 'Y' for "yes") compared with all other Australians (N). This was categorised according to respondents who reported living in Victoria and living in "metropolitan" compared with "regional/rural" areas. • By July 2021, many states/territories were going in and out of lockdown, and this question was added into the Poll and reported on directly with the question "are you currently under stay-at-home orders or restrictions due to COVID-19 (also known as 'lockdown')?", responses "yes" compared with "no".
Cumulative lockdown	Trichotomous variable based on total length of lockdown experienced by each state/territory. By 31 July 2021, the cumulative length of COVID-19 lockdown was greatest for the state of Victoria ("Vic", total 31 weeks); followed by the state of New South Wales ("NSW", total 15 weeks) and then all "other" states and territories (total range 10-12 weeks).
Mental health	
Caregiver mental health	6 items of the Kessler-6 (K6) assessing caregivers' self-reported anxiety and depressive symptoms encountered in the last 4 weeks. Scored on a 5-point Likert scale from 1 "none of the time" to 5 "all of the time". Summarized into (a) a continuous total score, and (b) a binary variable indicating "poor mental health" (total score 19 or more) compared with not (total score 6-18).[13]
Impact of the pandemic on mental health	A 5-point item adapted from UK Young Minds Matter Study,[14] describing the impact of COVID-19 on mental health, dichotomized into negative ("small negative/large negative") compared with positive ("none/small positive/large positive"). Reported by caregivers for (a) themselves and (b) each child.

RESULTS

In June 2020, n=2020 of 2697 (74.9%) caregivers approached provided data for themselves and 3411 children. In September 2020, 1434/1769 (81.1%) caregivers provided data for 2553 children. In July 2021, 2508/3925 (63.9%) provided data for 4327 children. Table 2 presents the sample characteristics for each survey. There were some differences between surveys in characteristics such as the proportion of respondents caring for young children, caregiver gender, sole caregiving, and SEIFA; characteristics that were used to create the sample weights (see *Analysis*).

For the three surveys overall, children's mean age was 9.3 years (standard deviation (SD) 5.0 years) and 47.8% were female. Caregivers' mean age was 42.2 years (SD 9.1 years), 54.0% were female, and each caregiver cared for 1.7 children on average (SD 0.8). One-quarter were sole caregivers, 56.6% had a university degree, 25.6% were born outside Australia, and 22.4% spoke a language other than English at home. Twenty-eight percent of caregivers reported job/income loss due to COVID-19; 17.2% reported low household income and 29.3% were unable to afford at least one essential item in the month prior. The proportions of respondents in lockdown were 0% (June 2020), 29.2% (September 2020) and 56.5% (July 2021).

Table 3 presents the weighted mental health experiences for each survey, separately by characteristics. When considering patterns over time, the proportions of respondents who reported poor mental health according to the K6 was similar between surveys. Negative mental health impacts for caregivers became more common between June 2020 and July 2021, although there were no consistent patterns between June 2020 and September 2020, or between September 2020 and July 2021. In contrast, there was a clear pattern of increasingly common negative mental health impacts for children across the three waves of data collection.

When considering differences between subgroups, negative mental health impacts for caregivers and children were more common when reported by female than male caregivers, and for older compared with younger children, but were similar between child genders. Poor mental health (K6) was more common for groups typically experiencing more social adversity, such as sole caregiving compared with multi-caregiver households, having a home language other than English (but not for country of birth), and several financial and lockdown characteristics (lower SEIFA and income, couldn't afford essentials, and cumulative or current lockdown). In contrast with the K6, negative mental health impacts for caregivers and children were more common for families experiencing greater social advantage, such as multi-caregiver (compared with sole caregiver) households and higher (compared with lower) education, SEIFA, and income. The proportions of respondents reporting all three negative mental health measures increased for families who had lost jobs or income due to the pandemic, and with cumulative lockdown.

Figure 1 shows that the three measures of negative mental health experiences increased for Victorian and NSW families by July 2021 compared with June 2020. Increases in the proportions of caregivers reporting poor mental health (K6) aligned with the second extended lockdowns; for Victorians this was September 2020 and for NSW it was July 2021 (supplementary data (a)). In contrast, there was weak evidence of change in the proportions of caregivers in Other states/territories reporting poor mental health (K6) over time. The proportions of caregivers and children experiencing negative mental health impacts due to the pandemic increased from June 2020 to July 2021 (supplementary data (b and c)). Increased proportions of Victorians reported negative impacts in September 2020 and again in July 2021. More caregivers from NSW and the Other states/territories self-reported negative impacts in July 2021 compared with June or September 2020.

Table 2. Sample characteristics by survey, described with the number of respondents (proportion).

Characteristic	Subgroup	Number of respondents (%)		
		June 2020	**Sept 2020**	**July 2021**
Child		**N=3411**	**N=2553**	**N=4327**
Age in years	0-4	628 (18.4)	520 (20.4)	1291 (29.8)
	5-11	1356 (39.8)	1153 (45.2)	1705 (39.4)
	12-17	1427 (41.8)	880 (34.5)	1331 (30.8)
Gender	F	1664 (48.7)	1242 (48.7)	2013 (46.5)
	M	1747 (51.2)	1311 (51.4)	2314 (53.5)
Caregiver		**N=2020**	**N=1434**	**N=2508**
Gender	F	1003 (49.7)	758 (52.9)	1458 (58.1)
	M	1017 (50.4)	676 (47.1)	1050 (41.9)
Sole carer	Yes	462 (22.9)	282 (19.7)	704 (28.1)
	No	1558 (77.1)	1152 (80.3)	1804 (71.9)
Education [a]	Y12	326 (16.1)	239 (16.7)	422 (16.8)
	Cert.	518 (25.6)	390 (27.2)	693 (27.6)
	Uni.	1176 (58.2)	805 (56.1)	1393 (55.5)
Born outside Australia	Yes	546 (27.4)	329 (23.4)	622 (25.4)
	No	1444 (72.6)	1078 (76.6)	1827 (74.6)
Home language other than English	Yes	413 (20.5)	324 (22.6)	600 (23.9)
	No	1607 (79.6)	1110 (77.4)	1908 (76.1)
Socio-economic				
SEIFA quintile (1=most disadvantage, 5=least disadvantage)	1	213 (10.6)	172 (12.0)	327 (13.0)
	2	273 (13.5)	245 (17.1)	391 (15.6)
	3	370 (18.3)	250 (17.4)	511 (20.4)
	4	457 (22.7)	319 (22.3)	571 (22.8)
	5	705 (34.9)	448 (31.2)	707 (28.2)
Low income (<$1000) [c]	Yes	254 (14.7)	212 (16.9)	427 (19.2)
	No	1480 (85.3)	1040 (83.1)	1782 (80.7)
Couldn't afford essential items [d]	Yes	527 (26.1)	400 (27.9)	822 (32.8)
	No	1493 (73.9)	1034 (72.1)	1686 (67.2)
Pandemic experience				
Job/income loss [b]	Yes	559 (27.7)	381 (26.6)	711 (28.4)
	No	1461 (72.3)	1053 (73.4)	1797 (71.7)
Current lockdown [e]	Yes	0 (0)	419 (29.2)	1416 (56.5)
	No	2020 (100)	1015 (70.8)	1092 (43.5)
State as a proxy for cumulative lockdown (Vic=most, Other=least) [e]	Vic	612 (30.3)	473 (33.0)	746 (29.7)
	NSW	617 (30.5)	449 (31.3)	791 (31.5)
	Other	791 (39.2)	512 (35.7)	971 (38.7)

Vic: Victoria, NSW: New South Wales. Data were weighted using national demographic distributions for caregiver age, gender, family structure (sole-caregiving, number of children and any under 5 years), state/territory and Socio-Economic Indexes for Areas Index of Relative Disadvantage (SEIFA).

[a] Highest education coded as Year 12 or less (up to the end of high/secondary school; Y12); vocational training certificate ("cert."); or university degree ("uni").

[b] Job loss by one or two adults, or reduction in income, due to COVID-19.

[c] Low-income defined according to Australian definitions of income-poverty. Missing 767 caregivers who preferred not to report income.

[d] Any one or more of mortgage or rent; electricity, gas, water bills; food; healthcare; prescription medicines; home or car insurance; mobile phone bills; internet, in the last month.

[e] No Australians were in lockdown for the June 2020 Poll, so the number/proportion is for the whole cohort. For the September 2020 Poll, only residents of metropolitan Melbourne in the state of Victoria were in lockdown (Y) compared with all other Australians (N). In July 2021, many states/territories were going in and out of

lockdown, and this question was asked directly (Y/N). Overall, Victorians experienced the most (cumulative) lockdown, followed by NSW, followed by Other states and territories.

Table 3. Mental health experiences by sample characteristic and survey, described with the number of respondents (weighted proportion).

Characteristic	Subgroup	Poor caregiver mental health (K6)[f]			Negative impact on caregiver[g]			Negative impact on child[g]		
		June 2020 N=2020	Sept 2020 N=1434	July 2021 N=2508	June 2020 N=2020	Sept 2020 N=1434	July 2021 N=2508	June 2020 N=3411	Sept 2020 N=2553	July 2021 N=4327
Overall	All	264 (17.4)	247 (19.5)	541 (18.9)	949 (47.4)	708 (50.3)	1490 (60.0)	1055 (25.6)	884 (33.1)	1807 (44.0)
Child										
Age in years	0-4	-	-	-	-	-	-	81 (12.8)	101 (19.9)	330 (24.2)
	5-12	-	-	-	-	-	-	440 (32.3)	414 (37.3)	789 (46.9)
	13-17	-	-	-	-	-	-	534 (38.2)	369 (43.7)	688 (51.5)
Gender	F	-	-	-	-	-	-	525 (25.1)	438 (32.8)	876 (45.5)
	M	-	-	-	-	-	-	530 (26.0)	446 (33.4)	931 (42.7)
Caregiver										
Gender	F	153 (18.3)	110 (18.0)	338 (20.5)	489 (49.6)	410 (53.9)	920 (63.7)	512 (25.9)	521 (35.1)	1099 (45.7)
	M	111 (15.8)	137 (21.8)	203 (16.6)	460 (44.1)	298 (44.7)	570 (54.7)	543 (25.1)	363 (30.0)	708 (41.6)
Sole carer	Yes	137 (36.1)	100 (36.5)	268 (32.6)	199 (42.7)	99 (36.2)	335 (44.3)	190 (24.3)	144 (27.1)	397 (30.9)
	No	127 (10.2)	147 (14.8)	273 (15.0)	750 (49.2)	609 (54.1)	1155 (64.5)	865 (26.0)	740 (34.8)	1410 (47.9)
Education[a]	Y12	55 (21.7)	61 (26.9)	124 (15.8)	141 (40.2)	98 (42.9)	211 (47.3)	148 (21.2)	115 (24.3)	293 (39.2)
	Cert.	73 (21.4)	65 (19.9)	152 (20.0)	244 (48.7)	204 (56.1)	418 (64.6)	293 (28.6)	264 (34.9)	524 (51.1)
	Uni.	136 (15.8)	121 (16.9)	265 (17.5)	564 (48.8)	406 (49.6)	861 (61.5)	614 (25.3)	505 (34.9)	990 (41.8)
Born outside Australia	Yes	58 (12.5)	40 (11.7)	124 (16.9)	243 (42.4)	169 (49.2)	379 (60.0)	253 (23.0)	184 (32.0)	396 (40.2)
	No	198 (18.9)	198 (21.0)	405 (19.6)	694 (49.4)	530 (50.8)	1080 (60.1)	789 (26.8)	695 (33.7)	1375 (45.3)
Home language other than English	Yes	80 (20.9)	79 (22.5)	176 (25.1)	172 (40.6)	154 (47.6)	337 (58.9)	178 (21.7)	181 (31.7)	352 (39.5)
	No	184 (16.3)	168 (18.6)	365 (17.0)	777 (49.4)	554 (51.1)	1153 (58.9)	877 (26.8)	703 (33.5)	1455 (45.4)
Socio-economic										
SEIFA quintile (1=most disadvantage, 5=least disadvantage)	1	41 (22.1)	35 (20.7)	97 (26.1)	86 (41.4)	77 (47.4)	172 (56.0)	115 (27.1)	90 (31.3)	223 (42.0)
	2	50 (20.5)	53 (25.6)	99 (21.4)	125 (46.1)	117 (51.3)	215 (60.6)	133 (21.1)	157 (32.6)	262 (45.1)
	3	49 (16.2)	41 (19.3)	114 (17.9)	186 (53.1)	129 (51.9)	295 (58.7)	191 (27.3)	170 (34.5)	350 (40.8)
	4	66 (17.9)	54 (20.8)	108 (16.0)	205 (44.1)	149 (49.6)	350 (62.4)	242 (26.6)	172 (31.0)	392 (42.8)
	5	58 (9.4)	64 (11.1)	123 (16.5)	347 (51.7)	236 (51.0)	458 (67.1)	374 (26.3)	295 (35.7)	580 (52.9)
Low income (<$1000)[c]	Yes	69 (27.5)	58 (31.3)	171 (29.5)	120 (53.4)	89 (45.6)	230 (51.2)	111 (26.1)	133 (33.3)	271 (35.3)
	No	165 (15.5)	167 (17.3)	326 (16.6)	686 (45.2)	507 (49.2)	1064 (61.0)	774 (24.8)	631 (32.4)	1292 (44.6)

Characteristic	Subgroup	Poor caregiver mental health (K6) [f]			Negative impact on caregiver [g]			Negative impact on child [g]		
		June 2020 N=2020	Sept 2020 N=1434	July 2021 N=2508	June 2020 N=2020	Sept 2020 N=1434	July 2021 N=2508	June 2020 N=3411	Sept 2020 N=2553	July 2021 N=4327
Couldn't afford essential items [d]	Yes	359 (37.4)	156 (38.4)	341 (35.5)	263 (50.3)	165 (44.8)	469 (59.2)	270 (26.5)	236 (33.9)	586 (42.6)
	No	96 (8.8)	91 (12.3)	200 (11.2)	686 (46.2)	543 (52.4)	1021 (60.4)	785 (25.1)	648 (32.8)	1221 (44.7)
Pandemic experience										
Job/income loss [b]	Yes	96 (23.7)	93 (27.5)	218 (28.8)	306 (51.0)	224 (59.5)	486 (71.1)	305 (25.0)	297 (40.7)	613 (55.8)
	No	168 (15.1)	288 (16.4)	323 (15.3)	643 (46.1)	484 (46.8)	1004 (55.9)	750 (25.7)	587 (30.2)	1194 (39.7)
Current lockdown [e]	Yes	n/a	83 (24.0)	353 (23.5)	n/a	240 (58.1)	867 (62.9)	n/a	355 (44.1)	1114 (49.6)
	No	264 (17.4)	164 (17.7)	188 (13.3)	949 (47.4)	468 (47.3)	623 (56.4)	1055 (25.6)	529 (28.7)	693 (37.3)
State as a proxy for cumulative lockdown (Vic=most, Other=least) [e]	Vic	85 (17.7)	93 (23.3)	200 (25.1)	284 (48.9)	266 (57.5)	483 (66.4)	321 (23.8)	389 (42.5)	626 (51.4)
	NSW	96 (21.6)	78 (18.6)	188 (22.0)	307 (49.2)	213 (49.0)	477 (60.8)	306 (25.9)	267 (34.4)	603 (48.9)
	Other	83 (12.4)	76 (16.6)	153 (11.2)	358 (43.9)	229 (44.1)	530 (54.4)	428 (26.9)	228 (21.0)	578 (33.7)

Vic: Victoria, NSW: New South Wales. proportions were weighted using national demographic distributions for caregiver age, gender, family structure (sole-caregiving, number of children and any under 5 years), state/territory and Socio-Economic Indexes for Areas Index of Relative Disadvantage (SEIFA). .

[a] Highest education coded as Year 12 or less (up to the end of high/secondary school; Y12); vocational training certificate ("cert."); or university degree ("uni").

[b] Job loss by one or two adults, or reduction in income, due to COVID-19.

[c] Low-income defined according to Australian definitions of income-poverty. Missing 767 caregivers who preferred not to report income.

[d] Any one or more of mortgage or rent; electricity, gas, water bills; food; healthcare; prescription medicines; home or car insurance; mobile phone bills; internet, in the last month.

[e] No Australians were in lockdown for the June 2020 Poll, so the proportion is for the whole cohort (see 'Overall' at top of Table). For the September 2020 Poll, only residents of metropolitan Melbourne in the state of Victoria were in lockdown (Y) compared with all other Australians (N). In July 2021, many states/territories were going in and out of lockdown, and this question was asked directly (Y/N). Overall, Victorians experienced the most (cumulative) lockdown, followed by NSW, followed by Other states and territories.

[f] Kessler-6 (K6) dichotomized into a binary "poor mental health" (total score 19 or more) compared with not (total score 6-18).

[g] Dichotomised into negative ("small negative/large negative") compared with positive ("none/small positive/large positive").

DISCUSSION

In the only nationally representative and repeated cross-sectional survey of Australian

families over 12 months of the COVID-19 pandemic, there were unequal distributions of

mental health impacts based on caregiver gender, child age, socioeconomic and pandemic-

related characteristics. As hypothesised, negative mental health impacts were more common

for female compared with male caregivers, and older children compared with younger

children. However, poor caregiver mental health (K6) was similar between female and male

genders. Job or income loss due to the pandemic was detrimental for caregiver and child

mental health. While poor caregiver mental health (K6) was more common for families

experiencing greater social adversity, reported negative impacts were more common for more

socially advantaged groups. Caregivers in states/territories who experienced the least

cumulative lockdown reported similar K6 scores over time. Otherwise, poor mental health

and negative mental health impacts increased over the course of the three surveys and with

cumulative lockdown.

Comparable data on the mental health experiences specific to caregivers of children during

the pandemic are limited. Aknin and colleagues' review of mental health experiences during

COVID-19 suggest that levels of psychological distress in general adult populations may

have declined from an early peak in the first months of the pandemic.[4] In our study, a similar

improvement was found for the mental health (K6) of Australian caregivers in the

states/territories that experienced the least lockdown. However, our data demonstrated

evidence that negative mental health experiences increased over time and were strongly

related to cumulative lockdown. These findings align with and extend data from the Pulse of

the Nation (TPTN), a weekly, cross-sectional survey of representative Australian adult

samples throughout the pandemic.[15] Like our study, TPTN data indicated that mental distress

in adults (measured with a single item that highly correlated with the K6) increased during

2020, which the authors describe as driven by increasing financial stress.[15] Notably, increases in mental distress were highest for parents, tripling from 8% pre-pandemic to 24% during the pandemic.[15] While the RCH Poll lacked pre-pandemic data, substantially more caregivers reported poor mental health on the K6 (average 18%) in the Polls throughout the pandemic than representative Australian adult data collected pre-pandemic (8% in 2017) or during the first national lockdown (11%).[16] Our findings of increasing negative mental health impacts for children are consistent with the emerging reviews of the published evidence.[9]

Strengths of our study included the large cross-sectional and nationally representative surveys, which employed strong methodology (surveys piloted and included the validated K6); collected data on caregiver and child mental health; surveyed female and male caregivers; and achieved good response rates. In other Polls, indicators (frequency/prevalence) across a range of topics are almost universally consistent with more traditionally obtained estimates, providing support for the sample selection and survey administration methods. By July 2021, we found that negative mental health impacts were reported for half of children and two-thirds of adults. This highlights how general pandemic stress, and disruption to family life including employment, routines, and social interactions, can adversely affect caregivers and children. Despite Australia's low incidence of COVID-19, the increases in psychological distress are consistent with data from countries with high infection rates and the increases in distress are sizable.[4 6 8] This pattern has been described for previous health crises; following the H1N1 pandemic, a quarter of parents and 30% of children who were required to isolate met criteria for PTSD symptoms.[17]

This study also had limitations. The reliance on caregiver-report, from only one caregiver per household, means the child rating may be biased by caregiver perception. We did not collect a validated measure of children's mental health that would provide a measure of clinical

impact. The finding of different social gradients for the K6 compared with the negative impact items may represent differing expectations and emphasises the need for validated measures for anticipating the longer-term consequences of lockdown, which ideally would be collected from all caregivers and children directly. The RCH Poll only samples adults who are 18 years and older, so younger caregivers are missing, and a significant proportion of caregivers did not report family income. However, the weighted and adjusted analyses support the generalisability of findings to the broader population of Australian families raising children and young people.

Australia's reliance on lockdown has meant that caregivers had to concurrently parent, work, and facilitate remote schooling. Our findings show that, despite the low disease incidence, the pandemic experience has undermined caregivers' and children's mental health. Understanding specific risk factors for poorer mental health over time was beyond the scope of the cross-sectional data. However, our findings support the inequities evident in existing research. As researchers have suggested,[4 6 8] understanding and prioritising population risk groups should be the subject of ongoing research. Clearly, pandemic response and recovery planning must prioritise and incorporate mental health supports. To this end, we endorse the recommendations proposed by Aknin and colleagues.[6] These include prioritizing safe access to childcare and schools; greater investment in evidence-informed mental health services including telehealth, and its integration with social care systems; and mental health promotion across communities such as work and schools. They also emphasise the importance of monitoring of mental health. As our findings demonstrate, mental health is changing during the pandemic. Understanding families' experiences is necessary to inform policy effort with a greater level of precision, so that we can respond to the evolving mental health needs of children and their families.

FIGURE LEGEND

Figure 1. Estimated probabilities over time (by survey) and by state/territory (as a proxy for cumulative lockdown) adjusted for potential confounders for (a) caregiver K6, and negative mental health impacts for the (b) caregiver and (c) children.

ACKNOWLEDGMENTS AND FUNDING

We thank all families who took part in the Royal Children's Hospital National Child Health Polls. The National Child Health Polls are funded by The Royal Children's Hospital Foundation. This work was supported by The Healthier Wealthier Families project, which was supported by the Helen Macpherson Smith Trust Impact Grant #9523 and The Corella Fund. The Murdoch Children's Research Institute (MCRI) administered research grants for the work and provided infrastructural support (as study sponsor) to its staff but played no role in the conduct or analysis of the research. Research at the MCRI is supported by the Victorian Government's Operational Infrastructure Support Program. Author AP was supported by The Erdi Foundation Child Health Equity (COVID-19) Scholarship. Author SG was supported by a NHMRC Practitioner Fellowship (#1155290).

COMPETING INTERESTS

None to declare.

AUTHOR CONTRIBUTIONS

Anna Price: Conceptualization, Validation, Formal analysis, Investigation, Resources, Data curation, Writing – Original Draft, Writing - Review & Editing, Visualization, Supervision, Project administration, Funding acquisition.

Mary-Anne Measey: Conceptualization, Methodology, Validation, Investigation, Data curation, Writing - Review & Editing.

Monsurul Hoq: Methodology, Validation, Formal analysis, Investigation, Data curation, Writing - Review & Editing.

Anthea Rhodes: Conceptualization, Methodology, Investigation, Resources, Writing - Review & Editing, Supervision, Project administration, Funding acquisition.

Sharon Goldfeld: Conceptualization, Methodology, Investigation, Resources, Writing - Review & Editing, Supervision, Funding acquisition.

AR and SG share senior authorship.

DATA SHARING STATEMENT

The data underlying the results presented in the study are available from the RCH Child Health Poll, please contact child.healthpoll@rch.org.au.

REFERENCES

1. Hale T, Angrist, Noam. , Goldszmidt R, Kira B, Petherick A, Phillips T, et al. A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). Nature Human Behaviour. https://doi.org/10.1038/s41562-021-01079-8. Retrieved from: https://ourworldindata.org/grapher/covid-stringency-index [Online Resource] (2021). 2021.

2. Australian Department of Health. Coronavirus (COVID-19) at a glance - 31 July 2021. https://www.health.gov.au/sites/default/files/documents/2021/08/coronavirus-covid-19-at-a-glance-31-july-2021_0.pdf.

3. Mathieu E, Ritchie H, Ortiz-Ospina E, et al. A global database of COVID-19 vaccinations. *Nature human behaviour* 2021:1-7.

4. Aknin L, De Neve J-E, Dunn E, et al. Mental health during the first year of the COVID-19 pandemic: A review and recommendations for moving forward. *Perspectives on Psychological Science* 2021

5. Centre for Mental Health. Addressing inequities in mental health research exacerbated by Covid-19. A report by the Mental Health Research Group. 6 July 2021 [https://www.centreformentalhealth.org.uk/publications/fit-for-purpose, last accessed 24 Nov 2021].

6. Samji H, Wu J, Ladak A, et al. Review: Mental health impacts of the COVID-19 pandemic on children and youth – a systematic review. *Child and Adolescent Mental Health*;n/a(n/a) doi: https://doi.org/10.1111/camh.12501

7. Jones EAK, Mitra AK, Bhuiyan AR. Impact of COVID-19 on Mental Health in Adolescents: A Systematic Review. *Int J Environ Res Public Health* 2021;18(5) doi: 10.3390/ijerph18052470 [published Online First: 2021/04/04]

8. Racine N, McArthur BA, Cooke JE, et al. Global Prevalence of Depressive and Anxiety Symptoms in Children and Adolescents During COVID-19: A Meta-analysis. *JAMA Pediatrics* 2021;175(11):1142-50. doi: 10.1001/jamapediatrics.2021.2482

9. Meherali S, Punjani N, Louie-Poon S, et al. Mental Health of Children and Adolescents Amidst COVID-19 and Past Pandemics: A Rapid Systematic Review. *International Journal of Environmental Research and Public Health* 2021;18(7):3432.

10. Nikolaidis A, Paksarian D, Alexander L, et al. The Coronavirus Health and Impact Survey (CRISIS) reveals reproducible correlates of pandemic-related mood states across the Atlantic. *Sci Rep* 2021; 11(1). (accessed 2021/04//).

11. Davidson P, Saunders P, Bradbury B, et al. Poverty in Australia 2020: Part 1, Overview. ACOSS/UNSW Poverty and Inequality Partnership Report No. 3, Sydney: ACOSS.; 2020.

12. Wilkins R, Lass I. The Household, Income and Labour Dynamics in Australia Survey: Selected Findings from Waves 1 to 16. Melbourne Institute: Applied Economic & Social Research, University of Melbourne., 2018.

13. Furukawa TA, Kessler RC, Slade T, et al. The performance of the K6 and K10 screening scales for psychological distress in the Australian National Survey of Mental Health and Well-Being. *Psychol Med* 2003;33(2):357-62. doi: 10.1017/s0033291702006700 [published Online First: 2003/03/08]

14. Young Minds. Coronavirus: Impact on young people with mental health needs. https://youngminds.org.uk/media/3708/coronavirus-report_march2020.pdf. 2020.

15. Broadway B, Payne A, A, Salamanca N. Coping with COVID-19: Rethinking Australia. Melbourne Institute: Applied Economic & Social Research, the University of Melbourne., 2020.

16. Edwards B, Biddle N, Gray M, et al. Initial impacts of COVID-19 on mental health in Australia. 2020

17. Sprang G, Silman M. Posttraumatic stress disorder in parents and youth after health-related disasters. *Disaster medicine and public health preparedness* 2013;7(1):105-10.

Supplementary data: estimated probabilities over time (by survey) and by state/territory (as a proxy for cumulative lockdown) for the three mental health measures (graphed in Figure 1)

(a) Poor caregiver mental health (Kessler-6)

```
Generalized linear models                      Number of obs   =      5,854
Optimization     : MQL Fisher scoring          Residual df     =      5,840
                   (IRLS EIM)                   Scale parameter =          1
Deviance         =  9280572.574                (1/df) Deviance =   1589.139
Pearson          =  10441040.84                (1/df) Pearson  =   1787.849

Variance function: V(u) = u*(1-u)                       [Bernoulli]
Link function    : g(u) = u                             [Identity]

                                               BIC             =    9229911
-------------------------------------------------------------------------------
               |              Semirobust
  parentk6tot_d | Risk diff.   std. err.      z    P>|z|    [95% conf. interval]
---------------+---------------------------------------------------------------
       state_r |
           NSW |   .0718525   .0258502     2.78   0.005     .021187    .1225179
           Vic |   .0494101    .023712     2.08   0.037    .0029355    .0958847
               |
          poll |
  W2 sept 2020 |   .0450288   .0269215     1.67   0.094   -.0077365    .097794
  W3 july 2021 |   .0080103   .0161058     0.50   0.619   -.0235565   .0395772
               |
   state_r#poll |
NSW#W2 sept 2020 |  -.0623205   .0408931    -1.52   0.128   -.1424695   .0178284
NSW#W3 july 2021 |   .0292478   .0336068     0.87   0.384   -.0366202   .0951158
Vic#W2 sept 2020 |   .0310635    .043261     0.72   0.473   -.0537265   .1158535
Vic#W3 july 2021 |   .0784145   .0331736     2.36   0.018    .0133954   .1434336
               |
           sex |   .0010905   .0125578     0.09   0.931   -.0235223   .0257033
           q2a |   .2219676   .0232243     9.56   0.000    .1764489   .2674863
   education_c3 |   .0207678   .0094677     2.19   0.028    .0022114   .0393241
         Score |  -.0002257   .0000974    -2.32   0.020   -.0004166  -.0000348
          cald |   .0236198   .0130602     1.81   0.071   -.0019777   .0492173
         _cons |   .2722676   .1000927     2.72   0.007    .0760895   .4684457
-------------------------------------------------------------------------------
```

(b) Negative impacts of the pandemic on caregiver mental health

```
Generalized linear models                      Number of obs   =      5,854
Optimization     : MQL Fisher scoring          Residual df     =      5,840
                   (IRLS EIM)                   Scale parameter =          1
Deviance         =  13913646.93                (1/df) Deviance =   2382.474
Pearson          =  10441040.98                (1/df) Pearson  =   1787.849

Variance function: V(u) = u*(1-u)                       [Bernoulli]
Link function    : g(u) = u                             [Identity]

                                               BIC             =    1.39e+07
-------------------------------------------------------------------------------
               |              Semirobust
      w16q52_d | Risk diff.   std. err.      z    P>|z|    [95% conf. interval]
---------------+---------------------------------------------------------------
       state_r |
           NSW |   .0600002   .0401823     1.49   0.135   -.0187556   .1387561
           Vic |   .0568247   .0407174     1.40   0.163   -.0229799   .1366293
               |
          poll |
  W2 sept 2020 |  -.0051744    .043287    -0.12   0.905   -.0900155   .0796666
  W3 july 2021 |   .1272643   .0357338     3.56   0.000    .0572273   .1973012
               |
   state_r#poll |
NSW#W2 sept 2020 |  -.0053991   .0603809    -0.09   0.929   -.1237435   .1129454
NSW#W3 july 2021 |  -.0234892   .0514006    -0.46   0.648   -.1242325   .0772541
Vic#W2 sept 2020 |   .0756745   .0606002     1.25   0.212   -.0430996   .1944487
Vic#W3 july 2021 |   .0442841   .0514682     0.86   0.390   -.0565917   .1451599
               |
           sex |  -.1035235   .0187399    -5.52   0.000   -.1402531   -.066794
           q2a |  -.1378773   .0240192    -5.74   0.000    -.184954  -.0908005
   education_c3 |  -.0199496   .0130221    -1.53   0.126   -.0454725   .0055733
         Score |   .0002641   .0001445     1.83   0.068    -.000019   .0005472
          cald |   -.059999   .0193952    -3.09   0.002   -.0980129  -.0219852
         _cons |   .2869052    .15113     1.90   0.058   -.0093041   .5831144
```

(c) Negative impacts of the pandemic on child mental health

```
Generalized linear models                    Number of obs    =      10,109
Optimization     : MQL Fisher scoring        Residual df      =      10,093
                   (IRLS EIM)                Scale parameter  =           1
Deviance         =  13214524.01              (1/df) Deviance  =    1309.276
Pearson          =  10717625.13              (1/df) Pearson   =    1061.887

Variance function: V(u) = u*(1-u/1)          [Binomial]
Link function    : g(u) = u                  [Identity]

                                             BIC              =    1.31e+07

                          (Std. err. adjusted for 4,877 clusters in responseid)
--------------------------------------------------------------------------------
                   |             Semirobust
         w16q54_d  | Risk diff.   std. err.     z    P>|z|    [95% conf. interval]
-------------------+------------------------------------------------------------
          state_r  |
             NSW   | -.0091262   .0346798    -0.26   0.792   -.0770974    .058845
             Vic   | -.0249992   .0323287    -0.77   0.439   -.0883623   .0383639
                   |
            poll   |
    W2 sept 2020   | -.0644975   .0345009    -1.87   0.062    -.132118   .0031231
    W3 july 2021   |  .0793999   .0316666     2.51   0.012    .0173345   .1414652
                   |
    state_r#poll   |
 NSW#W2 sept 2020  |  .1487946   .0523181     2.84   0.004    .0462531   .2513362
 NSW#W3 july 2021  |  .1413132    .046868     3.02   0.003    .0494535   .2331728
 Vic#W2 sept 2020  |  .2357508   .0501326     4.70   0.000    .1374927   .3340089
 Vic#W3 july 2021  |  .1847532   .0441666     4.18   0.000    .0981883   .2713181
                   |
            q1c_   | -.0048253   .0134087    -0.36   0.719   -.0311058   .0214552
             sex   | -.0455852   .0167225    -2.73   0.006   -.0783606  -.0128098
     parentk6tot_d |  .0411459   .0249218     1.65   0.099      -.0077   .0899918
             q2a   | -.0709401   .0208751    -3.40   0.001   -.1118546  -.0300255
     education_c3  | -.0062353    .011399    -0.55   0.584    -.028577   .0161063
           Score   |   .000286   .0001372     2.08   0.037    .0000171   .0005549
            cald   | -.0375271   .0178451    -2.10   0.035   -.0725028  -.0025514
           _cons   |  .0359978   .1433662     0.25   0.802   -.2449948   .3169904
--------------------------------------------------------------------------------
```