

1 **COVID-19 vaccine hesitancy among Algerian medical students: a cross-sectional study**  
2 **in five universities.**

3 Mohamed Amine KERDOUN <sup>a,b</sup>, Abdellah Hamza HENNI <sup>c</sup>, Assia YAMOUN <sup>d</sup>,  
4 Amine RAHMANI <sup>b</sup>, Rym Messaouda KERDOUN <sup>e</sup>, Nazia ELOUAR <sup>f</sup>.

5 a: Department of Medicine, Faculty of medical sciences, Kasdi Merbah University,  
6 Ouargla, 30000, Algeria.

7 b: Mohamed Boudiaf Public Hospital, Ouargla, 30000, Algeria.

8 c: Laboratory of dynamic interactions and reactivity of systems, Kasdi Merbah  
9 University, Ouargla, 30000, Algeria.

10 d: Department of Medicine, Faculty of medical sciences, Abdelhamid Mira  
11 University, Bejaia, 06000, Algeria.

12 e: Laboratory of biology and environment, Constantine 1 University, Constantine,  
13 25000, Algeria.

14 f: EPSP Bechir Mentouri, Constantine, 25000, Algeria.

15

16

17

18

19

20

21




22

23 **Graphical abstract**

**COVID-19 vaccine hesitancy among Algerian medical students: a cross-sectional study in five universities**

? Vaccine hesitancy is a limiting factor in global efforts to contain the current pandemic, wreaking havoc on public health. As today's students are the doctors of tomorrow, it is critical to understand their attitudes toward the COVID-19 vaccine.

Our work is the first national Algerian study that examines medical students' attitudes toward the SARS-CoV-2 vaccine.



★ Conclusion

Vaccine acceptance's key barriers were concerns regarding information, safety, efficacy, and potential adverse effects, consistent with studies in other countries among different population groups. The influence of the Vaccine Acceptant students should not be neglected with a possible role as ambassador to hesitant and resistant students

24  
25  
26  
27  
28  
29

30 **Abstract**

31 Vaccine hesitancy is a limiting factor in global efforts to contain the current  
32 pandemic, wreaking havoc on public health. As today's students are tomorrow's  
33 doctors, it is critical to understand their attitudes toward the COVID-19 vaccine. To  
34 our knowledge, this study was the first national one to look into the attitudes of  
35 Algerian medical students toward the SARS-CoV-2 vaccine using an electronic  
36 convenience survey.

37 383 medical students from five Algerian universities were included, with a mean age  
38 of 21.02. 85.37% (n=327) of respondents had not taken the COVID-19 vaccine yet  
39 and were divided into three groups; the vaccine acceptance group (n=175, 53.51%),  
40 the vaccine-hesitant group (n=75, 22.93%), and the vaccine refusal group (n=77,  
41 23.54%).

42 Gender, age, education level, university, and previous experience with COVID-19  
43 were not significant predictors for vaccine acceptance. The confirmed barriers to the  
44 COVID-19 vaccine concern available information, effectiveness, safety, and adverse  
45 effects.

46 This work highlights the need for an educational strategy about the safety and  
47 effectiveness of the COVID-19 vaccine. Medical students should be educated about  
48 the benefits of vaccination for themselves and their families and friends.

49 The Vaccine acceptant students' influence should not be neglected with a possible  
50 ambassador role to hesitant and resistant students.

51 **Keywords:** COVID-19, medical students, vaccine hesitancy, Algeria, public health.

52

53

54

55

## 56 **1. Introduction**

57 The COVID-19 crisis is a major health pandemic first detected in November 2019  
58 and is a severe acute respiratory disease caused by SARS-CoV-2 [1]. The infection  
59 spread quickly in Wuhan, then throughout China and other countries, including  
60 Algeria, the largest country of Africa [2, 3]. As of the end of July 2021, 170,189  
61 confirmed cases and 4,219 deaths had been reported in this country.

62 There are no particular antiviral treatments for COVID-19, and only a few of the  
63 already used therapies have demonstrated the ability to decrease mortality in critical  
64 patients (e.g., corticosteroids). Additionally, compliance with social distancing and  
65 prolonged use of face masks is not guaranteed. Thus, vaccination has been critical in  
66 potentially putting an end to the COVID-19 epidemic [4].

67 Quickly, after the SARS-CoV-2 virus discovery, an extraordinary amount of work  
68 has been accomplished. The scientific community has initiated over 300 vaccines  
69 projects. 40 vaccines are now being evaluated in clinical trials; several have already  
70 received approval from the Food and Drug Administration (FDA) [5] and others are  
71 being used in many countries [6, 7].

72 Although vaccination is a major advance in terms of public health and reduction in  
73 mortality, it has been linked from its origins to vaccine hesitancy [8]. Vaccine  
74 hesitation refers to behaviors of refusal of certain vaccines or their delivery with  
75 delay in a context where vaccination services are functioning [9].

76 Thus, despite its availability, the public acceptability of the COVID-19 vaccines  
77 created in a short time remains uncertain [10]. This hesitation is found worldwide.  
78 Concerns about the vaccine's efficacy or safety, the country of manufacture,  
79 antivaccine movements, and the belief in rushed vaccine research were vaccination  
80 hesitation causes, in addition to rumors and misinformation [11-13].

81 In Algeria, the government has approved the Sinopharm and Sinovac Chinese  
82 vaccines and the Sputnik Russian vaccine. The national vaccination campaign  
83 started by the end of January 2021 for healthcare workers and vulnerable groups.  
84 The current local policy targets all the population of the country.

85 A recent study published in July 2021 described the profile of COVID-19 vaccine  
86 acceptability among the Algerian population (49% of participants were healthcare  
87 workers) and found that 33.5% agreed to get the vaccination. They determined that  
88 two-thirds of Algerians are unlikely to be engaged in COVID-19 vaccine uptake,  
89 making them one of the most resistant populations to voluntary vaccination in  
90 Arabic countries. [14]. However, the vaccine campaign's success depends on the  
91 vaccination rate [15].

92 Today's medical students are tomorrow's doctors, and they are the future providers  
93 of health care and essential influencers of individuals and communities.  
94 Additionally, they will be responsible for making vaccination recommendations and  
95 counseling vaccine-positive individuals.

96 At the heart of this COVID-19 pandemic, we need to know and better understand  
97 the motivations and hesitations of medical students in the face of vaccination.  
98 Documenting the perception and attitude of students towards a vaccination,  
99 therefore, appears essential.

100 The current study targeted medical students from five Algerian universities to  
101 explore the level of COVID-19 vaccine hesitancy and determine the factors and  
102 barriers that may affect their vaccination decision-making.

## 103 **2. Materials and methods**

### 104 **2.1. Study Design and Settings**

105 An observational cross-sectional study was conducted among medical students of  
106 five universities in Algeria (Ouargla, Algiers, Annaba, Setif, and Tizi Ouzou) from  
107 23 to 31 July 2021 with a convenience sample and then closed when not receiving  
108 any new responses for 24 h.

109 The questionnaires were electronically distributed via social networks. Volunteer  
110 students filled in an anonymous online questionnaire. The inclusion criteria were  
111 medical students aged 18 years or older and willing to participate in the study.

112

## 113 **2.2. Questionnaire**

114 A literature review informed the development of the questionnaire. The  
115 questionnaire addressed the following data: personal characteristics, gender, age,  
116 residence, college, academic year, and COVID-19 infection. Lastly, vaccination or  
117 not, level of acceptance or hesitancy about the COVID-19 vaccine for respondents  
118 who do not vaccinate yet, beliefs regarding COVID-19 vaccination like side effects,  
119 efficacy, and information about the vaccine. The internal consistency was assessed  
120 by calculating the Cronbach's alpha as 0.804. Most of the questions were assigned to  
121 be mandatory answer items to avoid incompleteness and missing data.

## 122 **2.3. Statistical analysis of data**

123 Data were extracted from the form to an Excel sheet and statistically analyzed using  
124 IBM SPSS software version 23. Descriptive statistics, one-way ANOVA, t-test, and  
125 chi-square test were used for the respondents. A significance level of 0.05 was  
126 adopted for all statistical analyzes performed.

## 127 **3. Results**

128 A total of 383 medical students [80.67% (n = 302) female and 19.32% (n = 74)  
129 male] from five Algerian universities were included, with a mean age of 21.02  
130 (SD=2.15). The students were divided between 67.62% (n = 259) junior and 32.37%  
131 (n=124) senior students.

132 Table 1 summarizes the students' characteristics and the factors influencing their  
133 uptake of the COVID-19 vaccine.

**Table 1. Characteristics of study participants and comparison between vaccine uptake group and non-vaccine group (n=383).**

<b>Factors</b>	<b>Overall respondents (n=383) (%)</b>	<b>COVID 19 Vaccine group (n = 56)</b>	<b>COVID 19 not Vaccine yet group (n =327)</b>	<b><i>p</i></b>
Age, years mean (SD)	21.02 (2.15)	21.25 (1.78)	21.00 (2.21)	0.771
Gender				
Male	74 (19.32)	16 (21.62)	58 (17.77)	0.082
Female	309 (80.67)	40 (71.42)	269 (82.26)	
Study level				
Junior Students	259 (67.62)	32 (57.14)	227 (69.41)	0.097
Senior Students	124 (32.37)	24 (44.64)	100 (30.58)	
University				
Kasdi Merbah - Ouargla	105 (27.41)	22 (20.95)	83 (79.04)	0.062
Badji Mokhtar - Annaba	86 (22.45)	8 (9.30)	72 (90.69)	
Ferhat Abbas – Setif	72 (18.79)	12 (16.66)	60 (84.34)	
Benyoucef Benkhedda – Algiers	65 (16.97)	11 (16.92)	54 (83.08)	
Mouloud Mammeri - Tizi Ouzou	55 (14.36)	3 (5.45)	52 (94.54)	
COVID 19 Positivity				
Close entourage	224 (58.48)	35 (15.65)	189 (84.37)	0.804
Distant entourage	98 (25.58)	13 (13.26)	85 (86.73)	
No one	61 (15.92)	8 (13.11)	53 (86.88)	
Available information for the new vaccine				
Sufficient information	90 (23.49)	28 (50.00)	62 (18.96)	<0.001
Neutral opinion	71 (18.54)	11 (19.64)	60 (18.34)	

Insufficient information	222 (57.96)	17 (30.03)	205 (62.69)	
<b>Side effect of the vaccine</b>				
The vaccine is safe without side effects	53 (13.83)	17 (30.35)	36 (11.01)	<0.001
Neutral opinion	152 (39.95)	24 (42.86)	128 (39.14)	
The vaccine present multiple side effects	178 (46.47)	15 (26.78)	163 (49.84)	
<b>Efficacy of the vaccine</b>				
The vaccine is effective	128 (33.42)	34 (60.71)	94 (28.74)	<0.001
Neutral opinion	153 (39.94)	16 (28.57)	137 (41.49)	
The vaccine is not effective	102 (26.63)	6 (8.92)	96 (29.36)	

136


137

138

139

140 According to Figure 1, 14.62 % (n=56) of respondents have received the SARS-  
141 Cov-2 vaccine, while 85.37 % (n=327) have not yet. The non-vaccinated  
142 respondents were divided into three groups; the acceptance group (n= 175, 53.51%),  
143 the hesitant group (n=75, 22.93%), and the refusal group (n=77, 23.54%).

144



145

146 **Figure 1. COVID-19 vaccine uptake and acceptance among the respondents.**

147 Demographic variables and experience with COVID-19 infection were not  
148 predictive of COVID-19 vaccine uptake, and there was no significant difference  
149 concerning age, gender, experience with covid, year of study, or university location.

150 Confidence in the information, efficacy, and safety of vaccines shows that 57.96%  
151 (n=222) of respondents think they have insufficient information, 13.83% (n=53)  
152 think that the vaccine is safe, and 33.42 % (n=128) think that the vaccine is  
153 inefficient.

154 Respondents who have received the vaccine feel more informed about the vaccine  
155 (50.00% vs. 18.96%), think that the vaccine is effective (60.71% vs. 28.74%), and  
156 have no severe side effects (30.35% vs. 11.01%) (p<0.001).

157 Table 2 shows the characteristics of the non-vaccinated students and factors  
158 associated with the COVID-19 vaccine decision.

159

160

161

**Table II. Characteristics of unvaccinated respondents (n=327).**

<b>Factors</b>	<b>COVID 19 vaccine acceptance group (n=175)</b>	<b>COVID 19 vaccine hesitant group (n=75)</b>	<b>COVID 19 vaccine refusal group (n=77)</b>	<b><i>p</i></b>
Age, years mean (SD)	21.06 (4.40)	20.97 (3.40)	20.99 (6.79)	0.871
Gender				
Male	32 (18.29)	12 (16.00)	14 (18.18)	0.902
Female	143 (81.71)	63 (84.00)	63 (81.81)	
Study level			1	
Junior Student	118 (51.98)	53 (23.34)	56 (24.67)	0.671
Senior Student	57 (57.00)	22 (22.00)	21 (21.00)	
University				
Kasdi Merbah - Ouargla	40 (48.19)	20 (24.09)	23 (27.71)	0.144
Badji Mokhtar - Annaba	39 (50.00)	15 (19.23)	24 (30.76)	
Ferhat Abbas – Setif	38 (63.33)	18 (30.00)	4 (6.67)	
Benyoucef Benkhedda - Algiers	30 (55.55)	10 (18.51)	14 (25.92)	
Mouloud Mammeri - Tizi Ouzou	28 (53.84)	12 (23.07)	12 (23.07)	
COVID 19 Positivity				
Close entourage	109 (57.67)	41 (21.69)	39 (20.63)	0.162
Distant entourage	45 (52.94)	17 (20.00)	23 (27.06)	
No one	21 (39.62)	17 (32.07)	15 (28.30)	
Available information for the new vaccine				

Sufficient information	49 (28.00)	4 (5.33)	9 (11.68)	
Neutral opinion	46 (26.28)	8 (10.66)	6 (13.63)	<0.001
Insufficient information	80 (45.71)	63 (84.00)	62 (82.66)	
<b>Side effect of the vaccine</b>				
The vaccine is safe without side effects	34 (19.42)	2 (2.66)	0 (11.01)	
Neutral opinion	87 (49.71)	30 (40.00)	11 (14.28)	<0.001
The vaccine present multiple side effects	54 (32.57)	43 (57.33)	66 (85.71)	
<b>Efficacy of the vaccine</b>				
The vaccine is effective	69 (39.42)	18 (24.00)	7 (9.09)	
Neutral opinion	72 (41.14)	37 (49.33)	28 (36.36)	<0.001
The vaccine is not effective	34 (19.42)	20 (26.66)	42 (54.54)	

162

163

164 Medical students are concerned about vaccine hesitancy; only 53.51% are likely to  
165 be vaccinated. Being a male and senior student was associated with vaccine  
166 acceptance, however, no statistically significant difference has been recorded.

167 No available information, severe side effects, and inefficacy of vaccine were  
168 significantly associated with a higher risk of vaccine hesitancy and refusal ( $p <$   
169 0.001).

## 170 **4. Discussion**

### 171 **4.1. Vaccine uptake**

172 The urgency for vaccination is growing by the day, with increasing numbers of cases and reports  
173 of the virus variants in the United Kingdom, Brazil, and South Africa spreading to other  
174 countries [16]. As of 13 August 2021, 31.1% of the world population has received at least one  
175 dose of a COVID-19 vaccine, and 23.4% is fully vaccinated. 4.66 billion doses have been  
176 administered globally, and 35.85 million are now administered each day. However, only 1.2%  
177 of people in low-income countries have received at least one dose.

178 Vaccine hesitancy is a limiting factor in global efforts to contain the current  
179 pandemic, wreaking havoc on public health and socioeconomic systems [17, 18]. As  
180 today students are tomorrow's doctors; it is critical to understand their attitudes  
181 toward the COVID-19 vaccine and work to increase their acceptance to plan an  
182 effective post-pandemic strategy. To our knowledge, our study is the first national  
183 one to look into the attitudes of Algerian medical students toward the SARS-CoV-2  
184 vaccine.

185 The results of our research revealed that only 14.62% of students got the COVID-19  
186 vaccine. The proportion of the population that must be vaccinated against COVID-  
187 19 to initiate herd immunity is unknown. However, a modeling study determined  
188 that vaccines must have an efficacy rate of at least 80% and coverage of at least 75%  
189 to end this ongoing epidemic [19].

190 The low vaccination rates for students can be transposed to the general Algerian  
191 population and demonstrate the need for additional efforts to achieve immunization.  
192 A recent study conducted in April 2021 to describe the existing profile of potential

193 COVID-19 vaccine acceptance among the Algerian population (healthcare workers  
194 constituted 49% of participants), reported that 33.5% accept to receive the vaccine  
195 [20].

196 To increase vaccination coverage, it is necessary to ensure the availability of SARS-  
197 Cov-2 vaccines and vaccination promotion. Systematic strategies should be  
198 implemented to improve COVID-19 vaccine acceptance and uptake among  
199 university students.

200 According to previous studies, a few combined interventions, such as education and  
201 training sessions, easy access to vaccines, and post-vaccination rewards, could  
202 significantly increase vaccination uptake [21, 22].

#### 203 **4.2.Factors associated with vaccination**

204 This study examined vaccine uptake predictors in great detail. Gender, age,  
205 education level, university, and previous experience with COVID-19 were not  
206 significant predictors. Our findings indicated that women and junior students were  
207 less likely to receive the COVID-19, but these findings were not conclusive ( $p >$   
208 0.05).

209 A global population reports from Kuwait [23], Qatar [24], KSA [25], Jordan [26],  
210 and Algeria [20] show that men were more likely to accept the Covid-19 vaccine  
211 compared to females [26]. According to a systematic review, women were less  
212 likely to be vaccinated during the pandemic [27]. This could be because men engage  
213 in riskier behaviors than women [28].

214 However, past research data showed conflicting results about gender. A global survey including  
215 13,426 individuals in 19 countries with a high COVID-19 burden showed that men were  
216 relatively less likely to have a positive attitude towards vaccination than women [29]. Another  
217 study showed that women in Russia and Germany had higher acceptance rates of the SARS-  
218 CoV-2 vaccines than men [30]. This phenomenon has been named “*the Covid-19 gender*  
219 *paradox*”[31].

220 Despite their critical role in vaccine promotion and patient counseling, medical  
221 students are also concerned about vaccine hesitancy. Of the 327 respondents who  
222 had not yet received the vaccine, more than half (53.51%) were willing to take it, a  
223 quarter was hesitant (22.93%), and a quarter was resistant (23.54%).

224 The acceptance rate of our work is lower than a survey of American (75%) [32] and  
225 Italian students (86%) [33] conducted in December 2020, comparable to that of  
226 French students (57%) in January 2021 [34], and higher than that of Egyptian  
227 students (35%) in January 2021 [35]. Also, several studies show that Vaccine  
228 hesitancy and refusal among healthcare workers is a veritable concern and mainly  
229 originate from the safety and effectiveness of COVID-19 vaccines [36-38].

230 The respondents who chose not to be vaccinated did not differ significantly in  
231 demographic and social characteristics from the rest. However, senior students have  
232 more intention of getting vaccinated. According to Barelo et al. [33], vaccination  
233 attitudes are influenced by students' knowledge of health issues; students in  
234 advanced years are more likely to be vaccinated because of their responsibility to  
235 contribute to future waves or peaks.

236 The majority of participants in this study expressed concerns about vaccine  
237 information, effectiveness, safety, and adverse effects. Similar concerns have been  
238 raised globally about health workers worldwide [35, 39, 40].

239 With 57.96%, the primary vaccination barriers identified were a lack of information,  
240 and explain why, despite students' recognition of the COVID-19 vaccine's  
241 importance, they continue to express hesitancy and refusal due to a lack of certainty  
242 about vaccination safety and unknown potential adverse effects, in addition to  
243 misinformation gleaned from social media [41, 42].

244 The evidence suggests that it is critical to establish trust in COVID-19 vaccines [43].  
245 Vaccination education during medical school could be improved by using methods  
246 based on practical learning, like case-based learning and clinical placements [34].  
247 The positive influence of social networks could also be improved by medical

248 influencers that support the dissemination of scientific insights, including issues  
249 related to vaccines and their safety.

250 Other significant reasons for vaccine refusal and hesitancy include fear of vaccine  
251 side effects and doubts about vaccine efficacy. Similarly, Lucia et al. [32], stated  
252 that vaccine hesitancy was influenced by concerns about serious side effects and a  
253 lack of trustworthy information. These concerns are understandable, given the rapid  
254 pace of vaccine development, the novelty of the mRNA technology used in some  
255 vaccines, and the public mediatization of vaccine side effects [44, 45]. Many  
256 hesitant people about vaccination are expected to accept it after being reassured and  
257 provided with credible information about the vaccine's safety and effectiveness [45].

258 As future health care workers, medical students should be vaccinated, as vaccination  
259 of healthcare professionals and students is critical for preventing health-related  
260 infections associated with close contact with high-risk patients [35]. They also  
261 should be educated about the benefits of vaccination for themselves and their family  
262 and friends [46].

263 The Vaccine Acceptant students' influence should not be neglected with a possible  
264 ambassador role to hesitant and resistant students. They should be set as an example  
265 by encouraging other students and patients to get vaccinated.

## 266 **5. Limitations**

267 This study has some limitations, including an online sample with no random  
268 selection and weak generalizability. The percentage of women and junior student  
269 were elevated. The study findings are concerned with five universities and thus  
270 cannot be generalized. Also, this work was carried out during the third peak in  
271 Algeria and may influence COVID-19 vaccine acceptance. Additional research  
272 should be based on an equal representation of gender status and include studies  
273 across academic disciplines for comparison purposes.

## 274 **6. Conclusion**

275 This is the first study of vaccine hesitancy among medical students in Algeria. The  
276 COVID-19 vaccine acceptance was 57%, with considerably high hesitancy and  
277 refusal. COVID-19 vaccine acceptance's key barriers were concerns regarding  
278 information, safety, efficacy, and potential adverse effects, consistent with studies in  
279 other countries among different population groups. The influence of the Vaccine  
280 Acceptant students should not be neglected with a possible role as ambassador to  
281 hesitant and resistant students.

## 282 **Ethics Statement**

283 All procedures performed in this study involving human participants were following  
284 the ethical standards of the University's Research Ethics Board and with the 1975  
285 Helsinki Declaration.

286 The institutional review board of Ouargla University has approved this work.

287 All necessary participant consent has been obtained and the appropriate institutional  
288 forms have been archived.

## 289 **Conflict of Interest**

290 The authors declare that the research was conducted without any commercial or  
291 financial relationships that could be construed as a potential conflict of interest.

## 292 **Acknowledgments**

293 The first author would like to thank all the students who participated in this study,  
294 particularly Kasdi Merbah University. "I am very proud to be one of our teachers".

## 295 **References**

- 296 1. Jahangir MA, Muheem A, Rizvi MFJJPP, 10.31531/ d. Coronavirus  
297 (COVID-19): history, current knowledge and pipeline medications. 2020;3080(2).
- 298 2. Kadi N, Khelfaoui MJBotNRC. Population density, a factor in the spread of  
299 COVID-19 in Algeria: statistic study. 2020;44(1):1-7.

- 300 3. Lounis MJEJoM, Technologies E. COVID-19 in Algeria: Chronology and  
301 evaluation of preventive actions. 2020;13(1):em2001.
- 302 4. Dong Y, Shamsuddin A, Campbell H, Theodoratou EJJogh. Current COVID-  
303 19 treatments: Rapid review of the literature. 2021;11.
- 304 5. Kesselheim AS, Darrow JJ, Kulldorff M, Brown BL, Mitra-Majumdar M,  
305 Lee CC, et al. An Overview Of Vaccine Development, Approval, And Regulation,  
306 With Implications For COVID-19: Analysis reviews the Food and Drug  
307 Administration's critical vaccine approval role with implications for COVID-19  
308 vaccines. 2021;40(1):25-32.
- 309 6. Burki TKJTLRM. The Russian vaccine for COVID-19. 2020;8(11):e85-e6.
- 310 7. Dal-Ré R, Caplan AL, Gluud C, Porcher R. Ethical and scientific  
311 considerations regarding the early approval and deployment of a COVID-19  
312 vaccine. American College of Physicians; 2021.
- 313 8. Machingaidze S, Wiysonge CS. Understanding COVID-19 vaccine  
314 hesitancy. Nature Medicine. 2021.
- 315 9. Razai MS, Chaudhry UAR, Doerholt K, Bauld L, Majeed A. Covid-19  
316 vaccination hesitancy. 2021;373:n1138.
- 317 10. Machingaidze S, Wiysonge CSJNM. Understanding COVID-19 vaccine  
318 hesitancy. 2021:1-2.
- 319 11. Dubé E, MacDonald NE. Vaccine acceptance: barriers, perceived risks,  
320 benefits, and irrational beliefs. The Vaccine Book: Elsevier; 2016. p. 507-28.
- 321 12. Godlee F. What should we do about vaccine hesitancy? : British Medical  
322 Journal Publishing Group; 2019.
- 323 13. Schwarzingler M, Watson V, Arwidson P, Alla F, Luchini SJTLPH. COVID-  
324 19 vaccine hesitancy in a representative working-age population in France: a survey  
325 experiment based on vaccine characteristics. 2021;6(4):e210-e21.
- 326 14. Kacimi SEO, Klouche-Djedid SN, Riffi O, Belaoui HA, Yasmine F, Taouza  
327 FA, et al. Determinants of SARS-CoV-2 Vaccine Engagement in Algeria: A  
328 Population-based Study with Systematic Review of Studies from Arab Countries of  
329 the MENA Region. 2021.
- 330 15. Thunstrom L, Ashworth M, Finnoff D, Newbold SJAaS. Hesitancy towards a  
331 COVID-19 vaccine and prospects for herd immunity. 2020.
- 332 16. Kelekar AK, Lucia VC, Afonso NM, Mascarenhas AK. COVID-19 vaccine  
333 acceptance and hesitancy among dental and medical students. The Journal of the  
334 American Dental Association. 2021.

- 335 17. Salmon DA, Dudley MZ, Glanz JM, Omer SBJV. Vaccine hesitancy: causes,  
336 consequences, and a call to action. 2015;33:D66-D71.
- 337 18. Lo NC, Hotez PJJp. Public health and economic consequences of vaccine  
338 hesitancy for measles in the United States. 2017;171(9):887-92.
- 339 19. Bartsch SM, O'Shea KJ, Ferguson MC, Bottazzi ME, Wedlock PT, Strych U,  
340 et al. Vaccine efficacy needed for a COVID-19 coronavirus vaccine to prevent or  
341 stop an epidemic as the sole intervention. 2020;59(4):493-503.
- 342 20. Kacimi SEO, Klouche-Djedid SN, Riffi O, Belaoui HA, Yasmine F, Taouza  
343 FA, et al. Determinants of SARS-CoV-2 Vaccine Engagement in Algeria: A  
344 Population-based Study with Systematic Review of Studies from Arab Countries of  
345 the MENA Region. medRxiv. 2021.
- 346 21. Robbins SCC, Ward K, Skinner SRJV. School-based vaccination: a  
347 systematic review of process evaluations. 2011;29(52):9588-99.
- 348 22. Bonanni P, Bonaccorsi G, Lorini C, Santomauro F, Tiscione E, Boccalini S,  
349 et al. Focusing on the implementation of 21st century vaccines for adults.  
350 2018;36(36):5358-65.
- 351 23. AlAwadhi E, Zein D, Mallallah F, Haider NB, Hossain A. Monitoring  
352 COVID-19 Vaccine Acceptance in Kuwait During the Pandemic: Results from a  
353 National Serial Study. Risk Management and Healthcare Policy. 2021;14:1413.
- 354 24. Alabdulla M, Reagu SM, Al-Khal A, Elzain M, Jones RM. COVID-19  
355 vaccine hesitancy and attitudes in Qatar: A national cross-sectional survey of a  
356 migrant-majority population. Influenza and other respiratory viruses.  
357 2021;15(3):361-70.
- 358 25. Alfageeh EI, Alshareef N, Angawi K, Alhazmi F, Chirwa GC. Acceptability  
359 of a COVID-19 vaccine among the Saudi population. Vaccines. 2021;9(3):226.
- 360 26. El-Elimat T, AbuAlSamen MM, Almomani BA, Al-Sawalha NA, Alali FQ.  
361 Acceptance and attitudes toward COVID-19 vaccines: a cross-sectional study from  
362 Jordan. Plos one. 2021;16(4):e0250555.
- 363 27. Wang Q, Yang L, Jin H, Lin L. Vaccination against COVID-19: A  
364 systematic review and meta-analysis of acceptability and its predictors. Preventive  
365 Medicine. 2021;150:106694.
- 366 28. Goldman RD, Yan TD, Seiler M, Cotanda CP, Brown JC, Klein EJ, et al.  
367 Caregiver willingness to vaccinate their children against COVID-19: Cross sectional  
368 survey. 2020;38(48):7668-73.
- 369 29. Lazarus JV, Ratzan SC, Palayew A, Gostin LO, Larson HJ, Rabin K, et al. A  
370 global survey of potential acceptance of a COVID-19 vaccine. Nature medicine.  
371 2021;27(2):225-8.

- 372 30. Lazarus JV, Wyka K, Rauh L, Rabin K, Ratzan S, Gostin LO, et al. Hesitant  
373 or not? The association of age, gender, and education with potential acceptance of a  
374 COVID-19 vaccine: A country-level analysis. *Journal of Health Communication*.  
375 2020;25(10):799-807.
- 376 31. Galasso V, Profeta P, Foucault M, Pons V. COVID-19 Vaccine's Gender  
377 Paradox. *medRxiv*. 2021.
- 378 32. Lucia VC, Kelekar A, Afonso NMJJoPH. COVID-19 vaccine hesitancy  
379 among medical students. 2020.
- 380 33. Barello S, Nania T, Dellafiore F, Graffigna G, Caruso RJEJoe. 'Vaccine  
381 hesitancy' among university students in Italy during the COVID-19 pandemic.  
382 2020;35(8):781-3.
- 383 34. Tavolacci MP, Dechelotte P, Ladner JJV. COVID-19 Vaccine Acceptance,  
384 Hesitancy, and Resistancy among University Students in France. 2021;9(6):654.
- 385 35. Saied SM, Saied EM, Kabbash IA, Abdo SAEFJJomv. Vaccine hesitancy:  
386 Beliefs and barriers associated with COVID-19 vaccination among Egyptian  
387 medical students. 2021;93(7):4280-91.
- 388 36. Dror AA, Eisenbach N, Taiber S, Morozov NG, Mizrahi M, Zigran A, et al.  
389 Vaccine hesitancy: the next challenge in the fight against COVID-19.  
390 2020;35(8):775-9.
- 391 37. Sallam MJV. COVID-19 vaccine hesitancy worldwide: a concise systematic  
392 review of vaccine acceptance rates. 2021;9(2):160.
- 393 38. Kose S, Mandiracioglu A, Sahin S, Kaynar T, Karbus O, Ozbel YJIJoCP.  
394 Vaccine hesitancy of the COVID-19 by health care personnel. 2021;75(5):e13917.
- 395 39. Kwok KO, Li K-K, Wei WI, Tang A, Wong SYS, Lee SSJjons. Influenza  
396 vaccine uptake, COVID-19 vaccination intention and vaccine hesitancy among  
397 nurses: A survey. 2021;114:103854.
- 398 40. Troiano G, Nardi AJPH. Vaccine hesitancy in the era of COVID-19. 2021.
- 399 41. Cinelli M, Quattrocioni W, Galeazzi A, Valensise CM, Brugnoti E,  
400 Schmidt AL, et al. The COVID-19 social media infodemic. 2020;10(1):1-10.
- 401 42. Islam AN, Laato S, Talukder S, Sutinen EJTF, Change S. Misinformation  
402 sharing and social media fatigue during COVID-19: An affordance and cognitive  
403 load perspective. 2020;159:120201.
- 404 43. Schwartz JLJNEJoM. Evaluating and deploying COVID-19 vaccines—the  
405 importance of transparency, scientific integrity, and public trust. 2020;383(18):1703-  
406 5.

407 44. Willis DE, Andersen JA, Bryant-Moore K, Selig JP, Long CR, Felix HC, et  
408 al. COVID-19 vaccine hesitancy: Race/ethnicity, trust, and fear. 2021.

409 45. Rief W, editor Fear of Adverse Effects and COVID-19 Vaccine Hesitancy:  
410 Recommendations of the Treatment Expectation Expert Group. JAMA Health  
411 Forum; 2021: American Medical Association.

412 46. Afonso N, Kavanagh M, Swanberg SJV. Improvement in attitudes toward  
413 influenza vaccination in medical students following an integrated curricular  
414 intervention. 2014;32(4):502-6.

415

416

417

418

419

420

421

