Magnetic Resonance Enterography indices for ileal Crohn's disease assessment:

Comparison between classical and machine-learning-based indices

Itai Guez, BSc[1], Gili Focht, MBA[2], Mary-Louise C. Greer, MBBS[3], Ruth Cytter-Kuint, MD[4], Li-tal Pratt, MD[5],

Denise A. Castro, MD[5], Dan Turner, MD, PhD[2], Anne M. Griffiths, MD[6], Moti Freiman, PhD[1*]

[1]Technion - Israel Institute of Technology, Haifa, Israel

[2]Shaare Zedek Medical Center, Juliet Keidan Institute of Paediatric Gastroenterology and Nutrition, The

Hebrew University of Jerusalem, Jerusalem, Israel

[3]Department of Diagnostic Imaging, The Hospital for Sick Children, Department of Medical Imaging, University

of Toronto, Toronto, Canada

[4]Shaare Zedek Medical Center, Paediatric Radiology Unit, Jerusalem, Israel

[5]Kingston Health Sciences Centre, Queen's University, Department of Diagnostic Imaging, Kingston, Canada

[6]The Hospital for Sick Children, Gastroenterology, Hepatology and Nutrition, Toronto, Canada

Corresponding author: Itai Guez, MSc candidate, Technion - Israel Institute of Technology, Technion City, Haifa

3200003, Israel, +972-77-887-4147, itai.guez@campus.technion.ac.il

[Type here]

16   **Abstract:**

Rationale and Objectives: Magnetic Resonance Enterography (MRE) is a non-invasive examination to assess

Crohn's Disease (CD) activity. Several MRE-based indices were developed for standardized evaluation. The

goal of this study was to compare non-linear machine-learning models to linear models in evaluating terminal

ileum (TI) inflammation severity from MRE data of pediatric CD patients.

Materials and Methods: We developed Random-Forest machine-learning (RF) and baseline multiple linear-

regression (MLR) models to evaluate CD severity at the TI from MRE data with the Magnetic Resonance Index

of Activity (MaRIA), it's simplified version (sMaRIA), and the Pediatric Inflammatory Crohn's MRE Index (PICMI)

variables. We assessed the accuracy of the MRE-based TI CD severity evaluation with ileocolonoscopy-based

Simple Endoscopic Score for CD (SES-CD) as the reference standard using a stratified 2-fold validation

experimental setup, repeated 100 times.

Results: A total of 163 children who had all relevant MRE variables and available TI SES-CD score were included

in this study (51% in remission [TI SEC-CD<3], 25% moderate [3≤ TI SEC-CD≤6], 24% high [TI SEC-CD ≥7]). RF-

PICMI achieved the best R-squared coefficient on validation set compared to other models (Median value over

200 folds: RF-MaRIA 0.35 vs. MLR-MaRIA 0.29, RF-sMaRIA 0.37 vs. MLR-sMaRIA 0.27, and RF-PICMI 0.39 vs.

MLR-PICMI 0.33, p<1e-5). Receiver operation characteristics (ROC) area under curve (AUC) for classification of

mucosal healing [TI SEC-CD<3] was broadly similar in all models.

Conclusion: Machine-learning models with MRE variables as inputs might be more accurate than their linear-

regression counterpart models in evaluating CD activity from MRE data.

Keywords: Crohn's disease, Pediatrics, Diagnostic performance, Magnetic resonance enterography, Machine

learning.

[Type here]

1. Introduction

Treatment goals for Crohn's disease (CD) have shifted in the past two decades from symptom control towards

more objective goals in accordance to a "treat-to-target" philosophy.[1,2] Specifically, healing of inflammatory

lesions defined as endoscopic healing (EH)[3] is currently considered a primary objective treatment goal in

clinical trials.[4] The Simple Endoscopic Score for CD (SES-CD)[5] assessed by ileocolonoscopy is considered the

most established endoscopic score for EH evaluation.[6–8] A threshold value which is selected empirically by

experts or a reduction >50% from a baseline SES-CD score is considered as EH.[5]

During the past decade, Magnetic resonance enterography (MRE) has emerged as an imaging modality for

evaluating bowel healing.[3,9] MRE-based evaluation of healing in CD commonly takes the term "transmural

healing" (TH) as a synonym for complete healing of all the bowel layers (including the mucosa). Nevertheless,

there is still no validated definition of what constitutes TH in CD.[10] Current consensus for selecting therapeutic

targets in IBD (STRIDE-II) is that TH and histological healing are not formal targets but should be assessed as

measures of the remission depth, which means assessing healing from cross-sectional imaging is considered an

adjunct and not a treatment target.[11]

Several MRE-based indices were developed for standardized evaluation of TH from MRE data.[12] The most

common MRE index of CD activity in adults is the Magnetic Resonance Index of Activity (MaRIA) which includes

wall-thickness, ulcers, relative contrast enhancement (RCE) and wall edema.[13] Recently, a simplified version of

the MaRIA (sMaRIA) was introduced.[14] This version does not require RCE and thereby avoids intravenous

gadolinium-based contrast agents and includes fat stranding instead. More recently, the Pediatric

Inflammatory Crohn's MRE Index (PICMI) which also does not require RCE and avoids intravenous gadolinium-

based contrast agents and includes instead wall restricted diffusion, comb sign and mesenteric edema, has

been developed and validated specifically to assess TH in pediatric CD patients.[15]

Previous MRE index development and validation reference standards were primarily based on EH assessment

[Type here]

through correlation with endoscopic scores such as SES-CD[16] and Crohn's Disease Endoscopic Index of Severity

(CDEIS)[17] as these indexes are considered reflective of the current medical gold standard while the TH

definition has no consensus.[10]

Recently non-linear machine-learning models demonstrated their potential in improving inflammatory bowel

disease (IBD) assessment in various clinical applications by leveraging complex non-linear correlations between

the input variables and outcomes.[17–19] Yet, previous MRE-based indices development relied upon linear-

regression models, assuming only a linear correlation between MRE items and the endoscopic score.

Therefore, the goal of this study was to systematically compare diagnostic performances of machine-learning

methods over classical methods for the development of MRE-based indices for ileal Crohn's disease

assessment.

The presented models are available to the community through a dedicated website at: https://tcml-

bme.github.io/ML_SESCD.html. Parts of this work were presented at the RSNA 2021 and the WSPGHAN 2021

conferences. A preliminary version of this work is available as a pre-print at

https://doi.org/10.1101/2021.08.29.21262424

[Type here]

2. Materials and Methods

2.1 Ethical Approval

The imageKids study has been approved by the Helsinki committee of the Israeli Ministry of Health. Since all identifying elements related to patient privacy have been removed from the database used in the research presented in this manuscript, it does not need to provide an additional approval of the ethics committee.

2.2 Study population

This is a substudy of the multicenter ImageKids study (NCT01881490[20]) in which a total of 240 children who fulfilled the eligibility criteria of being 6 to 18 years old and having an established diagnosis of CD enrolled to the study. Enrolled participants underwent an ileocolonoscopy and MRE together with biosampling and comprehensive collection of demographics, clinical and laboratory data within 14 days without any change in treatment. Detailed description of the ImageKids study patient population and data collection is provided by Focht et al.[15]

ImageKids study patients' data records with incomplete colonoscopies, prior bowel surgeries, non-intubated TI during ileocolonoscopy, missing MRE information were excluded from our substudy.

2.3 MRE protocol and radiological evaluation

The ImageKids MRE protocol included a localizer sequence and a motility sequence in the coronal plane, followed by a series of coronal and/or axial rapid T2-weighted, T1-weighted gradient echo sequences and diffusion-weighted imaging (DWI). T1-weighted sequences were performed pre- and post-intravenous gadolinium-based contrast agent (GBCA) injection. An intravenous antispasmodic agent (glucagon or hyoscine butylbromide) was administered following the motility sequence. The MRE was performed without an enema as it is less feasible in pediatrics.

[Type here]

Each MRE was read twice by independent two independent central radiologists using an explicit case report

form who were chosen from a group of four pediatric radiologists: MLG, DC, RCK and LP, each with a minimum

of 10 years of experience, from four different sites, two in Canada and two in Israel). The radiologists were

blinded to the clinical and endoscopic data. Radiologists reported, among others, the MaRIA,[13] sMARIA,[14] and

the PICMI variables.[15] The TI was defined in the Imagekids study as the first 10cm segment of the ileum

measured from the ileocecal valve. Table supp1 in the supplementary materials summarizes the variables for

each index.

2.4 Endoscopic report and gastroenterologist assessment

The SES-CD evaluates four items (ulcer size, proportion of the surface area that is ulcerated, proportion of the

surface area affected, narrowing) by scoring each item on an ordinal scale from 0 to 3 in five bowel segments

(TI, right colon, transverse colon, left colon, rectum), where a higher score indicates a more severe disease

defined by a higher percentage of mucosal inflammation. The maximum score in a passable segment is 11

while a score of 12 can be obtained only in the last evaluated segment when there is impassable stenosis. The

total SES-CD score, which is the sum of the five segments scores, ranges from 0 to 56. No adjustment is made

to account for non-visualized segments (due to resection, technical difficulty, or stenosis).[5] The SES-CD items

were scored by a site gastroenterologist during ileocolonoscopy for each bowel segment. In addition,

Gastroenterologist global assessment (GGA) of the degree of inflammation was scored on a 100-mm visual

analogue score (VAS) and on a Likert scale from 1 (i.e. Complete deep remission) to 5 (i.e. Severe disease

activity). The site gastroenterologists were un-blinded to clinical data. For the purpose of this study only the TI

portion of the SES-CD score was used and not the rest of the score.

2.5. Development of Machine-Learning-based MRE indices for CD

We developed three non-linear machine-learning-based MRE indices for CD severity assessment using the

random-forest algorithm,[21] each using a different set of variables. The first index used the MaRIA variables (RF-

[Type here]

MaRIA), the second index used the sMaRIA variables (RF-sMaRIA), and the third index used the PICMI variables

(RF-PICMI). In addition, we constructed three multiple-linear-regression-based indices, each using the same

three sets of variables as input features (MLR-MaRIA, MLR-sMaRIA, MLR-PICMI). The linear regression models

produce a continuous score which is then rounded to nearest integer value, while the random-forest models

produce an ordinal score. The models' scores were normalized to a 0-1 range by dividing it by the maximum

possible SES-CD value of 12.

The TI SES-CD score was used as the reference for indices' development and validation. Therefore, we used

only patients with an available TI SES-CD score which had all MRE variables of both indices. Table supp2

summarizes the full parameter list used for configuration and optimization of the machine-learning algorithms.

The models were written in Python [version 3.6.4] using the open-source Scipy [version 1.5.4] library and the

open-source Statsmodels [version 0.11.0] library. Models ran on an Intel i3 CPU.

2.5 Statistics

We compared diagnostic performance of the machine-learning-based MRE indices to this of their counterpart

multiple-linear-regression-based indices using the TI SES-CD as the reference-standard. We defined distinct

groups of patients for development (50%) and validation (50%). We mitigated the bias of the selection of

patients to development and validation sets and properly compare the set of variables and the chosen

algorithm by using stratified 2-fold validation approach repeated 100 times, with a total of 200 folds.

We assessed indices accuracy of the different models by means of the R-squared coefficient with the reference

endoscopic TI SES-CD. We determined which set of variables (PICMI, MaRIA, or sMARIA) provides the most

accurate predictions in comparison to the reference endoscopic TI SES-CD measurements by means of the

highest R-squared. We used the non-parametric Wilcoxon non-parametric test with Bonferroni correction to

control the family wise error rate (FWER) with $p<0.01$ to assess the statistical significance of the differences in

the R-squared. Finally, we evaluated the capacity of the different indices to distinguish between patients with

    [Type here]

143 and without MH in comparison with reference MH assessments defined as reference SES-CD<3,[4,22] using the

144 area under the receiver operating characteristic (ROC) curves (AUC).

3. Results

3.1 Study population

Figure 1 summarizes the patients selection process for the study. Of the 240 children included in the original ImageKids study, a total of 163 children were included in this sub-study, while 77 were excluded (5 due to incomplete colonoscopies, 13 due to prior bowel surgeries, 33 due to non-intubated TI and 26 due to missing one or more MRE items [wall restricted diffusion (n=16), RCE (n=22)]).

Of the 163 children included in this sub-study, 51% were in remission [TI SES-CD<3], 25% moderate [$3 \leq$ TI SES-CD≤ 6], 24% high [TI SES-CD ≥ 7]). 107 out of the 163 children (65%) had some degree of inflammation according to radiologists reports while 56 (35%) did not have any signs of inflammation seen in the TI based on MRE (all MRE features were zeroes). only 6 of the 56 without any MRE signs of inflammation had a TI SES-CD>3. Table supp3 summarizes the demographic and variables values for the patients included in this sub-study.

4.2 TI SES-CD prediction from MRE

The stratified 2-fold validation evaluation scheme divided the set of 163 patients into 81/82 for development/validation 200 times. Table 1 summarizes the distribution of the R-squared between the models' CD severity assessment and reference TI SES-CD for the different models on the validation sets. The machine-learning-based models achieved a better R-squared compared to their multiple-linear-regression counterparts for each set of variables. The PICMI-based models achieved a better R-squared over MARIA/sMARIA-based models for both the machine-learning-based models and the multiple-linear-regression models. Fig. 2 depicts the R-squared distribution of the different validation sets. All differences were statistically significant.

[Type here]

Figure 1: Patients selection from the entire cohort. MRE: Magnetic Resonance Enterography

R-squared distribution	Median	Q1	Q3
MLR-MaRIA	0.29	0.22	0.33
MLR-sMaRIA	0.27	0.22	0.33
MLR-PICMI	0.33	0.26	0.38
RF-MaRIA	0.35	0.29	0.4
RF-sMaRIA	0.37	0.3	0.41
RF-PICMI	**0.39**	**0.33**	**0.42**

170   Table 1: Validation set R-squared distribution statistics over the folds per model, RF: random-forest, MLR:

171   multiple-linear-regression. Best result in bold.

[Type here]

Figure 2: R-squared distribution over the validation folds

174 Fig. 3 presents the averaged ROC curves of the different models for the differentiation between patients who

175 achieved MH (TI SES-CD<3) and patients who did not achieve MH (TI SES-CD≥3) according to the reference

176 ileocolonscopy-based TI SES-CD score. The RF-PICMI achieved the best distribution of AUC over the different

177 folds. However, the improvement was not statistically significant.

[Type here]

180   Figure 3: Average ROC curves over the different folds for all models.

4. Discussion

The development of MRE biomarkers, MRE indices and the definition of TH play an important role in

objective assessment of CD severity with MRE.[23] Yet, the reference standard for evaluating MRE indices for

CD severity assessment remains endoscopic scores such as the SES-CD.[24] However, current MRE indices

achieved only a moderate correlation with the reference endoscopic scores. In this study, we hypothesized

that the moderate correlation to the TI SES-CD achieved by MRE indices may be attributed, at least in part,

to the utilization of classical linear regression models to characterize the correlation between the MRE-

based variables and EH which limited their ability to determine the potentially complex, non-linear

correlation.[25]

Our results demonstrated that non-linear machine-learning models were better in characterizing the

correlation between the MRE items and EH compared to their linear counterparts. The machine-learning

models improved the correlation between the CD severity assessment by each MRE-based index (MaRIA,

[Type here]

sMaRIA and PICMI) and the reference TI SES-CD. The combination the PICMI MRE variables and the

machine-learning-based model yielded the highest correlation between the model's CD severity

assessment and the reference TI SES-CD.

The better performance of the machine-learning-based models compared to their multiple- linear-

regression counterpart was expected as the machine-learning-based models have a better capacity to

characterize complex, non-linear correlations between input variables and outcomes compared to linear

models.[25] Specifically, we chose to use the RF model[21] for the machine-learning models development as it

is a well-established model useful for small datasets and structured data. The RF ensemble mechanism

controls overfitting by built-in feature selection and especially suits discrete target as it had a fixed set of

possible values covered by the dataset.

Yet, the improved correlation between the models' predictions and the reference TI SES-CD did not lead to

an improvement in classification between patients who achieved MH and patients who did not achieve

MH by means of the AUC. This may be explained as a result of improved correlation demonstrated mostly

on patients with SES-CD higher than 2 rather than in the entire range of scores due to limited availability of

data. We expect that training the models on larger amount of data will result in improved classification

accuracy as well.

While, to the best of our knowledge, we are the first to propose using non-linear machine-learning models

for the development of MRE-based indices for CD severity assessment, the utilization of machine-learning

models in the domain of IBD is not new. Specifically, Dhaliwal et al. used a machine-learning model to

distinguish pediatric colonic IBD subtypes based on clinical, endoscopic, radiologic, and histologic data[26]

and Waljee et al. demonstrated the added value of a machine-learning model to predict disease course

from demographic data and clinical records compared to a linear-regression model.[27] Additional works

aimed to predict endoscopic scores of CD severity from ilecolonoscopy data.[28,29]

[Type here]

In addition, our results demonstrated an improved agreement between the MRE-based assessment and

reference TI SES-CD when using the PICMI variables over the MaRIA and sMaRIA variables in the model.

Although previous literature questioned the need to develop a specific MRE index of activity for the

pediatric population,[3] the improved agreement achieved by the PICMI based models suggest an added-

value to MRE-indices for CD severity assessment developed specifically to the pediatric population.

Finally, our best-performing model (RF-PICMI) used a DWI sequence to assess wall restricted diffusion

variable rather than a post-gadolinium T1-weighted sequence. This finding suggests that the diagnostic

accuracy of DWI, included in PICMI and not MaRIA and sMaRIA, is not overly estimated as was previously

hypothesized.[14] Further, it provides some evidence that the post-contrast sequence can be omitted on

MRE in children to reduce use of GBCA although it is not yet standard practice.[30]

There are several limitations to our study. First, the machine-learning models in this study were developed

with PICMI and MaRIA variables obtained by expert central readings. Further, the PICMI and MaRIA

variables were obtained from MRE data acquired with an optimized and standardized acquisition protocol.

Models' predictions using PICMI and MARIA variables obtained by radiologists' readings of MRE data that

was not necessarily standardized according to the ImageKids MRE protocol should be interpreted with

caution. Second, the development of machine-learning models is assumed to require more data compared

to the development of linear models due to their complexity, we expect better results when the models

are trained on the entire dataset, however then we cannot define properly held out validation set.

Therefore, the presented models' performances might underestimate the added-value of the machine-

learning models.

A future study to completly reproduce our results would require external validation by means of a fully

independent test dataset.

[Type here]

Finally, CD is a transmural disease rather than a mucosal disease. It is not clear yet whether mucosal or

transmural healing should be the primary treatment goal.[10] While MRE has the ability to detect CD lesions

beyond the mucosa, its ability to assess low severity mucosal lesions is much more limited.[3] Therefore,

combined assessment with MRE and ileocolonscopy is ideal to fully assess the extent of the disease.

Conclusions

Our study demonstrates the important role of the combination of non-linear machine-learning models and

the selection of specific MRE-based variables to improve our ability to assess the severity of CD from MRE

data. Such machine-learning methods have the potential to improve MRE-based IBD assessment in

additional applications.

PICMI-based models had better results than MaRIA/sMaRIA-based models using both the random-forest

and multiple-linear-regression models. Further, our evaluation approach can serve as a research tool for

the clinical community to determine the most predictive features in clinical challenges that are still lacking

consensus about the most appropriate data acquisition, and to guide more efficient data collection in large

prospective studies of machine-learning applications in healthcare. The presented models to CD severity

from MRE-based variables are available to the community through a dedicated website at: https://tcml-

bme.github.io/ML_SESCD.html.

[Type here]

5. References

1.   Peyrin-Biroulet L, Sandborn W, Sands BE, et al. Selecting Therapeutic Targets in Inflammatory Bowel

Disease (STRIDE): Determining Therapeutic Goals for Treat-to-Target. *Am J Gastroenterol.*

2015;110(9):1324-1338. doi:10.1038/ajg.2015.233

2.   Colombel JF, D'Haens G, Lee WJ, Petersson J, Panaccione R. Outcomes and Strategies to Support a

Treat-to-target Approach in Inflammatory Bowel Disease: A Systematic Review. *J Crohn's Colitis.*

2020;14(2):254-266. doi:10.1093/ecco-jcc/jjz131

3.   Panés J, Rimola J. Is the Objective of Treatment for Crohn's Disease Mucosal or Transmural Healing?

Clin Gastroenterol Hepatol. 2018;16(7):1037-1039. doi:10.1016/j.cgh.2018.03.034

4.   Turner D, Griffiths AM, Wilson D, et al. Designing clinical trials in paediatric inflammatory bowel

diseases: a PIBDnet commentary. *Gut.* 2020;69(1):32-41. doi:10.1136/gutjnl-2018-317987

5.   Daperno M, D'Haens G, Van Assche G, et al. Development and validation of a new, simplified

endoscopic activity score for Crohn's disease: The SES-CD. *Gastrointest Endosc.* 2004;60(4):505-512.

doi:10.1016/S0016-5107(04)01878-4

6.   Dubcenco E, Zou G, Stitt L, et al. Effect of standardised scoring conventions on inter-rater reliability in

the endoscopic evaluation of Crohn's disease. *J Crohn's Colitis.* 2016;10(9):1006-1014.

doi:10.1093/ecco-jcc/jjw120

7.   Khanna R, Bouguen G, Feagan BG, et al. A systematic review of measurement of endoscopic disease

activity and mucosal healing in Crohn's disease: Recommendations for clinical trial design. *Inflamm*

Bowel Dis. 2014;20(10):1850-1861. doi:10.1097/MIB.0000000000000131

[Type here]

279   8.   Khanna R, Zou G, D'Haens G, et al. Reliability among central readers in the evaluation of endoscopic

280   findings from patients with Crohn's disease. *Gut*. 2016;65(7):1119-1125. doi:10.1136/gutjnl-2014-

281   308973

282   9.   Walsh AJ, Bryant R V., Travis SPLL. Current best practice for disease activity assessment in IBD. *Nat Rev*

283   *Gastroenterol Hepatol*. 2016;13(10):567-579. doi:10.1038/nrgastro.2016.128

284   10.   Serban ED. Treat-to-target in Crohn's disease: Will transmural healing become a therapeutic endpoint?

285   *World J Clin Cases*. 2018;6(12):501-513. doi:10.12998/WJCC.V6.I12.501

286   11.   Turner D, Ricciuto A, Lewis A, et al. STRIDE-II: An Update on the Selecting Therapeutic Targets in

287   Inflammatory Bowel Disease (STRIDE) Initiative of the International Organization for the Study of IBD

288   (IOIBD): Determining Therapeutic Goals for Treat-to-Target strategies in IBD. *Gastroenterology*.

289   2021;160(5):1570-1583. doi:10.1053/j.gastro.2020.12.031

290   12.   Rozendorn N, Amitai MM, Eliakim RA, Kopylov U, Klang E. A review of magnetic resonance

291   enterography-based indices for quantification of Crohn's disease inflammation. *Therap Adv*

292   *Gastroenterol*. 2018;11(1):1-21. doi:10.1177/1756284818765956

293   13.   Rimola J, Ordás I, Rodriguez S, et al. Magnetic resonance imaging for evaluation of Crohn's disease:

294   Validation of parameters of severity and quantitative index of activity. *Inflamm Bowel Dis*.

295   2011;17(8):1759-1768. doi:10.1002/ibd.21551

296   14.   Ordás I, Rimola J, Alfaro I, et al. Development and Validation of a Simplified Magnetic Resonance Index

297   of Activity for Crohn's Disease. *Gastroenterology*. 2019;157(2):432--439.e1.

298   doi:10.1053/j.gastro.2019.03.051

299   15.   Focht G, Kuint RC, Greer M-LC, et al. Development, validation and evaluation of the Pediatric

300   Inflammatory Crohn's Magnetic Resonance Enterography Index (PICMI) from the ImageKids study.

[Type here]

Gastroenterology. 2022;0(0). doi:10.1053/J.GASTRO.2022.07.048

16.   Zheng X, Li M, Wu Y, et al. Assessment of pediatric Crohn's disease activity: validation of the magnetic

resonance enterography global score (MEGS) against endoscopic activity score (SES-CD). *Abdom Radiol*.

2020;45(11):3653-3661. doi:10.1007/s00261-020-02590-8

17.   Olivera P, Danese S, Jay N, Natoli G, Peyrin-Biroulet L. Big data in IBD: a look into the future. *Nat Rev*

Gastroenterol Hepatol. 2019;16(5):312-321. doi:10.1038/s41575-019-0102-5

18.   Le Berre C, Sandborn WJ, Aridhi S, et al. Application of Artificial Intelligence to Gastroenterology and

Hepatology. *Gastroenterology*. 2020;158(1):76-94.e2. doi:10.1053/j.gastro.2019.08.058

19.   Seyed Tabib NS, Madgwick M, Sudhakar P, Verstockt B, Korcsmaros T, Vermeire S. Big data in IBD: Big

progress for clinical practice. *Gut*. 2020;69(8):1520-1532. doi:10.1136/gutjnl-2019-320065

20.   https://clinicaltrials.gov/ct2/show/study/NCT01881490.

21.   Breiman L. Random Forests. *Mach Learn*. 2001;45:5-32.

22.   Weiss B, Turner D, Griffiths A, et al. Simple Endoscopic Score of Crohn Disease and Magnetic Resonance

Enterography in Children: Report from ImageKids Study. *J Pediatr Gastroenterol Nutr*. 2019;69(4):461-

465. doi:10.1097/MPG.0000000000002404

23.   Bruining DH, Zimmermann EM, Loftus E V., et al. Consensus recommendations for evaluation,

interpretation, and utilization of computed tomography and magnetic resonance enterography in

patients with small bowel Crohn's disease. *Radiology*. 2018;286(3):776-799.

doi:10.1148/radiol.2018171737

24.   Roseira J, Ventosa AR, de Sousa HT, Brito J. The new simplified MARIA score applies beyond clinical

trials: Asuitable clinical practice tool for Crohn's disease that parallels a simpleendoscopic index and

[Type here]

fecal calprotectin. *United Eur Gastroenterol J*. 2020;8(10):1208. doi:10.1177/2050640620943089

25. Shalev-Shwartz S, Ben-David S. *Understanding Machine Learning: From Theory to Algorithms*. Vol

9781107057135. Cambridge University Press; 2013. doi:10.1017/CBO9781107298019

26. Dhaliwal J, Erdman L, Drysdale E, et al. Accurate Classification of Pediatric Colonic Inflammatory Bowel

Disease Subtype Using a Random Forest Machine Learning Classifier. *J Pediatr Gastroenterol Nutr*.

2021;72(2):262-269. doi:10.1097/MPG.0000000000002956

27. Waljee AK, Lipson R, Wiitala WL, et al. Predicting Hospitalization and Outpatient Corticosteroid Use in

Inflammatory Bowel Disease Patients Using Machine Learning. *Inflamm Bowel Dis*. 2018;24(1):45-53.

doi:10.1093/ibd/izx007

28. Klang E, Grinman A, Soffer S, et al. Automated Detection of Crohn's Disease Intestinal Strictures on

Capsule Endoscopy Images Using Deep Neural Networks. *J Crohn's Colitis*. 2021;15(5):749-756.

doi:10.1093/ecco-jcc/jjaa234

29. Klang E, Barash Y, Margalit RY, et al. Deep learning algorithms for automated detection of Crohn's

disease ulcers by video capsule endoscopy. *Gastrointest Endosc*. 2020;91(3):606-613.e2.

doi:10.1016/j.gie.2019.11.012

30. Moy MP, Sauk J, Gee MS. The role of MR enterography in assessing Crohn's disease activity and

treatment response. *Gastroenterol Res Pract*. 2016;2016. doi:10.1155/2016/8168695

[Type here]

343 ## 6. Supplementary material

Variable	PICMI	MaRIA	sMARIA	Range	Definition
Wall-thickness	V	V	V	Positive Number (mm)	Measure the thickest part within the TI segment
Ulcers	V	V	V	No - 0, Yes - 1	Mucosal indentation smaller than the full thickn of the bowel wall
Perienteric fat stranding (assessed by mesenteric T2 hyperintensity)	V	-	V	No - 0, Yes - 1	T2 hyperintense signal of the mesentery adjacer to the same bowel loop assessed
Mural edema	-	V	V	No - 0, Yes - 1	T2 hyperintensity relative to psoas muscle
Relative Contrast Enhancement (RCE)	-	V	-	Unbounded number	Increased signal intensity of the bowel wall (WSI on post-contrast T1-weighted sequences, judged subjectively relative to adjacent normal bowel loops and compared with pre-contrast T1-weighted images. WSI is calculated by the average of three wall enhancement measurements.
Wall restricted diffusion	V	-	-	0 - None/Mild, 1 – Moderate/Marked	A subjective measure of hyperintensity on highe b-value trace image compared to normal bowel
Comb sign	V	-	-	No - 0, Yes - 1	Vascular engorgement related to the bowel loop assessed

345 Table supp1: Variables of the MRE-based indices

Parameter	Value
bootstrap	True
ccp_alpha	0
criterion	Mse
max_depth	2
max_features	Auto
max_leaf_nodes	None
max_samples	None
min_impurity_decrease	0
min_impurity_split	None
min_samples_leaf	1

[Type here]

min_samples_split	2
min_weight_fraction_leaf	0
n_estimators	100
n_jobs	None
oob_score	True
random_state	0
verbose	0
warm_start	False

Table supp2: Random forest algorithm parameters

[Type here]

	Subsample n=163	TI remission(MH) TI SES-CD < 3 n=83	No TI remission TI SES-CD >= 3 n=80	P
Male (%)	88 (54)	44 (53)	44 (55)	-
Disease duration, years	2.2 (0.45-4)	2.2 (1.1-3.95)	2.5 (0-4.8)	0.46
Age at enrollment, years	14.24±2.44	14.4±2.45	14.4±2.42	0.22
L1 (TI and cecum) *	37 (22)	13 (16)	24 (30)	-
L2 (Colonic)*	26(16)	23(27)	3(4)	-
L3 (Ileocolonic)*	96 (59)	43 (52)	53 (66)	-
L4 (Isolated upper GI)*	4(2)	4(5)	0(0)	-
Fecal Calprotectin (FC)	629 (151-1319) available in 132	393(70-806) available in 70	828 (532-1601) available in 62	7.9e-5
C-Reactive Protein (CRP)	4.95 (1.4-20.13) available in 152	4 (1-8.45) available in 75	9.5 (2.9-30.6) available in 77	0.003
Weighted Pediatric Crohn's Disease Activity Index (PCDAI)	22.5(10-40) available in 150	17.5 (0-37) available in 74	26.25 (15-41) available in 76	0.07
PICMI	21.5±10.43	15.56±10.46	24.64±9.02	2.7e-5
MaRIA	22.55±8	17.64±7.61	25.14±6.93	7.3e-6
GGA VAS (0-100mm)	31 (16-59)	19 (5-53)	40 (23-60)	6.1e-5
GGA Likert 1- Complete deep remission**	19(12)	19(23)	0(0)	-

[Type here]

GGA Likert 2 - Clinical remission**	30(18)	16(20)	14(18)	-
GGA Likert 3- Mild disease activity	46(28)	21(25)	25(31)	-
GGA Likert 4 - Moderate disease activity	50(31)	19(23)	31(39)	-
GGA Likert 5 - Severe disease activity	18(11)	8(10)	10(12)	-

370   Table supp3: Basic characteristics of the group with all relevant MRE variables and Ileocolonoscopy. Mean +/-

371   standard deviation, median (interquartile range), and frequency (%) are displayed as appropriate.

372   GGA=gastroenterologist global assessment; VAS=visual analogue scale; TI SES-CD = terminal ileum simple

373   endoscopic score of Crohn's disease; * according to Montreal classification; ** clinically and also

374   endoscopically (as much as is known).

[Type here]