Combined PD-L1 and TIM-3 blockade improves the expansion of fit human CD8+ antigen-specific T cells for adoptive immunotherapy

Shirin Lak[1], Valérie Janelle[1], Anissa Djedid[2], Gabrielle Boudreau[1], Ann Brasey[1,3], Véronique Lisi[2], Cédric Carli[1], Lambert Busque[1,3,4,5], Vincent-Philippe Lavallée[2,6,7], Jean-Sébastien Delisle[1,4,5]

[1] Centre de recherche de l'Hôpital Maisonneuve-Rosemont, Montréal, Qc, Canada

[2] Centre de recherche du CHU Sainte-Justine, Montréal, Qc, Canada

[3] Biomarker Unit, Centre C3i, Montréal, Qc, Canada

[4] Department of Medicine, Université de Montréal, Montréal, Qc, Canada

[5] Hematology-Oncology and Cell therapy Division, Hôpital Maisonneuve-Rosemont, Montréal, Qc, Canada

[6] Department of Pediatrics, Université de Montréal, Montréal, Qc, Canada

[7] Hematology-Oncology Division, CHU Sainte-Justine CHU Sainte-Justine

Correspondence to:

Jean-Sébastien Delisle MD, FRCPC, PhD
Centre de recherche de l'Hôpital Maisonneuve-Rosemont
5415, Boul de L'Assomption
Montréal, Qc H1T 2M4
Canada
js.delisle@umontreal.ca

Word count
    Abstract: 343
    Main text: 4948

Declarations

Ethics approval and consent to participate. Informed consent was obtained from all cell donors as detailed in the Method section.

Consent for publication. All authors have reviewed the final version and agreed with the submission. The corresponding author agrees to pay the publication charges.

Availability of data and material. As detailed in the Methods section, the genomic data have been deposited for public access and the authors agree to share protocols. All materials used are commercially available.

Competing interest. The authors have no relevant conflict of interest to disclose.

Funding. This work was funded by the Leukemia/Lymphoma Society of Canada (grants #622735 and #430053) to JSD.

Authors' contribution. SL and JSD conceived the study. SL, VJ, AB, CC performed experiments. SL, VJ, AB, AD, GB, VL analyzed data. LB, VPL, JSD oversaw data analysis. SL, VJ and JSD wrote the manuscript. All authors reviewed the manuscript.

Acknowledgments. The authors are grateful to the volunteer blood donors and Héma-Québec for the leukocytes reduction chamber procurement and handling. We also acknowledge the valuable contribution of Martine Dupuis (flow cytometry and sorting) and the Genome Québec staff for the single-cell RNA-sequencing. SL and VJ are respectively former Cole Foundation and Fonds de recherche du Québec-Santé (FRQS) studentship awardees, JSD and VPL hold FRQS clinician-scientist career awards and JSD is a member of the ThéCell network and of the Canadian Donation and Transplant Research Program (CDTRP).

List of Abbreviations

ACT	Adoptive Cell Therapy
APC	Antigen Presenting Cell
CCR7	C-C chemokine receptor type 7
CD	Cluster of differentiation
CDR	Complementarity Determining Region
CTLA-4	Cytotoxic T-lymphocyte-associated protein 4
CTLs	Cytotoxic T-lymphocytes
CTV	Cell Trace Violet
CTY	Cell Trace Yellow
DC	Dendritic Cell
DMSO	Dimethylsulfoxide
EBV	Epstein-Barr virus
ELISPOT	Enzyme-linked immune absorbent spot
FBS	Fetal Bovine Serum
FOXP3	Forkhead Box P3
Gal-9	Galectin-9
GM-CSF	Granulocyte-Macrophage Colony-Stimulating Factor
GrzB	GranzymeB
HLA	Human Leukocyte Antigen
IFN	Interferon
IL	Interleukin
KLRG1	Killer cell Lectin-like Receptor G1
LAG-3	Lymphocyte-Activation Gene 3
LRSC	Leukocyte Reduction System Chamber
mAB	Monoclonal Antibody
MHC	Major Histocompatibility Complex
NGS	Next-generation Sequencing
PBMC	Peripheral Blood Mononuclear Cell
PBS	Phosphate-buffered saline
PD-1	Programmed Cell Death-1
PD-L1	Programmed Death Ligand-1
PGE2	Prostaglandin E2
PHA	T-cell mitogen phytohemagglutinin
PMA	Phorbol 12-myristate 13-acetate
RT	Room Temperature
sc-RNA	Single Cell RNA

87	TAA	Tumor-Associated Antigen
88	Tcm	Central Memory T cell
89	TCR	T-cell Receptor
90	Teff	Effector T cell
91	Tem	Effector Memory T cell
92	TIM-3	T-cell Immunoglobulin and Mucin-domain Containing-3
93	TMB	3,3′,5,5′-Tetramethylbenzidine
94	TNF	Tumor Necrosis Factor
95	TSA	Tumor-Specific Antigen
96	WT1	Wilms' tumor suppressor gene1
97		

Abstract

Background. The stimulation and expansion of antigen-specific T cells *ex vivo* enables the targeting of a multitude of cancer antigens. However, clinical scale T-cell expansion from rare precursors requires repeated stimulations *ex vivo* leading to T-cell terminal effector differentiation and exhaustion that adversely impact therapeutic potential. We leveraged immune checkpoint blockade relevant to antigen-specific CD8+ human T cells to improve the expansion and function of T cells targeting clinically relevant antigens.

Methods. A clinically-compliant protocol relying on peptide-pulsed monocyte-derived dendritic cells and cytokines was used to expand antigen-specific CD8$^+$ targeting the oncogenic Epstein-Barr virus (EBV) and the tumor associated antigen (TAA) Wilms Tumor 1 (WT1) protein. The effects of antibody-mediated blockade of immune checkpoints applied to the cultures (T-cell expansion, phenotypes and function) were assessed at various time points. Genomic studies including single cell RNA (scRNA) sequencing and T-cell receptor sequencing were performed on EBV-specific T cells to inform about the impact of immune checkpoint blockade on the clonal distribution and gene expression of the expanded T cells.

Results. Several immune checkpoints were expressed early by *ex vivo* expanded antigen-specific CD8$^+$ T cells, including PD-1 and TIM-3 with co-expression matching evidence of T-cell dysfunction as the cultures progressed. The introduction of anti-PD-L1 (expressed by the dendritic cells) and anti-TIM-3 antibodies in combination (but not individually) to the culture led to markedly improved antigen-specific T-cell expansion based on cell counts, fluorescent multimer staining and functional tests. This was not associated with evidence of T-cell dysfunction when compared to T cells expanded without immune checkpoint blockade. Genomics studies largely confirmed these results, showing that double blockade does not impart specific transcriptional programs or patterns on TCR repertoires. However, our data indicate that combined blockade may nonetheless alter gene expression in a minority of clonotypes and have donor-specific impacts.

Conclusions. The manufacturing of antigen-specific CD8$^+$ T cells can be improved in terms of yield and functionality using blockade of TIM-3 and the PD-L1/PD-1 axis in combination. Overcoming the deleterious effects of multiple antigenic stimulations through PD-L1/TIM-3 blockade is a readily applicable approach for several adoptive-immunotherapy strategies.

Background

The vast majority of potentially actionable cancer antigens are major histocompatibility (MHC)-bound peptides. These include tumor-specific antigens (TSA) resulting from mutated proteins or aberrantly expressed regions of the genome, tumor associated antigens (TAA) that come from abnormally expressed self-proteins and finally, non-self peptides best described in the setting of virus-associated cancers or allogeneic stem cell transplantation[1][2]. While it has been possible to isolate TSA or TAA-specific T-cell receptors (TCR) and manufacture TCR-transgenic T cells *ex vivo* followed by their injection in patients, such adoptive immunotherapy approaches face numerous technical hurdles and are currently available only for a minority of relevant cancer antigens[3]. Several effective strategies rely on the *ex vivo* expansion of native antigen-specific T cells, enabling the targeting of a vast array of antigens. However, the expansion of large numbers of antigen-specific T cells requires repeated antigen exposure (through co-culture with antigen-presenting cells – APC) and stimulatory cytokines, potentially leading to T-cell dysfunction (terminal effector differentiation and exhaustion) and poor performance following adoptive transfer[4-6].

Alternatively, the endogenous cancer-reactive T-cell repertoire can be mobilized through the systemic administration of antibodies that prevent signaling from inhibitory co-signaling receptors present on the surface of exhausted T cells[7][8]. Such "immune-checkpoint" blockade, targeting most commonly Cytotoxic T-lymphocyte-associated protein 4 (CTLA-4) and Programmed death-1 (PD-1) on T cells (or its corresponding primary ligand, Programmed death-ligand 1 - PD-L1), is now the cornerstone of therapeutic regimes against several types of neoplasia including advanced melanoma and lung cancer[9]. As dysfunctional cancer-reactive T cells often express multiple negative co-signaling molecules, a strategy has been to use combined approaches with the caveat that increased response may come with more immune-related toxicities[10]. *Ex vivo* expanded T cells express inhibitory receptors, and PD-L1/PD-L2 silenced antigen-presenting dendritic cells have been shown to improve the expansion and function of antigen-specific T cells *ex vivo*[11], thus providing a solid rationale to leverage immune checkpoint blockade to improve T-cell manufacturing for adoptive immunotherapy.

We show herein that the *ex vivo* expansion of CD8+ T cells specific for an oncogenic virus antigen and a

TAA is enhanced by using the combined blockade of PD-L1 and the immune checkpoint T-cell

immunoglobulin and mucin-containing protein-3 (TIM-3), while single blockade of either receptor failed to

improve T-cell expansion. Increased antigen-specific T-cell expansion under dual immune checkpoint

blockade was not associated with phenotypic and functional evidence of exhaustion or terminal effector

differentiation. This was corroborated with single-cell RNA (sc-RNA) and VD(J) sequencing, which

nonetheless revealed that dual immune checkpoint blockade may impact T-cell gene expression in a donor

and clonotype-dependent fashion. Taken together, our results show that dual PD-L1/TIM-3 blockade during

ex vivo expansion can yield large quantities of fit human antigen-specific T cells for adoptive

immunotherapy.

Methods

Donors and cellular procurement

Peripheral blood mononuclear cells (PBMCs) from HLA-A0201 expressing volunteer donors were isolated using Ficoll-Hypaque gradient (STEMCELL Technologies) from fresh whole blood (collected by venipuncture) or leukoreduction system chambers provided by Héma-Québec (LRSCs) as previously described[12][13]. All donors provided written informed consent, and all the experiments were approved by the Héma-Québec and Hôpital Maisonneuve-Rosemont Research Ethics Committees. Recovered PBMCs were either used immediately for experiments or cryopreserved for future use in 90% fetal bovine serum and 10% dimethyl sulfoxide (DMSO).

Dendritic Cells (DCs) differentiation and antigen pulsing

Monocyte isolation and DC differentiation have been generated as previously described[4]. Briefly, monocytes were obtained using the adherence method whereby PBMCs were plated in adherent plastic plates (Sarstedt) in media (X-Vivo 15 medium (LONZA) supplemented with 5% human serum, 2 mM L-glutamine and 1mM Sodium Pyruvate (Gibco), 1000 U/ml (100 ng/ml) IL-4 and 800 U/ml (50ng/ml) GM-CSF (Both from STEMCELL technologies)) and incubated in a CO_2, 37°C incubator for 7 days. On day 4, media was replaced with fresh media supplemented with IL-4 and GM-CSF. On day 7, DCs were matured with maturation media containing 1000 U/ml (100 ng/ml) IL-4, 800 U/ml (50ng/ml) GM-CSF (Both from STEMCELL technologies), 10 ng/ml TNF-α (STEMCELL technologies), 1 µg/ml PGE2 (SIGMA), 10 ng/ml IL-1β (Feldan), 100 ng/ml IL-6 (Miltenyi Biotec) and loaded with desired peptide (1 µg/mL Latent membrane protein 2 - LMP2$_{426-434}$ (CLGGLLTMV) or 1 µg/mL Wilm's Tumor 1 - WT1$_{37-45}$ (VLDFAPPGA), both from JPT Peptides. Lastly, DC media was supplemented with IFN-γ 1000U/ml (Feldan) for the last 24 hours of maturation.

Ex Vivo expansion of antigen-specific T cells

Antigen-specific T cells were stimulated using a clinically compliant protocol as previously described ([4] and www.clinicaltrials.gov NCT03091933) from 15x10^6 PBMCs and expanded through multiple weekly stimulations using irradiated (40 Gy) autologous, peptide-loaded monocyte-derived DCs at a 1:10

(DC:PBMC) in a G-Rex6® Well Plate vessel (Wilson Wolf Manufacturing, New Brighton, MN). Our complete

T-lymphocyte culture (CTL) media (Advanced RPMI 1640, 10% human serum, 1X L-glutamine (Gibco))

was supplemented with the following cytokines; week 1: IL-21 (30 ng/mL) and IL-12 (10 ng/ mL) (Both from

Feldan), week 2: IL-21, IL-2 (100 IU/mL), IL-7 (10 ng/mL) and IL-15 (5 ng/mL) (STEMCELL Technologies),

subsequent weeks: IL-2, IL-7 and IL-15. Medium, including cytokines, was refreshed every 3 to 4 days, and

antigen re-stimulation was done every week with peptide-loaded monocyte-derived DCs. Cell concentration

was adjusted to 1:10 ratio each week. When indicated, cultures were supplemented with 20 µg/ml of anti-

PD-L1 blocking mAb (BioXcell, 29E.2A3) or/and 10 µg/ml of anti-TIM3 blocking mAb (Biolegend, F38-2E2)

with all media changes. Respective isotype antibodies, mouse IgG1 (BioXcell, MOPC-21) and InVivoMAb

mouse IgG2b (BioXcell, MPC-11), were used at the same concentration where indicated. All cell cultures

were performed in monitored incubators (37°C in 5% CO_2 and 5% air humidity). Cell viability and cell counts

were assessed by the countess automated cell counter (Invitrogen) using trypan blue (Invitrogen) at a 1:1

ratio with the cellular suspension in cell counting chamber slides (Invitrogen-C10283).

Flow cytometry

The phenotype of mature DCs was assessed by cell surface expression of the following markers (antibody

clone in parenthesis) CD80 (L307.4), CD86 (2331, FUN1), HLA-ABC (W6-32), CD11c (3.9), CD19 (HIB19)

(all from BD Biosciences)), HLA-DR (LN3), and CD83 (HB15e) (both from Invitrogen). To detect antigen-

specific CD8+ T cells, up to 10^6 cells were suspended in phosphate-buffered saline (PBS) plus 2% fetal

bovine serum (FBS) and stained with appropriate allophycocyanin (APC) labeled MHC-Dextramers A*0201

(Immudex,), for 10 minutes in the dark at 4°C. For additional cell surface markers and phenotyping, cells

were stained with following antibodies: CD3 (SKY7), CD3 (UCHT1), CD8 (SK1), CD45RO (UCHL1),

CD45RA (5H9), CCR7 (150503), LAG3 (T47.530), and CD4 (RPA-T4) all from BD Biosciences; CD62L

(DREG-56), TIM3 (F38-2E2), PD-1 (EH12.2H7), KLRG1 (2F1/KLRG1), CD57 (HCD57), and PD-L1

(29E.2A3) from Biolegend and CD8 (RPA-T8) and CD244 (eBioDM244) from Invitrogen. Staining was

performed at room temperature (RT) in the dark for 30 minutes. Intracellular staining was done with the

Foxp3 / Transcription Factor Staining Buffer Set as recommended by the manufacturer (eBioscience).

Before the fixation step, up to 10^6 cells were stimulated to produce cytokines with the indicated peptide (0.5

µg/ml) (test condition) or PMA (50 ng/mL)-ionomycin (500 ng/mL) (Sigma-Aldrich) (positive control) and a

pool of irrelevant peptides (from the cytomegalovirus pp65 protein - PepMix™ HCMVA, JPT peptides) for

4 h at 37°C. Cells are suspended in CTL media plus brefeldin A (Biolegend) to block the secretion of

cytokines during the stimulation period. Cells were then harvested and stained for cell-surface markers,

including CD3, CD4, and CD8 in 4 °C for 20 minutes. Next, fixed and permeabilized cells were stained with

intracellular cytokine detection antibodies; IFN-γ (4S.B3), IL-2 (MQ1-17H12), and TNF-α (Mab11) all

purchased from BD Biosciences) and Ki67 (ki-67) from Biolegend at RT for 20 minutes. Cell acquisition

was performed on a Fortessa or LSR II flow cytometer (BD Biosciences), and data were analyzed with

Flowlogic Software (inivai).

IFN-γ Enzyme-linked immune-spot (ELISpot)

227 ELISPOT assays were performed using Human IFN- ELISpot[PLUS] kit (MABTECH), following the

228 manufacturer's instruction. Cultured cells were added ($5{\times}10^4$) to wells in duplicates and then stimulated

229 with anti-CD3 mAb (positive control), irrelevant peptides (specificity control), test peptide or vehicle only

230 (DMSO). Spots were counted using ELISpot reader (vSpot Reader Spectrum, AID).

Flow Cytometry-based Cytotoxicity assay

Flow cytometry-based cytotoxicity assay was performed using either Cell Trace Violet (CTV) or Cell Trace

Yellow (CTY) (Invitrogen Life Technologies) labeled LMP2$_{426-434}$ pulsed as target cells as described

before[14]. Briefly, target cells were prepared by stimulating autologous PBMCs with the T-cell mitogen

phytohemagglutinin (PHA) (3×10^6 PBMCs/mL were incubated in T-cell media with 20 µg/mL PHA) for 3

days at 37 °C and 5% CO_2. Peptide pulsed (or not pulsed) target cells were co-cultured with LMP2-specific

T cells at various ratios for 4 hours in CTL plus 10% horse serum. Unpulsed target cells alone were used

as control. After incubation, cells were harvested, stained with LIVE/DEAD Fixable Aqua Dead Cell Stain

Kit, Life Technologies) according to manufacturers' instructions and viable target, cells were quantified by

flow cytometry using Flow Count Beads (Beckman Coulter). Cytotoxicity was calculated by comparing the

percentage of viable target cells in test conditions relative to control (100 - (Target cell alive/ Target cell

alone) × 100).

Single-cell RNA sequencing and High-throughput TCR sequencing

At indicated time points, T cells were sorted based on MHC-Dextramer and CD8 staining using FACS Aria

III (DB Biosciences). Post sorted purity of the sorted population was 94%. A fraction of the sorted antigen-

specific T cells was subjected to both VDJ and transcriptome sc-RNA sequencing at the Genome Quebec

facilities. Briefly, cells were counted and viability assessed using a hemocytometer and Trypan Blue. The

targeted cell recovery was set at 6000 cells and libraries were prepared using the kits Chromium Next GEM

Single Cell 5' Kit v2 and Chromium Single Cell Human TCR Amplification Kit (10x Genomics inc.), as per

the manufacturer's recommendations. Libraries were quantified using the Kapa Illumina GA with Revised

Primers-SYBR Fast Universal kit (Kapa Biosystems). Average size fragment was determined using a

LabChip GX (PerkinElmer) instrument. The 10X Single cells 5' libraries were sequenced on Illumina HiSeq

4000 PE28x98 while the 10X Single Cells VDJ (Human T) libraries were sequenced on Illumina HiSeq 4000

PE150. The Illumina control software was HCS HD 3.4.0.38, the real-time analysis program was RTA v.

2.7.7. bcl2fastq2 v2.20 was then used to demultiplex samples and generate fastq files. Reads were aligned

to GRCh38 genome assembly, gene expression matrices generated, and clonotype identification

performed using CellRanger v3.0.2. The resulting gene expression matrices were normalised by total UMI

counts per cell multiplied by the median UMI count per cell and natural-log transformed using Scanpy v1.4[15].

PhenoGraph[16] clustering was applied with k=20 and clusters with a median mitochondrial fraction greater

than 0.2 were filtered out. Clonotypes were defined by nucleotide sequences of alpha and beta chains. In

order to remove potential doublets, we restricted our analysis to cells in which exactly one alpha and one

beta chain were identified. Furthermore, we retained only the most abundant clonotype for each alpha and

beta chain. The number of retained cells per sample is provided in Table S1. After filtering, individual count

matrices were combined and normalized and log transformed, as described above, for downstream

analyses. Pathway expression is defined as the log2 of the sum of the normalized gene expression of

individual genes. Genes included with each pathway are listed in Table S2 (taken from[17]). Significance of

the changes in pathway expression between the control (CTRL) and delayed double blockade (DDB)

conditions was assessed using a Mann-Whitney U test. To set the threshold at which p-values are

considered significant, we compared the p-values obtained with the pathway of interest to those obtained

with random sets of genes. For each pathway of interest, we generated 1000 random gene sets with the

271 same number of genes drawn from the same expression distribution. On each of these random gene sets,

272 we performed the CTRL vs DDB comparison and recorded the p-values obtained from the Mann-Whitney

273 U test and report the percentile ranking of the p-value in the pathway compared to the p-values obtained

274 from the random gene sets.

276 Bulk TCR repertoire profiling was performed from sorted multimer positive and negative CD8[+] T-cell RNA

277 preserved in Trizol reagent (Invitrogen) using Next-Generation Sequencing (NGS) targeting the

278 hypervariable complementarity-determining region 3 (CDR3) of the T cell receptor beta chain (TCRβ).

279 TRIzolTM extracted RNA in combination with the PureLink RNA mini and microcolumn system (Thermo

280 Fisher), quantified by UV spectrophotometry (Tecan), and QC was completed on Bioanalyzer chip (Agilent).

281 TCRb amplicon libraries were prepared from 25 ng total RNA with the Oncomine™ TCR Beta-SR Assay

282 for RNA (ThermoFisher Scientific). The TCRb libraries obtained were quantified on the ViiA 7 Real-Time

283 PCR System with the Ion Library Taqman Quantitation kit (Thermo Fisher). NGS was completed on the Ion

284 S5 semiconductor platform using an Ion 540 chip, prepared with the Ion Chef System (all from Thermo

285 Fisher). TCRβ repertoire analysis was completed using the Ion Reporter Software (Thermo Fisher) and

286 Immunarch package[18] analyses using R software. The data for all genomic analyses can be found in the

287 Gene Expression Omnibus (GSE182537 and GSE181682).

288 **Statistical Analysis**

289 Statistical significance was analyzed were with the R software, version 4.0.4. Multiple group comparisons

290 were performed using one-way ANOVA and Turkey post hoc test or the Kruskal-Wallis test with the Holm

291 procedure to correct for multiple testing (if ANOVA requirements were not met). Paired Wilcoxon-Mann-

292 Whitney tests were performed for two-group comparisons. p values of 0.05 or less were considered

293 significant.

Results

Multiple stimulations are detrimental to antigen-specific T-cell expansion *ex vivo*

Antigen-specific T cells were stimulated using moDCs loaded with the peptide LMP2$_{426-434}$ (CLGGLLTMV),
which is an HLA-A0201 restricted antigen from the oncogenic Epstein-Barr virus (EBV). Weekly *ex vivo*
stimulations were performed in cytokine supplemented media, and antigen-specific expansion was
assessed before each stimulation. While overall T-cell expansion progressed following each of four
stimulations (albeit at limited pace after day 14 and two stimulations), the percentage and absolute number
of antigen-specific CD8 T cells, as identified by fluorescent HLA-0201/ LMP2$_{426-434}$ multimer, stagnated
despite additional stimulations (Figure 1A-C). Further, this halted growth was matched with a gradual
change in phenotype. The predominance of central memory (Tcm) differentiation profile at day 14 evolved
towards an effector memory (Tem) and effector (Teff) differentiation phenotype on days 21 and 28 (Figure
1D). These results were anticipated as serial T-cell stimulations have been associated with the development
of effector T cells that gradually lose their capacity to expand and persist after adoptive transfer[6]. The
expression of the inhibitory receptors related to T-cell exhaustion (PD-1, TIM-3, and to a lesser extent LAG3
and 2B4) was substantial on a significant fraction of antigen-specific T cells at day 14, with little modulation
over time in culture (Figure 1E). However, the fraction of cells showing dual expression of PD-1 and TIM-3
increased with repeated stimulation (Figure 1F) in line with the development of CD8+ T-cell exhaustion[19-21]. Phenotyping of the moDCs revealed prevalent expression of corresponding PD-1 and TIM-3 ligands,
(respectively PD-L1 and CEACAM-1/Galectin 9) (Figure 1G). These results confirm and extend previous
data, suggesting that the early and persistent expression of PD-1 and TIM-3 by CD8+ T cells, along with
the expression of their ligands by stimulating DCs, may represent a significant hurdle for the expansion of
antigen-specific T cells for immunotherapy[19 20].

Combination of PD-L1/PD-1 axis and TIM-3 blockade significantly increases antigen-specific CD8+
T-cell expansion

In order to assess if immune checkpoint blockade during *ex vivo* expansion may improve antigen-specific
T-cell yield, anti-PD-L1, anti-TIM-3, or both were added to the culture media at the beginning of culture and
with all media changes. Cell counts at day 14, and even more strikingly at day 21, revealed that the double

blockade condition significantly increased T-cell expansion relative to control (no checkpoint blockade) and

the single blockade groups (Figure 2A). This translated into a marked increase in LMP2$_{426\text{-}434}$-specific T-

cell yield in the dual blockade condition, most evident at day 21 of culture (Figure 2B-C). In contrast, the

single blockade of PD-L1 or TIM-3 offered no advantage at any time point. The phenotypic assessment of

LMP2$_{426\text{-}434}$-specific T cells revealed no statistically significant difference in the percentages of Tcm, Tem,

and Teff and no difference in PD-1 or TIM-3 expression (Figure 2D-E). However, the percentage of T cells

expressing the inhibitory receptors LAG3 and 2B4 was lower in the combined relative to the control

condition. Hence, dual PD-L1 and TIM-3 blockade increased T-cell growth without conferring phenotypic

changes associated with increased T-cell dysfunction.

It is known that TIM-3 has a dual function. TIM-3 is transiently upregulated at intermediate levels on

activated T cells and confers activation signals[22]. However, in settings of chronic stimulation in the presence

of its ligands, TIM-3 inhibits T-cell activation and behaves as a *bona fide* immune checkpoint. We therefore,

slightly modified our protocol to introduce TIM-3 blockade at day 7, a week after the first stimulation

(hereafter designated as delayed double blockade-DDB), with the expectation that it would further improve

antigen-specific T-cell yield. Compared to the dual blockade started on day 0, DDB slightly increased total

T-cell expansion at day 21 and more convincingly following an additional antigenic stimulation (day 28)

(Figure 3A). Moreover, the DDB approach increased the percentage of LMP2$_{426\text{-}434}$-specific T cells at all

time points relative to the control condition (no blocking antibodies used), including at day 14 which was

not the case when PD-L1 and TIM-3 blockade were both applied at day 0 (Figure 3B). Absolute antigen-

specific counts were also increased in the DDB relative to the control condition (significant on days 21 and

28) (Figure 3C). Finally, DDB did not impact T-cell differentiation or exhaustion marker expression (Figure

3D-E). Independent cultures using isotype control antibodies confirmed the specific effects of anti-PD-L1

and anti-TIM-3 antibodies on the antigen-specific T-cell expansion (Supplementary Figure 1). We conclude

that combined blockade of the PD-L1/PD-1 axis and TIM-3 can be incorporated in *ex vivo* cultures to

increase antigen-specific CD8$^+$ T-cell yield for adoptive immunotherapy without altering T-cell phenotypes.

Double immune checkpoint blockade generates functional antigen-specific T cells

Dual PD-L1 and TIM-3 blockade increases antigen-specific CD8+ T-cell expansion in culture without

altering T-cell phenotypes suggesting comparable functionality. Intracellular cytokine secretion

measurements and ELISpot assays at day 28 confirmed that a higher proportion of T cells were reactive

upon LMP2$_{426-434}$ peptide re-exposure in the DDB condition relative to control (no checkpoint inhibition)

(Figure 4A-B). This was also generally the case when DDB was compared to double immune checkpoint

blockade administered at day 0 (statistically significant in ELISPOT data at day 28). We thus pursued our

functional assessments comparing DDB with the control condition. As further indication of increased

functionality, a greater fraction of antigen-specific T cells expanded in the DDB condition relative to control

expressed the proliferation marker Ki-67 and upon peptide re-exposure, more DDB exposed T cells

displayed evidence of cytotoxic potential (surface CD107a and intracellular granzyme B expression) (Figure

4C-D). This was corroborated with cytotoxicity assays showing that T cells from the DDB condition were

highly effective, especially at low effector:target ratios (Figure 4E-F). Importantly, in all our assays,

combined PD-L1 and TIM-3 blockade did not lead to increased non-specific (off-target or spontaneous)

cytokine release or cytotoxicity. Thus, dual immune checkpoint inhibition expands T-cell products with

specific and augmented antigen reactivity.

PD-L1/TIM-3 blockade imparts no consistent gene expression signatures to expanded antigen-

specific CD8+ T cells

To gain more insight into the biological impacts of double immune checkpoint blockade on antigen-reactive

T cells, we submitted sorted day 28 multimer-positive T cells from three donors to paired transcriptome and

TCR alpha-beta single-cell RNA sequencing (scRNA-Seq). Gene expression was compared between

donors, conditions (no checkpoint blockade vs. DDB) and clonotypes. Global gene expression patterns

across the three donors revealed strong donor-specific clustering (Figure 5A). The impact of DDB on gene

expression for each donor antigen-specific T cells, was assessed using published human T-cell gene sets[17].

This enabled a detailed assessment of T-cell activation, proliferation, terminal differentiation, exhaustion

and metabolism-associated gene expression as these processes are likely to be impacted by DDB. Overall,

when all T cells from each condition were compared, no consistent change of gene expression signature

could be identified across all donors (Figure 5B). We next evaluated whether clonotype-specific signatures

could be identified. VD(J) sequencing revealed that the LMP2$_{426-434}$ specific T cells were oligoclonal in all

donors and conditions (1-4 clones representing more than 80% of all cells – Figure 5C and Supplementary

Table 3). Most dominant clonotypes were shared between experimental conditions (albeit in some

instances at different frequencies) and a few clonotypes were shared between donors (e.g. clonotype 1 and

3 common to donors 1 and 2, Supplementary Table 3). We next analyzed the expression of several

pathways in the abundant clonotypes, defined as those identified in at least 10 cells and representing at

least 1% of the repertoire in both the DDB and control conditions from the same donor. Similar to

comparisons involving all clonotypes, no consistent pattern was found across donors when comparing the

DDB versus control condition on a per clonotype basis (Figure 5D). Interestingly, our data unveiled

divergent gene expression patterns for one clonotype (Clonotype 3) which was shared by Donor 1 and 2.

Relative to its counterpart in the control condition, Donor 1 Clonotype 3 in the DDB condition expressed

higher levels of genes related to T-cell exhaustion/differentiation and had a lowered expression of genes

associated with T-cell proliferation. The same clonotype in the DDB condition from Donor 2 had, on the

contrary, an increased expression of the genes associated with T-cell activation without any transcriptional

changes related to T-cell dysfunction (T-cell exhaustion, terminal differentiation, low proliferation). Although

obtained from a limited number of cells and donors, these results suggest that donor rather than clonotype

related features may determine T-cell outcomes following DDB.

To assess whether clonotype-specific transcriptional signatures may impact their expansion and clonal

hierarchy within the cultures, we used bulk mRNA collected from sorted multimer negative and positive

CD8+ T cells at day 21 and 28 from the same donors. RNA of suitable quality was obtained for 23 out of a

potential of 24 samples (the multimer-positive fraction of the control condition from Donor 3 had to be

excluded) and subjected to complementary determining region 3 (CDR3) region sequencing of the T-cell

receptor-beta chain (TCRβ). Day 28 multimer-positive T-cell CDR3 sequencing matched very well paired

VDJ sequencing of single cells (Supplementary Figure 2), confirmed the oligoclonality of the multimer-

positive T cells at both day 21 and 28 and revealed that TCR repertoire diversity in the multimer-negative

T cells was not affected by DDB (Figure 6A). Further, the multimer positive and negative fractions had

limited overlap (Supplementary Figure 3), suggesting that sorting effectively separated most CD8+ antigen-

specific cells from the rest of the T cells in culture. Clonotype hierarchy among multimer-positive T cells at

day 21 and day 28 showed no consistent pattern of evolution in DDB relative to the control condition (Figure

6B-C). However, the proportion of certain clonotypes in these fractions (both DDB and control) varied

importantly (>20%) between day 21 and 28. This was notably the case for Clonotype 3 from Donor 1 in the

DDB condition, which declined markedly from day 21 to day 28, whereas this clonotype's frequency

changed only slightly between day 21 to 28 of DDB exposure in Donor 2 (Figure 6B). We then explored

whether the clonotypes with altered abundance between day 21 and day 28 in the same donor and condition

displayed a specific transcriptional profile at day 28. In the DDB condition, clonotype 3 of Donor 1, which

showed a decreased abundance from day 21 to day 28, had a higher expression of genes related to T-cell

activation, differentiation and exhaustion relative to other clonotypes from the same donor in the same

condition (Figure 6D). Although less striking, we also noticed a weak trend of increased expression of genes

related to exhaustion in clonotypes with decreased abundance at day 28 and a rise in the expression of

genes related to proliferation in clonotypes with an increased abundance at day 28, irrespective of

experimental conditions (Figure 6D). Taken together these data show that DDB has limited impact on the

clonal diversity of the expanded T cells overtime. The proportion of the various clonotypes in time can

nonetheless fluctuate in both culture conditions and that the gene expression signatures at day 28 offers

possible explanations for such fluctuations. Globally, these results suggest that DDB confers no consistent

transcriptional features to expanded antigen-specific CD8$^+$ antigen-specific T cells but may alter

activation/dysfunction and cell cycle-related processes in a clonotypes and donor-dependent fashion.

The benefits of dual PD-L1 and TIM-3 blockade extend to TAA-specific T cells

The EBV-derived LMP2$_{426-434}$ antigen stimulates a memory T-cell repertoire in more than 90% of adults[23].

The *ex vivo* expansion of naïve T cells is generally considered more challenging for several reasons such

as size of the repertoire and amount of stimulation required. We sought to determine the impact of dual PD-

L1/TIM-3 blockade on the priming and expansion of naïve T cells. It was previously shown that CD8+ T

cells specific against Wilm's tumor 1 (WT1) derived peptides, a clinically relevant tumor-associated antigen

(TAA), are phenotypically naïve in healthy individuals[24] but can be expanded from a majority of such

donors[25 26]. Volunteer donor T cells were stimulated with an HLA-0201 restricted WT1 peptide (WT1$_{37-45}$)

using the same stimulation/expansion conditions for LMP2$_{426-434}$ specific T cells. WT1-specific T cells

expanded following a similar pattern as LMP2$_{426-434}$-specific T cells (Figure 7A-C). The expansion,

percentages and number of antigen-specific cells as well as their cytokine secretion were improved in the

DDB condition relative to cultures without immune checkpoint blockade, again without altering T-cell

differentiation patterns or exhaustion marker expression (Figure 7D-H). We conclude that our results

obtained with LMP2$_{426-434}$ extend beyond virus-specific memory T-cell expansion and that double PD-

L1/TIM-3 immune checkpoint blockade can improve the expansion of functional TAA-specific CD8$^+$ T cells

from naïve repertoires.

Discussion

The stimulation and expansion of antigen-specific T cells for adoptive immunotherapy is an attractive strategy to target a wide variety of cancer antigens. However, T-cell expansion is limited by the expression of inhibitory receptors and the development of T-cell dysfunction. We confirmed and extended previous studies revealing that immune checkpoint receptors are expressed on repeatedly activated T cells and that their corresponding ligands can be present on antigen-presenting cells used *ex vivo*. Co-expression of PD-1 and TIM-3 has been used to describe highly exhausted CD8[+] T cells, and combined blockade of these receptors in murine models has previously been shown to improve tumor control relative to single blockade[19 20 27]. We show that antibody-mediated immune checkpoint blockade targeting PD-1 or TIM-3 alone is insufficient to improve CD8[+] T-cell expansion while the combination improves T-cell expansion and antigen-specific reactivity. These data confirm the synergistic potential of immune checkpoint blockade and the particular relevance of TIM-3 and PD-1 as inhibitory receptors in CD8[+] T cells[20]. The reasons for this synergistic effect might be due to the mobilization of different signaling intermediates by the two receptors. While PD-1 relies on the recruitment of SHP phosphatases to mediate its effects (as for several other negative co-stimulatory molecules), TIM-3 uses Bat3 and Fyn for its stimulatory and inhibitory effects, respectively. We reasoned that delaying TIM-3 blockade by one week after the start of the culture even in the presence of TIM-3 ligands on APC would be beneficial, given previous experimental evidence that TIM-3 provides activation signals following a first T-cell stimulation[28 29]. This contrasts with PD-1, which conveys inhibitory signals in both exhausted T cells and early after activation[30], justifying the use of anti-PD-L1 at culture initiation. We realize that several refinements may be required to fully leverage the potential agonistic/antagonistic effects of TIM-3 on T-cell activation and expansion and determine whether anti PD-L1/PD-1 blockade may also be improved by altering the timing of the inhibition. In addition, whether dual PD-L1/TIM-3 blockade improves T-cell expansion and function across multiple types of T-cell manufacturing protocols should be tested, just like further combinations with blockade of additional inhibitory receptors. Nonetheless, our work provides a new strategy for T-cell expansion and offers novel biological insights on the effects of PD-L1/PD-1 axis and TIM-3 blockade in human antigen-specific T-cell clonotypes submitted to multiple antigenic stimulations.

Our data support that sustained dual PD-L1/TIM-3 blockade through several rounds of antigenic

stimulations provides ongoing benefits without exacerbating T-cell dysfunction or curbing expansion as

inferred by previous studies of using inhibitory receptor gene deletion[31]. This was assessed through

phenotyping, functional assays, and gene expression in single cells. Our results indicate that antigen-

specific CD8[+] T cells expanded under combined TIM-3/PD-L1 blockade are functional as well as specific

and that their clonal composition is generally stable in time and relative to their counterpart not exposed to

immune checkpoint blockade. It has been reported that immune checkpoint blockade *in vivo* can shape the

T-cell repertoire[32-34]. Thus, from a perspective of T-cell therapy, it was essential to define whether an

intervention during *ex vivo* T-cell expansion alters the clonal identity of the cellular product. While we did

not observe consistent effects of PD-L1/TIM-3 blockade on T-cell transcriptome or clonotype distribution,

altered transcriptional profiles in a clonotype and donor-dependent manner nonetheless suggests that

immune checkpoint blockade in adoptive T-cell therapy may require personalization.

Further work will be required to understand why some clonotypes may be more susceptible to

differentiation/exhaustion following immune checkpoint blockade at different time-points (previous

activation/proliferation history, differentiation status at the beginning of checkpoint blockade, etc).

Nevertheless, our data suggest that even after 28 days in culture, the antigen-specific T cells are highly

functional, with a third expressing Tcm markers. This indicates that antigen-specific T cells generated in

high numbers after dual immune checkpoint blockade could further expand after transfer and likely respond

to further immune checkpoint blockade administered *in vivo*.

We conclude that dual PD-L1/TIM-3 blockade is a readily applicable strategy to improve the expansion of

functional antigen-specific CD8[+] T cells expansion *ex vivo* for adoptive immunotherapy.

References

1. Janelle V, Rulleau C, Del Testa S, et al. T-Cell Immunotherapies Targeting Histocompatibility and Tumor Antigens in Hematological Malignancies. *Front Immunol* 2020;11:276. doi: 10.3389/fimmu.2020.00276

2. Chrusciel E, Urban-Wojciuk Z, Arcimowicz L, et al. Adoptive Cell Therapy-Harnessing Antigen-Specific T Cells to Target Solid Tumours. *Cancers (Basel)* 2020;12(3) doi: 10.3390/cancers12030683

3. Oppermans N, Kueberuwa G, Hawkins RE, et al. Transgenic T-cell receptor immunotherapy for cancer: building on clinical success. *Ther Adv Vaccines Immunother* 2020;8:2515135520933509. doi: 10.1177/2515135520933509

4. Janelle V, Carli C, Taillefer J, et al. Defining novel parameters for the optimal priming and expansion of minor histocompatibility antigen-specific T cells in culture. *Journal of translational medicine* 2015;13:123. doi: 10.1186/s12967-015-0495-z

5. Gattinoni L, Klebanoff CA, Restifo NP. Paths to stemness: building the ultimate antitumour T cell. *Nat Rev Cancer* 2012;12(10):671-84. doi: nrc3322 [pii]

10.1038/nrc3322 [published Online First: 2012/09/22]

6. Janelle V, Delisle JS. T-Cell Dysfunction as a Limitation of Adoptive Immunotherapy: Current Concepts and Mitigation Strategies. *Cancers (Basel)* 2021;13(4) doi: 10.3390/cancers13040598

7. Wei SC, Duffy CR, Allison JP. Fundamental Mechanisms of Immune Checkpoint Blockade Therapy. *Cancer Discov* 2018;8(9):1069-86. doi: 10.1158/2159-8290.CD-18-0367

8. Chamoto K, Hatae R, Honjo T. Current issues and perspectives in PD-1 blockade cancer immunotherapy. *Int J Clin Oncol* 2020 doi: 10.1007/s10147-019-01588-7

9. Vaddepally RK, Kharel P, Pandey R, et al. Review of Indications of FDA-Approved Immune Checkpoint Inhibitors per NCCN Guidelines with the Level of Evidence. *Cancers (Basel)* 2020;12(3) doi: 10.3390/cancers12030738 [published Online First: 2020/04/05]

10. Schnell A, Bod L, Madi A, et al. The yin and yang of co-inhibitory receptors: toward anti-tumor immunity without autoimmunity. *Cell Res* 2020;30(4):285-99. doi: 10.1038/s41422-020-0277-x

11. Hobo W, Maas F, Adisty N, et al. siRNA silencing of PD-L1 and PD-L2 on dendritic cells augments expansion and function of minor histocompatibility antigen-specific CD8+ T cells. *Blood* 2010;116(22):4501-11. doi: 10.1182/blood-2010-04-278739

12. Boudreau G, Carli C, Lamarche C, et al. Leukoreduction system chambers are a reliable cellular source for the manufacturing of T-cell therapeutics. *Transfusion* 2019;59(4):1300-11. doi: 10.1111/trf.15121

13. Orio J, Carli C, Janelle V, et al. Early exposure to interleukin-21 limits rapidly generated anti-Epstein-Barr virus T-cell line differentiation. *Cytotherapy* 2015;17(4):496-508. doi: 10.1016/j.jcyt.2014.12.009

14. Dahmani A, Janelle V, Carli C, et al. TGFbeta Programs Central Memory Differentiation in Ex Vivo-Stimulated Human T Cells. *Cancer Immunol Res* 2019;7(9):1426-39. doi: 10.1158/2326-6066.CIR-18-0691

15. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol* 2018;19(1):15. doi: 10.1186/s13059-017-1382-0

16. Levine JH, Simonds EF, Bendall SC, et al. Data-Driven Phenotypic Dissection of AML Reveals Progenitor-like Cells that Correlate with Prognosis. *Cell* 2015;162(1):184-97. doi: 10.1016/j.cell.2015.05.047

17. Azizi E, Carr AJ, Plitas G, et al. Single-Cell Map of Diverse Immune Phenotypes in the Breast Tumor Microenvironment. *Cell* 2018;174(5):1293-308 e36. doi: 10.1016/j.cell.2018.05.060

18. Nazarov VI, Pogorelyy MV, Komech EA, et al. tcR: an R package for T cell receptor repertoire
advanced data analysis. *BMC Bioinformatics* 2015;16:175. doi: 10.1186/s12859-015-0613-1
19. Zhou Q, Munger ME, Veenstra RG, et al. Coexpression of Tim-3 and PD-1 identifies a CD8+ T-cell
exhaustion phenotype in mice with disseminated acute myelogenous leukemia. *Blood*
2011;117(17):4501-10. doi: 10.1182/blood-2010-10-310425
20. Sakuishi K, Apetoh L, Sullivan JM, et al. Targeting Tim-3 and PD-1 pathways to reverse T cell
exhaustion and restore anti-tumor immunity. *J Exp Med* 2010;207(10):2187-94. doi:
10.1084/jem.20100643 [published Online First: 2010/09/08]
21. Tan J, Yu Z, Huang J, et al. Increased PD-1+Tim-3+ exhausted T cells in bone marrow may influence
the clinical outcome of patients with AML. *Biomark Res* 2020;8:6. doi: 10.1186/s40364-020-
0185-8
22. Gorman JV, Starbeck-Miller G, Pham NL, et al. Tim-3 directly enhances CD8 T cell responses to acute
Listeria monocytogenes infection. *J Immunol* 2014;192(7):3133-42. doi:
10.4049/jimmunol.1302290
23. Taylor GS, Long HM, Brooks JM, et al. The immunology of Epstein-Barr virus-induced disease. *Annu*
Rev Immunol 2015;33:787-821. doi: 10.1146/annurev-immunol-032414-112326
24. Schmied S, Gostick E, Price DA, et al. Analysis of the functional WT1-specific T-cell repertoire in
healthy donors reveals a discrepancy between CD4(+) and CD8(+) memory formation.
Immunology 2015;145(4):558-69. doi: 10.1111/imm.12472
25. Chapuis AG, Ragnarsson GB, Nguyen HN, et al. Transferred WT1-Reactive CD8+ T Cells Can Mediate
Antileukemic Activity and Persist in Post-Transplant Patients. *Sci Transl Med*
2013;5(174):174ra27. doi: 5/174/174ra27 [pii]

10.1126/scitranslmed.3004916 [published Online First: 2013/03/01]
26. Doubrovina E, Carpenter T, Pankov D, et al. Mapping of novel peptides of WT-1 and presenting HLA
alleles that induce epitope-specific HLA-restricted T cells with cytotoxic activity against WT-1(+)
leukemias. *Blood* 2012;120(8):1633-46. doi: blood-2011-11-394619 [pii]

10.1182/blood-2011-11-394619 [published Online First: 2012/05/25]
27. Sun F, Guo ZS, Gregory AD, et al. Dual but not single PD-1 or TIM-3 blockade enhances oncolytic
virotherapy in refractory lung cancer. *J Immunother Cancer* 2020;8(1) doi: 10.1136/jitc-2019-
000294
28. Lee J, Su EW, Zhu C, et al. Phosphotyrosine-dependent coupling of Tim-3 to T-cell receptor signaling
pathways. *Mol Cell Biol* 2011;31(19):3963-74. doi: 10.1128/MCB.05297-11
29. Das M, Zhu C, Kuchroo VK. Tim-3 and its role in regulating anti-tumor immunity. *Immunol Rev*
2017;276(1):97-111. doi: 10.1111/imr.12520
30. Ahn E, Araki K, Hashimoto M, et al. Role of PD-1 during effector CD8 T cell differentiation. *Proc Natl*
Acad Sci U S A 2018;115(18):4749-54. doi: 10.1073/pnas.1718217115
31. Odorizzi PM, Pauken KE, Paley MA, et al. Genetic absence of PD-1 promotes accumulation of
terminally differentiated exhausted CD8+ T cells. *J Exp Med* 2015;212(7):1125-37. doi:
10.1084/jem.20142237
32. Friese C, Harbst K, Borch TH, et al. CTLA-4 blockade boosts the expansion of tumor-reactive CD8(+)
tumor-infiltrating lymphocytes in ovarian cancer. *Sci Rep* 2020;10(1):3914. doi: 10.1038/s41598-
020-60738-4
33. Rudqvist NP, Pilones KA, Lhuillier C, et al. Radiotherapy and CTLA-4 Blockade Shape the TCR
Repertoire of Tumor-Infiltrating T Cells. *Cancer Immunol Res* 2018;6(2):139-50. doi:
10.1158/2326-6066.CIR-17-0134

34. Han J, Duan J, Bai H, et al. TCR Repertoire Diversity of Peripheral PD-1(+)CD8(+) T Cells Predicts
Clinical Outcomes after Immunotherapy in Patients with Non-Small Cell Lung Cancer. *Cancer*
Immunol Res 2020;8(1):146-54. doi: 10.1158/2326-6066.CIR-19-0398

Figures

Figure 1

Figure 1. Repeated antigenic encounters ex vivo leads to antigen-specific T-cell exhaustion. (A)

Representative fluorescent multimer staining of LMP2$_{426-434}$-specific CD8$^+$ T cells (LMP2) and total cellular

expansion (expressed as fold change relative to culture input – 15x10^6 cells). (B) Percentages, as well as

(C) calculated number of antigen-specific T cells at the indicated time points (n=10 different donors). (D)

Representative staining to determine naïve (Tn), central memory (Tcm), effector memory (Tem), and

effector T cells (Teff) and the percentage of HLA-A02/LMP2$_{426-434}$ multimer positive CD8$^+$ T cells within

these categories at different culture time points (n= 12 different donors). (E) Representative histograms of

immune checkpoint expression evaluated relative to fluorescence minus one (FMO) control in HLA-A0201-

LMP2$_{426-434}$ (LMP2) multimer-stained T cells and compiled results from 12 different donors. (F)

Representative staining of PD-1 and TIM-3 co-expression and compiled results from 9-13 different donors.

(H) Percentage expression of PD-1 and TIM-3 ligands on dendritic cells used to stimulate T cells (n=5

different donors). Significant p values are indicated on figure panels. All error bars represent standard

deviation to the mean (SEM).

Figure 2. Dual but not single PD-L1 and TIM-3 blockade improves T-cell expansion. (A) Total cell

expansion relative to input at the beginning of the culture in function of time and culture condition; no

blocking antibodies (ctrl), anti-PD-L1 (α-PD-L1), anti-TIM3 (α-TIM3) or both (double blockade - DB).

Absolute count (B) and percentage (C) of HLA-A0201-LMP2$_{426-434}$ (LMP2) multimer positive T cells in the

same conditions and at the same time-points. (D) T-cell differentiation phenotypes; naïve (Tn), central

memory (Tcm), effector memory (Tem) and effector T cells (Teff) and (E) percentages of LMP2 multimer

positive T cells expressing immune checkpoints in the function of culture conditions and time-points (n=4-

10 different donors). Significant (<0.05) p-values are indicated on the figure panels. All error bars represent

SEM.

Figure 3

Figure 3. Delayed double blockade (DDB) further improves T-cell yield. (A) Total T-cell expansion

relative to input at day 0 in the function of time and culture condition; no blocking antibodies (Ctrl), double

anti-PD-L1 and anti-TIM-3 applied at culture initiation (double blockade – DB), anti-PD-L1 introduced at day

0 and anti-TIM-3 introduced at day 7 (delayed double blockade – DDB). (B) Percentages and (C) absolute

counts of HLA-A0201-LMP2$_{426-434}$ (LMP2) multimer positive T cells from the same cultures. (D) T-cell

differentiation phenotypes and (E) immune checkpoint expression of LMP2 multimer positive cells in the

same culture conditions and time-points (n=8 different donors). Significant (<0.05) p-values are indicated

on the figure panels. All error bars represent SEM.

Figure 4. DDB expands a high proportion and number of functional antigen-specific T cells. (A)
Representative dot plots showing intracellular cytokine secretion following LMP2$_{426-434}$ antigenic re-
stimulation at day 28 and compiled results from 5-6 independent donor cultures demonstrating the
percentage of CD8$^+$ T cells secreting IFNγ, TNFα, IL-2 and multiple cytokines; expansion with no blocking

antibodies (CTRL), anti-PD-L1 and anti-TIM3 antibodies both introduced at the beginning of the culture

(DB) or both antibodies but anti-TIM3 introduced at day 7 (DDB). (B) IFNγ ELISpot results using 50,000

cells per condition harvested from the cultures at the indicated time-points and using the following

stimulating conditions: vehicle only (DMSO), anti-CD3 (CD3, positive control), irrelevant peptides (Irr.),

LMP2$_{426\text{-}434}$ peptide (LMP2). (C) Percentage of Ki-67 staining at day 28 among CD8^{+} T cells from the ctrl

vs DDB cultures (n=5 different donors) and (D) co-expression of surface CD107a and intracellular

granzyme-b (GzB) as surrogate for degranulation following LMP2$_{426\text{-}434}$ exposure. (E) Fluorescence-based

cytotoxicity assay (cell tracer yellow – CTY or cell tracer violet – CTV) showing loss of targets loaded

(LMP2+) or not (LMP2-) with the LMP2$_{426\text{-}434}$ antigen (numbers indicate percentages of total events) and

(F) compiled results at different target:T-cell ratio (from 3 different donors). The significant p-values are

indicated on the panels and error bars indicate SEM.

Figure 5

Figure 5. Single-cell RNA-sequencing of antigen-specific T cells after double immune checkpoint inhibition.

(A) T-stochastic neighbor embedding (t-SNE) of normalized single-cell gene expression after dimensionality reduction from control (CTRL) and delayed double blockade (DDB) conditions, color coded by donor and experimental condition on the left and by cluster on the right. The barplot represents the percentage of each sample (colored as above left) in each cluster labeled on the x-axis and color-coded at the top of the graph. (B) Dot plot representing the change in the expression of genes related to pathways of interest in the DDB condition compared to the CTRL condition. The color of the dots represents the log2 fold-change of the genes in the pathway and the size of the dot is representative of the percentile ranking of the comparison in random gene sets (see Methods). (C) Clonotype frequencies in the CTRL and DDB

657   conditions for each donor. (D) Similar to (B) comparing the cells of selected clonotypes/patients between

658   the two experimental conditions.

Figure 6. Impact of DDB on clonal diversity and stability in time.

(A) Estimate of TCR repertoire diversity using the Gini-Simpson index among HLA-multimer negative and

positive T cells at day 21 and 28. (B) Clonal hierarchy and clonal relatedness among

HLA-A0201-LMP2$_{426-434}$ multimer positive T cells between day 21 and 28 using the day 28 repertoire as

reference in the DDB (B) or control condition (C) as determined by bulk CDR3 sequencing. The control

condition in Donor 3 was not assessed due poor RNA quality at day 28. (D) Dot plot representing the change

in the expression of genes related to pathways of interest when comparing a clonotype of interest in a

donor/experimental condition to all the other clonotypes in the same donor/condition. The clonotypes

selected are those whose abundance vary by more than 20% between day 21 and 28.

Figure 7

Figure 7. Improved expansion of functional WT1-specific T cells with DDB. (A) Representative HLA-

A0201-WT1$_{37-45}$ multimer staining (WT1) of CD8$^+$ T cells. (B) Cell expansion expressed as fold increase

relative to cell input at the beginning of the culture (15×10^6) in ctrl (no blocking antibodies) or delayed double

anti-PD-L1 and anti-TIM3 blockade (DDB) condition, as well as percentage (C) and absolute numbers (D)

of WT1-specific T cells at different time-points in culture. (E) Proportion of intracellular cytokine expression

(IFNγ, TNFα, IL-2, and multiple cytokines) following WT1 exposure at day 28 of the culture in CD8[+] cells

and (F) IFN-γ ELISpot results at days 14, 21, and 28 of culture following exposure to vehicle alone (DMSO),

peptides not used in the culture (irrelevant, Irr) as negative controls, anti-CD3ε (positive control) and the

targeted peptide (WT1). (G) T-cell differentiation; naïve (Tn), central memory (Tcm), effector memory

(Tem), and effector T cells (Teff) and (H) percentage of immune checkpoint surface expression on WT-1

specific T cells. Four (4) different donors, p-values are indicated on the figures, error bars indicate SEM.

683