Article

A transnational and transregional study of the impact and effectiveness of social distancing for COVID-19 mitigation

Tarcísio M. Rocha Filho[1],* , **Marcelo A. Moret[2]** , **José F. F. Mendes[3],†**

1   Instituto de Física and International Center for Condensed Matter Physics, Universidade de Brasília, Campus Universitário Darcy Ribeiro, Asa Norte, 70919-970 - Brasília, Brazil; marciano@fis.unb.br
2   Centro Universitário SENAI CIMATEC, Salvador, BA – Brazil and Universidade do Estado da Bahia, Salvador - Brazil
3   Departamento de Física & I3N, Universidade de Aveiro, 3810-193 Aveiro, Portugal
*   Correspondence: marciano@fis.unb.br (T.M.R.F)
†   Conceptualization: T.M.R.F. Interpretation, writing, review & editing: all authors contributed equally

Abstract: We present an analysis of the relationship between SARS-CoV-2 infection rates and a social distancing metric from data for all the states and most populous cities in the United States and Brazil, all the 22 European Economic Community countries and the United Kingdom. We discuss why the infection rate, instead of the effective reproduction number or growth rate of cases, is a proper choice to perform this analysis when considering a wide span of time. We obtain a strong Spearman's rank order correlation between the social distancing metric and the infection rate in each locality. We show that mask mandates increase the values of Spearman's correlation in the United States, where a mandate was adopted. We also obtain an explicit numerical relation between the infection rate and the social distancing metric defined in the present work.

Keywords: COVID-19; Social isolation; epidemiological model

Citation: Rocha Filho, T. M.; Moret, M. A.; Mendes, J. F. F. Impact and effectiveness of social distancing for COVID-19 mitigation. *Journal Not Specified* **2021**, *1*, 0. https://doi.org/

Received:
Accepted:
Published:

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

1. Introduction

The current COVID-19 pandemic is the main health crisis in the world in a century, with over 220 million cases and 4.5 million deaths [1]. It began in China at the end of 2019, and has since expanded to every country in the world, with waves occurring at different times in each location. A number of interventions were implement in most countries, such as travel ban, social distancing and mandatory mask use [2,3], and its effects have been discussed in different works, which generally concluded that they were effective in reducing the growth of cases and deaths [4–9]. Possibly the more effective measures being lock-downs, workplaces and business closing and school closing, i. e. the social distancing policies [10], with travel restrictions expected to have modest effects in reducing transmission when there is a high circulation of the virus [11].

In order to quantify and qualify the degree of social distancing and its effects, some different approaches have been proposed: by survey questionnaires in the population in order to assess adherence to social distancing and to compare it to the growth of cases, or deaths [12], or by using mobility data from different sources [13–19]. In the latter case, a mobility or social distancing metric is compared to the growth rate of cases (or deaths) of COVID-19, or to the effective reproduction number R_t. As we discuss below, this introduces a limitation in the analysis due to the fact that the interpretation of both the growth rate and R_t at the beginning of the pandemic, when most of the population is still susceptible to the virus, is different to that at latter stages, when a non-negligible proportion of the population has already been infected, of has already been vaccinated. A more informative parameter, that better represents information on the circulation of the SARS-CoV-2 virus, is the average infection rate $\overline{\beta}$, which is proportional to R_t divided by the proportion of the susceptible population (see Eq. 3 below). This explains particularly the result by Gatalo et al. [20] who obtained a strong Pearson correlation

between phone mobility data and COVID-19 growth rates at earlier stages, but a weaker
correlation at later stages, for 25 counties in the United States.

We present here an analysis of the effect of social distancing for 22 European
countries and for the 50 and 27 states of the United States and Brazil, and the most
populous cities and municipalities for the latter two, respectively. These localities have
different situations and histories of the pandemic. For instance, as mask use became
mandatory at different moments for American states, we were able to obtain quantitative
evidence on its effect on enhancing social distancing policies.

Our main goal is to evidence a monotonous relationship between social distancing
data and the value of the infection rate, and to quantify it explicitly.

2. Material and methods

47 *2.1. Effective reproduction number*

48   The effective reproduction number $R_t(i)$ at day i, estimated from the generation
49 time distribution w_j with j the number of days between infections, is given by [21]:

$$R_t = \frac{I(t)}{\sum_{t'} w(t')I(t-t')},\tag{1}$$

50 with $I(t)$ the number (or proportion) of infected individual at day i. The effective
51 reproduction number can also be estimated from the series of deaths by first determining
52 the number of infected individuals as:

$$I(t) = \frac{1}{\bar{\theta}} \sum_{t'} N_{\text{deaths}}(t+t')u(t'),\tag{2}$$

where $u(t)$ is the distribution of the number t of days (taken as discrete) between first
symptom and death [22], $N_{\text{deaths}}(t)$ the number of deaths at day t and $\bar{\theta}$ is the average
infection fatality ratio [23], computed from the demographic structure in each locality.
We then use Eq. (1 to determine R_t at a given day.

2.2. Infection rate

The infection rate can be estimated as [24]:

$$\bar{\beta} = \frac{\gamma R_t}{S(t)},\tag{3}$$

59 with $S(t)$ the proportion of susceptible individuals in the population at day t, R_t the
60 time dependent effective reproduction number, and γ the recovery rate from infection
61 with the value reported in the literature. [25]. We can also write that

$$\bar{\beta} = P_c C,\tag{4}$$

where C is the average number of contacts of one individual per day, and P_c the prob-
ability of contagion of a susceptible individual from a single contact with an infected
individual. Social distancing acts by reducing the number of contacts C, while other
non-pharmaceutical interventions reduce the value of P_c.

2.3. Social distancing metric

As a proxy for the "amount" of social distancing, we define a metric quantifying the
deviation from a baseline representing the pre-pandemic normality. Many possibilities
exist, and different mobility data are available from different sources [26–29]. We require
that data is freely available, with coverage up to the city level. For these sources only
Google mobility trends satisfies these two criteria, providing data on the following six
categories of locations: retail and recreation (D_1); grocery and pharmacy (D_2); parks (D_3);
transit stations (D_4); workplaces (D_5) and residential (D_6), as percentages of variation
of time spent in each type of place, with respect to a baseline defined for the period of

January 3 to February 6, 2020. The symbols between parenthesis represent the numeric
value of the time series for each type of data. An increase in the time spent at residence
is expected to decrease the value of the infection rate $\overline{\beta}$, and is considered as a negative
contribution to the metric, while an increase in the remaining five categories are expected
to increase $\overline{\beta}$ and thus contribute with a positive sign. The social distancing metric is
then defined as a weighted average of the data for each category, with the specified sign,
with weights given by an (arbitrarily) estimated average proportion of the duration of a
day spent in each type of location, and given by

$$M \equiv 100 + \frac{0.5}{24}D_1 + \frac{1}{24}D_2 + \frac{0.5}{24}D_3 + \frac{1}{24}D_4 + \frac{9}{24}D_5 - \frac{12}{24}D_6, \tag{5}$$

83 where the value of 100 is added such that the baseline is close to this value, and has
84 no effect of the value of the Spearman's correlation. The resulting metric M for each
85 Brazilian and American state are shown in Figs. 1A and 1B, respectively, with a similar
86 behavior for the other localities considered here (not shown). This definition is such that
87 a smaller value of M represents a more beneficial situation.

Figure 1. Social distancing metric M in Eq. (1) for A) Brazil states and B) USA states.

88 *2.4. Spearman's rank-order correlation*

89   Spearman's rank-order correlation $r_s(A, B)$ between two time series $A = (A_1,$
90 $\ldots, A_{N_{\text{data}}})$ and $B = (B_1, \ldots, B_{N_{\text{data}}})$, of length N_{data}, with A_i the value of the series at
91 the i-th data value, is defined as [30]:

$$r_s[A, B] = 1 - 6\frac{\sum_{i=1}^{N_{\text{data}}} d_i^2}{N_{\text{data}}(N_d^2 - 1)}, \tag{6}$$

such that $-1 \geq r_s \geq 1$ and d_i is the difference in paired ranks of the two series A
and B, i. e. the difference in position of the i-th data point for the two data sets when
ordered in ascending order. The coefficient r_s measures the strength of how two variables
are monotonically related, by an increasing or decreasing relation if $r_s > 0$ or $r_s < 0$,
respectively.

In order to show the importance to account for the decreasing number of susceptible
individuals with time, we show in Fig. 2A the time evolution of R_t and $\overline{\beta}$ for the Los
Angeles county in the United States. As the proportion of susceptible individuals
decreases over time, R_t and $\overline{\beta}$ diverge slowly. By computing the Spearman's correlation
between M and R_t and between M and $\overline{\beta}$, for a period of $N_{\text{data}} = 150$ days for the same
data, we see from Fig. 2B that a small difference between M and $\overline{\beta}$ has a significant effect
on the value of r_s. The Spearman's correlation between M and R_t is close to zero at
later times while clearly positive for M and $\overline{\beta}$. This is explained by the fact that, from
Eq. (2), that the same value of R_t can correspond to different values of the infection
rate $\overline{\beta}$ which is directly related to the circulation of the virus, as it measures the rate at
which susceptible individuals are infected, and thus more closely related to the different
mitigation policies implemented. We conclude that using R_t to represent the stage of
the pandemic can lead to misleading results at later stages in assessing the effectiveness

₁₁₀ of social distancing, as the number of susceptible individuals decreases, and that of
₁₁₁ vaccinated individuals increase.

Figure 2. A) Time variation of $\overline{\beta}(t)$ and R_t for Los Angeles county. B) Spearman's rank-order correlation $r_s[M, \overline{\beta}]$ between the social distancing metric M and the infection rate $\overline{\beta}$ and $r_s[M, R_t]$ between M and the effective reproduction number R_t for Los Angeles county in the United States.

₁₁₂ *2.5. Data sources*

₁₁₃   The following data sources were employed in the present work:

₁₁₄ •   Population by age for Europe: World Population Prospects - United Nations –
₁₁₅   https://population.un.org/wpp.
₁₁₆ •   Time series of deaths and cases by country: World Health Organization –
₁₁₇   https://www.who.int/emergencies/diseases/novel-coronavirus-2019.
₁₁₈ •   Time series of cases and deaths by US counties and states: New York Times COVID-
₁₁₉   19 Tracker data set –
₁₂₀   https://raw.githubusercontent.com/nytimes/covid-19-data/master/us-counties.csv.
₁₂₁ •   Population by age group in US counties and states: United States Census Bureau –
₁₂₂   https://www.census.gov/data/tables/time-series/demo/popest/2010s-counties-
₁₂₃   detail.html.
₁₂₄ •   Data on days of mask mandate in the US: Center for Disease Control and Prevention
₁₂₅   – https://data.cdc.gov/Policy-Surveillance/U-S-State-and-Territorial-Public-Mask-
₁₂₆   Mandates-Fro/62d6-pm5i.
₁₂₇ •   Population by age group for Brazilian municipalities and states: Brazilian Institute
₁₂₈   for Geography and Statistics – https://brasilemsintese.ibge.gov.br/populacao.
₁₂₉ •   Time series for cases and deaths by COVID-19 by municipality and state in Brazil:
₁₃₀   Brazilian Ministry of Health – https://covid.saude.gov.br.
₁₃₁ •   Detailed data on vaccination in Brazil: Brazilian Ministry of Health –
₁₃₂   https://opendatasus.saude.gov.br/dataset/covid-19-vacinacao.

₁₃₃ **3. Results and discussions**

₁₃₄   The localities that are analyzed here are:

₁₃₅ •   All 50 United States states, from the first reported case up to December 20, 2021.
₁₃₆ •   The 24 United States counties with a population of at least one million and at least
₁₃₇   1000 deaths in 2020 (Nassau was not considered due to inconsistent data for the
₁₃₈   number of deaths), from the first reported case in each county up to December 20,
₁₃₉   2021.
₁₄₀ •   All 27 Brazilian states from February 26, 2020 to June 14, 2021.
₁₄₁ •   The 22 Brazilian cities (municipalities) with a population of at least 750 thousand
₁₄₂   from February 26, 2020 to June 14, 2021.
₁₄₃ •   All countries in the European Economic Community and the United Kingdom with
₁₄₄   Google Mobility data and at least one thousand deaths by COVID-19 in 2020, from
₁₄₅   March, 1^{st} 2020 to December 31, 2020, with a total of 22 countries.

₁₄₆ The span of time of the data was chosen to avoid the effect of vaccination in the United
₁₄₇ States and Europe, while for Brazil detailed and publicly available anonimized data
₁₄₈ on each vaccine shot delivered allows modeling the time evolution of the pandemic

for a longer period. For estimations of susceptible population in Eq. (3 we use the epidemiological model described in [31] to determine the attack rate in each locality and is described in Appendix A. Serological surveys also provide such estimates, but are not available for every locality and for the required time window and, where available, data do not have the required time resolution.

The results of the Spearman's rank-order correlation between the social distancing metric M and the infection rate $\overline{\beta}$ for each locality are show in Fig. 3. In order to assess the effect of mandatory mask use in each United States county and state we compute r_s for two periods: for the whole period, indicating in the corresponding graphic the percentage of time with a mask mandate, and for the period with a mask mandate, for those counties with a mandate for at least 50% of the days since the beginning of the pandemic, while for the remaining counties we consider the whole period and display the corresponding histogram in black. We also computed the Spearman's correlation separately for each of the six mobility data reported by Google, with results shown in Figs. 4 and 5. The average of $\overline{\beta}/\gamma$, over the time period considered for each locality, versus the total number of deaths at the end of each period is shown in Fig. 6, where an approximately linear relation is clearly visible, with the exception of a few cases in Brazil.

In order to established a numeric relationship between $\overline{\beta}$ and M let us assume the linear relation

$$\overline{\beta}(t) = \alpha M(t), \tag{7}$$

with α a constant, and consider only the time window that allows to an accurate estimation of R_t. The distributions of values of the ratio $\alpha/\gamma = \overline{\beta}(t)/\gamma M(t)$ for the Brazilian states, Brazilian municipalities, European countries, United States states and counties are shown in Figs 7A–E, with values for α/γ (CI 95%) given by 0.015 (0.0096–0.023), 0.019 (0.0081–0.042), 0.014 (0.0089–0.021), 0.015 (0.0091–0.027) and 0.014 (0.0084–0.024), respectively. We also show the best fit with a log-normal distribution for values of α/γ in Figs. 7F–J.

While vaccination reduces the proportion of susceptible individuals in the population, it does not alter the relationship of the infection rate $\overline{\beta}$ with social distancing policies with M as a proxy, and was explicitly taken into account in our analysis by using an epidemiological model with vaccination compartments. The approach presented here allowed to evidence a monotonous relationship between the infection rate in each locality and the social distancing metric M. It also allowed to explicitly obtain a numeric relationship between $\overline{\beta}$ and a metric for social distancing. Behavioral changes can also have a significant impact on the evolution of any epidemic, and are difficult to include in the current analysis. Nevertheless, the significant values obtained for the Spearman's correlation indicate the important role that social distancing has played up to now. This is particularly clear in Belgium ($r_s = 0.75$), Spain ($r_s = 0.8$) and the United Kingdom ($r_s = 0.88$), three countries with a high attack rate. The correlation is somewhat smaller for other localities, but nevertheless with significant positive values, clearly indicating an approximately monotonous relationship between the two variables.

For Brazil and the European countries the results for Spearman's correlation are quite similar: the variation in time spent at residence is negatively correlated with the infection rate, i. e. the more time spent at home the smaller the value of $\overline{\beta}$, while other categories are positively correlated. For the United States, due to a much greater variety of mitigation policies implemented [13], we see a slightly different picture. In general, time at residence is negatively correlated with $\overline{\beta}$ while time at workplace is positively correlated with the transmission rate, as expected. For the remaining categories (grocery and pharmacy, park, retail and recreation and transit stations) we observe both negative and positive correlations according to the locality, indicating that the most relevant categories are those related to the increase of time spent at home and the decrease of time spent at work places. For the United States case there is a significant increase in the

value of r_s when considering only the time period with a mask mandate, which indeed
shows its effectiveness.

The values of the proportionality constant α/γ between $\overline{\beta}(t)/\gamma$ and $M(t)$ are sur-
prisingly close to one another, despite the great differences in the history and imple-
mented policies to mitigate the COVID-19 pandemic. We obtain a log-normal distribution
for the value of α/γ (and for α consequently) for all types of localities considered here,
with average values significantly closer, despite all the differences between countries,
implemented mitigation policies, and timings. This points to a universal efficacy of
social distancing, enhanced by a mandatory mask use. The explicit linear relation in Eq. (
7) with the value obtained for the proportionality constant α can be used, for instance, in
modeling studies with different scenarios for social isolation.

Of course not only social distancing affects the evolution of the infection rate,
causing the variation observed for the Spearman's correlation for the different localities.
We note that even a small increase in $\overline{\beta}$, and thus a small decrease in M, for a long period
of time, results in a significant increase in mortality, as can be seen from Fig. 6. Our
analysis does not grasp the impact of great gatherings of individuals and the possible
effect of the so-called superspreading events [20], or the implications of contact tracing.

4. Conclusion

A proper choice of a variable to represent the current circulation of the virus is
central to assessing the effects of mitigation policies. The infection rate as expressed
in Eq. (4) is affected by the reduction of social contacts through the average number
contacts C, and by other implemented protocols, such as mask wearing, that reduce the
probability of contagion per contact P_c. On the other hand, the effective reproduction
number R_t, or any other measure of growth rate of the pandemic, also depends on
the current attack rate, and confuses variables in the analysis. This is an important
point to consider as a more detailed analysis requires a large data set, and therefore a
larger time series, and therefore a significant variation in the proportion of susceptible
individuals. Computing Spearman's correlation, rather than Pearson correlation, for
instance, allows us to more clearly evidence a monotonous relationship between the
social distancing metric as defined here and the infection rate, and computed from
the whole time series for each locality. Future research considering socioeconomic and
demographic data would certainly provide valuable information on mitigation strategies
targeted at specific groups, such as elders and individuals with comorbidities, as well as
the impact of school closure, each considered separately from other factors [32].

We hope that the present work will contribute to a better assessment of the effects of
social distancing, and at least partially of mask mandates, on the still ongoing mitigation
interventions against the COVID-19 pandemic.

5. Acknowledgments

This work received financial support from the National Council of Technological
and Scientific Development - CNPq (grant number 305291/2018-1 MAM) and i3N (grant
numbers UIDB/50025/2020 & UIDP/50025/2020 JFFM) - Fundação para a Ciência e
Tecnologia/MEC (Portugal).

1. John Hopkins University. John Hopkins Coronavirus Resource Center, 2021. Last accessed 4
September 2021.
2. Perra, N. Non-pharmaceutical interventions during the COVID-19 pandemic: A review.
Physics Reports **2021**, *913*, 1. doi:10.1016/j.physrep.2021.02.001.
3. Chiesa, V.; Antony, G.; Wismar, M.; Rechel, B. COVID-19 pandemic: health impact of staying
at home, social distancing and 'lockdown' measures - a systematic review of systematic
reviews. *Journal of public health (Oxf.)* **2021**, *fdab102. Advance online publication*. doi:
10.1093/pubmed/fdab102.

4. Margraf, J.; Brailovskaia, J.; Schneider, S. Adherence to behavioral COVID-19 mitigation measures strongly predicts mortality. *PLoS ONE* **2021**, *16*, e0249392. doi:10.1371/journal.pone.0249392.

5. Zhang, J.; Litvinova, M.; Liang, Y.; Wang, Y.; Wang, W.; Zhao, S. Changes in contact patterns shape the dynamics of the COVID-19 outbreak in China. *Science* **2021**, *368*, 1481. doi:10.1126/science.abb8001.

6. Thu, T.P.B.; Ngoc, P.N.H.; Hai, N.M.; Tuan, L. Effect of the social distancing measures on the spread of COVID-19 in 10 highly infected countries. *Science of The Total Environment* **2020**, *742*, 140430. doi:10.1016/j.scitotenv.2020.140430.

7. Ayouni, I.; Maatoug, J.; Dhouib, W.; Zammit, N.; Fredj, S.B.; Ghammam, R.; Ghannem, H. Effective public health measures to mitigate the spread of COVID-19: a systematic review. *BMC Public Health* **2021**, *21*, 1015. doi:10.1186/s12889-021-11111-1.

8. Zhang, M.; Wang, S.; Hu, T.; Fu, X.; Wang, X.; Hu, Y.; Halloran, B.; Cui, Y.; Liu, H.; Liu, Z.; Bao, S. Human mobility and COVID-19 transmission: a systematic review and future directions. *medRxiv preprint* **2021**. doi:10.1101/2021.02.02.21250889.

9. Oh, J.; Lee, H.Y.; Khuong, Q.L.; et al. Mobility restrictions were associated with reductions in COVID-19 incidence early in the pandemic: evidence from a real-time evaluation in 34 countries. *Scientific Reports* **2021**, *11*, 13717. doi:10.1038/s41598-021-92766-z.

10. Mendez-Brito, A.; Bcheraoui, C.E.; Pozo-Martina, F. Systematic review of empirical studies comparing the effectiveness of non-pharmaceutical interventions against COVID-19. *Journal of Infection* **2021**, *83*, 281. doi:10.1016/j.jinf.2021.06.018.

11. Chinazzi, M.; Davis, J.T.; Ajelli, M.; abd Marcia Litvinova, C.G.; Merler, S.; y Pionti, A.P.; abd Luca ROssi, K.M.; Vespigniani, A. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science* **2020**, *368*, 395. doi:10.1126/science.aba9757.

12. Jarvis, C.I.; Van Zandvoor, K.; Gimma, A.; Prem, K.; working group, C.C..; Klepac, P.; Rubin, G.J.; Edmunds, W.J. Quantifying the impact of physical distance measures on the transmission of COVID-19 in the UK. *BMC Medicine* **2020**, *18*, 124. doi:10.1186/s12916-020-01597-8.

13. Badr, H.S.; Du, H.; Marshall, M.; Dong, E.; Squire, M.M.; Gardner, L.M. Association between mobility patterns and COVID-19 transmission in the USA: a mathematical modelling study. *The Lancet Infectious Diseases* **2020**, *20*, 1247. doi:10.1016/S1473-3099(20)30553-3.

14. Candido, D.S.; Claro, I.M.; Jesus, J.G.; Souza, W.M.; Moreira, F.R.R.; Dellicour, S. Evolution and epidemic spread of SARS-CoV-2 in Brazil. *Science* **2020**, *369*, 1255. doi:10.1126/science.abd2161.

15. Wellenius, G.A.; Vispute, S.; Espinosa, V.; Fabrikant, A.; Tsai, T.C.; Hennessy, J. Impacts of social distancing policies on mobility and COVID-19 case growth in the US. *Nature Communications* **2021**, *12*, 3118. doi:10.1038/s41467-021-23404-5.

16. Singh, B.B.; Lowerison, M.; Lewinson, R.T.; Vallerand, I.A.; Deardon, R.; Gill, J.P.S.; Sing, B.; W, B.H. Public health interventions slowed but did not halt the spread of COVID-19 in India. *Transboundary and Emerging Diseases* **2021**, *68*, 2171. doi:10.1111/tbed.13868.

17. Nouvellet, P.; Bhatia, S.; Cori, A.; Ainslie, K.E.C.; Baguelin, M.; Bhatt, S. Reduction in mobility and COVID-19 transmission. *Nature Communications* **2021**, *12*, 1090. doi:10.1038/s41467-021-21358-2.

18. Yilmazkuday, H. Stay-at-home works to fight against COVID-19: International evidence from Google mobility data. *Journal of Human Behavior in the Social Environment* **2021**, *31*, 210. doi:10.1080/10911359.2020.1845903.

19. Nova, A.C.; Ferreira, P.; Almeira, D.; Dionísio, A.; Quintino, D. Are Mobility and COVID-19 Related? A Dynamic Analysis for Portuguese Districts. *Entropy* **2021**, *23*, 786. doi:10.3390/e23060786.

20. Gatalo, O.; Tseng, K.; Hamilton, A.; Lin, G.; Klein, E. Associations between phone mobility data and COVID-19 cases. *The Lancet Infectious Diseases* **2020**, *21*, E111. doi:10.1016/S1473-3099(20)30725-8.

21. Fraser, C. Estimating Individual and Household Reproduction Numbers in an Emerging Epidemic. *PLOS One* **2007**, *2*, e758. doi:10.1371/journal.pone.0000758.

22. Russell, T.W.; Hellewell, J.; Jarvis, C.I.; van Zandvoort, K.; Abbott, S.; Ratnayake, R.; working group, C.C..; Flasche, S.; Eggo, R.M.; Edmunds, W.J.; Kucharski, A.J. Estimatins the infection and case fatality ratio for coronavirus disease (COVID-19) using age-adjusted data from the outbreak on the Diamond Princess cruise ship, February 2020. *Eurosurveillance* **2020**, *25*, 2000256. doi:10.2807/1560-7917.ES.2020.25.12.2000256.

23. Verity, R.; Okell, L.C.; Dorigatti, I. Estimates of the severity of coronavirus disease 2019:
A model-based analysis. *The Lancet Infectious Diseases* **2020**, *20*, 669. doi:10.1016/S1473-
3099(20)30243-7.
24. Kermack, W.O.; McKendrick, A.G. A contribution to the mathematical theory of epidemics.
Proceedings of the Royal Society A **1927**, *115*, 700. doi:10.1098/rspa.1927.0118.
25. Li, R.; Pei, S.; Chen, B.; Song, Y.; Zhang, T.; Yang, W.; Shaman, J. Substantial undocumented
infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science* **2020**,
368, 489. doi:10.1126/science.abb3221.
26. Google. Google COVID-19 Comunity Mobility Reports, 2021.
27. Teralytics. Teralytics Matrix (Zürich, Switzerland), 2021.
28. Unacast. Unacast Social Distancing Scorecard, 2021.
29. Apple. Apple Mobility Trends, 2021.
30. Myers, J.L.; Well, A.D., Research Design and Statistical Analysis; Lawrence Erlbaum: London,
2003.
31. Rocha-Filho, T.M.; Mendesc, J.F.F.; Chow, C.C.; Phillips, J.C.; Cordeiro, A.J.A.; Scorza, F.A.;
Almeidai, A.C.G.; AMoret, M. A data-driven model for COVID-19 pandemic – Evolution
of the attack rate and prognosis for Brazil. *Chaos, Solitons & Fractals* **2021**. In press, doi:
10.1016/j.chaos.2021.111359.
32. Lachassinne, E.; de Pontual, L.; Caseris, M.; Lorrot, M.; Guilluy, C.; Naud, A.; Dommergues,
M.A.; Pinquier, D. SARS-CoV-2 transmission among children and staff in daycare centres
during a nationwide lockdown in France: a cross-sectional, multicentre, seroprevalence
study. *The Lancet Child & Adolescent Health* **2021**, *5*, 256. doi:10.1016/S2352-4642(21)00024-9.
33. Jrahmandad, J.; Lim, T.Y.; Sterman, J. Behavioral dynamics of COVID-19: estimating under-
reporting, multiple waves, and adherence fatigue across 92 nations. *SSRN* **2021**. Forthcoming,
doi:10.2139/ssrn.3635047.

Appendix A Epidemiological model

336 In order to determine the proportion of susceptible individuals in a given locality
338 we use the approach described in [31] based on the SEIAHRV epidemiological model
339 with variables described in Table A1

Variable	Description
S_i	Susceptible individuals
E_i	Exposed individuals (non-contagious)
I_i	Infected symptomatic individuals (contagious)
A_i	Asymptomatic symptomatic individuals (contagious)
H_i	Hospitalized individuals
$V_i^{(l,k)}$	Vaccinated individual with l doses with vaccine of type k without primary vaccination failure
$U_i^{(l,k)}$	Vaccinated individual with l doses with vaccine of type k with primary vaccination failure
R_i	Recovered individuals.
$d_l^{(i,k)}$	individual vaccinated per unit of time with the l-th dose of vaccine of type k.
$e_l^{(k)}$	Efficacy of vaccine of type k withe l doses.

Table A1. Variables in the SEIAHRV model in Ref. [31]. All variables are proportions with respect to the initial population and the index i refers to the age-group.

340 The proportion S_i of susceptible individuals is obtained from the epidemiological
341 model described in [31], with model equations:

$$\frac{dS_i}{dt} = -\left[\lambda_i + \mu' + \nu_1 + \frac{1}{p_v}\sum_{k=1}^{K} d_1^{(i,k)}\right]S_i$$
$$+\kappa'\delta_{i,1} + \nu_{i-1}S_{i-1},$$

Figure 3. Spearman's correlation index r_s between the social distancing metric M and the infection rate $\overline{\beta}$ for: A) Main Brazilian municipalities with population over 750 thousand; B) Brazilian states; C) 22 European countries; d) US counties with at least one million inhabitants. D) All American states. Bar colors give the proportion of days with a mask mandate since the beginning of the pandemic in each location, up to December 20, 2020; E) Same as (D) but considering only the period with a mask mandate. States without a mask mandate in the period considered are marked in black. F) Same as (D) but for the main counties in the United states. G) Same as (E) but for the main counties in the United states.

Figure 4. Spearman's correlation index r_s between each mobility variable and the infection rate $\overline{\beta}$ for the main Brazilian municipalities, each Brazilian state and European countries.

Figure 5. Spearman's correlation index r_s between changes in each mobility category and the infection rate $\bar{\bar{\beta}}$ for A) Counties with more than one million inhabitants and one thousand deaths for the period from the first COVID-19 case up to December, 20 2020; B) Same as (A) but for the period with a mask mandate, except those counties with no mask mandate in 2020 (marked in black in figure 3E) for which the whole period is considered; C) Same as (A) for all US states; D) same as (B) for all US states.

Figure 6. Total number of deaths per 100 thousand inhabitants at the end of the considered period as a function of the average value of $\overline{\beta}/\gamma$ during the same time span.

Figure 7. Coefficient $\overline{\beta}/\gamma M = \alpha/\gamma$ for A) Brazilian states; B) Brazilian municipalities, C) US states; D) US counties and E) European countries. The normalized histogram (in red) and the log−normal distribution function (in black) for the values for α/γ: F) Brazilian states; G) Brazilian municipalities, H) US states; I) US counties and J) European countries. The values for α/γ (CI 95%) are 0.015 (0.0096–0.023), 0.019 (0.0081–0.042), 0.014 (0.0089–0.021), 0.015 (0.0091–0.027) and 0.014 (0.0084–0.024), respectively.

$$\frac{dE_i}{dt} = \lambda_i S_i + \lambda_i \sum_{k=1}^{K} U_i^{(k)} + \nu_{i-1} E_{i-1}$$
$$- \left[\sigma + \mu' + \nu_i + \frac{1}{p_v} \sum_{k=1}^{K} d_1^{(i,k)} e_1^{(k)}\right] R_i,$$

$$\frac{dI_i}{dt} = (1-\chi)\sigma E_i - \left[\gamma + \mu' + \nu_i\right] I_i + \nu_{i-1} I_{i-1}$$
$$-(1-\chi)\zeta_i \sigma E_i(t-\tau_1),$$

$$\frac{dA_i}{dt} = \chi\sigma E_i - \left[\gamma + \mu' + \nu_i + \frac{1}{p_v} \sum_{k=1}^{K} d_1^{(i,k)} e_1^{(k)}\right] A_i$$
$$+\nu_{i-1} A_{i-1},$$

$$\frac{dH_i}{dt} = -\left[\psi + \mu' + \nu_i\right] H_i + (1-\chi)\zeta_i \sigma E_i(t-\tau_1)$$
$$-(1-\chi)\theta_i \zeta_i \sigma E_i(t-\tau_2) + \nu_{i-1} H_{i-1},$$

$$\frac{dR_i}{dt} = \gamma I_i + \gamma A_i - \left[\mu' + \nu_i + \frac{1}{p_v} \sum_{k=1}^{K} d_1^{(i,k)} e_1^{(k)}\right] R_i$$
$$+\psi H_i + \nu_{i-1} R_{i-1},$$

$$\frac{dV_i^{(1,k)}}{dt} = -\left[\mu' + \nu_i\right] V_i^{(1,k)} + \nu_{i-1} V_{i-1}^{(1,k)}$$
$$+e_1^{(k)} d_1^{(i,k)} - e_1^{(k)} d_2^{(i,k)},$$

$$\frac{dV_i^{(2,k)}}{dt} = -\left[\mu' + \nu_i\right] V_i^{(2,k)} + \nu_{i-1} V_{i-1}^{(2,k)} + e_2^{(k)} d_2^{(i,k)},$$

$$\frac{dU_i^{(k)}}{dt} = -\left[\lambda_i + \mu' + \nu_i\right] U_i^{(k)} + \nu_{i-1} U_{i-1}^{(k)}$$
$$+\frac{(1-e_1^{(k)}) d_1^{(i,k)} S_i}{p_v} - (e_2^{(k)} - e_1^{(k)}) d_2^{(i,k)},$$

$$p_v = S_i + E_i + A_i + R_i. \tag{A1}$$

350 This is a non-linear delayed set of ODEs due ti the time delay between infection, hos-
351 pitalization and death. The different parameter values used in the model are given in
352 Table 1 of Ref. [31]. The force of infection in Eq. (A1) is given by

$$\lambda_i = \sum_{j=1}^{M} \beta_{i,j} \frac{I_j + \xi A_j}{n_i}, \tag{A2}$$

353 with $\beta_{i,j})$ the infection rate from an infected individual of age group j to infect an
354 individual of age-group i. The epidemiological model is calibrated using the time series
355 of deaths in order to avoid the significant under-notification of cases [33]. The value $\bar{\beta}$ in

356 Eq. (3) is an age-independent estimate obtained from the total proportion of susceptible
357 individuals obtained from

$$S(t) = \frac{1}{P_{\text{tot}}} \sum_i P_i S_i(t), \tag{A3}$$

358 where P_i is the population in age group i and P_{tot} the total population for the given
359 locality. The model is fitted from the time series of deaths as described in [31].