Secondary transmission of SARS-CoV-2 in educational settings in Northern Italy from September

2020 to April 2021: a population-based study

Authors:

Olivera Djuric[1,2], Elisabetta Larosa[3], Mariateresa Cassinadri[3], Silvia Cilloni[3], Eufemia Bisaccia[3], Davide

Pepe[3], Massimo Vicentini[1], Francesco Venturelli[3], Laura Bonvicini[1], Paolo Giorgi Rossi[1], Patrizio

Pezzotti[4], Alberto Mateo Urdiales[4], Emanuela Bedeschi[3], and the Reggio Emilia Covid-19 Working

Group.

Affiliations:

1) Epidemiology Unit, Azienda Unità Sanitaria Locale – IRCCS di Reggio Emilia, Reggio Emilia, Italy

2) Center for Environmental, Nutritional and Genetic Epidemiology (CREAGEN), Section of Public
Health, Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio
Emilia. Italy

3) Public health unit, Azienda Unità Sanitaria Locale – IRCCS di Reggio Emilia, Reggio Emilia, Italy

4) Department of Infectious Diseases, Istituto Superiore di Sanità, Rome, Italy.

The following are members of the Reggio Emilia Covid-19 Working Group:

Reggio Emilia Covid-19 Working Group: Massimo Costantini, Paolo Giorgi Rossi, Roberto Grilli,
Massimiliano Marino, Debora Formisano, Giulio Formoso, Emanuela Bedeschi, Cinzia Perilli, Ivano
Venturi, Eufemia Bisaccia, Elisabetta Larosa, Mariateresa Cassinadri, Silvia Cilloni, Cinzia Campari,
Francesco Gioia, Serena Broccoli, Marta Ottone, Pierpaolo Pattacini, Giulia Besutti, Valentina Iotti,
Lucia Spaggiari, Pamela Mancuso, Andrea Nitrosi, Marco Foracchia, Rossana Colla, Alessandro
Zerbini, Marco Massari, Anna Maria Ferrari, Mirco Pinotti, Nicola Facciolongo, Ivana Lattuada, Laura
Trabucco, Stefano De Pietri, Giorgio Francesco Danelli, Laura Albertazzi, Enrica Bellesia, Simone
Canovi, Mattia Corradini, Tommaso Fasano, Elena Magnani, Annalisa Pilia, Alessandra Polese, Silvia
Storchi Incerti, Piera Zaldini, Efrem Bonelli, Bonanno Orsola, Matteo Revelli, Carlo Salvarani,
Francesco Venturelli.

27
28 Corresponding author:
29
Olivera Djuric, MD, PHD
Epidemiology Unit
Azienda USL-IRCCS di Reggio Emilia
Via Amendola 2
42122 Reggio Emilia, Italy
Email: olivera.djuric@ausl.re.it
Tel: +39/ 0522335278

Abstract

Background: We aimed to quantify the risk of transmission of SARS-CoV-2 in the school setting by

type of school, characteristics of the index case and calendar period in the Reggio Emilia province

(RE), Italy. The secondary aim was to estimate the promptness of contact tracing.

Methods: A population-based analysis of surveillance data of all COVID-19 cases occurring in RE,

Italy, from September 1, 2020, to April 4th, 2021, for which a school contact and/or exposure was

suspected. Indicator of the delay in contact tracing was computed as the time elapsed since

positivity of the index case and the date on which the swab for classmates was scheduled (or most

were scheduled).

Results: Overall, 30,184 and 13,608 contacts among classmates and teachers/staff, respectively,

were identified and received recommendation for testing; 43,214 (98.7%) performed the test.

Secondary transmission occurred in about 40% of the investigated classes, and the overall secondary

case attack rate was 4%, slightly higher when the index case was a teacher, but with almost no

differences by type of school and stable during the study period. Promptness of contact tracing

increased during the study period, reducing the time from index case identification and testing of

contacts from 7 to 3 days, as well the ability to identify possible source of infection in the index case.

Conclusions: Despite the spread of the Alpha variant during the study period in RE, the secondary

case attack rate remained stable from school reopening in September 2020 until the beginning of

April 2021.

Key words: SARS-CoV-2, COVID-19, schools, attack rate, contact tracing

Background

Background

Since the first case of COVID-19 was described at the end of 2019, there have been almost 249

million cases and 5 million deaths reported globally up to October 2021.[1] In the period prior to the

implementation of mass vaccination programmes, social distancing and movement-restriction

measures were the main measures undertaken to prevent virus transmission. Closure of educational

institutions was one of the preventive measures considered and often adopted during the pandemic.

This was due to the concern about potential school-to-home transmissions of the virus from

students to more susceptible family members, although the overall risk of severe COVID-19 in

children and young people was shown to be very small.[2]

The role of school contacts in the spread of the virus as well the effectiveness of school closures in

controlling the epidemic have been debated, and inconsistent results have emerged.[3,4] Modelling

studies have provided scenarios with a small impact of school closure on virus transmission under

the conditions hypothesized by the authors.[5,6,7] A recent systematic review of empiric studies[8]

showed heterogenous results ranging from no effect to an important reduction in virus transmission

in the community, with studies with lower risk of bias showing a small effect.[9,10] Finally, three large

studies from the US, the UK and Italy suggest a limited role of school transmission in determining

mortality in the community.[11-13] It is worth noting that school opening may impact on virus

transmission because of the increase in public transport-related contacts and other non-school

activities, not only through contacts occurring in the classroom.

In our first study conducted between 1 September and 15 October 2020, non-negligible secondary

attack rates were detected, especially in secondary schools. After this first report, several changes

occurred that could have influenced the risk of transmission and its control in schools.[14] Starting

from 27 November 2020, the local health authority improved contact tracing protocols and

introduced immediate molecular tests for all contacts, whether symptomatic or asymptomatic, at

the beginning of quarantine, with the aim of identifying all possible sources of infection in

asymptomatic contacts, i.e., backward tracing (figure 1). Policies to reduce crowding especially in

high schools were introduced (reducing in-class time by 25% to 50%) as were several short closures

in the periods of highest incidence (figure 2). Finally, at the end of December 2020, the Alpha

variant started to circulate in our province, becoming predominant in February 2021.[15]

In this study we present the results of comprehensive epidemiological investigations conducted from

school reopening in September 2020 until the end of March 2021. Our aim was to quantify the risk

of infection transmission in the school setting by type of school, characteristics of the index case and

calendar period in the Reggio Emilia province by analysing surveillance data about all cases who had

contacts in the school setting. The second aim was to estimate the promptness of contact tracing.

Methods

Setting

There are approximately 95,000 inhabitants form 6 months-old to 19-year-old that can attend

infant-toddler centres (ages 0-3 years), preschools (ages 3-5), primary schools (ages 6-10), middle

schools (ages 11-13) and high schools (ages 14-19), and about 12,000 teachers/ school staff

members in Reggio Emilia province (531,751 inhabitants, Emilia Romagna, Northern Italy). During

the study period, there were two peaks of infections: in November 2020 and in February/March

2021 (figure 2). After the school reopening on 1 September 2020 for remedial courses and on 15

September 2020 for the regular school year, in-class learning was in place until 26 October 2020,

when most high schools moved to remote learning for at least 75% of scheduled lessons. In addition,

because of the to the high circulation of the virus, the Christmas school holidays were extended to

the second week of January, i.e., from 20 December to 11/15 January. Another lockdown led to the

closing of schools on 3 March 2021 (figure 2). Only infant-toddler centres and preschools, schools

that requires laboratory work and schools with pupils with disabilities or special needs continued in-

class didactic activities.

Design

An analysis of population-based surveillance data was conducted including all consecutive positive

cases confirmed with RT-PCR for SARS-COV-2 infection between 1 September 2020 and 4 April 2021

in Reggio Emilia province that led to an epidemiological investigation among children and

adolescents (0–19 years) who may have had exposure or contact with positive cases at school. Given

that reporting positivity of even one child led to a field investigation, contact tracing and, in some

cases, quarantine measures, all cases were included, regardless of the presence of an outbreak or

not. We excluded cases that occurred among children not attending schools in the period

investigated for contact tracing, i.e., in the period starting 48 h before symptom onset, and for

asymptomatic cases, 48 h before diagnosis or 48 h after contact with a certain case, whichever

occurred first.

Control measures

When a case occurred in primary and secondary schools, in-class activities were usually suspended,

but quarantine was not mandatory if all physical distancing measures were respected (sufficient

distance in the classroom, physical distance during all activities and utilization of face masks). All

students and school personnel were asked to perform a swab, with an appointment scheduled as

early as possible and communicated to the parents by means of a text message. If at least one

secondary case was found, quarantine was mandatory for all students in that class. For all close

contacts identified during the epidemiological investigation, a 10-day quarantine and an exit

negative test were required. In the infant-toddler and preschool settings, both the teachers and

children were considered close contacts if they had had contact with a case at school 48h before

symptom onset or a positive swab, in which case they were subjected to quarantine. In some cases,

based on the surveillance team's evaluation, the entire school was closed. Teachers were never

quarantined for school contacts, except if they taught in an infant-toddler centre or preschool.

Other control measures have been described more in detail elsewhere[14]; the major changes

occurring during the study period are described in figure 1.

Outcome definition and variables of interest

The main outcome was the secondary transmission rate. Overall attack rates were calculated by

dividing the number of cases by the population at risk, i.e., classmates, teachers and staff, who had

had close contact with the index case in a period starting from 48h before symptom onset of the

symptomatic index case and, for asymptomatic cases, 48h before diagnosis or 48h after contact with

a confirmed case, whichever occurred first. Mean attack rate was calculated as the mean value of all

class-level transmissions.

A classroom cluster was defined as the presence of two or more positive students or teachers/staff

members in the same class.

In this analysis we consider each single case or outbreak within a class as one statistical unit. For

each class an index case was identified, i.e., the first case who tested positive (considering the date

the swab was done). If more than one case in a class tested positive on the same day, the one with

the earliest symptom onset was considered the index case. The same class can be included more

than once because it may have been involved in more than one investigation during the study

period.

When more than one class was included in a between-class transmission or there was a single index

case for more than one class (usually, but not only, when the index case was a teacher), this was

considered a school cluster.

If a classmate was already in isolation prior to symptom onset or swab positivity of the index case,

due to contact with a positive person or re-entry from abroad, he/she was excluded from the

denominator. Any student or staff who refused to perform a swab was excluded from the

denominator. Cluster characteristics are reported; instead, information on the characteristics of

cases diagnosed outside the province or that were identified during some experimental screening

programme (2 classes) are missing.

Index cases were classified as student or teacher/staff and according to whether the case had a

known contact outside the school setting (household, social activity, sport activity, not identified,

and other/unspecified). The classes were classified according to the type of school (infant-toddler

centre and preschool, primary school, lower secondary school, upper secondary school) and whether

they were part of a multiclass cluster.

Delay in diagnosis of the index case, computed only for symptomatic cases, as the number of days

between symptom onset and the date of swab positivity. Delay in contact tracing was computed as

the time from swab positivity of the index case to the date on which the swab for (the majority of)

classmates was scheduled.

Descriptive statistic was used to analyse and present the data. Data were analysed using Microsoft

Excel and Stata v.13.1 (Statacorp, Tx).

Ethical statement

The study was approved by the Area Vasta Emilia Nord Ethics Committee (no. 2020/0045199). The

Ethics Committee authorised the use of subject data even in the absence of consent for people that

could not be reached, if every reasonable effort had been made to contact that subject.

Results

A total of 1213 student and 391 teacher/staff index cases were identified, corresponding to 1882

investigated classes, because some index cases, particularly teachers, had contacts in more than one

class. A total of 43,792 swabs were requested for contact tracing in these classes; 43,214 (98.7%)

were performed (table 1). The number of students and teachers/staff involved in contact tracing was

about slightly less than half of the number of total resident students and teachers/staff in the

province. However, this does not mean that epidemiological investigations involved this proportion

of the school population, because some classes might have been involved in more than one

investigation during the study period. Teacher index cases were more frequent in preschools and

primary schools than in secondary schools.

Student index cases were identified in 56.5% of cases as a contact of other cases, the vast majority

of which were within the same household (46.5%); a possible source of infection was identified for

only 26.6% of teacher index cases (Table 1).

Secondary transmission occurred in less than 40% of classes, and the overall attack rate was 3.9%

(95% CI 3.8-4.1%), 3.5% (95% CI 3.3-3.7%) for student index cases and 4.9% (95% CI 4.5-5.3%) for

teacher index cases (Table 1). The attack rate over the study period was 4.8% in September and

October, 2.6% in November, 3.9% in December and 4% from January to April (Table 2). The attack

rate was similar in all types of schools (Table 3).

The mean delay in testing was 6 days in student index cases and 4.8 in teachers (Table 1); this did

not change over the study period, except for a slight increase in December, and was similar across

types of schools (Table 3). On the contrary, the delay in contact tracing decreased from December

2020 on, from about 7 days to 3 days (Table 2), not showing any differences between type of index

case and type of school (Table 3).

Discussion

We confirm a modest secondary attack rate in schools, as already observed in previous studies.[16-19]

Secondary cases occurred in about 40% of classes that were exposed to an infected classmate or

teacher, and the overall attack rate was about 4%. Differently from our previous report[14] and other

studies reporting data on school cluster investigations,[19] the attack rate was similar in preschools,

primary schools and secondary schools. In fact, compared with the first two months of the study

period,[14] the attack rate increased in preschools from almost no transmission to 4%, while it

remained stable in secondary schools. It must be considered that remote learning and periodic

closures were implemented much more frequently in secondary schools, starting from 26 October

2020. Furthermore, an increase in the secondary case attack rate starting form late

December/January was expected because the spread of the Alpha variant, which is more infectious

than the previously circulating variants,[20] started in that period in our region.[15]

Surprisingly, the difference in attack rate between classes where the index case was a student and

those where the index case was a teacher was also small, despite the different kind of interactions

and length of exposure of student-student contact and teacher-student contacts in the school

setting. Unfortunately, we do not have any analytical information to define the quality and the

duration of interactions between index cases and contacts.

The effect of changes in contact tracing strategies introduced mainly after the second wave of the

pandemic, i.e., at the end of November 2020, was appreciable in the reduction in the delay in

contact tracing and in the increase in the proportion of index cases for which a contact, particularly a

contact in their household, was known. Indeed, a higher proportion of school investigations with a

known link to household clusters was precisely the anticipated effect of introducing immediate

testing of all asymptomatic contacts for all incident cases in the community, because this backward

tracing made it possible to identify asymptomatic or paucisymptomatic students and to link them to

their school contacts.

Limitations and strengths

The main limitation of this study is that any observed association between changes in secondary

attack rate and school or index case characteristics and especially actions implemented during the

study period could be confounded by other factors that we could not measure. Among these, the

most important was the spread of the Alpha variant, which probably started in our region around

Christmas and which became the dominant circulating variant in February. Unfortunately, as very

few cases were sequenced in Italy in that period,[15] we cannot determine which clusters were due to

the Alpha variant. Other variables that were missing or difficult to standardize were the duration and

intensity of contacts between the index case and other classmates and the control measures each

school independently put in place both before the occurrence of the case and during the outbreak.

The main strength of the study is its population-based nature; it included all cases for a which a

school contact or exposure was identified. For an analysis of routine surveillance data, the

completeness of field investigations is surprisingly high; only a few classes were not characterized,

and very few classmates refused to be tested. Each outbreak in each school was investigated and re-

checked for plausibility by the investigation team, thereby partially overcoming the shortcomings of

cross-sectional, retrospective and self-reported studies.

Conclusions

Overall, we confirmed a moderate risk of secondary transmission in the school setting (4.1% vs 3.2%

in the previous study up to 15th October 2020), similar across all types of schools. The attack rate

was also stable during the study period, despite the fact that the Alpha variant, which is much more

infectious than the previously circulating variants, became predominant in the last included months.

The promptness of contact tracing increased over the study period, reducing the time from index

case identification to testing of contacts from 7 to 3 days, as well the ability to identify possible

sources of infection in the index case.

Acknowledgements

We thank all the Public Health Department officers and COVID-19 reference persons in the schools

that collaborated in the investigations.

We thank Jacqueline M. Costa for the English language editing.

Conflict of interest

None declared.

Authors' contributions

PGR, EL, MV and PP designed the study. EL, MC, SC, EB, DP, FV and EB contributed to the collection

and management of the data. OD and LB analysed the data. OD and PGR interpreted the data and

wrote the manuscript. AMU contributed to data analysis and interpretation of results. All authors

revised the manuscript and approved the final version.

Funding

The study was conducted using exclusively institutional funds of the Azienda USL-IRCCS di Reggio

Emilia. There was no external funding source for this study.

What is already known on this topic

• Closure of educational institutions was one of the preventive measures considered and

often adopted during the pandemic.

• The role of school contacts in the spread of the virus as well the effectiveness of school

closures in controlling the epidemic is still debated.

• Potential effect of the more prompt (backward) contact tracing on secondary transmission in

educational setting has not been evaluated.

What this study adds

• Risk of transmission in educational setting is moderate with stable attack rates over time

and by type of school.

• Secondary attack rate is slightly higher if index case is a teacher.

• The tracing delay has halved after introducing the backward contact tracing, while ability to

identify possible sources of infection in the index case increased.

References

1. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time.

Lancet Infect Dis. 2020;20(5):533-534. doi: 10.1016/S1473-3099(20)30120-1.

2. Viner RM, Mytton OT, Bonnel C, Mellendez-Torres JG, Ward J, Hudson L, et al. Susceptibility to

SARS-CoV-2 Infection Among Children and Adolescents Compared With Adults. A Systematic Review

and Meta-analysis. JAMA Pediatr. 2021;175(2):143-156. Doi: 10.1001/jamapediatrics.2020.4573

3. Gaythorpe, KAM, Bhatia S, Mangal T, et al. Children's role in the COVID-19 pandemic: a systematic

review of early surveillance data on susceptibility, severity, and transmissibility. 2021;11(1):13903.

doi: 10.1038/s41598-021-92500-9.

4. Haug N, Geyrhofer L, Londei A, Dervic E, Desvars-Larrive A, Loreto V, Pinior B, Thurner S, Klimek P.

Ranking the effectiveness of worldwide COVID-19 government interventions. Nat Hum Behav.

2020;4(12):1303-1312. doi: 10.1038/s41562-020-01009-0.

5. Head JR, Andrejko K, Cheng Q, et al. The effect of school closures and reopening strategies on

COVID-19 infection dynamics in the San Francisco Bay Area: a cross-sectional survey and modeling

analysis. medRxiv [Preprint]. 2020. doi: 10.1101/2020.08.06.20169797.

6. Keeling MJ, Tildesley MJ, Atkins BD, et al. The impact of school reopening on the spread of COVID-

19 in England. Philos Trans R Soc Lond B Biol Sci. 2021;376(1829):20200261 doi:

10.1098/rstb.2020.0261.

7. Landeros A, Ji X, Lange K, et al. An examination of school reopening strategies during the SARS-

CoV-2 pandemic. PLoS One. 2021;16(5):e0251242. doi: 10.1371/journal.pone.0251242.

8. Walsh S, Chowdhury A, Russell S, et al. Do school closures reduce community transmission of

COVID-19? A systematic review of observational studies MedRxiv. 2021. doi:

https://doi.org/10.1101/2021.01.02.21249146.

9. Courtemanche C, Garuccio J, Le A, et al. Strong Social Distancing Measures In The United States

Reduced The COVID-19 Growth Rate. Health Aff (Millwood) 2020; 39: 1237–46. doi:

10.1377/hlthaff.2020.00608.

10. Hsiang S, Allen D, Annan-Phan S, et al. The effect of large-scale anti-contagion policies on the

COVID-19 pandemic. Nature 2020; 584: 262–7. doi: 10.1038/s41586-020-2404-8.

11. Ertem Z, Schechter-Perkins E, Oster E, et al. The Impact of School Opening Model on SARS-CoV-2

Community Incidence and Mortality: A Nationwide Cohort Study. Res Sq [Preprint]. 2021 Jul

15:rs.3.rs-712725. doi: 10.21203/rs.3.rs-712725/v1.

12. Forbes H, Morton CE, Bacon S, et al. Association between living with children and outcomes from

covid-19: OpenSAFELY cohort study of 12 million adults in England. BMJ. 2021;372:n628. doi:

10.1136/bmj.n628.

13. Marziano V, Guzzetta G, Rondinone BM, et al. Retrospective analysis of the Italian exit strategy

from COVID-19 lockdown. Proc Natl Acad Sci U S A. 2021;118(4):e2019617118. doi:

10.1073/pnas.2019617118.

14. Larosa E, Djuric O, Cassinadri M, et al. Secondary transmission of COVID-19 in preschool and

school settings in northern Italy after their reopening in September 2020: a population-based study.

Euro Surveill. 2020;25(49):2001911. doi: 10.2807/1560-7917.ES.2020.25.49.2001911.

15. Istituto Superiore di Sanità. Prevalenza e distribuzione delle varianti di SARS-CoV-2 di interesse

per la sanità pubblica in Italia. Rapporto n. 5 del 23 luglio 2021. July 2021. [Italian] Available at:

https://www.epicentro.iss.it/coronavirus/pdf/sars-cov-2-monitoraggio-varianti-rapporti-periodici-

23-luglio-2021.pdf

16. Ehrhardt J, Ekinci A, Krehl H, Meincke M, Finci I, Klein J, et al. Transmission of SARS-CoV-2 in

children aged 0 to 19 years in childcare facilities and schools after their reopening in May 2020,

Baden-Württemberg, Germany. Euro Surveill. 2020;25(36):2001587. doi: 10.2807/1560-

7917.ES.2020.25.36.2001587

17. Macartney K, Quinn HE, Pillsbury AJ, Koirala A, Deng L, Winkler N, et al. Transmission of SARS-

CoV-2 in Australian educational settings: a prospective cohort study. Lancet Child Adolesc Health.

2020;4(11):807-816. doi: 10.1016/S2352-4642(20)30251-0

18. Yoon Y, Kim KR, Park H, et al. Stepwise school opening online and off-line and an impact on the

epidemiology of COVID-19 in the pediatric population. MedRxiv.2020; (Preprint). doi:

10.1101/2020.08.03.20165589.

19. Aiano F, Mensah AA, McOwat K, Obi C, Vusirikala A, Powell AA, Flood J, Bosowski J, Letley L,

Jones S, Amin-Chowdhury Z, Lacy J, Hayden I, Ismail SA, Ramsay ME, Ladhani SN, Saliba V. COVID-19

outbreaks following full reopening of primary and secondary schools in England: Cross-sectional

national surveillance, November 2020. Lancet Reg Health Eur. 2021 Jul;6:100120. doi:

10.1016/j.lanepe.2021.100120. Epub 2021 May 19.

20. Davies NG, Abbott S, Barnard RC, et al. Estimated transmissibility and impact of SARS-CoV-2

lineage B.1.1.7 in England. Science. 2021;372(6538):eabg3055. doi: 10.1126/science.abg3055.

Figure legends

Figure 1. Timeline of the major changes in regulations and recommendations for management of

COVID-19 cases in educational settings in Reggio Emilia province

Figure 2. Weekly notification rates of new COVID-19 cases per 100,000 inhabitants aged 0-19 years,

by age class, Reggio Emilia province, 1 September 2020 – 31 March 2021. In the graph the main

changes in school opening and closure are also reported, and (in red) the proportion of Alpha variant

among sequenced cases as reported by the Italian National Institute of Health.

Tables

Table 1. Characteristics of the field investigation in 1884* classes for which a school contact with a COVID-19 case was suspected, by type of index case. Reggio Emilia, September 2020-March 2021

	Student		Teacher		Total*	
N of classes	**1224**		**658**		**1882**	
Type of school, n (%)						
Infant-toddler centre	194	15.9	156	23.7	350	18.6
Primary school	341	27.8	197	29.9	538	28.6
Middle school	310	25.3	186	28.3	496	26.3
High school	366	29.9	112	17.0	478	25.4
Other educational services	13	1.1	7	1.1	20	1.1
Classes with secondary transmission, n (%)	471	38.5	255	38.7	726	38.6
Classes that were part of a school cluster, n (%)	105	8.6	412	62.6	517	27.5
School clusters, n (%)	51		142		193	
Testing delay - days mean (SD)	6.1	4.8	4.8	4.2	5.6	4.6
Tracing delay - days mean (SD)	4.3	3.3	4.1	2.9	4.2	3.2
Number of contacts	30,184		13,608		43,792	
Number of tested contacts	29,802		13,412		43,214	
N of secondary cases	1047		658		1705	
Overall attack rate, % (95% CI)[¥][**]	3.5	3.3 - 3.7	4.9	4.5 - 5.3	3.9	3.8 - 4.1
Mean attack rate[^][**]	3.6	3.2-4.1	5.0	4.2-5.8	4.1	3.7-4.5
N of index cases	**1213**		**391**		**1604**	
Possible source of infection of index case, n (%)	686	56.6	104	26.6	790	49.2
Household outbreak	564	46.5	50	12.8	614	38.3
Social contact	19	1.6	7	1.8	26	1.6
Sport contact	18	1.5	0	0.0	18	1.1
Someone's contact	85	7.0	47	12.0	132	8.2
No contact reported	527	43.4	287	73.4	814	50.8

* In two classes there were no index cases due to screening

** Swabs not performed were excluded, as were 11 classes with unknown number of contacts

‡ Calculated by dividing the overall number secondary cases by the number of tested contacts

^ Mean of all classroom-level attack rates

Table 2. Characteristics of the field investigation in 1884 classes for which a school contact with a COVID-19 cases was suspected, by calendar period. Reggio Emilia, September 2020-March 2021

	September+October		November		December		January+February+March+April*	
N of classes	**248**		**263**		**316**		**1057**	
Type of school, n (%)								
Infant-toddler centre	39	15.7	72	27.4	53	16.8	186	17.6
Primary school	46	18.6	89	33.8	113	35.8	292	27.6
Middle school	52	21.0	82	31.2	104	32.9	258	24.4
High school	111	44.8	18	6.8	42	13.3	307	29.1
Other educational services	0	0.0	2	0.8	4	1.3	14	1.3
Classes with secondary transmission, n (%)	106	43.6	86	33.0	124	40.3	411	39.1
Classes that are part of a school cluster, n (%)	55	22.2	73	27.8	112	35.4	277	26.2
N of school clusters	22		27		37		107	
N of index cases	**226**		**220**		**250**		**908**	
Type of index case, n (%)								
Teacher	38	16.8	75	34.1	72	28.8	206	22.7
Student	188	83.2	145	65.9	178	71.2	702	77.3
Possible source of infection of index case	100	44.2	54	24.5	138	55.2	498	54.8
Household outbreak	59	26.1	45	20.5	113	45.2	397	43.7
Social contact	7	3.1	0	0.0	5	2.0	14	1.5
Sport contact	7	3.1	0	0.0	0	0.0	11	1.2
Unidentifiable contact	27	11.9	9	4.1	20	8.0	76	8.4
No contact reported	126	55.8	166	75.5	112	44.8	410	45.2
Number of contacts	6,327		6,279		6,860		24,314	
Number of tested contacts	6,252		5,964		6,827		24,204	
N of secondary cases	302		156		264		984	
Overall attack rate, % (95% CI)‡**	4.8	4.3 - 5.4	2.6	2.2 - 3.1	3.9	3.4 - 4.4	4.1	3.8 - 4.3
Mean attack rate^**	4.8	3.8-5.9	2.8	2.0-3.5	4.2	3.2-5.2	4.2	3.7-4.8
Testing delay - days mean (SD)	5.7 (4.3)		5.8 (3.8)		7.0 (5.1)		5.2 (4.6)	

| Tracing delay - days mean (SD) | 7.4 (3.8) | 7.4 (3.9) | 3.6 (1.7) | 2.9 (1.9) |

*Until 4 April 2021

**Swabs not performed were excluded, as were 11 classes without known number of contacts

¥ Calculated by dividing overall secondary cases by the number of tested contacts

^ Mean of all classroom-level attack rates

Table 3. Characteristics of the field investigation in 1884 classes for which a school contact with a Covid-19 cases was suspected, by type of school. Reggio Emilia, September 2020-March 2021

	Infant-toddler centre		Primary school		Middle school		High school		Other educational services	
N of classes	**350**		**540**		**496**		**478**		**20**	
Classes with secondary transmission	131	37.4	216	40.0	173	34.9	202	42.3	5	25.0
Classes that were part of a school cluster	25	7.1	162	30.0	207	41.7	116	24.3	7	35.0
N of school clusters	10		70		68		43		2	
N of index cases	**338**		**466**		**370**		**415**		**15**	
Type of index case, n (%)										
Teacher	144	42.6	126	27.0	66	17.8	51	12.3	4	26.7
Student	194	57.4	340	73.0	304	82.2	364	87.7	11	73.3
Number of contacts	7,767		11,709		12,106		11,936		306	
Number of tested contacts	7,652		11,575		11,923		11,799		298	
N of secondary cases	349		553		386		409		9	
Overall attack rate, % (95% CI)[¥][*]	4.6	4.1 - 5.1	4.8	4.4 - 5.2	3.2	2.9 - 3.6	3.5	3.1 - 3.8	3.0	1.4 - 5.7
Mean attack rate[^][*]	4.8	3.7-5.9	5.0	4.1-5.9	3.4	2.8-4.0	3.3	2.8-3.9	2.1	0.3-4.5
Testing delay - days mean (SD)	6.0	4.6	5.8	4.9	5.4	4.6	5.4	4.3	6.7	5.1
Tracing delay - days mean (SD)	4.4	3.5	4.1	3.1	4.1	3.3	4.4	3.0	3.3	2.3

*Swabs not performed were excluded, as were 11 classes without known number of contacts

[¥] Calculated by dividing the overall number of secondary cases by the number of tested contacts

[^] Mean of all classroom-level attack rates

Figure 1

21/9/2020

- Infant-toddler centres, preschools and elementary school (1st and 2nd grades): all present in the cluster 48h before symptom onset or swab positivity of the index case are close contacts and subject to 14-day quarantine and one swab.

- Elementary schools (3rd-5th grades), middle and high schools: only one swab without quarantine

Changed duration of isolation/quarantine and exit swab.

- For cases, 10 days, of which at least 3 days without symptoms + test: if negative, stop, if positive, 21 days of quarantine.

- For contacts, quarantine for 14 days from the last contact with positive person or 10 days from the last contact with positive person + negative test

12/10/2020

29/10/2020

Everyone over age 6 years should wear a mask at all times when in school facility or school bus

- One swab (including antigenic) should be performed as soon as possible for all close contacts

- Temporary suspension of attendance while awaiting results of the swab for close contacts in elementary and secondary schools

- The second swab after 14 days of quarantine for the close contacts who became cases

27/11/2020

31/01/2021

Extension of the period defining close contact for more than 48h

Figure 2