COVID-19 convalescent plasma and randomized clinical trials: rebuilding confidence by
explaining failures and finding signals of efficacy.

Daniele Focosi[1,#]

Massimo Franchini[2]

Liise-anne Pirofski[3]

Thierry Burnouf[4]

Nigel Paneth[5]

Michael J. Joyner[6]

Arturo Casadevall[7]

[1]North-Western Tuscany Blood Bank, Pisa University Hospital, Pisa, Italy.

[2]Division of Transfusion Medicine, Carlo Poma Hospital, 46100 Mantua, Italy;
massimo.franchini@asst-mantova.it

[3]Division of Infectious Diseases, Albert Einstein College of Medicine and Montefiore Medical Center,
New York, NY 10467, USA; l.pirofski@einsteinmed.org

[4]Graduate Institute of Biomedical Materials and Tissue Engineering & International PhD Program in
Biomedical Engineering, College of Biomedical Engineering, Taipei Medical University, Taiwan;
thburnouf@gmail.com

[5]Departments of Epidemiology & Biostatistics and Pediatrics & Human Development, College of
Human Medicine, Michigan State University, East Lansing, MI 48824, paneth@msu.edu

[6]Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, MN, USA;
joyner.michael@mayo.edu

[7]Department of Medicine, Johns Hopkins School of Public Health and School of Medicine, Baltimore,
MD, USA; acasade1@jhu.edu

[#]corresponding author: via Paradisa 2, 56124 Pisa, Italy. E-mail: daniele.focosi@gmail.com

Word count: abstract 250; body 5835.

Keywords: COVID-19; convalescent plasma; randomized clinical trial; propensity score-matched;

neutralizing antibodies; viral neutralization tests.

Abbreviations: BSC : best supportive care; CP : convalescent plasma; CCP : COVID-19

convalescent plasma; nAbs : neutralizing antibodies; VNT: viral neutralization test;

Author contributions: D.F. designed the paper, analyzed the data, and wrote the first draft. M.F.,

M.J.J., T.B., L.P., N.P. and A.C. revised the manuscript. All authors approved the final version.

Abstract

Convalescent plasma (CP) recurs as a frontline treatment in epidemics because it is available as
soon as there are survivors. The COVID-19 pandemic represented the first large-scale opportunity to
shed light into mechanisms of action, safety and efficacy of convalescent plasma using modern
evidence-based medicine approaches. Studies ranging from observational case series to randomized
controlled trials (RCT) have reported highly variable efficacy results for COVID-19 CP (CCP), resulting
in more doubt than certainty. Reasons for CCP success and failure may be hidden in study details,
which are usually difficult to explain to physicians and the public but provide fertile ground for
designing next-generation studies. In this paper we analyzed variables associated with efficacy such
as clinical settings, disease severity, CCP SARS-CoV-2 antibody levels and function, dose, timing of
administration (variously defined as time from onset of symptoms, molecular diagnosis, diagnosis of
pneumonia, or hospitalization, or by serostatus), outcomes (defined as hospitalization, requirement for
ventilation, clinical improvement or mortality), CCP provenance and time for collection, and criteria for
efficacy. Focusing only on the results from the 23 available RCT we noted that these were more likely
to show signals of efficacy, including reductions in mortality, if the plasma neutralizing titer was ≥ 160
and the time to randomization was ≤ 9 days, consistent with passive antibody therapy efficacy
requiring dosing with sufficient antibody. The fact that most studies revealed signals of efficacy
despite variability in CCP and its use suggest robust therapeutic effects that become apparent despite
the data noise.

Introduction

In the first 21 years of the 21st century humanity has experienced six major epidemics. The agents involved were SARS-CoV, MERS, influenza A(H$_1$N$_1$), Ebola, Zika and SARS-CoV-2 viruses. For five of these outbreaks the response included the use of convalescent plasma (CP) (reviewed in (1, 2)) and it was considered for the sixth (Zika virus). The attraction of CP is that it is readily available as soon as there are convalescing survivors, that unlike drugs or monoclonal antibodies it needs no development, and it is polyclonal, cheap and deployable even in resource poor countries. CP has been proposed as a first line response to new pandemics (3) and was deployed during the COVID-19 pandemic in March 2020 in countries that experienced the early waves of disease such as China (4, 5) and Italy (6).

While in early 2020 most clinical use was reported in case series or small phase II clinical trials (7), beginning in late March 2020 the US expanded access program (EAP) generated a large and robust treatment dataset, with insights on safety and optimal use. This database provided the first clear evidence that CP is safe, which was important given that early in the pandemic there were significant concerns about antibody-dependent enhancement (8). Later, an analysis of the first 3082 patents within the EAP database provided evidence that associated early administration of high titer CCP to non-ventilated hospitalized patients with reduced mortality (9). Before the FDA granted emergency use authorization (EUA), the US EAP provided CCP to as many as 94,287 patients. During the past year, many studies employing either randomized controls (RCT) or propensity score-matched (PSM) controls have been published. RCTs and PSM studies reported so far have had largely opposite outcomes, with most but not all RCTs finding little overall effect on mortality while the PSM and many smaller trials reporting mortality benefits. Several RCTs did not have mortality as a primary endpoint or it was part of a composite endpoint (5, 10-12). These disparate results have led to confusion for both the public and the clinicians, leading to reduced enthusiasm for the use of CP, in part because RCT data is more influential in affecting the opinion of many physicians, specialty societies and government regulators.

As with any other medical treatment, several key factors should be taken into account when evaluating a trial, including the indication (which can be estimated by timing or clinical severity), the therapeutic dose and the intended outcomes. The choices made by the trial designers determine whether the trial will demonstrate or conceal clinical benefit. While much attention is appropriately focused on the performance features of clinical trials (sample size, fidelity to randomization, appropriate analysis), the biological rationale for the hypothesis being tested is critically important but not always taken into account.

Methods

On September 7, 2021, we searched PubMed (which is also indexing the medrXiv prepublishing server) for clinical trials of CCP in COVID19, focusing on RCTs and PSM studies only. Each study was analyzed for the following variables: NCT identifier, recruitment, randomization strategy, type of

92   control arm, baseline patient status, median neutralizing antibody (nAb) titer in both recipients (before
93   CCP transfusion) and CCP units, type of viral neutralization test (VNT), primary endpoint, signals of
94   efficacy, and reasons for failure

95   At the same date, the ClinicalTrials.gov database was searched for CCP RCTs worldwide having as
96   status "completed", "active, not yet recruiting" or "recruiting".

Results

PubMed search retrieved 23 RCTs and 12 PSM studies about CCP, whose main variables are
summarized in Tables 2 and 3. The characteristics of the VNTs used are summarized in Table 1. The
variables were reconciled in 4 major topics, discussed in the following sections: the indication, the
therapeutic doses, the relevance of CCP to the viral variant, and the intended outcome.

ClinicalTrials.gov search retrieved 8 CCP RCTs completed but not yet prepublished or published, 7
active but not yet recruiting RCTs, and 10 RCTs which are still recruiting (summarized in Table 4).

The indication

While it would be desirable to have a single drug that works at any disease stage, it was not
reasonable to expect a silver bullet effect from neutralizing antibody-based treatments such as CCP in
later stages of disease. COVID-19 is now well-defined as a disease with two stages, an initial viral
phase characterized by flu-like and upper and lower respiratory symptoms, followed, in severe cases,
by an inflammatory phase that is characterized by inflammation-driven damage to multiple organ
systems, including the lungs that can impair gas exchange and cause life-threatening hypoxia and
damage to multiple organs, including the brain and blood vessels (13). Specific intact antibodies in
CCP are expected to neutralize SARS-CoV-2 in the intravascular system and, in some patients,
prevent progression from early to severe and life-threatening disease (as seen in animal models (14)),
but this antiviral therapy cannot be expected to reverse the inflammatory phase of the disease, nor
neutralize infectious viruses invading the extravascular system. Thus, COVID-19 is similar to
influenza, a disease in which antivirals are effective early in disease but have no effect in later stages
when the symptomatology stems largely from the inflammatory response.   The rationale for
administering CCP as early as possible in the course of COVID-19 stems from the neutralization
stoichiometry itself: the larger the number of actively replicating virions in the body, the higher the nAb
dose needed for neutralization (15). Some uncontrolled studies have reported a lack of association
between early intervention and outcomes (16, 17), but in these studies the level of neutralizing
antibody (nAb) or the overall anti-Spike antibody level in the infused CCP was unknown, leaving room
for alternative explanations.

At the beginning of the pandemic, some investigators and opinion leaders, riding the wave of CCP
successes in anecdotal reports in the media and small case series, introduced CCP to the general
public as a panacea for any patient with COVID-19, including life-threatening cases, leading to
confusing messaging: after reports of failure in severely ill patients emerged, opinions became
polarized and the debate became everything but scientific (18). In clinical trials, the indication (i.e., the

baseline clinical setting) has been variously defined by patient status (outpatient vs. presenting to the
emergency room vs. hospitalized vs. ICU-admitted), disease severity (using 5-category COVID-19
Outpatient Ordinal Outcome Scale (19), a 6-category ordinal scale (12), a 7-category COVID-19
severity scale (20), the WHO 8- (21) or 11-category (22) ordinal scales, or pneumological scores such
as SOFA), the time elapsed before recruitment (also variably defined as from molecular diagnosis,
from onset of hospitalization, from diagnosis of pneumonia, or from onset of symptoms), or by
serological status (presence of antibodies or the ability to neutralize SARS-CoV-2). This variability in
inclusion criteria for studies has resulted in marked heterogeneity in recruited patients.

An additional complexity in recruitment to CCP trials is time to treatment. Clinical trials involve
administrative requirements and consent procedures, and recruitment to a RCT further requires
randomization, which may produce delays in treatment. CCP therapy requires matching on blood
type, ordering the CCP, which may or may not be available on site, and setting up the transfusion.
This inherent delay from randomization to infusion means that RCTs may build in a disadvantage for
the CCP study arm, where controls may have received treatment earlier in the disease course (as, for
example, in the CP3O trial (23). ABO-compatible CCP units may be not readily available at the local
blood bank and recruited patients may have to wait for a compatible unit of CCP. These almost
inevitable delays from randomization mean that CCP may be provided later in the illness than is ideal,
and even if the trial intends to treat early, in practice it may not be possible.

During a pandemic, moreover, delays in treatment are magnified. The accrual of severely ill patients
in emergency departments and the overwhelmed or even collapsed health care systems can create
long delays from arrival in the emergency room to treatment. In the absence of quick (antigenic or
molecular) tests for SARS-CoV-2, the turnaround time for final confirmation of diagnosis with PCR,
which must often be run in batches, can take several hours. All of these factors are likely to impact the
efficacy of CCP treatment. To shorten such time, fully screened CCP collected from eligible donors
(24) could be safely administered within emergency departments shortly after admission and even
before the patient reaches the ward.

The therapeutic dose
Determining the effective dose of CCP is difficult in a pandemic because the antibody assays and
other tests needed to assess the potency of any antibody product take time to be developed. In
practice, the effective dose is the product of multiple factors, none of which is fully standardized. The
first factor is the concentration of the nAbs as measured by a VNT. At the beginning of the pandemic,
only a few BSL3 (or higher)-equipped virology laboratories could run VNT using authentic live SARS-
CoV-2 virus: the procedure was time-consuming (3-5 days) and the reports were operator-dependent.
Nowadays, the availability of Spike-pseudotyped viruses which can be managed under the more
widely available BSL-2 laboratories, or cell-free ACE-2 competition assays, combined with automated
(e.g., luminescence-based) readings, have standardized outcomes and shortened turnaround times
(25): however, harmonization between different assays is still a work in progress (26). The VNT differs
according to the type of replication-competent cell line, the viral isolate used for the challenge (which

is critically important when the virus is mutating rapidly as has been the case with emergence of
variants of concern), the multiplicity of infection (i.e., the ratio between the viral inoculum - referred
with different measuring units – and the number of replication-competent cells within each well), the
detection system (optic microscopy for cytopathic effect, immunostaining, quantitative PCR, or
luminometer for engineered pseudoviruses), and finally the threshold of neutralization (50% or 90%).
The DAWN-plasma RCT provides a clear example of such heterogeneity, with 4 different VNTs used
in at different participating laboratories. It was not until August 2020, when many trials were already
underway, that the FDA Emergency Use Authorization 26382 defined high-titer CCP on the basis of
correlation with a reference standard, the Broad Institute the live-virus, 5-dilution VNT as a 50%
inhibitory dilution (ID_{50}) of 1:250 or more (https://www.fda.gov/media/141481/download), and
exclusive use of high-titer CCP was formally recommended by the FDA only on March 9, 2021.

Table 1 summarizes the key variables in VNT employed to date in CCP RCTs. Published trials have
varied greatly in their approaches to antibody quantification whether in measured transfused CCP
units or in recipients. Many trials have relied on high-throughput semi-quantitative or qualitative
assays with a poor-to-moderate relationship with nAb titers. Although most trials performed a
correlation analysis between VNT and high-throughput serological assays, in many cases the CCP
units were tested only with the latter without validation, as was the case with 66% of the patients in
the PlasmAR trial (12). This procedure risks an incorrect evaluation of the neutralizing CCP activity.
Another cause for discrepancies in outcomes could be that although IgM, IgG, and IgA are all
capable of mediating neutralization, VNT titers correlate better with binding levels of IgM and IgA_1
than they do with IgG (27). Yet it is IgG that is routinely measured in high-throughput serological
assays, and these assays include non-neutralizing IgGs, the role of which in activity against SARS-
CoV-2 has not been established. Trials should preferentially use VNTs to assess serostatus of
transfused units and not rely on high-throughput serology.

As for any other medicinal product, CCP exhibits a dose-response relationship, which is also evident
when using high-throughput assays. In the subgroup analysis of the EAP, a gradient of mortality was
seen in relation to IgG antibody levels in the transfused CCP. In the subgroup of patients who were
not receiving mechanical ventilation, death within 30 days after CCP transfusion occurred in 81 of 365
patients (22.2%; 95% CI, 18.2 to 26.7) in the low titer group, 251 of 1297 patients (19.4%; 95% CI,
17.3 to 21.6) in the medium-titer group, and 50 of 352 patients (14.2%; 95% CI, 10.9 to 18.2) in the
high-titer group. Depending on the statistical model, the RR for 30-day mortality in high-titer CCP
compared to low-titer CCP recipients ranged from 0.64 – 0.67, with an upper 95% confidence bound
of 0.91 (8). Similarly, the large retrospective PSM study from HCA reported a 0.2% decreased risk of
mortality for every 1 unit of S/Co serology level (28).

The nAb titer (or total IgG levels as measured by surrogate assays) only describes one factor involved
in defining the real therapeutic dose in that it represents the concentration of just one (likely the main)
active ingredient. But CCP contains additional antibodies that mediate antibody-dependent cellular
cytotoxicity (ADCC), complement activation and phagocytosis of viral particles, functions that can
each contribute to its antiviral effects (29). At this time the relative importance of nAbs vs. the other

antibody activities is not understood, but, hopefully, retrospective analyses that correlate CCP efficacy
with these activities will reveal additional variables that need to be considered in choosing optimal
CCP units.

Despite these uncertainties, we can make estimates of likely effective doses based on the available
clinical experience thus far. The therapeutic dose of nAb is a product of its concentration in the
infused CP multiplied by the overall infused CP volume, adjusted to the recipient body weight to take
account of dilution into the blood volume and tissues. RCTs have varied in the provision of volume per
unit (200-300 ml), and most importantly in cumulative volume per patient (1-4 units) and in extent of
exposure to diverse antibodies from various CCP donors, and no published trials have adjusted levels
of nAbs by recipient body weight (or, when attempts have been performed, they referred to the old-
fashioned 10-15 ml/kg dose inferred from treatment of hemorrhagic coagulopathies (30)). A failure of
CCP to improve outcomes when 200-ml of 1:160 nAb-titer CCP is provided to a patient who weighs
120 kg represents quite a different scenario from failure of a 600-ml transfusion of 1:640 nAb-titer
CCP to produce improvement in a 60-kg patient. But these central issues in dosage have not been
considered in the RCTs published so far.

Several RCTs performed nAb titration, but with highly heterogenous methods which makes
comparability of doses across studies difficult. Table 1 attempts to reconcile doses across those trials,
showing that they actually differed more than was apparent by inspection of raw titers. The lack of
utility from low-titer (1:40) CCP in moderate COVID-19 was confirmed by the PLACID trial (10). As
long as a clear therapeutic dose is not identified, it seems prudent to transfuse units containing nAb
titers at least 10-fold higher than the nAb titer measured before transfusion in recipient serum.
Similarly, the ConCOVID RCT showed that CCP units having nAb titers similar to those of the
recipients (1:160) did not confer a clinical benefit (31). CCP units with an adequate nAb titer
(nowadays estimated at $\geq$1:160) are more easily found among older males who recovered from a
previous symptomatic COVID-19 requiring hospitalization (32, 33): unfortunately, such donors were
poorly represented in the first donation waves, which tended to obtain CCP from younger donors will
mild disease, and, presumably lower nAb titers (10).

Relevance of CCP to the viral variant

Albeit not formally demonstrated, CCP manufactured by pooling ABO-matched transfusion from many
different donors (e.g., in PlasmAr (12)) theoretically have greater polyclonality of nAbs than repeated
CCP doses from a single-donor (e.g. CAPSID (34)) and should grant higher efficacy against viral
variants. Nevertheless, pooling typically occurs among donors attending the same blood bank,
making donor exposure to different viral variants unlikely.

An analysis of potential variables associated with CCP efficacy associated near-sourcing with
reduced mortality, with the efficacy of CCP in reducing mortality falling sharply when the CCP source
was more than 150 miles from where it was used (35). This finding suggests that SARS-CoV-2
viruses vary enough in their antigenic composition in different geographic locations to create antibody

responses that differ by locale (36). Even though CCP is often standardized for nAb titer to the Spike
protein, the VNT could use a nonrelevant viral strain, or miss major functional differences for the
antibody response (29). This finding has implication for RCTs that use nationally sourced (centralized)
CCP, since the attempt to standardize the therapeutic units centrally could inadvertently reduce CCP
efficacy if hospitals use CCP obtained from distant loci. For example, in the C3PO RCT, which was
conducted in 21 USA states, 95% of the donor CCP was collected in either Chicago or Denver: since
only 4 of the 48 centers were in Illinois or Colorado, most CCP usage had to be from remote sources
(23). By contrast, the NCT04359810 RCT in New York and Brazil used CCP locally sourced in New
York, whose efficacy against P.1 was tested to ensure efficacy at the other recruiting center in Brazil
(11).

Although also not formally demonstrated during clinical trials, it is also reasonable to assume that
CCP collected during early pandemic waves could be less effective against currently circulating
variants of concern (37). RCTs whose recruitment was protracted across multiple pandemic waves
(e.g., ConPlas-19) and which relied on CCP collected and banked months earlier could have
inadvertently used CCP with reduced activity against the SARS-CoV-2 strains circulating the
community when the therapy was administered. Hence, both geography and time of collection of the
CCP are important variables when considering the efficacy of the treatment.

The intended outcomes
Most trials (CONTAIN, COMPILE, and PassItOn being exceptions) have used composite endpoints or
specialty scores (e.g., SOFA) rather than progression in the simple WHO ordinal scale or mortality,
and many were stopped because of apparent futility at a time when they may have been
underpowered to detect significant benefit. As represented in Figure 1, several studies have reported
overall negative results (panel A) despite the presence of positive signals of efficacy just barely
missing statistical significance (panels B and C). The significance level (i.e., $p= 0.05$) is largely a
socially constructed convention for rejecting the null hypotheses, but it has often been misinterpreted
as a measure of reality by many individuals not familiar with the nuances of statistics. For example,
some CCP studies have concluded that a difference that did not achieve a p value < 0.05 was an
absence of difference, even when mortality in the CCP arm was ~20-40% lower than in controls. This
reasoning played a central role in the polarized views of CCP efficacy and prevented subsequent
studies to drill down on positive effects that were observed. The dogged pursuit of statistical
significance, viewed as a measure of reality instead of the actual reality demonstrated by the data,
during a public health emergency dealt a serious blow to studies of CCP and created significant
confusion for clinicians. It is also important to understand that RCTs are powered to be less tolerant of
Type I error than Type II error, which are conventionally set at .05 and .20, meaning that a Type II
error is expected four times as often as a Type 1 error. This statistical convention can contribute to
the absence of significance in studies that were set up early in the pandemic when there was little
information on expected effects for the various patient populations studied and the patients were very
heterogenous such that only subgroups may have responded. Many studies were originally designed

to enroll patients at any disease stage, and it should be no surprise that subgroup analyses on the
groups that were later demonstrated more likely to benefit from CCP (e.g., early treated, seronegative
patients, those receiving high nAb titre) were underpowered to reach statistical significance, as shown
by orange color predominance in panel C of Figure 2. Nevertheless, favorable trends are a shared
feature across such trials. Lastly, rigid adherence to primary outcomes that were often fixed in the
early days of the pandemic when information about disease stage and quality of CCP associated with
efficacy were not understood. When these outcomes were not met, trials were considered failures
even though there were often signals of efficacy in the data that were not considered as valuable
since these had not been pre-specified, even when they made biological sense. For example, in the
New York-Brazil RCT cited above, CCP did not lower the primary end-point of clinical status on an
ordinal scale, but the statistically significant halving of mortality was acknowledged in the abstract.
Would it have made sense to ignore the strong effect of CCP on mortality in this trial just because
mortality was not selected as a primary outcome? Although we agree that subgroup analysis carries
the risk of 'cherry picking' data, such analyses are often important for hypothesis generation and
critically important during the emergency of a pandemic where neither viral pathogenesis nor
therapeutic variables are well understood. When sub-group analyses are based on firm biological
principles, such as focusing on those treated early in disease or lacking their own serological
response, the exercise is not cherry picking. To emphasize this point, Christopher Columbus missed
the pre-specified primary endpoint of his mission - reaching India - but no one considers his discovery
of the New World to be a failure! Turning to the clinical arena, most trials of anticoagulants in
myocardial infarction found reductions in mortality of about 20-25%, which was generally not
significant in these underpowered trials that declared the findings to be null, even though such a
mortality reduction would clearly be of value (38).

Another misunderstood endpoint is viral clearance, defined as the conversion of nasopharyngeal
swabs (NPS) from positive to negative for PCR evidence of SARS-CoV-2 in CCP-treated patients.
While there was early and robust evidence for this effect from CCP (4, 10), some RCTs failed to find
differences between arms just because they sampled NPS too late after CCP treatment, when the
endogenous immune response had also mounted in the control arm, and differences vanished.

Analyzing failures in individual RCTs.

We use the word 'failures' with care and considerable nuance, since negative trials can be very
important in teaching us about populations that do not benefit from CCP or variables that affect its
efficacy. Keeping the factors discussed above in mind, we have analyzed individual RCTs in detail.
At the very beginning, many historically or internally controlled observational studies showed clinical
benefit from CCP and this led the FDA to issue an EAP in March 2020 that was converted into an
emergency use authorization (EUA) in August 23, 2020. The largest observational study is the US
open-label EAP (NCT04338360) led from Joyner *et al*, which enrolled 105,717 hospitalized patients
with severe or life-threatening COVID-19 from April 3 to August 23, 2020 (39). In an analysis of the

effect of antibody in CCP performed independently of the results cited above (8) and using a nAb titer

in an overlapping but non-identical group of EAP patients, the FDA showed that the 7-day mortality in

non-intubated patients who were younger than 80 years of age and were treated within 72 hours after

diagnosis was 6.3% in those receiving high-titer CCP and 11.3% in those receiving low-titer CCP

(https://www.fda.gov/media/142386/download).

In a later analysis of a larger (N = 35,322) subset of EAP patients, (including 52.3% in the intensive

care unit (ICU) and 27.5% receiving mechanical ventilation), the 7-day mortality rate was 8.7% in

patients transfused within 3 days of diagnosis but 11.9% in patients transfused ≥4 days after

diagnosis. Similar findings again from the US EAP were observed in 30-day mortality (21.6% vs.

26.7%) (40). The major criticism of these results is that controls were neither randomized nor PSM:

hence a difference in the treatment outcome between treated and untreated groups may be caused

by a factor that predicts treatment rather than by the treatment itself. However, importantly, nAb titer

analysis was retrospectively done, both patients and physicians were unaware of the nAb content in

the CCP units used, the results are what would have been expected from the experience with

antibody therapy, and multivariate models were used to adjust for potential confounders (1).

Additionally, given the outline of an optimal use case with this data and the earlier underpowered RCT

by Li *et al* (5), it is unfortunate that due to (a) lack of awareness and (b) logistical burden associated

with protocol adjustments, involving repowering and new patients' recruitment criteria, later treatment

RCTs either continued or initiated without modifications to include newly available evidence.

The highest level of scientific evidence in primary clinical research stems from prospective PSM and

RCTs. PSM studies (Table 3) balance treatment and control groups on a large number of covariates

without losing a large number of observations. Unfortunately, no PSM study to date has investigated

nAb titers by VNT, and all times have been reported since hospitalization (excluding outpatients).

Nevertheless, in 2 retrospective PSM studies from 2 different hospitals in New York, trends for

improved outcomes in non-intubated and those treated within 7 days since hospitalization (HR 0.33)

were observed (41, 42). These findings were later confirmed in a prospective PSM study from

Houston (43, 44). Of interest, a retrospective PSM study from Providence did not show any benefit,

but patients were treated at a median of 7 days after onset of symptoms (45). Another PSM study

from Yale associated CCP with a 35% reduction in mortality (46). That study is notable in that it

included patients on mechanical ventilation who would not normally be expected to benefit from CCP

and the percentage of individuals receiving corticosteroids was very low since the study was

conducted in the early days of the pandemic in the USA. Another PSM from the Washington DC area

found a reduction in mortality with CCP use at both days 14 and 28, which reached statistical

significance at the earlier date (47). Finally, a very large study from 176 community hospitals affiliated

with Healthcare Corporation of America confirmed substantial mortality reduction in hospitalized

patients receiving CCP within 3 days from admission (48).

Since PSM only accounts for observed (and observable) covariates, and not latent characteristics,

RCT remains the gold standard for highest-level evidence (Table 2). In the PlasmAr RCT, the small

number of early arrivals (less than 72 hours) showed superior primary and secondary outcomes in the

CCP arm (n= 28) compared to the placebo arm (n=11), but the minimal contribution of this group to the overall cohort (228 CCP and 105 placebo) made the advantage disappear in the final outcomes at day 30 (12). In another Argentinean RCT on 160 outpatients older than 65 years of age with mild COVID-19 who were treated with CCP within 72 hours, progression to severe COVID-19 halved at day 30 (49). An RCT from India reported that patients younger than 67 treated at a median of 4 days after hospital admission manifested superior mitigation of hypoxia and survival in the CCP arm (50). Another RCT in Spain enrolling patients at less than 7 days of hospitalization showed four deaths in the control arm, none in the CCP arm (51). Given that conventional peer-review slows down during a pandemic, pre-publishing RCT results by the preprint mechanism should be encouraged to accelerate sharing of potentially life-saving therapeutic approaches and to provide pre-publication review that could improve the quality of the final published study.

Figure 1 graphically places the outcomes of RCTs and PSM studies on a Cartesian plot having timeliness and nAbs dose as variables (if values are disclosed in the reports): this makes immediately clear that the few successes at reaching the primary endpoints have gathered into the lower right corner (high nAb dose and early intervention), while the many "failures" have been scattered all around (panel A), reflecting lower antibody levels infused or late treatment, or both, with the latter being the commoner problem. Nevertheless, when we focus on mortality irrespective of statistical significance (panel B) or focusing on statistical significance (panel C), many more RCTs showed clear benefits.

We will focus here on "failures" as identified by title, abstract and/or press recognition. Narratively, we could group so-called "failures", with failure implying inability to demonstrate a favorable outcome to CCP use, into 4 categories, according to the main reasons:

1. Trials that transfused insufficient therapeutic doses of CCP due to either low total IgG levels or low nAb levels (e.g., PLACID)
2. Trials that transfused appropriate doses of CCP but too late, but which nevertheless reported signals of efficacy (e.g., RECOVERY, CAPSID, NCT04359810 and TSUNAMI)
3. Trials that were stopped too early to observe benefit or with inherent design flaws, and/or were underpowered such that likelihood of success was reduced (e.g., C3PO)
4. Trial in which CCP was used to treat a condition not amenable to antibody intervention, such as hypoxia that is caused by pulmonary inflammation

Stopping trials for futility is an occurrence that deserves special attention, because it represents wasted resources during a pandemic. Six RCTs so far have been halted for futility, namely RECOVERY, REMAP-CAP, CONCOR-1, C3PO, and NCT04361253, with the first one being to date the strongest evidence for futility (30), with its massive recruitment affecting the outcomes of systematic reviews (52). Instead of stopping trials for futility based on pre-set endpoints it makes more sense that DSMBs facing a high likelihood of lack of statistical significance provide advice on trial modifications that are likely to amplify the significance of signals of efficacy evident in these studies. This would seem a more responsible action than trial cessation given the paucity of therapeutic

alternatives in the pandemic emergency. Indeed, a Bayesian re-analysis of RECOVERY data with a
wide variety of priors (vague, optimistic, skeptical and pessimistic) calculated the posterior probability
for both any benefit or a modest benefit (number needed to treat of 100). Across all patients, when
analyzed with a vague prior, the likelihood of any benefit or a modest benefit was estimated to be
64% and 18% respectively. In contrast, in the seronegative subgroup, the likelihood of any benefit or
a modest benefit was estimated to be 90% and 74% (53). This finding of benefit accruing to specific
sub-groups, who were not determined post-hoc but because they were likely to benefit based on
understanding of principles of CP treatment is found in nearly every trial whose overall finding is
negative.

The inadequacy of meta-analyses.

With all the heterogeneity in key drivers discussed in the former paragraphs, it becomes clear that
secondary research (ranging from umbrella reviews to meta-analyses to systematic reviews), whereby
each study is considered at the same level, invariably ends up with biased and divergent conclusions.
This adds confusion to the already complex field of individual trial outcomes. Amazingly, as of August
24, 2021, PubMed has indexed 25 meta-analyses on CCP efficacy, more than the RCTs reported at
the same date. Until the beginning of 2021, meta-analyses (variably including observational studies)
were generally in favor of CCP (54), but began to be biased towards failure after publication of the
large RECOVERY trial (30), which, by enrolling as many as 11,448 patients, diluted all the other
divergent RCTs. Clear examples of this phenomenon come from a widely cited metanalysis from
Janiaud *et al* in *JAMA* (52) which included press release data from RECOVERY and from the living
systematic review by the Cochrane Group (55). This paper was surely unprecedented in the tradition
of meta-analysis, not only because it included a study based only on a news release (which proved to
differ in some important respects from the published paper), but because it allowed these data from a
news release to dominate the entire analysis. Several groups attempted to dissect the RECOVERY
trial and others by running subgroup analyses in their systematic reviews (53, 56, 57), but these
reviews were unable to restore confidence in CCP efficacy in the clinical community that had been
lost because of the publication of the overall negative findings of RECOVERY and PlasmAr (58). A
metanalysis of 22,591 patients (enrolled in 10 RCTs and 15 observational studies) showed that early
CCP significantly reduced mortality (RR 0.72, p<0.00001), but only in patients who were not suffering
severe or critical disease (59). On the other hand, another metanalysis of 18 peer-review clinical
trials, 3 preprints, and 26 observational studies actually found that CCP use was associated with
reduced risk of all-cause mortality in severe or critical COVID-19 patients (60). A recent umbrella
review of 29 metanalyses and systematic reviews found evidences for improvement in the CCP arms
for some outcomes (overall mortality, viral clearance at day 3,) but not for others (clinical
improvement, length of hospital stay (61).

Rather than pooling published RCTs, the Continuous Monitoring of Pooled International Trials of
Convalescent Plasma for COVID-19 Hospitalized Patients (COMPILE) study pooled individual patient

data from ongoing RCTs at two-week intervals. Unfortunately, with the single exception of CONTAIN,

participating RCTs largely shared late usage (DAWN-plasma, PLACID, ConCOVID, ConPlas-19,

NCT04421404, NCT04397757, and the Brasília Covid-19 Convalescent Plasma (BCCP)) (62).

Conclusions

While CCP contains a plethora of biologically active molecules (63), we now have very strong

evidence that appropriately vetted CCP from eligible convalescent donors is safe for patients (64, 65),

with no evidence of increased risks of transfusion-transmitted acute lung injury, antibody-mediated

enhancement concerns feared in the early days of the pandemic (66) nor is there evidence that CCP

induces accelerated SARS-CoV-2 evolution (11). Polyclonal antibodies such as CCP, or CCP-derived

hyperimmune globulins made from large donor pools, are likely to offer better protection against onset

of variants than monoclonal antibodies. Outcomes in immunocompromised patients treated with CCP

have been successful in the long-term, with minimal evidence for immune escape (67). There is

evidence that vaccinated convalescents may have even higher nAb titers than unvaccinated

convalescents offering the promise of expanded success in using CCP (68).

We have also learned that CCP is less likely to benefit patients requiring oxygen (i.e., from level 4 and

up on the 11-point WHO ordinal scale), and hence, ideally, the focus should be on outpatients and in

identifying that subset of patients who seek hospital care and are still sufficiently early in the course of

disease such that they can benefit from CCP. This finding parallels the finding with hyperimmune

serum and anti-Spike monoclonal antibodies, which at first failed in hospitalized patients (69, 70), but

later succeeded for ambulatory patients with mild to moderate COVID-19 (71) and were approved for

emergency use. However, at this moment clinical use in the US is restricted by the FDA to inpatients.

CCP usage per admission peaked after issuance of the EUA, with more than 40% of inpatients

estimated to have received CCP between late September and early November 2020. However,

following reports of RCTs that failed to show clear benefit from CCP, usage per admissions declined

steadily to a nadir of less than 10% in March 2021. A strong inverse correlation (Pearson correlation

coefficient of -0.5176 with P = 0.00242) was found between CCP usage/hospital admission and

deaths occurring 2 weeks after admission, and this finding was robust to examination of deaths taking

place 1, 2 or 3 weeks after admission. Changes in the number of hospital admissions, prevalence of

variants, and age of patients could not explain these findings. The authors estimated that the retreat

from CCP usage, a phenomenon they termed "plasma hesitancy", might have resulted in 29,000 to

36,000 excess deaths in the period from mid-November 2020 to February 2021 (72). The same

analysis estimated that USA had avoided 96,000 excess deaths from August 2020 to March 2021 by

its liberal deployment of CCP.

Several lines of evidence, ranging from the EAP to clinical trials employing RCT or PSM controls are

now indicating how CCP should be used in immunocompetent patients (73). The evidence supports

the initiation of CCP treatment as early as 44-72 hours within onset of symptoms (which largely

pertains to outpatients) and using CCP with a nAb titer > 1:160. Benefit within 1 week from onset of

symptoms (including in hospitalized patients) is less well understood, although a benefit from higher

therapeutic doses cannot be ruled out at this stage. Clinical benefit seems absent when administered
after 1 week from onset of symptoms or in patients requiring ventilation, or in those who receive CCP
with a low nAb titer. Nevertheless, chronically immunosuppressed patients benefit from CCP even at
later stages (67, 74, 75) : the best evidence for this scenario comes from a prospective PSM showing
a halving of mortality in ICU-admitted oncohematological COVID-19 patients who received CCP (76).
We note that while there have been concerns that use in immuncompromised can promote the
emergence of antibody-resistant variants, such variants have emerged from massive replication in
susceptible populations and not from treated patients, who in any case are isolated in hospitals where
mitigation efforts to reduce transmission are employed, and are thus very unlikely to transmit their
viruses further (77). Such simple concepts have been poorly communicated to the general public and
the clinical community, who should be better informed of the state of current evidence that support
CCP efficacy.

487 The future of CCP

CCP remains a relatively inexpensive therapy that is available throughout the world even in resource
poor areas that cannot afford expensive antiviral drugs or monoclonal antibody therapies. Much has
been learned about the variables that affect CCP efficacy even though, as recounted here, the clinical
efficacy data is mixed. Table 4 lists the RCTs whose outcomes have still to be reported after
completion or which are still recruiting patients. Unfortunately, little new can be expected given that
most of these RCTs were designed to enroll patients having symptoms for more than 7 days. Given
the heterogeneity of the product and the complex variables that contribute to efficacy it is remarkable
that many studies have reported reductions in mortality. This suggests a robust therapeutic effect that
allow signals of efficacy to break through all the noise imposed by variability in the product and its
clinical use. The positive evidence for CCP efficacy cannot be dismissed while negative results can
be explained. In the absence of good therapeutic options for COVID-19, CCP is likely to find a niche
in the early treatment of disease. Instead of looking for unlikely superiority outcomes, noninferiority
RCTs comparing monoclonal antibody versus CCP in early arrivals should be initiated. Such an RCT
is very unlikely to be sponsored by vendor companies, so public institutions should be sensitized to
funding it.

Given the experience accumulated with COVID-19, it is almost certain that CP will again be deployed
for the next epidemic and we are hopeful that lessons learned in this pandemic are heeded such that
use and trials focus on the very early use with high-titer CP.

We declare we have no conflict of interest to disclose.

Figure 1

Simplified graphical representation of CCP RCTs reported to date, plotted according to earliness of intervention and nAb titers in CCP. In **panel A**, green text indicates trials which met the primary endpoint with statistical significance; orange text indicates trials which failed to meet the primary endpoint but showed statistically nonsignificant trends in favor of CCP; red text indicates trials which failed to show and benefit from CCP in the primary endpoint. In **panel B**, green text indicates trials which showed overall mortality benefit from CCP; orange text indicates trials which showed mortality benefit from CCP in the subgroup of early arrivals or higher nAb titers; red text indicates trials which failed to show any mortality benefit from CCP. In **panel C**, green text indicates trials which showed statistically significant mortality benefit from CCP (overall or in the subgroup of early arrivals or higher nAb titers); orange text indicates trials which showed statistical trends towards mortality benefit from CCP (overall or in the subgroup of early arrivals); red text indicates trials which failed to show any mortality benefit trend from CCP in any subgroup. Underlined text indicates large trials which were not RCT and for which nAb levels was inferred from high-throughput serology, but are nevertheless reported as reference studies. Numbers in parenthesis indicate cumulative number of patients enrolled.

Table 1

520 Details of viral neutralization tests (VNT) employed in CCP RCTs. Information was retrieved from original article (including Supplementary Appendix).

521 Whenever not reported, the corresponding author was contacted (marked with *). If the details could not be retrieved the field is labelled "n.a." (not available).

522 IC: inhibitory concentration. NT: neutralization titer. PFU: plaque-forming unit.

	RCT (acronym/first author)	cell line	cells seeded per well	virus lineage	virus per well	multiplicity of infection (MOI)	length of incubation	assay read-out	threshold	protocol reference
authentic live SARS-CoV-2	NeuCoV-NET NCT04393727 (TSUNAMI)	Vero E6	12,000	SARS-CoV-2/Human/ITA/PAVIA10734/2020 (D614G)	100 $TCID_{50}$	0.01	until the cytopathic effect (CPE) became evident.	CPE	last serum dilution that inhibited SARS-CoV-2 CPE by 90%.	(78)
	NCT04433910 (CAPSID)	Vero E6	n.a.	n.a.	100 PFU	n.a.	3 days	CPE	$PRNT_{50}$	(79)
	Broad Institute on a high throughput platform (BROAD PRNT). Part of NCT04355767 (C3PO)	Vero E6-TMPRSS2	10,000	SARS-CoV-2 live virus (D614)	n.a.	n.a.	48 hours	N-protein ELISA	samples whose curves lay above 0.5 for all the data points were considered non-neutralizing, with ID_{50}=20, while samples whose curves fell below 0.5 were considered highly neutralizing and assigned an ID_{50}=10,240.	(80)
	NCT04359810 (O'Donnell)	Vero E6	10,000	2019-nCoV/USA-WA1-2020	100 $TCID_{50}$	0.01	48 hours	Triplex CII-SARS-CoV-2 rRT-PCR Test, EUA200510).	the highest CCP dilution that prevented virus growth (cycle threshold [Ct] was rated as neutralization titer.	(11)

	NCT04348656 (CONCOR-1)	Vero-E6	n.a.	Canada/ON_ON-VIDO-01-2/2020, EPI_ISL_42517	50 PFU	n.a.	72 hours	CPE	$PRNT_{50}$	(81)
	NCT04342182 (ConCOVID)	Vero-E6	n.a.	German isolate (GISAID ID EPI_ISL 406862)	400 PFU	n.a.	8 hours	CTL ImmunoSpot Image Analyzer	reciprocal of the highest dilution resulting in a reduction >50% of infected cells ($PRNT_{50}$)	(31)
	NCT04429854 (DAWN-plasma)	Vero E6	n.a.	BetaCov/Belgium/Sart-Tilman/2020/1	100 $TCID_{50}$	n.a.	5 days	CPE	$PRNT_{50}$	(82)
			18,000	2019-nCoV-Italy-INMI1	3 $TCID_{50}$	n.a.				(83)
			n.a.	Belgium/GHB-03021/2020	400 PFU	n.a.	4 days			(84)
			20,000	Belgium/S1871/2020	100 $TCID_{50}$	n.a.	2 days	anti-N staining	NT_{50}	(85, 86)
Spike pseudotyped viruses	Vitalant Research Institute (VRI) Pseudovirus Neutralization Part of NCT04355767 (C3PO)	ACE2 and TMPRSS2 expressing HEK293T cells	n.a.	VSV pseudotyped with Wuhan-Hu-1 Spike (D614G mutation and without 21 C-	n.a.	n.a.	18-24 hours	chemiluminescence reader	NT were calculated as a percentage of no-serum control and the NT_{50} was estimated from the dilution curve	(80)

								terminal aa)	
NCT04383535 (PlasmAr)	Vero-CCL81	20,000	VSV pseudotyped with Spike (CoV2pp) and carrying *Renilla* luciferase gene in place of its G glycoprotein (VSVΔG-rLuc).	n.a.	n.a.	18-22 hours	luminometer	IC_{50} is calculated as the midway point between the upper and lower plateaus of the curve. $absIC_{80}$ appeared to be a more stringent measure of nAb activity, as some sera that have respectable MN $absIC_{50}$ titers never achieve an $absIC_{80}$; this is due in part to the difference in the dynamic ranges between a luciferase-based assay (≥3 logs RLUs) and a MN assay (~1.5-log optical density [OD] values corresponding to the amount of viral protein detected).	(87)
CTRI/2020/04/024775 (PLACID)	Vero CCL-81 293 T/ACE2 cells	10,000	SARS-CoV-2 strain NIV2020770	n.a.	n.a.	36 hours	luminometer	n.a.	(88)
NCT04345523 (ConPlas-19)	Vero E6	5,000	lentivirus pseudotyped with Spike and luciferase	titrated at 10 ng p24 Gag	n.a.	48 hours	luminometer	ID_{50} expressed as the highest dilution of plasma (reciprocal dilution), which resulted in a 50% reduction of luciferase activity compared to control without serum. Sigmoid curves were generated and ID_{50} neutralization titers (NT_{50}) were calculated by non-linear regression	(20)

NCT04375098 (Elvira-Balcells)	HEK293T/hACE2	10,000*	HIV-1–SΔ19 pseudotyped with Spike (Genebank: QHU36824.1) and luciferase	n.a.	n.a.	48 hours*	luminometer	samples with a neutralizing activity of at least 50% at a 1:160 dilution were considered positive and used to perform titration curves and ID_{50} NT calculations	(89)
NCT04344535 (Bennett-Guerrero)	n.a.	n.a.	PRNT and pseudovirus	n.a.	n.a.	n.a.	n.a.	n.a.	(90)

Table 2

Randomized controlled trials (RCT) of COVID-19 convalescent plasma (CCP) reported to date. nAb: neutralizing antibodies. BSC: best supportive care. FFP: fresh frozen (nonconvalescent) plasma. n.a.: not assessed (i.e. antivirus antibodies were assessed only using high-throughput serology). IQR: interquartile range.

RCT identifier (acronym/ first author)	recruitment (out of expected) (randomization strategy)	control arm	median days before randomization	baseline recipient 8-point WHO score* (21)	transfused CCP volume (ml)	median nAb titer in CCP units	median pretransfusion nAb titer in recipient	main outcomes reported in abstract or conclusions	likely reason(s) for failure	signals of efficacy	ref
NCT04479163 (Libster)	160 (out of 210) (1:1)	normal saline	39.6 hours (from symptoms; and > 65 yrs)	2	250	n.a.	n.a.	progression to severe COVID-19 halved at day 30	no failure	-	(49)
BKH-CT-012	49 (1:1)	BSC	< 3 (from RCU admission)	5	400	n.a.	n.a.	duration of infection reduced by 4 days; mortality 1/21 in CCP arm vs 8/28	no failure	-	(91)
CTRI/2020/05/025209 (Raj)	80 (1:1)	BSC	4.2 (from hospital admission)	5	200+200	n.a.	n.a.	immediate mitigation of hypoxia, reduction in hospital stay as well as survival benefit was recorded in severe COVID-19 patients with ARDS aged less than 67 years	no failure	-	(50)

ChiCTR2000029757 (Li)	103 (out of 200) (1:1)	BSC	30 (from symptoms)	5-6	200	n.a.	n.a.	no significant difference in 28-day mortality (15.7% vs 24.0%) or time from randomization to day-28 discharge (51.0% vs 36.0%)	moderately late usage	reduced mortality at day 28 only in WHO score 5 patients (HR 2.5); negative conversion rate of viral PCR at 72 hours in 87.2% of the CCP group vs 37.5% of the control group (OR, 11.39)	(5)
NCT04342182 (ConCOVID)	86 (out of 426) (1:1)	BSC	10 from symptoms; 2 from hospitalization	5-6	300	1:320 (PRNT$_{50}$)	1:160 in 79% of recipients	no benefit at day 15	very late usage, high rate of seropositives	none	(31)
CTRI/2020/04/024775 (PLACID)	464 (1:1)	BSC	6 (from symptoms)	4-5	200+200	1:40	1:90	no benefit at day 28	moderately late usage; high rate of seropositives; extremely low nAb titre in CCP	none	(10)
NCT04345523 (ConPlas-19)	350 (1:1)	BSC	8 (from symptoms)	3 (25%) 4 (75%)	250-300	1:292	n.a.	no significant differences in primary endpoint (proportion of patients in categories 5, 6 or 7 (death) at 14 days	underpowered for mortality; primary endpoint set at just 15 days	primary endpoint significant at day +28. Trends for reduced overall mortality (p = 0.087) at day +28, expecially in aged > 75 years	(51)
NCT04375098 (Elvira-Balcells)	58 (1:1)	late CCP	6 (from symptoms)	3-4	200+200	≥ 1:160	59% < 1:160 (16% of patients enrolled before day 5 were ≥1:160 vs 60% of	no benefit at day 30 in death, mechanical ventilation or prolonged hospitalization compared to CCP administration only in case of clinical	underpowered, moderately late usage	none	(92)

							those enrolled after day 6	worsening or > 7 days after enrolment			
NCT04383535 (PlasmAr)	333 (2:1)	normal saline	8 (from symptoms)	5	500	1:300 IC_{80}	n.a.	no benefit at day 30 (16.2% vs. 31.2%)	moderately late usage	early arrivals (less than 72 hours) showed superior primary and secondary outcomes in the CCP arm (n= 28) compared to the placebo arm (n=11), but the minimal contribution of this group to the overall cohort (228 CCP and 105 placebo) made the advantage disappear in the final outcomes at day 30 (12).	(12)
NCT04356534 (AlQahtani)	40 (1:1)	BSC	n.a.	4 (95%) 5 (5%)	200+200	n.a.	n.a.	no difference in requirement for ventilation, white blood cell count, LDH, C-reactive protein (CRP), troponin, ferritin, D-dimer, procalcitonin, mortality rate at 28 days	underpowered; the CP group were a higher risk group with higher ferritin levels	primary outcome measure – ventilation – was required in 6 controls and 4 patients on CCP (risk ratio 0.67 95% CI 0.22 – 2.0, p=0.72); mean time on ventilation was 10.5 days in the control against 8.2 days in patients on CCP (p=0.81).	(93)
NCT04346446 (Bajpai)	29 (1:1)	FFP	< 3 (from symptoms)	4-5	250+250	n.a.	n.a.	no significant reduction in mortality or hospitalization	nAb measured with surrogate competitive assay (GenScript); beneficial factors in FFP used in control arm (63)	better median improvement in PaO_2/FiO_2 at 48-hours [42 vs 231] and at day 7	(94)
NCT04381	11558	BSC	9 from	4-7	275±75	n.a.	83%	no significant	late usage	the risk ratio for patients	(30)

936 (RECOVERY)	(1:1)		symptoms; 2 from hospitalization		(81% 2 units from different donors; 12%1 unit)		seronegative	difference in 28-day mortality, progression to invasive mechanical ventilation. Closed for futility		randomized within 7 days of symptom onset was 0.92 in favor of CCP versus 1.06 in patients randomized later. A reanalysis of seronegative patients (having 10% lower mortality) with a vague prior found that the likelihood of any or modest benefit was 90-74% (53).	
NCT04348656 (CONCOR-1)	940 (out of 1200) (2:1)	BSC	8 from symptoms	4-6	1-2 250-ml units	1:250	n.a.	closed for futility (even in the subgroup transfused within 3 days from diagnosis) in intubation or death by day 30	late usage (hypoxemic), sicker CCP arm (more abnormal CXR, more in ICU), varyng standard of care across 72 centres in 3 countries	each standard log increase in neutralization or ADCC independently reduced the potential harmful effect of CCP (OR=0.74), while anti-Spike IgG increased it (OR=1.53)	(81)
NCT02735707 (REMAP-CAP)	1084 (out of 7100?)	n.a.	≤ 3 from ICU hospitalization	5-6	1-2 units	n.a.	n.a.	no significant difference in organ support-free days at day 21 or in-hospital mortality. Closed for futility	very late usage	none	(95, 96)
NCT04355767 (C3PO)	511 (out of 900) with at least 1 risk factor associated with severe COVID-19	BSC	4 from symptoms, presented to the emergency department	2-3	1 250-ml unit	1:641 ID_{50}	n.a.	nonsignificant difference in risk difference (1.9%). Outcomes regarding worst illness severity and hospital-free days were similar in the	'all cause' outcome instead of COVID-19-related outcome; centralized CCP supply to distant sites likely affected by different SARS-CoV-2 variants (35) (since only 4 of the 48 centers were in	9.4% reduction in primary event endpoint in CCP group, which rises to 20% after exclusion on patients admitted on the index visit	(23, 98)

								two groups	Illinois or Colorado, most CCP usage had to be from remote sources); immunosuppressed individuals were nearly twice as common in the treatment group (12.8% vs 6.7%); designed to detect an absolute risk difference of 10% in disease progression (97)		
NCT04359810 (O'Donnell)	223 (2:1)	FFP	9 from symptoms	5-7	1 200-250 ml unit	1:160	n.a.	at 28 days, no significant improvement in clinical status	very late usage; beneficial factors in FFP used in control arm (63)	lower mortality (12.6% vs. 24.6%) compared to nonconvalescent plasma	(11)
NCT04381858 (Gonzalez)	190 (2:1)	IVIg	12 from symptoms	6-7 (85%)	2 200-ml units 24 hours apart	n.a. (29.5% received at least 1 unit of CCP with antibodies)	n.a.	no difference in mortality at day 28	very late usage; beneficial factors of IVIg used in control arm	none	(99)
NCT04344535 (Bennett-Guerrero)	74 (out of 500) (4:1)	FFP	9 from symptoms, 4 from hospitalization	n.a.	2 200-ml units	1:526	n.a.	no difference in ventilator-free days or mortality (27% vs. 33% (at day 28	very late usage; beneficial factors in FFP used in control arm (63)	all-cause mortality through 90 days was numerically lower in the CCP versus standard plasma groups (27% vs 33%; p = 0.63)	(90)
NCT04433910 (CAPSID)	105 (1:1)	BSC	7 from symptoms	4-7	3 units from same donor over 5	1:160 ($PRNT_{50}$)	1:160 ($PRNT_{50}$)	not significant difference in the primary outcome (dichotomous composite outcome	moderately late usage	median time to clinical improvement was 26 days in the CCP group and 66 days in the control group (p=0.27). Median time to	(34)

					days (850 ml)			of survival and no longer fulfilling criteria for severe COVID-19) and secondary outcomes			discharge from hospital was 31 days (IQR 16-n.r.) in the CCP and 51 days (IQR 20-n.r.) in the control group (p=0.24). In the subgroup that received a higher cumulative amount of nAbs the primary outcome occurred in 56.0% (versus 32.1%), with a shorter interval to clinical improvement, shorter time to hospital discharge and better survival compared to the control group	
NCT04547660 (PLACOVID)	160 (1:1)	BSC	10 from symptom	37% 5-6 66% 7	2 300-ml aliquots 2 days apart,	n.a.	>1:80 in 83%	no difference in 28-day mortality, days alive, days free of respiratory support, duration of invasive ventilatory support, inflammatory and other laboratorial markers values on days 3, 7 and 14	very late usage	none		(100)
NCT04429854 (DAWN-plasma)	320 (2:1)	BSC	7 from symptoms	3-5	2 200-250 ml aliquots within 12 hours followe	n.a.	≥1:320	no significant improvement proportion of patients that require mechanical ventilation or have died at day 15 or 30	late usage	none		(101, 102)

				d by 2 units within 36 hours						
NCT04393727 (TSUNAMI)	417 (1:1)	BSC	<10 from onset of symptoms	4-5	2 200-ml aliquots	each unit > 1:160)	n.a.	no statistically significant improvement in progression to ventilatory support or death	late usage	trends favoring CCP in basally seronegatives, P/F > 300 mmHg (p = 0.059),	(103)

527 * 0: no clinical or virological evidence of infection; 1 : no limitations of activities; 2 : limitations of activities ; 3 : hospitalized, no oxygen therapy; 4 : oxygen
528 by mask or nasal prongs ; 5 : non-invasive ventilation or high-flow oxygen; 6 : intubation and mechanical ventilation ; 7 : ventilation + additional organ
529 support - pressors, RRT, ECMO; 8 : death.

Table 3

531 Propensity score-matched (PSM) CCP studies reported to date. DPH: days post-hospitalization. None of these studies titered nAbs in either donor or
532 recipient using NT.

time	country	patients+ controls	median days post-hospitalization	baseline recipient WHO score (21)	transfused CCP volume (ml)	statistically significant outcomes	Reason(s) for failure	ref
retrospective	Mount Sinai, NY, USA	39 + 156	4	5 (87%) 6 (10%)	250+250	on day 14 oxygen requirements worsened in 17.9% of plasma recipients versus 28.2% of controls (aOR 0.86). Survival improved in plasma recipients (aHR 0.34)	No failure	(42)
	Providence, RI, USA	64 + 177	> 2 (< 10 from onset of symptoms: median 7)	4 (70%) 5 (30%)	n.a. (2 units)	no significant differences in incidence of in-hospital mortality (12.5% and 15.8%; aHR 0.93) or overall rate of hospital discharge (RR 1.28, although increased among patients > 65-years)	Late usage	(45)
	Montefiore Medical Center, NY, USA	90 + 258	< 3 (3-7 days from onset of symptoms)	5-6 (< 24 hrs mechanical ventilation)	200	anti-S IgG titer ≥1:2,430 (median 1:47,385) recipients <65 years had 4-fold lower mortality and 4-fold lower deterioration in oxygenation or mortality at day 28	No failure	(41)
	Washington, USA	263+263	< 14	n.a.	245 (median)	reduced 7-day (9.1 vs. 19.8%) and 14-day mortality (14.8 vs. 23.6%), but not 28-day mortality (P = 0.06), and longer hospital stay	Late usage; control cohort was treated, on average, 29 days prior to the CCP cohort	(47)
	USA (176 HCA Healthcare-affiliated community hospitals)	3774 + 10687	< 3 vs. 4-7	n.a.	n.a.	lower mortality (aHR = 0.71) and faster recovery. CCP within 3 days after admission, but not 4-7 days, was associated with a significant reduction in mortality risk (aHR = 0.53). CCP serology level was inversely associated with mortality when controlling for interaction with days to transfusion (HR = 0.998) but was not significant in a univariable analysis	No failure	(48)
	China	163 + 163	23	n.a.	300	hospital stay in CCP group was significantly longer than matched control group (P < 0.0001).	Very late usage; more advanced disease in the CCP group (23 days vs.	(104)

							15 since hospital admission	
	Greece	59 + 59	7	≥ 4	200-233 ml (days 1, 3, and 5)	significantly reduced risk of death [HR: 0.04, 3.4% vs. 13.6%], significantly better overall survival by Kaplan-Meir analysis, and increased probability of extubation [OR: 30.3]. Higher levels of antibodies (as measured with Euroimmun or pseudoVNT) in CCP were independently associated with significantly reduced risk of death.	No failure	(105)
	New Haven, USA	132 + 2551	< 6 vs > 6 days	moderate to severe		early CCP recipients, of whom 31 (40%) were on mechanical ventilation, had lower 14-day (15% vs 23%) and 30-day (38% vs 49%) mortality compared to a matched unexposed cohort, with nearly 50% lower likelihood of in-hospital mortality (HR 0.52). Early plasma recipients had more days alive and ventilator-free at 30 days (+3.3 days) and improved WHO scores at 7 days (-0.8) and hospital discharge (-0.9) compared to the matched unexposed cohort	No failure	(46)
	USA	143+823 (hematogical cancer)	n.a.	n.a.	n.a.	improved 30-day mortality (HR, 0.52; 95% CI, 0.29-0.92). Among the 338 patients admitted to the ICU, mortality was significantly lower in CCP recipients compared with nonrecipients (HR 0.40). Among the 227 patients who required mechanical ventilatory support, mortality was significantly lower in CCP recipients compared with nonrecipients (HR 0.32).	No failure	(76)
prospective	Houston, USA	136 + 251	n.a.	3 (9%) 4 (63%) 5 (18%) 6 (10%)	300 (1-2 units)	reduction in mortality within 28 days, specifically in patients transfused < 72 hours of admission with CCP with an anti-RBD titer ≥1:1350 (i.e., ~80% probability of a live virus *in vitro* neutralization titer of ≥1:160 (106))	No failure	(43)
		341 + 594	n.a.	7 (1%)	300 (1-2 units)	reduced 28-day (aHR=2.09 for controls) and 60-day (5.7% vs. 10.7%; aHR=1.82 for controls) mortality in those transfused with anti-RBD ≥1:1350 within 72 hours	No failure	(44)

					post-hospitalization. Optimal window of 44 hours to maximize benefit in 60 days mortality (4% vs 12.3%). 91% received CCP with an anti-RBD titer ≥1:1350. median S/CO ratio =24 using Ortho Vitros.			
	Poland	102 + 102	n.a.	n.a.	n.a.	lower mortality rate (13.7% vs. 34.3%; OR=0.25) related to time of first administration (12.2% at day 5, 21.5% at day 10), no significant differences in ICU stay, ventilator time, and hospitalization time. Earlier administration resulted in a ventilator being needed for a shorter length of time ($r = 0.41$)	No falure	(107)
	Colorado (16 hospitals)	188 + 188	n.a.	n.a.	1 unit if < 90 kg; 2 units if > 90kg	increased length of hospital stay in CCP-treated patients and no change in inpatient mortality compared to controls. In subgroup analysis of CCP-treated patients within 3 or 7 days of admission, there was no difference in length of hospitalization and inpatient mortality.	Covariate matching not achieved for subgroup receiving CCP < 3 days	(108)

Table 4

Summary of completed but not yet reported or ongoing RCTs of CCP, as registered on ClinicalTrials.gov as of August 26, 2021. BSC: best supportive care; FFP: fresh frozen plasma. Several studies were withdrawn (NCT04377568).

Status	NCT number	Patient subtype	Control arm	Study design	Number planned to enrol	Study start	Locations
Completed	NCT04332835	severe (SOFA <6)	BSC	single masking (outcomes assessor)	92	August 8, 2020	Colombia
	NCT04349410	any	10 different arms	single masking (investigator)	1800	April 11, 2020	USA
	NCT04397757	ordinal scale 5 to 7	BSC	open label	80	May 30, 2020	USA
	NCT04421404	within 3 days from hospitalization or 14 from symptoms	placebo	triple masking (participant, care provider, investigator)	34		USA
	NCT04374526	pneumonia, age > 65 and PaO2/FiO2 ≥300 mmHg and comorbidities	BSC	open label	29	May 27, 2020	Italy
	NCT04358783	hospitalized requiring supplemental oxygen	BSC	quadruple masking (participant, care provider, investigator, outcomes assessor)	30	April 27, 2020	Mexico
	NCT04405310	moderate to severe requiring supplemental oxygen	albumin	double (participant, care provider)	80	May 20, 2020	Mexico
	NCT04425915	on ventilator within 3 days from onset of symptoms	BSC	open label	400	June 14, 2020	India
Active, not recruiting	NCT04539275	ventilated and within 3 days from hospitalization	masked saline placebo	triple masking (participant, care provider, investigator)	702	November 16, 2020	USA
	NCT04374487	hospitalized and severe	BSC	open label	100	May 9, 2020	India
	NCT04323800	exposed within 96 h	FFP	triple masking (participant, care provider, investigator)	500	June 10, 2020	USA

		of enrollment and 120 h of receipt of CCP					
	NCT04364737 (CONTAIN)	hospitalized and within 7 days from symptoms	placebo	double masking (participant, investigator)	300	April 17, 2020	USA
	NCT04425837	high-risk	BSC	single masking (outcomes assessor)	236	July 2020	Colombia
	NCT04395170	hospitalized	2 arms (BSC; anti-COVID-19 IVIG)	open label	75	September 2020	Colombia
	NCT04391101	severe	BSC	open label	231	June 2020	Colombia
Recruiting	NCT04516811	moderate to severe	placebo	triple masking (participant, care provider, investigator)	600	September 21, 2020	South Africa
	NCT04388410	hospitalized, severe disease or risk for severe diseases	BSC	quadruple masking (participant, care provider, investigator, outcomes assessor)	410	August 25, 2020	Mexico
	NCT04385043	severe Infection	BSC	open label	400	May 1, 2020	Italy
	NCT04380935	acute respiratory distress syndrome	BSC	open label	60	May 18, 2020	Indonesia
	NCT04362176 (PassItOn)	hospitalized adults	placebo	triple masking (participant, care provider, outcomes assessor)	1000	April 24, 2020	USA
	NCT04390503	exposed within 7 days or mild symptoms within 5 days	albumin	double masking (participant, outcome assessor)	150	March 12, 2021	USA
	NCT04376034	severe or life-threatening	BSC	open label	240	March 30, 2021	USA
	NCT04373460	outpatients within 8 days from symptoms	FFP	triple masking (participant, care provider, outcomes assessor)	1344	June 3, 2020	USA
	NCT04366245	hospitalized ventilated	BSC	open label	72	April 23, 2020	Spain
	NCT04333251	hospitalized within 7 days from symptoms	BSC	open label	115	April 1, 2020	USA
	NCT04345991	mild and within 8	BSC	open label	120	April 15, 2020	France

	NCT04372979	hospitalized within 10 days from symptoms	FFP	triple masking (participant, care provider, outcomes assessor)	80	September 14, 2020	France
		days from symptoms					
	NCT04345289	adults with pneumonia	placebo	quadruple masking (participant, care provider, investigator, outcomes assessor)	1100	May 1, 2020	Denmark

References

1. Casadevall A, Pirofski LA, Joyner MJ. 2021. The Principles of Antibody Therapy for Infectious Diseases with Relevance for COVID-19. mBio 12:e03372-20.

2. Casadevall A, Pirofski L-a. 2020. The convalescent sera option for containing COVID-19. The Journal of Clinical Investigation 130.

3. Luke TC, Casadevall A, Watowich SJ, Hoffman SL, Beigel JH, Burgess TH. 2010. Hark back: passive immunotherapy for influenza and other serious infections. Crit Care Med 38:e66-73.

4. Duan K, Liu B, Li C, Zhang H, Yu T, Qu J, Zhou M, Chen L, Meng S, Hu Y, Peng C, Yuan M, Huang J, Wang Z, Yu J, Gao X, Wang D, Yu X, Li L, Zhang J, Wu X, Li B, Xu Y, Chen W, Peng Y, Hu Y, Lin L, Liu X, Huang S, Zhou Z, Zhang L, Wang Y, Zhang Z, Deng K, Xia Z, Gong Q, Zhang W, Zheng X, Liu Y, Yang H, Zhou D, Yu D, Hou J, Shi Z, Chen S, Chen Z, Zhang X, Yang X. 2020. Effectiveness of convalescent plasma therapy in severe COVID-19 patients. Proceedings of the National Academy of Sciences doi:10.1073/pnas.2004168117:202004168.

5. Li L, Zhang W, Hu Y, Tong X, Zheng S, Yang J, Kong Y, Ren L, Wei Q, Mei H, Hu C, Tao C, Yang R, Wang J, Yu Y, Guo Y, Wu X, Xu Z, Zeng L, Xiong N, Chen L, Wang J, Man N, Liu Y, Xu H, Deng E, Zhang X, Li C, Wang C, Su S, Zhang L, Wang J, Wu Y, Liu Z. 2020. Effect of Convalescent Plasma Therapy on Time to Clinical Improvement in Patients With Severe and Life-threatening COVID-19: A Randomized Clinical Trial. JAMA 324:460-470.

6. Perotti C, Baldanti F, Bruno R, Delfante C, Seminari E, Casari S, Percivalle E, Glingani C, Musella V, Belliato M, Garuti M, Meloni F, Frigato M, Di Sabatino A, Klersy C, De Donno G, Franchini M. 2020. Mortality reduction in 46 severe Covid-19 patients treated with hyperimmune plasma. A proof of concept single arm multicenter interventional trial. Haematologica 105:2834-2840.

7. Focosi D, Anderson AO, Tang JW, Tuccori M. 2020. Convalescent Plasma Therapy for COVID-19: State of the Art. Clin Microbiol Rev 33:e00072-20.

8. Joyner MJ, Wright RS, Fairweather D, Senefeld JW, Bruno KA, Klassen SA, Carter RE, Klompas AM, Wiggins CC, Shepherd JR, Rea RF, Whelan ER, Clayburn AJ, Spiegel MR, Johnson PW, Lesser ER, Baker SE, Larson KF, Ripoll JG, Andersen KJ, Hodge DO, Kunze KL, Buras MR, Vogt MN, Herasevich V, Dennis JJ, Regimbal RJ, Bauer PR, Blair JE, van Buskirk CM, Winters JL, Stubbs JR, Paneth NS, Verdun NC, Marks P, Casadevall A. 2020. Early safety indicators of COVID-19 convalescent plasma in 5,000 patients. J Clin Invest doi:10.1172/jci140200.

9. Joyner MJ, Carter RE, Senefeld JW, Klassen SA, Mills JR, Johnson PW, Theel ES, Wiggins CC, Bruno KA, Klompas AM, Lesser ER, Kunze KL, Sexton MA, Diaz Soto JC, Baker SE, Shepherd JRA, van Helmond N, Verdun NC, Marks P, van Buskirk CM, Winters JL, Stubbs JR, Rea RF, Hodge DO, Herasevich V, Whelan ER, Clayburn AJ, Larson KF, Ripoll JG, Andersen KJ, Buras MR, Vogt MNP, Dennis JJ, Regimbal RJ, Bauer PR, Blair JE, Paneth NS, Fairweather D, Wright RS, Casadevall A. 2021. Convalescent Plasma Antibody Levels and the Risk of Death from Covid-19. N Engl J Med 384:1015-1027.

10. Agarwal A, Mukherjee A, Kumar G, Chatterjee P, Bhatnagar T, Malhotra P, Latha B, Bundas S, Kumar V, Dosi R, Khambholja JK, de Souza R, Mesipogu RR, Srivastava S, Dube S, Chaudhary K, S S, Mattuvar K SA, Rajendran V, Sundararajaperumal A, Balamanikandan P, Maheswari RSU, Jayanthi R, Ragunanthanan S, Bhandari S, Singh A, Pal A, Handa A, Rankawat G, Kargirwar K, Regi J, Rathod D, Pathrose E, Bhutaka N, Patel MH, Verma RJ, Malukani K, Patel S, Thakur A, Joshi S, Kulkarni R, Suthar NN, Shah NM, Purohit HM, Shah CK, Patel MN, Shah S, Shah SH, Memon T, Beriwala VR, et al. 2020. Convalescent plasma in the management of moderate COVID-19 in India: open-label parallel-arm phase II multicentre randomized controlled trial (PLACID Trial). BMJ 371:m3939.

11. O'Donnell MR, Grinsztejn B, Cummings MJ, Justman J, Lamb MR, Eckhardt CM, Philip NM, Cheung YK, Gupta V, João E, Pilotto JH, Diniz MP, Cardoso SW, Abrams D, Rajagopalan K, Borden S, Wolf A, Sidi LC, Vizzoni A, Veloso VG, Bitan ZC, Scotto DE, Meyer BJ, Jacobson SD,

Kantor A, Mishra N, Chauhan LV, Stone E, Dei Zotti F, La Carpia F, Hudson KE, Ferrera SA,
Schwartz J, Stotler B, Lin W-H, Wontakal S, Shaz B, Briese T, Hod EA, Spitalnik SL, Eisenberger
A, Lipkin WI. 2021. A randomized, double-blind, controlled trial of convalescent plasma in
adults with severe COVID-19. J Clin Invest doi:10.1172/JCI150646:150646.

12.   Simonovich VA, Burgos Pratx LD, Scibona P, Beruto MV, Vallone MG, Vázquez C, Savoy N,
Giunta DH, Pérez LG, Sánchez MdL, Gamarnik AV, Ojeda DS, Santoro DM, Camino PJ, Antelo
S, Rainero K, Vidiella GP, Miyazaki EA, Cornistein W, Trabadelo OA, Ross FM, Spotti M,
Funtowicz G, Scordo WE, Losso MH, Ferniot I, Pardo PE, Rodriguez E, Rucci P, Pasquali J,
Fuentes NA, Esperatti M, Speroni GA, Nannini EC, Matteaccio A, Michelangelo HG, Follmann
D, Lane HC, Belloso WH. 2020. A Randomized Trial of Convalescent Plasma in Covid-19
Severe Pneumonia. N Engl J Med 384:619-629.

13.   Pirofski LA, Casadevall A. 2020. Pathogenesis of COVID-19 from the Perspective of the
Damage-Response Framework. mBio 11.

14.   Van Rompay KKA, Olstad KJ, Sammak RL, Dutra J, Watanabe JK, Usachenko JL, Immareddy R,
Roh JW, Verma A, Lakshmanappa YS, Schmidt BA, Germanio CD, Rizvi N, Stone M, Simmons
G, Dumont LJ, Allen AM, Lockwood S, Pollard RE, de Assis RR, Yee JL, Nham PB, Ardeshir A,
Deere JD, Patterson J, Jain A, Felgner PL, Iyer SS, Hartigan-O'Connor DJ, Busch MP, Reader
JR. 2021. Early post-infection treatment of SARS-CoV-2 infected macaques with human
convalescent plasma with high neutralizing activity reduces lung inflammation.
doi:10.1101/2021.09.01.458520 %J bioRxiv:2021.09.01.458520.

15.   Focosi D, Maggi F, Franchini M, Aguzzi A, Lanza M, Mazzoni A, Menichetti F. 2021. Patient-
blood management for COVID19 convalescent plasma therapy: relevance of affinity and
donor-recipient differences in concentration of neutralizing antibodies. Clin Microbiol Infect
doi:10.1016/j.cmi.2021.04.003.

16.   Alamgir J, Abid R, Garibaldi BT, Munir N, Setoguchi S, Hong SS, Chen X, Kocis PT, Yajima M,
Alexander GC, Mehta HB, Madhira V, Ergas R, O'Brien TR, Bozzette S. 2021. Lack of
association between convalescent plasma administration and length of hospital stay: a
hospital-day stratified multi-center retrospective cohort study. medRxiv [Preprint]
doi:10.1101/2021.05.04.21256627 %J medRxiv:2021.05.04.21256627.

17.   Kocayiğit H, Demir G, Karacan A, Süner K, Tomak Y, Yaylacı S, Dheir H, Kalpakci Y, Erdem AF.
2021. Effects on mortality of early vs late administration of convalescent plasma in the
treatment of Covid-19. Transfus Apher Sci doi:10.1016/j.transci.2021.103148:103148.

18.   Focosi D, Navarro D, Maggi F, Roilides E, Antonelli G. 2021. COVID-19 infodemics: the role of
mainstream and social media. Clin Microbiol Infect doi:10.1016/j.cmi.2021.08.003.

19.   Harrell F, Lindsell C. 2021. Statistical Design and Analysis Plan for Sequential Parallel-Group
RCT for COVID-19.

20.   Borobia AM, Carcas AJ, Pérez-Olmeda M, Castaño L, Bertran MJ, García-Pérez J, Campins M,
Portolés A, González-Pérez M, García Morales MT, Arana-Arri E, Aldea M, Díez-Fuertes F,
Fuentes I, Ascaso A, Lora D, Imaz-Ayo N, Barón-Mira LE, Agustí A, Pérez-Ingidua C, Gómez de
la Cámara A, Arribas JR, Ochando J, Alcamí J, Belda-Iniesta C, Frías J. 2021. Immunogenicity
and reactogenicity of BNT162b2 booster in ChAdOx1-S-primed participants (CombiVacS): a
multicentre, open-label, randomised, controlled, phase 2 trial. Lancet 398:121-130.

21.   WHO. 2020. R & D Blueprint - novel Coronavirus COVID19 Therapeutic Trial Synopsis.
https://www.who.int/blueprint/priority-diseases/key-action/COVID-
19_Treatment_Trial_Design_Master_Protocol_synopsis_Final_18022020.pdf?ua=1.
Accessed October 1.

22.   Anonymous. 2020. A minimal common outcome measure set for COVID-19 clinical research.
Lancet Infect Dis 20:e192-e197.

23.   Korley FK, Durkalski-Mauldin V, Yeatts SD, Schulman K, Davenport RD, Dumont LJ, El Kassar
N, Foster LD, Hah JM, Jaiswal S, Kaplan A, Lowell E, McDyer JF, Quinn J, Triulzi DJ, Van
Huysen C, Stevenson VLW, Yadav K, Jones CW, Kea B, Burnett A, Reynolds JC, Greineder CF,

Haas NL, Beiser DG, Silbergleit R, Barsan W, Callaway CW. 2021. Early Convalescent Plasma
for High-Risk Outpatients with Covid-19. N Engl J Med doi:10.1056/NEJMoa2103784.

24. Epstein J, Burnouf T. 2020. Points to consider in the preparation and transfusion of COVID-19
convalescent plasma. Vox Sang 115:485-487.

25. Focosi D, Mazzetti P, Pistello M, Maggi F. 2020. Viral infection neutralization tests: a focus on
SARS-CoV-2 with implications for convalescent plasma therapy. Rev Med Virol
doi:10.1002/rmv.2170:e2170.

26. Nguyen D, Simmonds P, Steenhuis M, Wouters E, Desmecht D, Garigliany M, Romano M,
Barbezange C, Maes P, Van Holm B, Mendoza J, Oyonarte S, Fomsgaard A, Lassaunière R,
Zusinaite E, Resman Rus K, Avšič-Županc T, Reimerink JH, Brouwer F, Hoogerwerf M,
Reusken CB, Grodeland G, Le Cam S, Gallian P, Amroun A, Brisbarre N, Martinaud C, Leparc
Goffart I, Schrezenmeier H, Feys HB, van der Schoot CE, Harvala H. 2021. SARS-CoV-2
neutralising antibody testing in Europe: towards harmonisation of neutralising antibody
titres for better use of convalescent plasma and comparability of trial data. Euro Surveill 26.

27. Klingler J, Weiss S, Itri V, Liu X, Oguntuyo KY, Stevens C, Ikegame S, Hung C-T, Enyindah-
Asonye G, Amanat F, Baine I, Arinsburg S, Bandres JC, Kojic EM, Stoever J, Jurczyszak D,
Bermudez-Gonzalez M, Simon V, Nádas A, Liu S, Lee B, Krammer F, Zolla-Pazner S, Hioe CE.
2020. Role of IgM and IgA Antibodies in the Neutralization of SARS-CoV-2. medRxiv
[Preprint] doi:10.1101/2020.08.18.20177303 %J medRxiv:2020.08.18.20177303.

28. Egloff SAA, Junglen A, Restivo JSA, Wongskhaluang M, Martin C, Doshi P, Schlauch D, Fromell
G, Sears LE, Correll M, Burris HA, LeMaistre CF. 2021. Convalescent plasma associates with
reduced mortality and improved clinical trajectory in patients hospitalized with COVID-19. J
Clin Invest doi:10.1172/JCI151788.

29. Natarajan H, Crowley AR, Butler SE, Xu S, Weiner JA, Bloch EM, Littlefield K, Wieland-Alter
W, Connor RI, Wright PF, Benner SE, Bonny TS, Laeyendecker O, Sullivan D, Shoham S, Quinn
TC, Larman HB, Casadevall A, Pekosz A, Redd AD, Tobian AAR, Ackerman ME. 2021. Markers
of Polyfunctional SARS-CoV-2 Antibodies in Convalescent Plasma. mBio 12.

30. Horby PW, Estcourt L, Peto L, Emberson JR, Staplin N, Spata E, Pessoa-Amorim G, Campbell
M, Roddick A, Brunskill NE, George T, Zehnder D, Tiberi S, Aung NN, Uriel A, Widdrington J,
Koshy G, Brown T, Scott S, Baillie JK, Buch MH, Chappell LC, Day JN, Faust SN, Jaki T, Jeffery
K, Juszczak E, Lim WS, Montgomery A, Mumford A, Rowan K, Thwaites G, Mafham M,
Roberts D, Haynes R, Landray MJ. 2021. Convalescent plasma in patients admitted to
hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial.
Lancet 397:2049-2059.

31. Gharbharan A, Jordans CCE, GeurtsvanKessel C, den Hollander JG, Karim F, Mollema FPN,
Stalenhoef JE, Dofferhoff A, Ludwig I, Koster A, Hassing R-J, Bos JC, van Pottelberge GR,
Vlasveld IN, Ammerlaan HSM, Segarceanu E, Miedema J, van der Eerden M, Papageorgiou G,
te Broekhorst P, Swaneveld FH, Katsikis PD, Mueller Y, Okba NMA, Koopmans MPG,
Haagmans BL, Rokx C, Rijnders B. 2021. Effects of potent neutralizing antibodies from
convalescent plasma in patients hospitalized for severe SARS-CoV-2 infection. Nat Comm
12:3189.

32. Del Fante C, Franchini M, Baldanti F, Percivalle E, Glingani C, Marano G, Mengoli C,
Mortellaro C, Viarengo G, Perotti C, Liumbruno GM. 2020. A retrospective study assessing
the characteristics of COVID-19 convalescent plasma donors and donations. Transfusion
61:830-838.

33. Mehew J, Johnson R, Roberts D, Harvala H. 2020. Convalescent plasma for COVID-19: male
gender, older age and hospitalisation associated with high neutralising antibody levels,
England, 22 April to 12 May 2020. Euro Surveill 25:2001754.

34. Koerper S, Weiss M, Zickler D, Wiesmann T, Zacharowski K, Corman VM, Gruener B, Ernst L,
Spieth P, Lepper PM, Bentz M, Zinn S, Paul G, Kalbhenn J, Dollinger M, Rosenberger P,
Kirschning T, Thiele T, Appl T, Mayer B, Schmidt M, Drosten C, Wulf H, Kruse JM, Jungwirth B,

Seifried E, Schrezenmeier H. 2021. Results of the CAPSID randomized trial for high-dose
convalescent plasma in severe COVID-19 patients. J Clin Invest
doi:10.1172/JCI152264:2021.05.10.21256192.

35. Kunze KL, Johnson PW, van Helmond N, Senefeld JW, Petersen MM, Klassen SA, Wiggins CC,
Klompas AM, Bruno KA, Mills JR, Theel ES, Buras MR, Golafshar MA, Sexton MA, Diaz Soto JC,
Baker SE, Shepherd JRA, Verdun NC, Marks P, Paneth NS, Fairweather D, Wright RS, van
Buskirk CM, Winters JL, Stubbs JR, Senese KA, Pletsch MC, Buchholtz ZA, Rea RF, Herasevich
V, Whelan ER, Clayburn AJ, Larson KF, Ripoll JG, Andersen KJ, Lesser ER, Vogt MNP, Dennis JJ,
Regimbal RJ, Bauer PR, Blair JE, Casadevall A, Carter RE, Joyner MJ. 2021. Mortality in
individuals treated with COVID-19 convalescent plasma varies with the geographic
provenance of donors. Nat Commun 12:4864.

36. Rochman ND, Wolf YI, Faure G, Mutz P, Zhang F, Koonin EV. 2021. Ongoing global and
regional adaptive evolution of SARS-CoV-2. Proc Natl Acad Sci U S A 118.

37. Junker D, Dulovic A, Becker M, Wagner TR, Kaiser PD, Traenkle B, Rothbauer U, Kienzle K,
Bunk S, Strümper C, Häberle H, Schmauder K, Malek N, Althaus K, Koeppen M, Bitzer M,
Göpel S, Schneiderhan-Marra N. 2021. Reduced serum neutralization capacity against SARS-
CoV-2 variants in a multiplex ACE2 RBD competition assay. medRxiv [Preprint]
doi:10.1101/2021.08.20.21262328 %J medRxiv:2021.08.20.21262328.

38. Chalmers TC, Matta RJ, Smith H, Jr., Kunzler AM. 1977. Evidence favoring the use of
anticoagulants in the hospital phase of acute myocardial infarction. N Engl J Med 297:1091-
6.

39. Senefeld JW, Johnson PW, Kunze KL, van Helmond N, Klassen SA, Wiggins CC, Bruno KA,
Golafshar MA, Petersen MM, Buras MR, Klompas AM, Sexton MA, Diaz Soto JC, Baker SE,
Shepherd JRA, Verdun NC, Marks P, van Buskirk CM, Winters JL, Stubbs JR, Rea RF,
Herasevich V, Whelan ER, Clayburn AJ, Larson KF, Ripoll JG, Andersen KJ, Vogt MNP, Dennis
JJ, Regimbal RJ, Bauer PR, Blair JE, Wright K, Greenshields JT, Paneth NS, Fairweather D,
Wright RS, Casadevall A, Carter RE, Joyner MJ. 2021. Program and patient characteristics for
the United States Expanded Access Program to COVID-19 convalescent plasma. medRxiv
[Preprint] doi:10.1101/2021.04.08.21255115 %J medRxiv:2021.04.08.21255115.

40. Joyner MJ, Senefeld JW, Klassen SA, Mills JR, Johnson PW, Theel ES, Wiggins CC, Bruno KA,
Klompas AM, Lesser ER, Kunze KL, Sexton MA, Diaz Soto JC, Baker SE, Shepherd JRA, van
Helmond N, van Buskirk CM, Winters JL, Stubbs JR, Rea RF, Hodge DO, Herasevich V, Whelan
ER, Clayburn AJ, Larson KF, Ripoll JG, Andersen KJ, Buras MR, Vogt MNP, Dennis JJ, Regimbal
RJ, Bauer PR, Blair JE, Paneth NS, Fairweather D, Wright RS, Carter RE, Casadevall A. 2020.
Effect of Convalescent Plasma on Mortality among Hospitalized Patients with COVID-19:
Initial Three-Month Experience. medRxiv [Preprint] doi:10.1101/2020.08.12.20169359 %J
medRxiv:2020.08.12.20169359.

41. Yoon Ha, Bartash R, Gendlina I, Rivera J, Nakouzi A, Bortz RH, Wirchnianski AS, Paroder M,
Fehn K, Serrano-Rahman L, Babb R, Sarwar UN, Haslwanter D, Laudermilch E, Florez C,
Dieterle ME, Jangra RK, Fels JM, Tong K, Mariano MC, Vergnolle O, Georgiev GI, Herrera NG,
Malonis RJ, Quiroz JA, Morano NC, Krause GJ, Sweeney JM, Cowman K, Allen S, Annam J,
Applebaum A, Barboto D, Khokhar A, Lally BJ, Lee A, Lee M, Malaviya A, Sample R, Yang XA,
Li Y, Ruiz R, Thota R, Barnhill J, Goldstein DY, Uehlinger J, Garforth SJ, Almo SC, Lai JR, Gil MR,
et al. 2021. Treatment of Severe COVID-19 with Convalescent Plasma in the Bronx, NYC. JCI
Insight doi:10.1172/jci.insight.142270:142270.

42. Liu STH, Lin H-M, Baine I, Wajnberg A, Gumprecht JP, Rahman F, Rodriguez D, Tandon P,
Bassily-Marcus A, Bander J, Sanky C, Dupper A, Zheng A, Nguyen FT, Amanat F, Stadlbauer D,
Altman DR, Chen BK, Krammer F, Mendu DR, Firpo-Betancourt A, Levin MA, Bagiella E,
Casadevall A, Cordon-Cardo C, Jhang JS, Arinsburg SA, Reich DL, Aberg JA, Bouvier NM. 2020.
Convalescent plasma treatment of severe COVID-19: a propensity score–matched control
study. Nature Medicine 26:1708-1713.

43. Salazar E, Christensen PA, Graviss EA, Nguyen DT, Castillo B, Chen J, Lopez BV, Eagar TN, Yi X,
Zhao P, Rogers J, Shehabeldin A, Joseph D, Leveque C, Olsen RJ, Bernard DW, Gollihar J,
Musser JM. 2020. Treatment of Coronavirus Disease 2019 Patients with Convalescent
Plasma Reveals a Signal of Significantly Decreased Mortality. Am J Pathol 190:2290-2303.
44. Salazar E, Christensen PA, Graviss EA, Nguyen DT, Castillo B, Chen J, Lopez BV, Eagar TN, Yi X,
Zhao P, Rogers J, Shehabeldin A, Joseph D, Masud F, Leveque C, Olsen RJ, Bernard DW,
Gollihar J, Musser JM. 2021. Significantly Decreased Mortality in a Large Cohort of
Coronavirus Disease 2019 (COVID-19) Patients Transfused Early with Convalescent Plasma
Containing High-Titer Anti-Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2)
Spike Protein IgG. Am J Pathol 191:90-107.
45. Rogers R, Shehadeh F, Mylona EK, Rich J, Neill M, Touzard-Romo F, Geffert S, Larkin J, Bailey
JA, Lu S, Sweeney J, Mylonakis E. 2020. Convalescent plasma for patients with severe COVID-
19: a matched cohort study. Clin Infect Dis ciaa1548:Epub ahead of print.
46. Briggs N, Gormally MV, Li F, Browning SL, Treggiari MM, Morrison A, Laurent-Rolle M, Deng
Y, Hendrickson JE, Tormey CA, Desruisseaux MS. 2021. Early but not late convalescent
plasma is associated with better survival in moderate-to-severe COVID-19. PLoS One
16:e0254453.
47. Shenoy AG, Hettinger AZ, Fernandez SJ, Blumenthal J, Baez V. 2021. Early mortality benefit
with COVID-19 convalescent plasma: a matched control study. Br J Haematol
doi:10.1111/bjh.17272.
48. Arnold Egloff SA, Junglen A, Restivo JSA, Wongskhauluang M, Martin C, Doshi P, Schlauch D,
Fromell G, Sears LE, Correll M, Burris HA, LeMaistre CF. 2021. Association of Convalescent
Plasma Treatment with Reduced Mortality and Improved Clinical Trajectory in Patients
Hospitalized with COVID-19 in the Community Setting. doi:10.1101/2021.06.02.21258190
%J medRxiv:2021.06.02.21258190.
49. Libster R, Pérez Marc G, Wappner D, Coviello S, Bianchi A, Braem V, Esteban I, Caballero MT,
Wood C, Berrueta M, Rondan A, Lescano G, Cruz P, Ritou Y, Fernández Viña V, Álvarez Paggi
D, Esperante S, Ferreti A, Ofman G, Ciganda Á, Rodriguez R, Lantos J, Valentini R, Itcovici N,
Hintze A, Oyarvide ML, Etchegaray C, Neira A, Name I, Alfonso J, López Castelo R, Caruso G,
Rapelius S, Alvez F, Etchenique F, Dimase F, Alvarez D, Aranda SS, Sánchez Yanotti C, De Luca
J, Jares Baglivo S, Laudanno S, Nowogrodzki F, Larrea R, Silveyra M, Leberzstein G, Debonis A,
Molinos J, González M, Perez E, et al. 2021. Early High-Titer Plasma Therapy to Prevent
Severe Covid-19 in Older Adults. N Engl J Med 384:610-618.
50. Ray Y, Paul SR, Bandopadhyay P, D'Rozario R, Sarif J, Lahiri A, Bhowmik D, Vasudevan JS,
Maurya R, Kanakan A, Sharma S, Kumar M, Singh P, Roy R, Chaudhury K, Maiti R, Bagchi S,
Maiti A, Perwez MM, Mondal A, Tewari A, Mandal S, Roy A, Saha M, Biswas D, Maiti C,
Chakraborty S, Sarkar BS, Haldar A, Saha B, Sengupta S, Pandey R, Chatterjee S, Bhattacharya
P, Paul S, Ganguly D. 2020. Clinical and immunological benefits of convalescent plasma
therapy in severe COVID-19: insights from a single center open label randomised control
trial.   medRxiv   [Preprint]   doi:10.1101/2020.11.25.20237883   %J
medRxiv:2020.11.25.20237883.
51. Avendano-Sola C, Ramos-Martinez A, Munez-Rubio E, Ruiz-Antoran B, Malo de Molina R,
Torres F, Fernandez-Cruz A, Callejas-Diaz A, Calderon J, Payares-Herrera C, Salcedo I, Romera
I, Lora-Tamayo J, Mancheno-Losa M, Paciello ML, Villegas C, Estrada V, Saez-Serrano I,
Porras-Leal ML, Jarilla-Fernandez MdC, Pano-Pardo JR, Moreno-Chulilla JA, Arrieta-Aldea I,
Bosch A, Belhassen-Garcia M, Lopez-Villar O, Ramos-Garrido A, Blanco L, Madrigal ME,
Contreras E, Muniz-Diaz E, Domingo-Morera JM, Casas-Flecha I, Perez-Olmeda M, Garcia-
Perez J, Alcami J, Bueno JL, Duarte RF. 2021. A multicenter randomized open-label clinical
trial for convalescent plasma in patients hospitalized with COVID-19 pneumonia. J Clin Invest
doi:10.1172/JCI152740:152740.

52.   Janiaud P, Axfors C, Schmitt AM, Gloy V, Ebrahimi F, Hepprich M, Smith ER, Haber NA,
Khanna N, Moher D, Goodman SN, Ioannidis JPA, Hemkens LG. 2021. Association of
Convalescent Plasma Treatment With Clinical Outcomes in Patients With COVID-19: A
Systematic Review and Meta-analysis. JAMA 325:1185-1195.
53.   Hamilton FW, Lee TC, Arnold DT, Lilford RJ, Hemming K. 2021. Is convalescent plasma futile
in COVID-19? A Bayesian re-analysis of the RECOVERY randomised controlled trial. Int J
Infect Dis S1201-9712:00523-3.
54.   Klassen SA, Senefeld JW, Johnson PW, Carter RE, Wiggins CC, Shoham S, Grossman BJ,
Henderson JP, Musser J, Salazar E, Hartman WR, Bouvier NM, Liu STH, Pirofski LA, Baker SE,
van Helmond N, Wright RS, Fairweather D, Bruno KA, Wang Z, Paneth NS, Casadevall A,
Joyner MJ. 2021. The Effect of Convalescent Plasma Therapy on Mortality Among Patients
With COVID-19: Systematic Review and Meta-analysis. Mayo Clin Proc 96:1262-1275.
55.   Piechotta V, Iannizzi C, Chai KL, Valk SJ, Kimber C, Dorando E, Monsef I, Wood EM, Lamikanra
AA, Roberts DJ, McQuilten Z, So-Osman C, Estcourt LJ, Skoetz N. 2021. Convalescent plasma
or hyperimmune immunoglobulin for people with COVID-19: a living systematic review.
Cochrane Database Syst Rev 5:Cd013600.
56.   Cruciani M, Bongiovanni G, Franchini M. 2021. High-titer convalescent plasma therapy for
coronavirus disease 2019 and mortality. Transfusion 61:1988-1990.
57.   Klassen SA, Senefeld JW, Senese KA, Johnson PW, Wiggins CC, Baker SE, van Helmond N,
Bruno KA, Pirofski LA, Shoham S, Grossman BJ, Henderson JP, Wright RS, Fairweather D,
Paneth NS, Carter RE, Casadevall A, Joyner MJ. 2021. Convalescent Plasma Therapy for
COVID-19: A Graphical Mosaic of the Worldwide Evidence. Front Med (Lausanne) 8:684151.
58.   Casadevall A, Grossman BJ, Henderson JP, Joyner MJ, Paneth NS, Pirofski LA, Shoham S.
2021.   Please   Reevaluate   the   Data   on   Convalescent   Plasma   for   COVID-19.
https://www.medpagetoday.com/opinion/second-opinions/94256?xid=nl_mpt_DHE_2021-
08-
30&eun=g1783322d0r&utm_source=Sailthru&utm_medium=email&utm_campaign=Daily%
20Headlines%20Top%20Cat%20HeC%20%202021-08-30&utm_term=NL_Daily_DHE_dual-
gmail-definition. Accessed August 31, 2021.
59.   de Candia P, Prattichizzo F, Garavelli S, La Grotta R, De Rosa A, Pontarelli A, Parrella R,
Ceriello A, Matarese G. 2021. Effect of time and titer in convalescent plasma therapy for
COVID-19. iScience doi:10.1016/j.isci.2021.102898:102898.
60.   Kloypan C, Saesong M, Sangsuemoon J, Chantharit P, Mongkhon P. 2021. Convalescent
Plasma for COVID-19: A Meta-analysis of Clinical Trials and Real-World Evidence. Eur J Clin
Invest doi:10.1111/eci.13663:e13663.
61.   Franchini M, Focosi D, Corsini F, Cruciani M. 2021. Safety and efficacy of convalescent
plasma in COVID-19: an overview of systematic reviews. Diagnostics.
62.   Anonymous. 2021. Continuous Monitoring of Pooled International Trials of Convalescent
Plasma   for   COVID-19   Hospitalized   Patients.   https://med.nyu.edu/departments-
institutes/population-health/divisions-sections-centers/biostatistics/research/continuous-
monitoring-pooled-international-trials-convalescent-plasma-covid19-hospitalized-patients.
Accessed August 31, 2021.
63.   Focosi D, Franchini M, Pirofski L-a, Burnouf T, Fairweather D, Joyner MJ, Casadevall A. 2021.
COVID-19 Convalescent Plasma Is More than Neutralizing Antibodies: A Narrative Review of
Potential Beneficial and Detrimental Co-Factors. Viruses 13:1594.
64.   Franchini M, Cruciani M. 2021. How Safe Is COVID-19 Convalescent Plasma? Mayo Clin Proc
96:2279-2281.
65.   Joyner MJ, Bruno KA, Klassen SA, Kunze KL, Johnson PW, Lesser ER, Wiggins CC, Senefeld JW,
Klompas AM, Hodge DO, Shepherd JR, Rea RF, Whelan ER, Clayburn AJ, Spiegel MR, Baker
SE, Larson KF, Ripoll JG, Andersen KJ, Buras MR, Vogt MN, Herasevich V, Dennis JJ, Regimbal
RJ, Bauer PR, Blair JE, van buskirk CM, Winters JL, Stubbs JR, van Helmond N, Butterfield BP,

Sexton M, Diaz Soto J, Paneth NS, Verdun NC, Marks P, Casadevall A, Fairweather D, Carter
RE, Wright RS. 2020. Safety Update: COVID-19 Convalescent Plasma in 20,000 Hospitalized
Patients. Mayo Clinic Proceedings 95:1888-95.

66. Lee WS, Wheatley AK, Kent SJ, DeKosky BJ. 2020. Antibody-dependent enhancement and
SARS-CoV-2 vaccines and therapies. Nature Microbiology 5:1185-1191.

67. Focosi D, Franchini M. 2021. Potential use of convalescent plasma for SARS-CoV-2
prophylaxis and treatment in immunocompromised and vulnerable populations. Expert Rev
Vaccines doi:10.1080/14760584.2021.1932475:1-8.

68. Vickers MA, Sariol A, Leon J, Ehlers A, Locher AV, Dubay KA, Collins L, Voss D, Odle AE, Holida
M, Merrill AE, Perlman S, Knudson CM. 2021. Exponential increase in neutralizing and spike
specific antibodies following vaccination of COVID-19 convalescent plasma donors.
Transfusion doi:10.1111/trf.16401.

69. Anonymous. 2021. CoVIg-19 Plasma Alliance Announces Topline Results from NIH-
Sponsored Clinical Trial of Investigational COVID-19 Hyperimmune Globulin Medicine.
https://www.takeda.com/newsroom/newsreleases/2021/covig-19-plasma-alliance-
announces-topline-results-from-nih-sponsored-clinical-trial-of-investigational-covid-19-
hyperimmune-globulin-medicine/. Accessed April 2, 2021.

70. Anonymous. 2020. ACTIV-3/TICO LY-CoV555 Study Group. A Neutralizing Monoclonal
Antibody for Hospitalized Patients with Covid-19. N Engl J Med 384:905-914.

71. Dougan M, Nirula A, Azizad M, Mocherla B, Gottlieb RL, Chen P, Hebert C, Perry R, Boscia J,
Heller B, Morris J, Crystal C, Igbinadolor A, Huhn G, Cardona J, Shawa I, Kumar P, Adams AC,
Van Naarden J, Custer KL, Durante M, Oakley G, Schade AE, Holzer TR, Ebert PJ, Higgs RE,
Kallewaard NL, Sabo J, Patel DR, Dabora MC, Klekotka P, Shen L, Skovronsky DM. 2021.
Bamlanivimab plus Etesevimab in Mild or Moderate Covid-19. NEJM
doi:10.1056/NEJMoa2102685.

72. Casadevall A, Dragotakes Q, Johnson PW, Senefeld JW, Klassen SS, Wright SR, Joyner MJ,
Paneth N, Carter R. 2021. Convalescent Plasma Use in the United States was inversely
correlated with COVID-19 Mortality: Did Convalescent Plasma Hesitancy cost lives? Elife
4:e69866.

73. Focosi D, Franchini M. 2021. COVID-19 convalescent plasma therapy: hit fast, hit hard! . Vox
Sang.

74. Focosi D, Franchini M. 2021. COVID-19 neutralizing antibody-based therapies in humoral
immune deficiencies: A narrative review. Transfus Apher Sci
doi:10.1016/j.transci.2021.103071:103071.

75. Senefeld JW, Klassen SA, Ford SK, Senese KA, Wiggins CC, Bostrom BC, Thompson MA, Baker
SE, Nicholson WT, Johnson PW, Carter RE, Henderson JP, Hartman WR, Pirofski LA, Wright
RS, Fairweather L, Bruno KA, Paneth NS, Casadevall A, Joyner MJ. 2021. Use of convalescent
plasma in COVID-19 patients with immunosuppression. Transfusion 61:2503-2511.

76. Thompson MA, Henderson JP, Shah PK, Rubinstein SM, Joyner MJ, Choueiri TK, Flora DB,
Griffiths EA, Gulati AP, Hwang C, Koshkin VS, Papadopoulos EB, Robilotti EV, Su CT, Wulff-
Burchfield EM, Xie Z, Yu PP, Mishra S, Senefeld JW, Shah DP, Warner JL. 2021. Association of
Convalescent Plasma Therapy With Survival in Patients With Hematologic Cancers and
COVID-19. JAMA Oncol doi:10.1001/jamaoncol.2021.1799.

77. Casadevall A, Henderson J, Joyner M, Pirofski L-a. 2021. SARS-Cov2 variants and
convalescent plasma: reality, fallacies, and opportunities. The Journal of Clinical
Investigation doi:10.1172/JCI148832.

78. Moscato G, Mazzetti P, Lucenteforte E, Rosellini A, Cara A, Quaranta P, Mainardi V, Villa P,
Focosi D, Lanza M, Bianco I, Mazzoni A, Falcone M, Menichetti F, Maggi F, Lai M, Freer G,
Pistello M. 2021. Assessment of automated high-throughput serological assays for prediction
of high-titer SARS-CoV-2 neutralizing antibody. Journal of Clinical Virology Plus 1:100016.

79.   Jahrsdörfer B, Kroschel J, Ludwig C, Corman VM, Schwarz T, Körper S, Rojewski M, Lotfi R,
893   Weinstock C, Drosten C, Seifried E, Stamminger T, Groß HJ, Schrezenmeier H. 2021.
894   Independent Side-by-Side Validation and Comparison of 4 Serological Platforms for SARS-
895   CoV-2 Antibody Testing. J Infect Dis 223:796-801.
896   80.   Di Germanio C, Simmons G, Kelly K, Martinelli R, Darst O, Azimpouran M, Stone M, Hazegh K,
897   Grebe E, Zhang S, Ma P, Orzechowski M, Gomez JE, Livny J, Hung DT, Vassallo R, Busch MP,
898   Dumont LJ. 2021. SARS-CoV-2 Antibody persistence in COVID-19 convalescent plasma
899   donors: Dependency on assay format and applicability to serosurveillance. Transfusion
900   doi:10.1111/trf.16555:2021.03.24.21254260.
901   81.   Bégin P, Callum J, Jamula E, Cook R, Heddle NM, Tinmouth A, Zeller MP, Beaudoin-Bussières
902   G, Amorim L, Bazin R, Loftsgard KC, Carl R, Chassé M, Cushing MM, Daneman N, Devine DV,
903   Dumaresq J, Fergusson DA, Gabe C, Glesby MJ, Li N, Liu Y, McGeer A, Robitaille N, Sachais BS,
904   Scales DC, Schwartz L, Shehata N, Turgeon AF, Wood H, Zarychanski R, Finzi A, Marceau D,
905   Huang A, Carr H, Lin Y, Lall R, Graham C, Arsenault C, Sales V, Sidhu D, Semret M, Hamm C,
906   Arhanchiague E, Solh Z, Srour N, Soliman K, Yee C, Laroche V, Nahirniak S, et al. 2021.
907   Convalescent plasma for hospitalized patients with COVID-19: an open-label, randomized
908   controlled trial. Nature Medicine doi:10.1038/s41591-021-01488-2.
909   82.   Misset B, Hoste E, Donneau A-F, Grimaldi D, Meyfroidt G, Moutschen M, Compernolle V,
910   Gothot A, Desmecht D, Garigliany M, Najdovski T, Laterre P-F. 2020. A multicenter
911   randomized trial to assess the efficacy of CONvalescent plasma therapy in patients with
912   Invasive COVID-19 and acute respiratory failure treated with mechanical ventilation: the
913   CONFIDENT trial protocol. BMC Pulmonary Medicine 20:317.
914   83.   Mariën J, Ceulemans A, Michiels J, Heyndrickx L, Kerkhof K, Foque N, Widdowson MA,
915   Mortgat L, Duysburgh E, Desombere I, Jansens H, Van Esbroeck M, Ariën KK. 2021.
916   Evaluating SARS-CoV-2 spike and nucleocapsid proteins as targets for antibody detection in
917   severe and mild COVID-19 cases using a Luminex bead-based assay. J Virol Methods
918   288:114025.
919   84.   Betrains A, Godinas L, Woei AJF, Rosseels W, Van Herck Y, Lorent N, Dierickx D, Compernolle
920   V, Meyfroidt G, Vanderbeke L, Vergote V, Lagrou K, Verhamme P, Wauters J, Vermeersch P,
921   Devos T, Maes P, Vanderschueren S. 2021. Convalescent plasma treatment of persistent
922   severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) infection in patients with
923   lymphoma with impaired humoral immunity and lack of neutralising antibodies. Br J
924   Haematol 192:1100-1105.
925   85.   Okba NMA, Müller MA, Li W, Wang C, GeurtsvanKessel CH, Corman VM, Lamers MM,
926   Sikkema RS, de Bruin E, Chandler FD, Yazdanpanah Y, Le Hingrat Q, Descamps D, Houhou-
927   Fidouh N, Reusken C, Bosch BJ, Drosten C, Koopmans MPG, Haagmans BL. 2020. Severe
928   Acute Respiratory Syndrome Coronavirus 2-Specific Antibody Responses in Coronavirus
929   Disease Patients. Emerg Infect Dis 26:1478-1488.
930   86.   Amanat F, White KM, Miorin L, Strohmeier S, McMahon M, Meade P, Liu W-C, Albrecht RA,
931   Simon V, Martinez-Sobrido L, Moran T, García-Sastre A, Krammer F. 2020. An In Vitro
932   Microneutralization Assay for SARS-CoV-2 Serology and Drug Screening. 58:e108.
933   87.   Oguntuyo KY, Stevens CS, Hung C-T, Ikegame S, Acklin JA, Kowdle SS, Carmichael JC, Chiu H-
934   p, Azarm KD, Haas GD, Amanat F, Klingler J, Baine I, Arinsburg S, Bandres JC, Siddiquey MN,
935   Schilke RM, Woolard MD, Zhang H, Duty AJ, Kraus TA, Moran TM, Tortorella D, Lim JK,
936   Gamarnik AV, Hioe CE, Zolla-Pazner S, Ivanov SS, Kamil JP, Krammer F, Lee B. 2020.
937   Quantifying absolute neutralization titers against SARS-CoV-2 by a standardized virus
938   neutralization assay allows for cross-cohort comparisons of COVID-19 sera. mBio 12.
939   88.   Anonymous. Wu, Fan and Wang, Aojie and Liu, Mei and Wang, Qimin and Chen, Jun and Xia,
940   Shuai and Ling, Yun and Zhang, Yuling and Xun, Jingna and Lu, Lu and Jiang, Shibo and Lu,
941   Hongzhou and Wen, Yumei and Huang, Jinghe, Neutralizing Antibody Responses to SARS-

CoV-2 in a COVID-19 Recovered Patient Cohort and Their Implications (3/28/2020). Available
at SSRN: https://ssrn.com/abstract=3566211 or http://dx.doi.org/10.2139/ssrn.3566211.

89.　Beltrán-Pavez C, Riquelme-Barrios S, Oyarzún-Arrau A, Gaete-Argel A, González-Stegmaier R,
Cereceda-Solis K, Aguirre A, Travisany D, Palma-Vejares R, Barriga GP, Gaggero A, Martínez-
Valdebenito C, Corre NL, Ferrés M, Balcells ME, Fernandez J, Ramírez E, Villarroel F, Valiente-
Echeverría F, Soto-Rifo R. 2021. Insights into neutralizing antibody responses in individuals
exposed to SARS-CoV-2 in Chile. 7:eabe6855.

90.　Bennett-Guerrero E, Romeiser JL, Talbot LR, Ahmed T, Mamone LJ, Singh SM, Hearing JC,
Salman H, Holiprosad DD, Freedenberg AT, Carter JA, Browne NJ, Cosgrove ME, Shevik ME,
Generale LM, Andrew MA, Nachman S, Fries BC. 2021. Severe Acute Respiratory Syndrome
Coronavirus 2 Convalescent Plasma Versus Standard Plasma in Coronavirus Disease 2019
Infected Hospitalized Patients in New York: A Double-Blind Randomized Trial. Crit Care Med
doi:10.1097/ccm.0000000000005066.

91.　Rasheed AM, Fatak DF, Hashim HA, Maulood MF, Kabah KK, Almusawi YA, Abdulamir AS.
2020. The therapeutic potential of convalescent plasma therapy on treating critically-ill
COVID-19 patients residing in respiratory care units in hospitals in Baghdad, Iraq. Infez Med
28:357-366.

92.　Balcells ME, Rojas L, Le Corre N, Martínez-Valdebenito C, Ceballos ME, Ferrés M, Chang M,
Vizcaya C, Mondaca S, Huete Á, Castro R, Sarmiento M, Villarroel L, Pizarro A, Ross P,
Santander J, Lara B, Ferrada M, Vargas-Salas S, Beltrán-Pavez C, Soto-Rifo R, Valiente-
Echeverría F, Caglevic C, Mahave M, Selman C, Gazitúa R, Briones JL, Villarroel-Espindola F,
Balmaceda C, Espinoza MA, Pereira J, Nervi B. 2020. Early Anti-SARS-CoV-2 Convalescent
Plasma in Patients Admitted for COVID-19: A Randomized Phase II Clinical Trial. PLoS
Medicine doi:10.1371/journal.pmed.1003415.

93.　AlQahtani M, Abdulrahman A, Almadani A, Alali SY, Al Zamrooni AM, Hejab AH, Conroy RM,
Wasif P, Atkin SL, Otoom S, Abduljalil M. 2020. Randomized controlled trial of convalescent
plasma therapy against standard therapy in patients with severe COVID-19 disease. Sci Rep
11:9927.

94.　Bajpai M, Kumar S, Maheshwari A, Chhabra K, kale P, Gupta A, Narayanan A, Gupta E,
Trehanpati N, Bihari C, Agarwal R, Gupta K, Gupta Uk, Bhardwaj A, Kumar G, Islam M, Singh
R, Yadav P, Maiwall R, Sarin SK. 2020. Efficacy of Convalescent Plasma Therapy compared to
Fresh Frozen Plasma in Severely ill COVID-19 Patients: A Pilot Randomized Controlled Trial.
medRxiv [Preprint] doi:10.1101/2020.10.25.20219337 %J medRxiv:2020.10.25.20219337.

95.　Angus DC, Berry S, Lewis RJ, Al-Beidh F, Arabi Y, van Bentum-Puijk W, Bhimani Z, Bonten M,
Broglio K, Brunkhorst F, Cheng AC, Chiche JD, De Jong M, Detry M, Goossens H, Gordon A,
Green C, Higgins AM, Hullegie SJ, Kruger P, Lamontagne F, Litton E, Marshall J, McGlothlin A,
McGuinness S, Mouncey P, Murthy S, Nichol A, O'Neill GK, Parke R, Parker J, Rohde G,
Rowan K, Turner A, Young P, Derde L, McArthur C, Webb SA. 2020. The REMAP-CAP
(Randomized Embedded Multifactorial Adaptive Platform for Community-acquired
Pneumonia) Study. Rationale and Design. Ann Am Thorac Soc 17:879-891.

96.　Estcourt LJ. 2021. Convalescent Plasma in Critically ill Patients with COVID-19. medRxiv
[Preprint] doi:10.1101/2021.06.11.21258760 %J medRxiv:2021.06.11.21258760.

97.　Anonymous. National COVID19 Convalescent Plasma Project. Commentary from the COVID-
19 Convalescent Plasma Project (CCPP19) Leadership Group on "Early Convalescent Plasma
for   High-Risk   Outpatients   with   Covid-19".   Accessed   online   at
https://ccpp19.org/news/review%20of%20NEJM%20US%20outpatient%20CP%20trial%208-
23.docx on August 25, 2021.

98.　Anonymous. March 2, 2021. NIH halts trial of COVID-19 convalescent plasma in emergency
department   patients   with   mild   symptoms.   https://www.nih.gov/news-events/news-
releases/nih-halts-trial-covid-19-convalescent-plasma-emergency-department-patients-
mild-symptoms. Accessed

99. Beltran Gonzalez JL, Gonzalez Gamez M, Mendoza Enciso EA, Esparza Maldonado RJ,
Hernandez Palacios D, Duenas Campos S, Ovalle Robles I, Macias Guzman MJ, Garcia Diaz AL,
Gutierrez Pena CM, Reza Escalera AL, Tiscareno Gutierrez MT, Galvan Guerra E, Dorantes
Morales MdR, Martinez Medina L, Monroy Colin VA, Arreola Guerra JM. 2021. Efficacy and
safety of convalescent plasma and intravenous immunoglobulin in critically ill COVID-19
patients. A controlled clinical trial. medRxiv [Preprint] doi:10.1101/2021.03.28.21254507 %J
medRxiv:2021.03.28.21254507.
100. Sekine L, Arns B, Fabro BR, Cipolatt MM, Machado RRG, Durigon EL, Parolo E, Pellegrini JAS,
Viana MV, Schwarz P, Lisboa TC, Dora JMS, Balsan AM, Schirmer FD, Franz JPM, da-Silveira
LM, Breunig RC, Petersen V, Sosnoski M, Mesquita NF, Volpato FCZ, Sganzerla D, Falavigna
M, Rosa RG, Zavascki AP. 2021. Convalescent plasma for COVID-19 in hospitalised patients:
an open-label, randomised clinical trial. Eur Respir J doi:10.1183/13993003.01471-2021.
101. Devos T, Van Thillo Q, Compernolle V, Najdovski T, Romano M, Dauby N, Jadot L, Leys M,
Maillart E, Loof S, Seyler L, Moonen M, Moutschen M, Van Regenmortel N, Ariën KK,
Barbezange C, Betrains A, Garigliany M, Engelen MM, Gyselinck I, Maes P, Schauwvlieghe A,
Liesenborghs L, Belmans A, Verhamme P, Meyfroidt G. 2021. Early high antibody-titre
convalescent plasma for hospitalised COVID-19 patients: DAWn-plasma. Eur Respir J
doi:10.1183/13993003.01724-2021.
102. Devos T, Geukens T, Schauwvlieghe A, Ariën KK, Barbezange C, Cleeren M, Compernolle V,
Dauby N, Desmecht D, Grimaldi D, Lambrecht BN, Luyten A, Maes P, Moutschen M, Romano
M, Seyler L, Nevessignsky MT, Vandenberghe K, van Griensven J, Verbeke G, Vlieghe E,
Yombi JC, Liesenborghs L, Verhamme P, Meyfroidt G. 2020. A randomized, multicentre,
open-label phase II proof-of-concept trial investigating the clinical efficacy and safety of the
addition of convalescent plasma to the standard of care in patients hospitalized with COVID-
19: the Donated Antibodies Working against nCoV (DAWn-Plasma) trial. Trials 21:981.
103. Menichetti F, Popoli P, Puopolo M, Spila Alegiani S, Tiseo G, Bartoloni A, De Socio G, Luchi S,
Blanc P, Puoti M, De Donno G, Toschi E, Massari M, Palmisano L, Marano G, Chiamenti M,
Martinelli L, Franchi S, Pallotto C, Suardi L, Luciani Pasqua B, Merli M, Fabiani P, Bertolucci L,
Borchi B, Modica S, Moneta S, Marchetti G, d'Arminio Monforte A, Stoppini L, Ferracchiato
N, Piconi S, Fabbri C, Beccastrini E, Saccardi R, Giacometti A, Esperti S, Pierotti P, Bernini l,
Bianco C, Benedetti S, Lanzi A, Bonfanti P, Massari M, Sani S, Saracino A, Castagna A, Trabace
l, Lanza M, Focosi D, et al. 2021. Effect of High-Titer Convalescent Plasma on Progression to
Severe Respiratory Failure or Death in Hospitalized Patients with COVID-19 Pneumonia: A
Randomized Clinical Trial. Submitted.
104. Jiang W, Li W, Xiong L, Wu Q, Wu J, He B, Shen J, Pang R, Luo T, Guo Y, Yang Y, Han Y, Dai W,
Zhu P, Xia X. 2020. Clinical efficacy of convalescent plasma therapy on treating COVID-19
patients: Evidence from matched study and a meta-analysis. Clin Transl Med 10:e259.
105. Pappa V, Bouchla A, Terpos E, Thomopoulos TP, Rosati M, Stellas D, Antoniadou A, Mentis A,
Papageorgiou SG, Politou M, Kotanidou A, Kalomenidis I, Poulakou G, Jahaj E, Korompoki E,
Grigoropoulou S, Hu X, Bear J, Karaliota S, Burns R, Pagoni M, Trontzas I, Grouzi E,
Labropoulou S, Stamoulis K, Bamias A, Tsiodras S, Felber BK, Pavlakis GN, Dimopoulos MA.
2021. A Phase II Study on the Use of Convalescent Plasma for the Treatment of Severe
COVID-19- A Propensity Score-Matched Control Analysis. Microorganisms 9:806.
106. Salazar E, Kuchipudi SV, Christensen PA, Eagar TN, Yi X, Zhao P, Jin Z, Long SW, Olsen RJ,
Chen J, Castillo B, Leveque C, Towers DM, Lavinder J, Gollihar JD, Cardona J, Ippolito GC,
Nissly RH, Bird IM, Greenawalt D, Rossi RM, Gontu A, Srinivasan S, Poojary IB, Cattadori IM,
Hudson PJ, Joselyn N, Prugar L, Huie K, Herbert A, Bernard DW, Dye J, Kapur V, Musser JM.
2020. Convalescent plasma anti-SARS-CoV-2 spike protein ectodomain and receptor-binding
domain IgG correlate with virus neutralization. J Clin Invest 130:6728-6738.
107. Tworek A, Jaroń K, Uszyńska-Kałuża B, Rydzewski A, Gil R, Deptała A, Franek E, Wójtowicz R,
Życińska K, Walecka I, Cicha M, Wierzba W, Zaczyński A, Król ZJ, Rydzewska G. 2021.

Convalescent plasma treatment is associated with lower mortality and better outcomes in
high risk COVID-19 patients - propensity score matched case-control study. Int J Infect Dis
105:209-215.
108.   Chauhan L, Pattee J, Ford J, Thomas C, Lesteberg K, Richards E, Loi M, Dumont LJ, Annen K,
Berg M, Zirbes M, Miller A, Jenkins TC, Bennett TD, Monkowski D, Boxer RS, Beckham JD.
2021. A Multi-center, Prospective, Observational-cohort controlled study of Clinical
Outcomes following COVID-19 Convalescent plasma therapy in hospitalized COVID-19
patients. medRxiv [Preprint] doi:10.1101/2021.06.14.21258910 %J
medRxiv:2021.06.14.21258910.

1053