



Implicit, Intrinsic, Extrinsic, and Host Factors Attributing the Covid-19 Pandemic. Part 1- Intrinsic Factor Nitrous Oxide Emissions: A Systematic Analysis

Nikita Thapliyal, Gaurav Joshi 0000-0002-7812-2871 and **Prashant Gahtori** ^{*} 0000-0002-2944-7856

School of Pharmacy, Graphic Era Hill University, Dehradun 248002, India

*Corresponding author: pgahtori@gehu.ac.in


NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Despite Nitrous oxide (N₂O) being the most widely used anesthetics in dental and other medical applications, it is associated with global warming and stratospheric ozone destruction. With globalization, a larger amount of N₂O emissions are articulated especially from human activities (30%, 6.7 Tg N per year), which are primarily dominated by agriculture that is even above the emissions of all oceans (26%). The synthesis of N₂O reflects the general chemistry and readily from a substrate Nitric oxide (NO) in the environment. The modeling of infectious disease dynamics covering common pathogen-transmission factors, for example intrinsic (or microbes nutrient supply) at a population level, is indeed imperative to curb the menace of any disease. Nonetheless, in areas where novel coronavirus disease (COVID-19) was at its worst, for example, Wuhan China, Mumbai India, Milan Italy, Washington USA etc., the reduction in N₂O emissions was well noticed. Nonetheless, viruses exhibit greater mobility than humans and hijack nutrients including nitrogen to complete their epidemiological cycle all due to limited sequence space of viral genomes, the high probability of genetic drift, extremely large population sizes, the high mutation and recombination rates. In consequence of drastic fall in N₂O emissions, lower human transport can not be an all alone contributor, but contrarily it may also be associated with coronavirus intrinsic factors. This prompted us to analyze freely accessible and large global data from two authenticated sources, the World Health Organization and World Bank. We hereby argue that intrinsic factor N₂O emissions fueling the COVID-19 progression significantly. Entire predictions were found consistent with the recently observed shreds of evidence. These insights enhanced scientific ability to interrogate viral epidemiology and recommended a 7-points framework covering all-natural lifestyle and dietary supplements for COVID-19 prevention before the arrival of a front-line therapeutic(s) or preventable vaccine.

Introduction

Over the years, the oxides of nitrogen have become increasingly important in contemporary science and are subjected to many reviews [1,2]. Among them three oxides, Nitrous oxide (N_2O) popularly known as laughing gas, Nitrogen dioxide (NO_2) and a free radical molecule Nitric oxide (NO) are the most abundant in the air [3]. N_2O oxidation by ozone can occur at any temperature and considered as an ozone depleting substance with a long half-life (100 to 150 years). The NO ($k \sim 10^7 \text{ lit mol}^{-1} \text{ sec}^{-1}$) is very rapidly oxidized to NO_2 by ozone. The NO_2 ($k \sim 3 \times 10^4 \text{ lit mol}^{-1} \text{ sec}^{-1}$) is still rapidly but slowly than NO oxidized to NO_3 by ozone.



The NO is produced in many tissues from L-arginine and molecular oxygen by four distinct isoforms of Nitric Oxide Synthase (NOS): (i) neuronal NOS-1 (nNOS), (ii) inducible NOS-2 (iNOS), (iii) endothelial NOS-3 (eNOS) and (iv) mitochondrial NOS (mtNOS). Nitric oxide reductases (NOR) play a predominant role in the detoxification of NO and likely to increase fitness of many Gram-positive and –negative pathogens during their associations with invertebrate and vertebrate hosts. The presence of an unpaired electron with uncharged molecule turns NO as an ideal messenger in human with a very short half-life of 2 to 30 seconds [4]. These characteristics of NO also allow its free diffusion across biomembranes and spontaneous signal transmission to perform a wide variety of physiological functions including an olfactory signal procession; platelet clotting; smooth muscle relaxation and vasodilation; blood flow regulation in the brain, heart, lung, gastrointestinal tract, and kidney; blood pressure regulation; cardiac contractility; and killing or inhibition of pathogens etc.[5,6] Despite having large physiological functions, the Global warming potential of 1 gram of N_2O is 265-298 times of same carbon dioxide for a 100-year timescale. A recent report published in Nature by Tian and coworkers highlighted that N_2O emissions from human activities such as agriculture, fuel combustion, wastewater management, and industrial processes have ballooned 30% over the past forty years [7]. Here agriculture constitutes the largest source of N_2O gas emissions, 100-, 8-, and 5-folds in comparison to transport, industry, and international bunkers respectively. Needless to say, that a well-balanced ecosystem is a cornerstone of human health and well-being[8,9]. However, research into how natural environments relates to human

health remains limited in important respects.

The COVID-19 disease surge caused by the novel corona (SARS-COV2) virus is posing enormous health challenges since its first detection in China during December 2019, and subsequently its declaration as a pandemic by the World Health Organization (WHO) on dated March 11, 2020 [10]. Recent MEDLINE records of COVID-19 accessed via PubMed shares 0.155931 million results during the writing of the manuscript as on July 13, 2021, which further emphasize several successful scientific accomplishments across its virology, biology, prevention, therapeutics, diagnostics, etc. [11] Despite these success stories, still the eradication of COVID-19 is limited to 3Ts plus P i.e. Tracing (immediate patient isolation), Testing (early diagnosis), Treatment (supportive respiratory care) and Prevention (vaccination). However, according to a recent estimate, there has been a total of 185.29153 million confirmed COVID-19 cases, resulting in 4.010834 million deaths and 0.458355 million new cases are still appearing worldwide on daily basis. [12] The COVID-19 patients have mild to severe symptoms, most of are seeming closely linked with the physiological functions of NO such as loss of olfactory sensation; platelet clotting; difficulty in breathing; multi-organ brain, heart, lung, gastrointestinal tract, and kidney failure etc. [13] In fact, recently the FDA approved an inhaled nitric oxide (iNO plus) for COVID-19 emergency use to a biotherapeutics company, Bellerophon Therapeutics. [14] In addition, a fall in N₂O gas emissions was noticed in local affected areas after the surge of the SARS-CoV-2 virus. Consequently, these real evidences convince us about the physiological role of substrate NO in COVID-19 management and huge spread of SARS-CoV-2 virus may be due to atmospheric imbalance of N₂O. Furthermore, there is considerable evidence that intrinsic, extrinsic (or environment), implicit, and host factors are key regulators of any pathogen transmission [15] and an intrinsic factor both operationally and conceptually have yet to be reached for full annihilation of COVID-19. Suppressing N₂O emissions seems the most viable pathway for attaining the great health challenges posed by COVID-19 pandemic. This prompted us to perform a wider sampling of global COVID-19 infectious cases with both nitric oxide and carbon dioxide emissions in understanding the SARS-COV2 transmission in atmosphere and human population.

Methods

A large authenticated data set with both Covid-19 cases and nitrous oxide/ carbon dioxide emissions consisting 97 countries/ territories is freely accessible, and retrieved from two reliable sources, the World Health Organization (WHO) and the World Bank (n= 97). All COVID-19 infectious cases (n= 97) were collected from five WHO situation reports- dated 09-Jul-2021, 22-Dec-2020, 13-Mar-2020, 19-Mar-2020, and 28-Mar-2020 [16]. All recent value pertaining to both nitrous oxide and carbon dioxide

emissions countries/ territories- wise is also publically accessible and supplied by the World Bank [17]. Carbon dioxide emissions listed in per capita were multiplied with country's population and finally converted into thousand metric tons. Here, 97 Countries/ Territories randomly fitted to top variable indices of correlation coefficient for unbiased cause-effect relationships (Table 1).

Table 1. Cumulative COVID-19 Infected Cases and Nitric oxide/ Carbon dioxideemissionsWorldwide

	Country/ Territory/Area	CUMULATIVE COVID-19 INFECTED CASES					Nitrous Oxide Emissions (Thousand metric tons of CO ₂ equivalent)	Carbon Dioxide Emissions (Thousand metric tons)
		09-Jul- 2021	22-Dec- 2020	13-Mar- 2020	19-Mar- 2020	28-Mar- 2020		
1	Albania	132565	52542	23	59	197	1,100	5,582
2	Algeria	143652	94781	25	72	367	12,110	157,498
3	Argentina	4593763	1531374	31	79	589	47,930	180,206
4	Armenia	226135	153825	1	84	372	1,170	5,572
5	Australia	30905	28128	140	510	3635	76,760	394,624
6	Austria	647111	334629	361	1646	7697	3,620	64,365
7	Azerbaijan	336788	199127	11	34	147	4,160	32,662
8	Bahrain	266797	90062	195	256	473	90	33,339
9	Bangladesh	1000543	499560	3	10	48	29,240	84,459
10	Belarus	424554	171579	12	46	94	12,010	59,096
11	Belgium	1092477	625928	314	1486	7284	4,460	94,800
12	Bhutan	2258	446	1	1	3	190	1,411
13	Bolivia	449687	149149	3	12	61	8,770	23,350
14	Brazil	18909037	7162978	77	291	2915	179,200	434,020
15	Brunei Darussalam	266	152	12	56	115	150	7,282
16	Bulgaria	422353	191029	7	92	293	4,050	40,682
17	Burkina Faso	13505	4954	2	26	146	9,850	4,519
18	Cameroon	80858	25849	2	10	75	62,710	9,075
19	Canada	1418632	495346	138	569	4018	43,120	584,891
20	Chile	1579591	583354	33	238	1610	6,410	88,410
21	China	119141	95716	80991	81174	82230	538,790	10,658,497
22	Colombia	4426811	1482072	9	93	491	21,720	81,446
23	Costa Rica	377091	157472	22	50	231	1,940	8,416
24	Cote d Ivoire	48693	21772	1	9	92	3,030	10,555
25	Croatia	360680	194962	16	81	586	1,790	16,651
26	Cyprus	81581	17476	6	58	162	370	7,340
27	Czech Republic	1668891	624140	116	522	2279	5,170	103,242
28	Denmark	297543	131606	674	1044	2046	4,740	33,372
29	Dominican Republic	331826	159064	5	21	581	3,390	25,642
30	Ecuador	465878	205920	17	155	1595	4,950	40,823
31	Egypt	282582	124891	67	196	536	22,320	256,045
32	Estonia	131396	21794	13	258	575	1,350	16,055
33	Finland	97651	32582	109	359	1025	4,670	44,563

34	France	5686066	2418439	2860	9043	32542	38,030	301,514
35	French Polynesia	19026	16182	1	3	30	38	781
36	Georgia	374836	208638	25	38	85	2,090	9,468
37	Germany	3734468	1494009	2369	8198	48582	32,900	717,056
38	Greece	433021	130485	98	418	966	4,450	63,405
39	Guyana	20645	6076	1	4	8	970	2,464
40	Honduras	270689	116212	2	9	67	3,210	10,093
41	Hungary	808437	302989	16	58	343	5,790	45,843
42	Iceland	6675	5621	61	250	890	400	2,128
43	India	30752950	10031223	74	151	724	253,790	2,459,313
44	Indonesia	2455912	657948	34	227	1046	95,690	589,547
45	Iran	3327526	1152072	10075	17361	32332	37,580	646,159
46	Iraq	1406289	583118	70	164	458	5,420	196,897
47	Ireland	276104	78776	70	292	2121	10,060	37,647
48	Israel	845123	371373	75	427	3460	2,360	60,385
49	Italy	4267105	1938083	15113	35713	86498	17,900	325,065
50	Jamaica	50497	12135	1	13	26	410	8,588
51	Japan	813976	195880	675	873	1499	18,010	1,109,045
52	Jordan	754927	272797	1	52	235	1,250	25,289
53	Kuwait	369227	147775	80	142	235	750	92,341
54	Latvia	137797	30297	16	71	280	1,860	7,468
55	Lebanon	546366	156570	66	133	391	740	27,573
56	Lithuania	279024	112359	3	26	358	3,530	11,262
57	Luxembourg	71879	44067	17	210	1605	300	9,596
58	Malaysia	808658	91969	129	673	2161	11,120	245,989
59	Maldives	74848	13474	8	13	13	30	1,910
60	Malta	30851	11621	9	48	139	40	1,412
61	Mexico	2558369	1301546	12	93	589	42,250	482,399
62	Mongolia	131962	953	1	5	12	13,190	21,320
63	Morocco	537253	415226	6	49	358	9,430	68,311
64	Nepal	652859	253184	1	1	3	8,330	12,476
65	Netherlands	1701911	676589	614	2051	8603	7,790	150,321
66	New Zealand	2409	1760	5	20	416	15,040	31,695
67	North Macedonia	155760	77949	7	36	219	550	7,375
68	Norway	132572	42775	489	1423	3581	3,190	38,119
69	Oman	281688	127019	18	33	152	780	77,580
70	Pakistan	967633	454673	20	241	1235	60,950	212,628
71	Panama	411226	206310	14	86	674	1,310	10,475
72	Paraguay	432801	98296	5	11	52	10,070	8,634
73	Peru	2071637	993760	22	145	580	9,360	55,947

74	Philippines	1455585	458044	52	187	803	13,640	142,241
75	Poland	2880670	1202700	49	287	1389	21,480	311,685
76	Portugal	899295	370787	41	642	4268	3,090	49,358
77	Qatar	222918	141858	262	442	562	880	94,296
78	Republic of Korea	165344	49665	7979	8413	9478	10,770	626,745
79	Republic of Moldova	257272	134578	4	36	199	1,160	12,781
80	Russian Federation	5733218	2848377	34	147	1264	58,610	1,623,065
81	Saudi Arabia	497773	360848	21	238	1104	5,750	531,565
82	Senegal	44790	17670	10	36	119	7,820	10,413
83	Slovakia	391813	151336	21	105	295	1,970	33,078
84	Slovenia	257619	105013	57	286	632	790	14,084
85	South Africa	2135246	912477	17	116	1170	18,820	444,616
86	Spain	3915313	1797236	2965	13716	64059	20,350	258,103
87	Sri Lanka	269946	36049	3	42	106	2,240	21,374
88	Sweden	1092308	367120	620	1279	3046	4,920	35,731
89	Switzerland	701473	402264	858	3010	12104	2,140	38,098
90	Thailand	317506	4331	75	212	1136	19,950	259,240
91	Togo	14176	3396	1	1	25	1,870	2,372
92	Tunisia	464914	119151	7	29	227	3,320	30,637
93	Turkey	5465094	1189947	1	191	5698	34,380	422,996
94	Ukraine	2240246	964448	3	16	311	25,310	181,678
95	United Kingdom	5022897	2004223	594	2630	14547	28,290	1,411,859
96	United States of America	33451965	17314834	1264	7087	85228	250,060	1,786,987
97	Vietnam	24810	1411	39	66	169	24,080	262,698
Correlation	Nitrous Oxide Emissions	0.6	0.5	0.7	0.7	0.6		
	Carbon Dioxide Emissions	0.2	0.2	0.9	0.9	0.6		

*Top with red, Bottom with green

Results and Discussion

We ascertained a large data set involving a total of ninety seven Countries/ Territories randomly who fulfilled the unbiased study criteria used in the assessment of role of nitrous oxide emissions over Covid-19 pandemic. There was a large variation among both N₂O emissions (variation= 4.53 X 10⁹) and cumulative COVID-19 cases (variation ranges from 2.48X10¹³ to 2.88X10⁸ dated from 28-March-2020 to 09- July-2021) across all Countries/ Territories (Table 1). Here with the large variation, we observed a good and consistent correlation ($r^2= 0.6 \pm 0.06$) between N₂O emissions and Cumulative covid-19 cases from last one year plus (Table 2). Moreover, the Z-test with a probability, P values falls below the level of significance at 0.05 (5%) (one-tail ranges from 0.0002 to 0.0034 and two-tail ranges from 0.0004 to 0.0068). This reveals that there is a significant relationship between N₂O emissions and Cumulative COVID-19 cases. The result of the analysis shows the calculated values of the Z-test of 3.2776 (22-Dec-2020)and 3.5593 (09- July-2021) which are higher than the Z-critical two-tail value of 1.9600 revealing that there is a positive significant relationship between N₂O emissions and cumulative covid-19 cases. Other hands, in the case of CO₂ emissions, a weak and inconsistent correlation ($r^2= 0.2 \pm 0.23$) exist, suggesting that COVID-19 cases are independent of CO₂ utilization. These statistical findings were overall supportive of N₂O emission's impact on enhancing SARS-COV2 spread in the human population.

Table 2. z-Test: Two sample for means (Cumulative COVID-19 infected cases and Nitrous oxide emissions)

	CUMULATIVE COVID-19 INFECTED CASES			Nitrous Oxide Emissions
	09-Jul-2021	22-Dec-2020	28-Mar-2020	
Mean	1825200.4	740951.0309	5815.2062	24886.4
Known Variance	2.48 X 10 ¹³	4.63 X 10 ¹²	2.88 X 10 ⁸	4.53 X 10 ⁹
Observations	97	97	97	97
Hypothesized Mean Difference	0	0	0	
z	3.5593	3.2776	-2.7060	
P(Z<=z) one-tail	0.0002	0.0005	0.0034	
z Critical one-tail	1.6449	1.6449	1.6449	
P(Z<=z) two-tail	0.0004	0.0010	0.0068	
z Critical two-tail	1.9600	1.9600	1.9600	

Notably, the top five N₂O emissions countries China, India, United States of America (US), Brazil, and Indonesia. [18] According to the WHO, currently the US, India and Brazil are the top three countries with the highest number of infections in the world. Yet standing at the 15th global rank of COVID-19, in our opinion Indonesia is still at high risk. Other hand, the Brunei Darussalam, Maldives, French Polynesia, Malta, Bahrain and Bhutan are among the least N₂O emissions Countries/ Territories. All these

are with the least COVID-19 outbreaks. Overall, these findings are further supported by real global evidences and suggest N₂O gas emissions significantly increase SARS-COV2 virus transmission.

Statistical analyses of epidemiological data summarize the way by which COVID-19 spread in the human population. According to the Oxford University report, agriculture constitutes the largest source of N₂O gas emissions 100-folds in comparison to transport vehicles, and both synthetic fertilizers/pesticides usually increase the negative environmental footprint of agriculture by contributing to greenhouse N₂O emissions. Surprisingly, a fall in N₂O gas emissions was noticed in affected areas after the surge of the SARS-CoV-2 virus [19] and is supportive of a natural virus evolution utilizes the major N₂O emissions.

By science value, there is a growing awareness that any infectious disease modulate the interplay between pathogens, hosts and environment. As stated by the National Institutes of Health that once any pathogen withstand the environment outside its host for a long period of time, an indirect contact occurs and this type of indirect transmission is already well established in case of gastrointestinal diseases, for example cholera, rotavirus infection, cryptosporidiosis, and giardiasis [20]. Hence, these sporadic COVID-19 instances are in full agreement with intrinsic N₂O emissions and suggest that these emissions is emerging as SARS-COV2 virus feed supplements.

Nature remains silent killer of harmful pathogens, and at the core of its response is the fundamental realization that nature' effect against COVID-19 epidemic is varied around the globe. Consequently, policy and strategy response covering suppression of N₂O emissions underlying scientific interventions for effective tackling of COVID-19 pandemic before arrival of front-line therapeutics and preventive vaccine.

The COVID-19 attributing factors with recently observed evidence, for instance, (i) a fall in N₂O gas emissions after the surge of COVID-19 in China, N₂O itself is a biochemical precursor of NO (↓NO level); (ii) High blood sugar- a reduced ability to produce NO (↓NO level); (iii) High blood pressure- NO inactivation (↓NO level); (iv) Obesity inhibit NO (↓NO level); (v) Smoking- inhibit biomolecular signaling molecule NO via nicotinic acetylcholine receptors (↓NO level); (vi) concurrent drug use (↓NO level); and (viii) a high salt intake attenuates NO production (↓NO level). It has been corroborated that the SARS-COV2 virus is surviving using N₂O energy source in atmosphere and in human population. Once again, COVID-19 pathophysiology and common symptoms fever, apnoea, cough, tiredness, and highly vulnerable groups, such as hypertensive, diabetics, smoker, and obese people, all observed reduced bioactive NO after the COVID-19 viral entry. [21]

NO is a gaseous secondary messenger known to be involved in numerous intra- and intercellular signaling events[22-24], for example, regulation of non-specific immune defense, body temperature, blood sugar, inflammation, vasodilation, airway tone, etc. More specifically, reports by Dantonio et al.

suggested systemic inhibition of NO synthesis attenuates bacterial endotoxin-induced fever[25], Hori et al. highlighted NO is involved in an ovalbumin-induced increase in cough reflex sensitivity[26], Prado et al. outlined role of nitric oxide in the various pathologic mechanisms of allergic asthma[27]. As a result of NO inhibition, common symptoms fever, cough, tiredness, and difficulty in breathing were appearing after COVID infection. The NO inducible agents, for example, hydroxychloroquine is being clinically relevant in COVID-19 treatment. A review of Nitric oxide enhancement strategies emphasizes many dietary strategies for more safe and effective NO based solutions than NO donating therapeutics[28].

The NO inactivation concept has evolved from the current approach and, endorse novel SARS-CoV2 coronavirus is greatly dependent upon local places with high N₂O emissions in past and CURRENT N₂O is turning low. It's surprising but most infected places, Wuhan, New York, Mumbai, Milan etc. are perfect falling in this line (Intrinsic- factors). In response, this study also proposed a framework of all-natural supplements for COVID-19 reduction (not prevention) by means of lifestyle and dietary supplements as follows:

1. Smoker, Hypertensive, Diabetics, and Obese people are among highly vulnerable groups, require counselling and strict attention (As reported above).
2. Indoor fitness exercises including meditation/ yoga are one of the front-line COVID-19 reduction markers. (Increase NO production).
3. People are advised to consume a very minimum amount of sugar and salt. (Both salt and sugar decreases NO production).
4. Patients with poly-pharmacy or concurrent drugs, for example, statins, fibrates, thiazolidinediones, metformin, antioxidant vitamins, aspirin, n-3 polyunsaturated fatty acids, and plant flavonoids should remain attentive. (Poly-pharmacy attenuate NO production).
5. Consumption of natural L-Arginine-rich sources, for example, dairy products, soybeans, peanuts, spirulina, lentils, eggs, etc. (L- Arginine is a precursor of NO production) and Citrulline supplementation using natural sources, for example, watermelon, pumpkins, cucumber, etc. (Citrulline stimulates de novo Arginine production)
6. Moderate consumption of nitrate-rich natural sources, such as beetroot, spinach, lettuce, chervil, radish, celery, pomegranate, etc. (Nitrate-nitrite-NO is second metabolic pathway).

In line with modern science multi-factorial approach, this study is endorsing nature's role in direct pathogen invasion and to limit the transmission of SARS-COV-2 pathogen. Thus, similar approach help researchers think undoubtedly about the causation of COVID-19 disease. We hope the insights presented

above along with disease reduction framework will be a game-changer in fighting against the COVID-19 pandemic before the launch of a front-line therapeutic(s) or preventable vaccine.

Acknowledgements

The authors are thankful to the World Health Organization and Worldbank for keeping data free to use. The authors are also thankful to GEHU India for providing necessary facilities and technical support.

Disclosure statement

The authors declare no competing financial interest.

Author contributions

PG devised the project and the main conceptual ideas. NT drafted the manuscript. GJ critically revised the manuscript and all authors approved the final version for submission.

References

1. Vitousek, P.M., Aber, J.D., Howarth, R.W., Likens, G.E., Matson, P.A., Schindler, D.W., Schlesinger, W.H. and Tilman, D.G., 1997. Human alteration of the global nitrogen cycle: sources and consequences. *Ecological applications*, 7(3), pp.737-750.
2. Snyder, C.S., Bruulsema, T.W., Jensen, T.L. and Fixen, P.E., 2009. Review of greenhouse gas emissions from crop production systems and fertilizer management effects. *Agriculture, Ecosystems & Environment*, 133(3-4), pp.247-266..
3. Jevtović-Todorović, V., Todorović, S.M., Mennerick, S., Powell, S., Dikranian, K., Benshoff, N., Zorumski, C.F. and Olney, J.W., 1998. Nitrous oxide (laughing gas) is an NMDA antagonist, neuroprotectant and neurotoxin. *Nature medicine*, 4(4), pp.460-463.
4. Lowenstein, C.J., Dinerman, J.L. and Snyder, S.H., 1994. Nitric oxide: a physiologic messenger. *Annals of internal medicine*, 120(3), pp.227-237.
5. Trieger, N., Loskota, W.J., Jacobs, A.W. and Newman, M.G., 1971. Nitrous oxide—a study of physiological and psychomotor effects. *The Journal of the American Dental Association*, 82(1), pp.142-150.
6. Bruckdorfer, R., 2005. The basics about nitric oxide. *Molecular aspects of medicine*, 26(1-2), pp.3-31.
7. Tian, H., Xu, R., Canadell, J.G., Thompson, R.L., Winiwarter, W., Suntharalingam, P., Davidson, E.A., Ciais, P., Jackson, R.B., Janssens-Maenhout, G. and Prather, M.J., 2020. A comprehensive quantification of global nitrous oxide sources and sinks. *Nature*, 586(7828), pp.248-256.
8. Smith, K.A., McTaggart, I.P. and Tsuruta, H., 1997. Emissions of N₂O and NO associated with nitrogen fertilization in intensive agriculture, and the potential for mitigation. *Soil use and management*, 13, pp.296-304.
9. Del Grosso, S.J., Wirth, T., Ogle, S.M. and Parton, W.J., 2008. Estimating agricultural nitrous oxide emissions. *EOS, Transactions American Geophysical Union*, 89(51), pp.529-529.
10. Ciotti, M., Ciccozzi, M., Terrinoni, A., Jiang, W.C., Wang, C.B. and Bernardini, S., 2020. The COVID-19 pandemic. *Critical reviews in clinical laboratory sciences*, 57(6), pp.365-388.
11. Pubmed Available from: <https://pubmed.ncbi.nlm.nih.gov/?term=covid-19> Retrieved on 13-July-2021.

12. Wilder-Smith, A., Chiew, C.J. and Lee, V.J., 2020. Can we contain the COVID-19 outbreak with the same measures as for SARS?. *The lancet infectious diseases*, 20(5), pp.e102-e107.
13. Ogen, Y., 2020. Assessing nitrogen dioxide (NO₂) levels as a contributing factor to coronavirus (COVID-19) fatality. *Science of the Total Environment*, 726, p.138605.
14. Alvarez, R.A., Berra, L. and Gladwin, M.T., 2020. Home nitric oxide therapy for COVID-19.
15. Hochberg, M.E., Hassell, M.P. and May, R.M., 1990. The dynamics of host-parasitoid-pathogen interactions. *The American Naturalist*, 135(1), pp.74-94.
16. Covid-19 situation reports available from <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>
17. Worldbank data available from <https://data.worldbank.org/indicator/EN.ATM.NOXE.KT.CE> Retrieved on 15-July-2020.
18. Han, M., Zhang, B., Zhang, Y. and Guan, C., 2019. Agricultural CH₄ and N₂O emissions of major economies: consumption-vs. production-based perspectives. *Journal of Cleaner Production*, 210, pp.276-286.
19. Otmani, A., Benchrif, A., Tahri, M., Bounakhla, M., El Bouch, M. and Krombi, M.H., 2020. Impact of Covid-19 lockdown on PM₁₀, SO₂ and NO₂ concentrations in Salé City (Morocco). *Science of the total environment*, 735, p.139541.
20. National Institutes of Health (US); Biological Sciences Curriculum Study. NIH Curriculum Supplement Series [Internet]. Bethesda (MD): National Institutes of Health (US); 2007. Understanding Emerging and Re-emerging Infectious Diseases. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK20370/>
21. Tay, M.Z., Poh, C.M., Rénia, L., MacAry, P.A. and Ng, L.F., 2020. The trinity of COVID-19: immunity, inflammation and intervention. *Nature Reviews Immunology*, 20(6), pp.363-374.
22. Bender, D. and Schwarz, G., 2018. Nitrite-dependent nitric oxide synthesis by molybdenum enzymes. *FEBS letters*, 592(12), pp.2126-2139.
23. Doel, J.J., Benjamin, N., Hector, M.P., Rogers, M. and Allaker, R.P., 2005. Evaluation of bacterial nitrate reduction in the human oral cavity. *European journal of oral sciences*, 113(1), pp.14-19.
24. Luiking, Y.C., Engelen, M.P. and Deutz, N.E., 2010. Regulation of nitric oxide production in health and disease. *Current opinion in clinical nutrition and metabolic care*, 13(1), p.97.
25. Dantonio, V., Batalhão, M.E., Fernandes, M.H., Komegae, E.N., Buqui, G.A., Lopes, N.P., Gargaglioni, L.H., Carnio, É.C., Steiner, A.A. and Bicego, K.C., 2016. Nitric oxide and fever: immune-to-brain signaling vs. thermogenesis in chicks. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 310(10), pp.R896-R905.
26. Hori, A., Fujimura, M., Ohkura, N. and Tokuda, A., 2011. Involvement of nitric oxide (NO) in cough reflex sensitivity between non-sensitized and OVA-sensitized guinea pigs. *Cough*, 7(1), pp.1-9.
27. Prado, C.M. and Martins, M., 2011. a, and Tibério, IFLC (2011). *Nitric oxide in asthma physiopathology*. *ISRN allergy*, 832560.
28. Bryan, N.S., 2015. Nitric oxide enhancement strategies. *Future science OA*, 1(1).